

To the Editor: Although he stresses the usefulness of the serum-ascites albumin gradient, Runyon claims that measurement of fibronectin or cholesterol in ascitic fluid is not useful for the differential diagnosis of ascites. We have a different view. A test for fibronectin¹ or cholesterol² in ascitic fluid can accurately discriminate between malignant and nonmalignant ascites and is superior to protein measurement for this purpose. The serum-ascites albumin gradient may be inferior to ascitic-fluid cholesterol measurement in the differential diagnosis of ascites^{3,4,5}. For these reasons, we advocate the measurement of ascitic-fluid cholesterol.

A.L. Gerbes, M.D.
University of Munich
81377 Munich, Germany

J. Scholmerich, M.D.
University of Regensburg
93053 Regensburg, Germany

References

1. Scholmerich J, Volk BA, Kottgen E, Ehlers S, Gerok W. Fibronectin concentration in ascites differentiates between malignant and nonmalignant ascites. *Gastroenterology* 1984;87:1160-1164. [\[Medline\]](#)
2. Jungst D, Gerbes AL, Martin R, Paumgartner G. Value of ascitic lipids in the differentiation between cirrhotic and malignant ascites. *Hepatology* 1986;6:239-243. [\[CrossRef\]](#) [\[Medline\]](#)
3. Gerbes AL, Jungst D, Xie YN, Permanetter W, Paumgartner G. Ascitic fluid analysis for the differentiation of malignancy-related and nonmalignant ascites: proposal of a diagnostic sequence. *Cancer* 1991;68:1808-1814. [\[CrossRef\]](#) [\[Medline\]](#)
4. Prieto M, Gomez-Lechon MJ, Hoyos M, Castell JV, Carrasco D, Berenguer J. Diagnosis of malignant ascites: comparison of ascitic fibronectin, cholesterol, and serum-ascites albumin difference. *Dig Dis Sci* 1988;33:833-838. [\[CrossRef\]](#) [\[Medline\]](#)
5. Misra SP, Dwivedi M, Misra V, Gupta R, Agarwal SK. Diagnostic paracentesis. *Gastroenterology* 1990;99:902-903. [\[Medline\]](#)