

Editors: K. Fellingner and R. Höfer

# FURTHER ADVANCES IN THYROID RESEARCH

VOL 2



Verlag der Wiener Medizinischen Akademie

N 172/19055



1971

Verlag der Wiener Medizinischen Akademie

A-1090 Wien, Alser Straße 4

Druck: G. Gistel & Cie., Wien III

## Table of Contents

	Page
K. FELLINGER and R. HÖFER, Vienna	
Introduction . . . . .	11
K. FELLINGER, Vienna, Austria	
Adress of the President . . . . .	11
ROSALIND PITT-RIVERS, London, Great Britain	
Some Biological Reactions of Iodine . . . . .	15
TH. WINSHIP, Washington (D. C.), U.S.A.	
The History of the International Thyroid Conferences— American Thyroid Association . . . . .	25
F. DELANGE, J. M. HERSHMAN and A. M. ERMANS, Brussels, Belgium, and Birmingham (Ala.), U.S.A.	
Blood TSH Level in Idjwi Island: Modifications Related to Regional Prevalence of Goitre and Not to Individual Thyroidal Size and Function . . . . .	35
E. A. PRETELL, M. WAN and P. PALACIOS, Lima, Peru	
Fetal Thyroid Function in Endemic Goiter . . . . .	45
I. H. BUTTFIELD, P. O. D. PHAROAH and B. S. HETZEL, Melbourne, Australia, and New Guinea	
Evidence of Prevention of Neurological Defect in New Guinea Children by Iodised Oil Injection of Mothers Prior to Pregnancy . . . . .	53
R. FIERRO-BENITEZ, I. RAMIREZ, E. ESTRELLA, A. QERIDO and J. B. STANBURY, Quito, Ecuador, Leiden, Netherlands, and Cambridge (Mass.), U.S.A.	
The Effect of Goitre Prophylaxis with Iodized Oil on the Prevention of Endemic Cretinism . . . . .	61
D. A. KOUTRAS, G. A. RIGOPOULOS, A. D. PHARMAKIOTIS, J. SFONTOURIS, J. MANTZOS, G. VLASSIS and B. MALAMOS, Athens, Greece	
The Metabolic Effects of Iodized Oil in Endemic Goitre . . . . .	79
A. M. ERMANS and M. CAMUS, Brussels, Belgium	
Evidence of a Regulation of the Thyroid Activity by the Level of Dietary Iodine in Some Thyroid Diseases . . . . .	85
NOBUO Ue, Maebashi, Japan	
Studies of Hog Thyroglobulin by Isoelectric Focusing . . . . .	91
H. BORNET, A. SCHNEIDER and H. EDELHOCH, Bethesda (Md.), U.S.A.	
Newly Synthesized Thyroglobulin: Effect of Temperature and 6M Guanidine . . . . .	97
L. CRAMAROSSA, M. ANDREOLI, G. SCUNCIO, M. D'ARMIENTO and C. CASSANO, Rome, Italy	
Effects of Temperature and Methyl-Mercaptoimidazole on the Subunit Structure of Human and Bovine Thyroglobulin . . . . .	105
P. G. MALAN and H. EDELHOCH, Bethesda (Md.), U.S.A.	
Reactivity of Tyrosyl Residues in Bovine and Human Goitre Thyroglobulins with Tetranitromethane . . . . .	113
L. WARTOFSKY and S. H. INGBAR, Washington (D.C.) and Boston (Mass.), U.S.A.	
A Method for Assessing the Latency, Potency and Duration of Action of Anti-thyroid Agents in Man . . . . .	121
R. HÖFER, P. LAURINGER, G. PAUMGARTNER, E. VORMITTAG and W. VORMITTAG, Vienna, Austria	
Effect of 1-methyl-2-mercaptoimidazole on Liver Alkaline Phosphatase . . . . .	137
B. E. W. BROWNLIE, B. MARCHANT and W. D. ALEXANDER, Glasgow, United Kingdom	
Placental Transfer of <sup>35</sup> S-Propylthiouracil and <sup>35</sup> S-Methimazole in the Rat . . . . .	143
P. PELTOLA and F.-E. KRUSIUS, Helsinki, Finland	
The Effect of Small Doses of 1-5-vinyl-2-thioxazolidone on the Human Thyroid Function During Long-Term Treatment . . . . .	149
N. MICHALOVSKIJ and P. LANGER, Bratislava, Czechoslovakia	
Chemical Identification and Goitrogenic Activity of L-5-vinyl-4-hydroxy-2-thioxazolidone— a Metabolite of Naturally Occurring L-5-vinyl-2-thioxazolidone . . . . .	155

T. HJORT, U. B. LAURIDSEN and I. B. PERSSON, Copenhagen, Denmark Thyroglobulin-Like Substances with Low Iodine Content in the Serum of Patients Receiving Antithyroid Therapy . . . . .	163
P. DE NAYER, M. LABRIQUE and M. DE VISSCHER, Louvain, Belgium Cell-Free Synthesis of Thyroglobulin: Covalent Association of Subunits — Effect of 2-Mercaptoethanol . . . . .	173
P. OLIN, S. ALMQVIST and R. EKHOLM, Göteborg, Sweden The Appearance of Immunoreactive Thyrotrophin in the Human Fetal Pituitary Gland Related to Iodide Accumulation, Biosynthesis of Thyroglobulin and Ultra- structure of the Thyroid Gland . . . . .	181
S. LISSITZKY, M. ROQUES, J. TORRESANI and A. JOST, Marseille and Paris, France Relationship Between Thyroglobulin Biosynthesis, Thyroid Hormone Formation and Histogenesis of the Fetal Rabbit Thyroid. In vivo Studies . . . . .	189
L. LAMAS, I. COVELLI, H. EDELHOCH, F. CORTESE and G. SALVATORE, Naples, Italy Relationships Between Structure of Thyroglobulin and Synthesis of Thyroxine . .	201
A. TAUROG, Dallas (Tex.), U.S.A. Thyroid Peroxidase-Catalyzed Iodination and Thyroxine Formation in Various Proteins . . . . .	211
H. KOHLER, H. STUDER, A. JUNG and H. BÜRGI, Berne, Switzerland Thyroglobulin Heterogeneity and its Impact on Iodine Kinetics: A Link in the Chain of Thyroid Control Mechanisms . . . . .	225
C. S. PITTMAN, J. B. CHAMBERS JR. and V. H. READ, Birmingham (Ala.), U.S.A. The Extrathyroidal Conversion Rate of Thyroxine to Triiodothyronine in Man . .	233
J. H. DUSSAULT and D. A. FISHER, Torrance and Los Angeles (Cal.), U.S.A. Absence of In Vivo Extrathyroidal Conversion of Thyroxine to Triiodothyronine in the Serum of Sheep . . . . .	237
L. E. BRAVERMAN, A. VAGENAKIS, P. DOWNS, A. E. FOSTER, K. STERLING and S. H. INGBAR, Brighton (Mass.), Boston (Mass.) and New York (N.Y.), U.S.A. The Effects of Stable Thyroxine (T <sub>4</sub> ) Replacement on the Peripheral Metabolism of T <sub>4</sub> and Triiodothyronine (T <sub>3</sub> ) . . . . .	243
G. HENNEMANN, C. BECKERS, R. DOCTER, A. DOLMAN and PH. DE NAYER, Rotter- dam, Netherlands, and Louvain, Belgium Observations Concerning the Relation Between Total Thyroxine (TT <sub>4</sub> ) and Absolute Free Thyroxine (AFT <sub>4</sub> ), and the Influence of AFT <sub>4</sub> on Serum TSH Levels and Thyroxine Disposal in Humans . . . . .	255
C. H. BASTOMSKY, Montreal, Canada The Mechanism of Theophylline-Induced Goiter . . . . .	263
J. H. DUSSAULT, C. J. HOBEL and D. A. FISHER, Torrance and Los Angeles (Cal.), U.S.A. Thyroxine Secretion in Fetal Sheep . . . . .	273
P. V. N. MURTHY, P. R. ADIGA and J. M. MCKENZIE, Montreal, Canada Effects of Poly-L-Ornithine on Stimulation of Thyroid Function . . . . .	281
P. ROCMANS, C. WILLEMS, P. NÈVE, J. OTTEN, F. RODESCH and J. E. DUMONT, Brus- sels, Belgium Thyroid Secretion and its Regulation . . . . .	291
M. A. PISAREV, L. J. DEGROOT and J. F. WILBER, Buenos Aires, Argentina, and Chicago, (Ill.), U.S.A. Regulation of Protein Synthesis in the Thyroid: Role of Cyclic Nucleotides . . .	297
R. E. LECOQ, F. M. LAMY, E. M. KEYHANI, G. VASSART and J. E. DUMONT, Brussels, Belgium Regulation of Protein Synthesis in the Thyroid . . . . .	307
J. KNOPP, V. STOLC and W. TONG, Pittsburgh (Pa.), U.S.A. The Dynamics of Certain Responses of Bovine Thyroid Cells to the Addition and the Withdrawal of TSH . . . . .	311
G. C. SCHUSSLER, Buffalo (N. Y.), U.S.A. Tissue Distribution of Radioactivity Following Single Injection and Continuous In- fusion of <sup>131</sup> I-Thyroxine and <sup>125</sup> I-Triiodothyronine. . . . .	319

R. R. CAVALIERI, M. STEINBERG and G. L. SEARLE, San Francisco (Cal.), U.S.A. Metabolism of Triiodothyronine (T <sub>3</sub> ) in Graves' Disease . . . . .	327
J. MCCONNON, V. V. ROW and R. VOLPÉ, Toronto, Canada Simultaneous Comparative Studies of Thyroxine (T <sub>4</sub> ) and Triiodothyronine (T <sub>3</sub> ) Production Rates in Health and Disease . . . . .	335
A. A. ZANINOVICH, O. DEGROSSI and V. PECORINI, Buenos Aires, Argentina Changes of Triiodothyronine Kinetics Produced by Variations of Serum Thyroxine- Binding Globulin Capacity in Subjects Without Thyroxine . . . . .	343
J. J. DiSTEFANO and R. F. CHANG, Los Angeles (Cal.), U.S.A. Computer Simulation of the Kinetics of Binding, Distribution and Metabolism of Thyroxine and Triiodothyronine in Man . . . . .	351
R. G. SPIRO, MARY J. SPIRO and T. ARIMA, Boston (Mass.), U.S.A. Studies on the Carbohydrate Units of Human Thyroglobulin . . . . .	363
SIMONE BOUCHILLOUX, ODILE CHABAUD and M.-MICHEL-BECHET, Marseille, France Subcellular Localization of Thyroid Glycosyltransferases . . . . .	371
R. EKHOLM and U. BJÖRKMAN, Göteborg, Sweden Cell-Free Incorporation of Labeled Monosaccharides into Thyroid Proteins . . . .	381
ANNETTE HERSCOVICS, P. WHUR, A. HADDAD and MEREDITH SMITH, Montreal, Canada Biosynthesis of the Carbohydrate of Thyroglobulin in Rat Thyroid . . . . .	387
H. HAIBACH, I. KOBAYASHI and M. A. GREER, Portland (Or.), U.S.A. Comparison of the Relative Rates of Endogenous Enzymatic Hydrolysis of Im- mature and Mature Thyroglobulin . . . . .	399
A. GORDON, O. SPIRA and J. GROSS, Jerusalem, Israel Evidence for the Presence of an Iodide Binding Peptide in the Thyroid and Salivary Gland . . . . .	407
L. J. DEGROOT and A. NAGASAKA, Chicago (Ill.), U.S.A. Biosynthesis of Thyroid Hormone: Polymeric Structure of Enzyme and Mechanism of H <sub>2</sub> O <sub>2</sub> Generation for Peroxidation . . . . .	413
N. M. ALEXANDER, New Haven (Conn.), U.S.A. Thyroid Peroxidase: Nature of the Heme Binding to Apoperoxidase . . . . .	423
J. POMMIER, J. NUNEZ and L. SOKOLOFF Enzymatic Iodination of Proteins . . . . .	433
H. OGAWARA and H. J. CAHNMANN, Bethesda (Md.), U.S.A. Nonenzymic Synthesis of Thyroxine in Thyroglobulin by Coupling with 4-Hy- droxy-3,5-Diiodophenylpyruvate . . . . .	441
J. ROBBINS, Bethesda (Md.), U.S.A. Abnormal Thyroglobulin in a Rat Thyroid Tumour . . . . .	451
C. CASSANO, F. MONACO, S. FONTANA, B. G. SALABÉ and M. ANDREOLI, Rome, Italy Physicochemical and Immunological Characterization of Abnormal Thyroglobulin from Human Congenital Goitre . . . . .	457
P. P. VAN JAARSVELD, L. SENA, B. VAN DER WALT and A. VAN ZYL, Bellville, South Africa, and Naples, Italy Abnormal Iodoproteins in a Congenital Bovine Goitre . . . . .	465
I. R. FALCONER, Nottingham, Great Britain Iodoproteins in a Congenital Goitre Lacking the Ability to Synthesize Normal Thyroglobulin . . . . .	481
S. SHULMAN and M. GHAYASUDDIN, New York (N.Y.), U.S.A. Isolation and Antigenic Characterization of Thyroglobulin Fragments . . . . .	493
Z. LEWITUS and Y. SHAHAM, Tel-Aviv, Israel An Extrathyroidal Iodine Concentration Pool as a First Stage in the Thyroidal Iodine Uptake in the Lizard . . . . .	503
R. RIVIÈRE, D. COMAR, G. F. CAMUZZINI, M. GIRAULT and C. KELLERSHOHN, Orsay, France, and Merida, Venezuela Determination of the Specific Radioactivity of Intrafollicular Iodine in Human Thyroid . . . . .	511

H. STUDER, H. BÜRGI and H. KOHLER, Berne, Switzerland Differential Sensitivity of Recently Iodinated and Aged Thyroglobulin to TSH In Vivo . . . . .	519
L. VAN MIDDLESWORTH, Memphis (Tenn.), U.S.A. Development of Relatively Stagnant Iodine Pools in Thyroids of Rats . . . . .	525
G. A. BRAY and STELLA MOTHON, Boston (Mass.), U.S.A. Studies on the Turnover and Concentration of Thyroid in Old and Young Rats . . . . .	533
W. H. FLORSHEIM, Long Beach (Cal.), U.S.A. On Several Mechanisms Determining LATS Responses in the MCKENZIE Assay . . . . .	549
D. H. SOLOMON, G. N. BEALL and I. J. CHOPRA, Torrance and Los Angeles (Cal.), U.S.A. Mobilization of Tissue Thyroxine in Mice by Thyroxine-Binding Proteins: Studies with a Double-Isotopic MCKENZIE Assay . . . . .	557
I. J. CHOPRA, G. N. BEALL and D. H. SOLOMON, Torrance and Los Angeles (Cal.), U.S.A. Demonstration of a Soluble Lipoprotein LATS-Inhibitor in Human Thyroid Extract . . . . .	565
B. R. SMITH, D. S. MUNRO and K. J. DORRINGTON, Sheffield and Cambridge, Great Britain Purification and Characterisation of LATS-Binding Protein from Human Thyroid Tissue . . . . .	571
T. MORI and J. P. KRISS, Stanford (Cal.), U.S.A. Rapid Competitive Binding Radioassay of Serum Anti-Microsomal and Anti-Thyroglobulin Antibodies: Measurements in Graves' Disease . . . . .	577
R. J. WINAND, J. SALMON and P. H. LAMBERT, Liège, Belgium Characterization of the Exophthalmogenic Factor Isolated from the Serum of Patients with Malignant Exophthalmos . . . . .	583
I. R. HART, Ottawa, Canada The Lipolytic Effect of LATS-IgG and Graves' Disease Sera . . . . .	595
D. V. BECKER, W. M. MCCONAHEY, B. M. DOBYNS, EDYTHALENA TOMPKINS, G. E. SHELINE and J. B. WORKMAN, New York (N.Y.), Rochester (Minn.), Cleveland (O.), Washington (D. C.), San Francisco (Cal.) and Baltimore (Md.), U.S.A. The Results of Radioiodine Treatment of Hyperthyroidism . . . . .	603
R. N. SMITH, D. S. MUNRO and G. M. WILSON, Sheffield and Glasgow, United Kingdom Two Clinical Trials of Different Doses of Radio-Iodine ( <sup>131</sup> I) in the Treatment of Thyrotoxicosis . . . . .	611
W. R. GREIG, H. W. GRAY, I. R. MCDUGALL, J. F. B. SMITH, F. C. GILLESPIE, J. A. THOMSON and E. M. MCGIRR, Glasgow, United Kingdom <sup>125</sup> I Therapy for Thyrotoxicosis: Results of Treatment of 50 Patients Followed for at Least 1 Year after Therapy . . . . .	619
Z. LEWITUS, E. LUBIN, J. RECHNIC, M. BEN-PORATH, Y. FEIGE and J. LAOR, Beilinson, Israel Treatment of Thyrotoxicosis with Small Doses of <sup>125</sup> I . . . . .	643
H. K. IBBERTSON, R. B. HUNTON, B. MCL. WHITE and P. D. GLUCKMANN, Auckland, New Zealand Early Thyroid Clearance Measurement for the Assessment of Carbimazole Therapy in Thyrotoxicosis . . . . .	653
A. W. G. GOOLDEN, M. BROWN and E. D. WILLIAMS, London, Great Britain Measurement of the Thyroid Trap with <sup>99m</sup> Tc as a Guide to the Medical Treatment of Thyrotoxicosis . . . . .	663
J. WOLFF, Bethesda (Md.), U.S.A. Enzymatic Properties of Thyroid Membranes . . . . .	669
J. A. WILLIAMS and J. WOLFF, Bethesda (Md.), U.S.A. Thyroid Secretion In Vitro: Effects of Membrane Stabilizers . . . . .	677
M. L. MAAYAN, R. J. SHAPIRO and S. H. INGBAR, Boston (Mass.), U.S.A. Metabolic Functions of Thyroid Cell "Ghosts" . . . . .	683

M. PAVLOVIC-HOURNAC, L. RAPPAPORT and J. NUNEZ, Paris, France Protein Synthesis in Thyroid Gland from Hypophysectomized Rats . . . . .	693
N. D. ESHCHENKO, F. E. PUTILINA and R. R. RACHEV, Leningrad, U.S.S.R., and Sofia, Bulgaria Effect of Thyroid Hormones on the Content and Redox Potential of Pyridine Nucleotides in Animal Tissues . . . . .	701
P. LANGER, K. GSCHWENDTOVA and L. PAZMANOVA, Bratislava, Czechoslovakia Short-term Changes of Pituitary Protein Metabolism Following a Single Adminis- tration of Antithyroid Drug in Rats . . . . .	705
M. SUZUKI and K. SHIBASAKI, Maebashi, Japan Early Effect of Thyroxine on Synthesis of Growth Hormone and Prolactin in the Adenohypophysis of Thyroidectomized Rat . . . . .	711
J. MANTZOS and L. CHIOTAKI, Athens, Greece Effect of Thyroid Hormones on Biosynthesis of Myelin Lipids . . . . .	723
R. MICHEL, J. BOUHNIC and O. MICHEL, Paris, France Action of Thyroid Hormones on the Mixed Function Oxidases of Isolated Beef Adrenal Cortex Mitochondria . . . . .	731
M. MORRISON, D. J. DANNER and G. S. BAYSE, Memphis (Tenn.), U.S.A. Subcellular Distribution and Catalytic Activity of Thyroid Peroxidase . . . . .	741
S. OHTAKI and I. N. ROSENBERG, Boston (Mass.), U.S.A. Some Effects of Monoamine Oxidase Inhibitors upon the Thyroid . . . . .	749
A. NAGASAKA and H. HIDAKA, Chicago (Ill.) and Nutley (N. J.), U.S.A. Inhibition of Iodide Peroxidase by Iproniazid . . . . .	765
B. BÉNARD and J. BRAULT, Sherbrooke, Canada Hydrogen Peroxide Production in Thyrotropin Stimulated Thyroid . . . . .	771
A. MELANDER, L. E. ERICSON, CH. OWMAN and F. SUNDLER, Göteborg and Lund, Sweden Colloid Droplet Formation and Release of Thyroid Hormone by Catecholamines and Indoleamines, and Their Inhibition by Adrenergic Blocking Drugs . . . . .	779
P. HEIMANN and B. THOLIN, Göteborg, Sweden Ultrastructure of Human Thyroid Tumours . . . . .	787
M. MICHEL-BECHET, L. VALENTA and F. KYNCL, Marseille, France, and Prague, Czechoslovakia Ultrastructure of Thyroid Carcinoma . . . . .	797
A. A. AL-SAADY, Ann Arbor (Mich.), U.S.A. The Effect of Host Condition on the Development and Progression of Iodine Deficiency Induced Transplanted Thyroid Tumours . . . . .	807
J. B. FIELD, U. ZOR, T. KANEKO, K. YAMASHITA and A. DEKKER, Pittsburgh (Pa.), U.S.A. Comparison of Effects of TSH, LATS and Prostaglandins on Dog Thyroid Slice Adenyl Cyclase, Cyclic AMP, Colloid Droplet Formation and Intermediary Metabolism . . . . .	817
C. S. AHN and I. N. ROSENBERG, Boston (Mass.), U.S.A. Oxidation of <sup>14</sup> C-Formate in Thyroid Slices: Effects of TSH, Dibutyl Cyclic 3', 5'-AMP (dbcAMP) and Prostaglandin E <sub>1</sub> (PGE <sub>1</sub> ) . . . . .	825
R. H. LINDSAY, A. G. CASH and J. B. HILL, Birmingham (Ala.), U.S.A. The Mechanism of a TSH Stimulation of Pyrimidine Pathways Leading to RNA Synthesis . . . . .	839
B. D. WILSON and R. L. WRIGHT, Bellville, South Africa Actions of Thyrotropin and Dibutyl Cyclic AMP on Ribonucleic Acid Synthesis in Isolated Thyroid Cells . . . . .	849
M. ZAKARIJA and C. H. BASTOMSKY, Montreal, Canada Thyroid Cyclic AMP Hydrolysis in Relation to Thyroid Gland Activity . . . . .	863
K. OSSOINIG, Vienna, Austria Echo-Orbitography—a Reliable Method for the Differential Diagnosis of Endo- crine Exophthalmos . . . . .	871

J. LECLÈRE, J. ROBERT, A. BERTRAND and P. HARTEMANN Ultrasound in the Diagnosis of Thyroid Diseases . . . . .	879
W. L. ROBISON and L. VAN MIDDLESWORTH, Livermore (Cal.) and Memphis (Tenn.), U.S.A. Determination of Calcium, Iodine, and Phosphorus Distributions in Thyroid Glands by Electron Probe Microanalysis . . . . .	891
H. FRITZSCHE, R. HÖFER and H. SCHATZ, Vienna, Austria Investigations on Thyroid Thin Needle Biopsies by Immunofluorescence . . . .	907
J. KAPITOLA, M. SCHÜLLEROVA and O. SCHREIBEROVA, Prague, Czechoslovakia Blood Flow Through the Thyroid Gland of Rats . . . . .	913
H. E. TRIANTAPHYLLIDIS and C. GUICHARD, Paris, France Effect of Casein and N-Acetyl-Neuraminic-Acid on Thyroid Function . . . . .	919
L. O. LUTHERER, M. J. FREGLY, A. H. BURNS and B. C. LUTHERER, Gainesville (Flor.), U.S.A. Effect of Ultracentrifugal Fractions of Rat Renal Homogenates on Uptake and Binding of <sup>131</sup> I by Porcine Thyroid Slices in Vitro and Rat Thyroid Glands in Vivo . . . . .	943
L. L. ROSENBERG, Berkeley (Cal.), U.S.A. Some Thyroidal Effects of Hydrocortisol in Hypophysectomized Rats . . . . .	957
W. L. GREEN, Seattle (Wash.), U.S.A. Iodine Metabolism During Treatment with the Tyrosine Dehalogenase Inhibitor, 3-Nitro-L-Tyrosine . . . . .	965
K. INOUE, M. A. GREER and A. TAUROG, Dallas (Tex.) and Portland (Or.), U.S.A. The Role of Protein Synthesis in the Utilization of Administered Iodide by Thyroid Glands in Rats . . . . .	973
J. A. WEAVER, R. C. LOWRY, D. R. HADDEN and D. A. D. MONTGOMERY, Belfast, North Ireland Thyroid Suppressibility — Follow-Up for Two Years After Antithyroid Treatmen C. E. CASSIDY, Boston (Mass.), U.S.A. Long-Term Follow-Up of Patients with Hyperthyroidism in Whom a Thyroid Suppression Test Was Performed at the End of Treatment . . . . .	989
D. W. SLINGERLAND, E. S. DELL and B. A. BURROWS, Boston (Mass.), U.S.A. The Spectrum of Thyroid Function After Radioiodine Treatment . . . . .	993
J. NOLTE, D. PETTE, B. BACHMAIER, P. KIEFHABER, H. SCHNEIDER and P. C. SCRIBA, Munich, Western Germany Enzyme Activities in Liver and Muscle Biopsy Specimens from Thyrotoxic and Hypothyroid Patients . . . . .	1005
Y. SHISHIBA, T. SHIMIZU, T. SAITO and K. SHIZUME, Tokyo, Japan Elevation of Immunoreactive Insulin Concentration Preceding an Attack of Thyrotoxic Periodic Paralysis . . . . .	1013
J. HERRMANN and H. L. KRÜSKEMPER, Hannover, Western Germany Observations on Treatment by Peritoneal Dialysis of Patients with Thyroid Storm . . . . .	1021
C. Y. BOWERS, A. V. SCHALLY, A. WEIL, G. A. REYNOLDS and K. FOLKERS, Austin (Tex.), U.S.A. Chemical and Biological Identity of Thyrotropin Releasing Hormone (TRH) of Bovine and Human Origin . . . . .	1029
J. WILBER, Chicago (Ill.), U.S.A. Stimulation of Thyrotropin Biosynthesis by Thyrotropin Releasing Hormone and 59 mM Potassium Chloride . . . . .	1041
D. D. ADAMS, T. H. KENNEDY and R. D. UTIGER, Dunedin, New Zealand, and Philadelphia (Pa.), U.S.A. Serum Thyrotropin (TSH) Concentrations: Measurements by Bioassay and Immunoassay in Iodine Deficiency and Other States . . . . .	1049



B. R. WEBSTER, B. E. FURNIVAL and J. C. PAICE, Toronto, Canada, and London, Great Britain Reference Preparations of Human Thyrotropin: Comparative Stability of Biological and Immunological Potencies . . . . .	1057
T. LEMARCHAND-BÉRAUD, M. GRIESSEN and B. R. SCAZZIGA, Lausanne, Switzerland Comparison of Bovine and Human TSH Radioimmunoassay for Human Plasma TSH Determination and Correlation with Bioassay . . . . .	1069
TH. COUTSOFTIDES and A. GORDON, Jerusalem, Israel The Effect of pH on Thyroid Hormone Binding by Serum Proteins . . . . .	1085
S. REFETOFF, N. I. ROBIN and C. A. ALPER, Chicago (Ill.) and Boston (Mass.), U.S.A. Four Phenotypes of Thyroxine Binding Globulin (TBG) in Man: Possible Mutations of a Single Regulatory Gene Locus . . . . .	1089
B. N. PREMACHANDRA, St. Louis (Mo.), U.S.A. Slow-Moving Human Thyroxine Binding Globulin (TBG): Similarity to TBG of Sialidase Treated Human Sera . . . . .	1097
B. L. BROWN, R. P. EKINS, S. M. ELLIS and E. S. WILLIAMS, London, Great Britain The Radioimmunoassay of Tri-Iodothyronine . . . . .	1107
J. BENOTTI, R. GRIMALDI, S. PINO and F. MALOOF, Waltham and Boston (Mass.), U.S.A. A Modified Method for Total Triiodothyronine (T <sub>3</sub> ) by Competitive Protein Binding . . . . .	1121
W. A. HARLAND and J. S. ORR, Glasgow, United Kingdom The Whole-Body Counter and Thyroxine Turnover . . . . .	1127
M. D. GARCÍA and G. MORREALE DE ESCOBAR, Madrid, Spain Radioimmuno and MCKENZIE Assays for Rat TSH . . . . .	1137
J. M. HERSHMAN, H. P. HIGGINS and W. R. STARNES, Birmingham (Ala.), U.S.A. and Toronto, Canada Differences Between a Thyroid Stimulator in a Hydatidiform Mole and Other Human Thyrotropins . . . . .	1155
D. M. COOK, J. W. KENDALL and E. M. NICHOLS, Portland (Ore.), U.S.A. Studies on the Role of CSF Circulation in Feedback Control of TSH Secretion . . . . .	1165
J. T. NICOLOFF, H. A. GROSS and M. D. APPLEMAN, JR., Los Angeles (Cal.), U.S.A. Inhibition of Thyroid Release by Estradiol . . . . .	1173
S. A. D'ANGELO and N. R. WALL, Philadelphia (Pa.), U.S.A. Maternal-Fetal Endocrine Interrelations: Effects of Goitrogens, Adrenal Inhibitors and Cyclic AMP in the Pregnant Rat . . . . .	1183
F. DE RUBERTIS, K. YAMASHITA, A. DEKKER and J. B. FIELD, Pittsburgh (Pa.), U.S.A. Dissociation of In Vivo and In Vitro Responsiveness to Thyroid Stimulating Hormone in Thyroid Adenomas . . . . .	1197
V. MACCHIA, M. F. MELDOLESI and M. CHIARIELLO, Naples, Italy Effect of TSH and TSH-like Substances on Some Properties of a Transplantable Thyroid Tumour of the Rat . . . . .	1205
K. SHIMAOKA and J. E. SOKAL, Buffalo (N. Y.), U.S.A. Comparison of L-Thyroxine (T <sub>4</sub> ) and L-Triiodothyronine (T <sub>3</sub> ) in Suppressive Therapy of Nontoxic Nodular Goitre . . . . .	1215
J.-G. LJUNGGREN, P. OLIN and BERTA JEREB, Stockholm, Sweden On the Biosynthesis of Thyroglobulin in Thyroid Carcinomas . . . . .	1221
G. V. FOSTER, M. B. CLARK, B. M. NATHANSON, D. GRAHAME-SMITH, L. GALANTE, R. HORTON and T. V. GUDMUNDSSON, London, Great Britain The Diagnosis of Medullary Carcinoma of the Thyroid by Bioassay and Radioimmunoassay . . . . .	1233

C. S. HILL, JR., A. H. TASHJIAN, JR., M. L. IBANEZ, N. A. SAAMAN and R. L. CLARK, Houston (Tex.) and Boston (Mass.), U.S.A.  
Diagnostic Value of Plasma Calcitonin Levels in Patients with Medullary Carcinoma of the Thyroid . . . . . 1245  
Authors and Co-Authors . . . . . 629, 1253

## **Enzyme Activities in Liver and Muscle Biopsy Specimens from Thyrotoxic and Hypothyroid Patients**

by J. NOLTE, D. PETTE, B. BACHMAIER, P. KIEFHABER, H. SCHNEIDER and P. C. SCRIBA \*)

### **Abstract**

Diagnosis of thyroid disorders was based on clinical status, radioiodine studies,  $PB^{127}I$  and  $T_3$  in vitro test. Needle biopsy samples from livers were immediately frozen (liquid  $N_2$ ) and enzyme activities were determined in supernatant and sediment fractions of total homogenates of each biopsy sample.

Phosphoglucumutase (E.C.2.7.5.1) is diminished in thyrotoxicosis (N = 10), as compared to controls (N = 19), indicating low glycogenolysis, and glyceraldehyde DH (E.C.1.2.1.12) was increased (augmented glycolytic rate). Hexokinase (E.C.2.7.1.1) and glucose-6-phosphate DH (E.C.1.1.1.49) remained unchanged by thyrotoxicosis or hypothyroidism (N = 5).

The activity of PEP-carboxykinase (4.1.1.31) is elevated in thyrotoxicosis and lowered in hypothyroidism, presumably with equivalent alterations of gluconeogenesis. The latter result has to be discussed with respect to the demonstration of an increased level of free cortisol in serum, resp. of diminished insulin efficiency in our thyrotoxic patients.

Enzymes of citric acid cycle and connected pathways—condensing enzyme (E.C.4.1.3.7), NADP-isocitrate DH (E.C.1.1.1.42), glutamate DH (E.C.1.4.1.2), aspartate amino transferase (E.C.2.6.1.1), malate DH (E.C.1.1.1.37)—were unchanged by thyrotoxicosis or hypothyroidism.

Malic enzyme (E.C.1.1.1.40) was increased in livers of thyrotoxic patients, 3-Hydroxyacyl-CoA DH (E.C.1.1.1.35) was not changed, whereas carnitine acetyl-transferase (E.C.2.3.1.7) was elevated in thyrotoxicosis. Mitochondrial  $\alpha$ -glycerol phosphate DH (E.C.1.1.99.5) showed no increase in thyrotoxic patients. This was unexpected in view of the known induction of this enzyme in the liver from rats treated with thyroid hormones (LARDY).

These results are discussed with respect to species differences and to mechanism of action of thyroid hormones.

### **Extrait**

Le diagnostic des troubles thyroïdiens était basé sur l'état clinique, l'exploration par le radioiode, le  $PB^{127}I$  et le test à la  $T_3$  in vitro. Les échantillons biopsiques des foies ont immédiatement été congelés ( $N_2$  liquide) et les activités enzymatiques déterminées dans le surnageant et les fractions de sédimentation des homogénats totaux de chaque échantillon biopsique.

La phosphoglucumutase (E.C.2.7.5.1) est diminuée dans la thyrotoxicose (N = 10), en comparaison aux témoins (N = 19), ce qui indique une basse glycogénolyse, et glyceraldehydphosphate DH (E.C.1.2.1.12) était augmentée (taux glycolytique accru). L'hexokinase (E.C.2.7.1.1) et le glucose-6-phosphate DH (E.C.1.1.1.49) n'ont pas été modifiés par la thyrotoxicose ou par l'hypothyroïdie (N = 5).

L'activité de la PEP-carboxykinase (4.1.1.31) est élevée dans la thyrotoxicose et abaissée dans l'hypothyroïdie, probablement avec des altérations équivalentes de la gluconéo-

\*) Supported by Deutsche Forschungsgemeinschaft SFB 51.

génése. Ce dernier résultat doit être discuté en rapport avec la manifestation d'un taux accru de cortisol libre dans le sérum et d'une efficacité insulinique diminuée chez nos malades thyrotoxiques.

Les enzymes du cycle de l'acide citrique et les voies connectées — citrate synthase (E.C.4.1.3.7), NADP-isocitrate DH (E.C.1.1.1.42), glutamate DH (E.C.1.4.1.2), aspartate amino-transférase (E.C.2.6.1.1), malate DH (E.C.1.1.1.37) — n'ont pas été changés par la thyrotoxicose ou l'hypothyroïdie.

L'enzyme malique (E.C.1.1.1.40) était augmenté dans les foies des malades thyrotoxiques. 3-hydroxyacyl-CoA DH (E.C.1.1.1.35) n'a pas été modifié, alors que la carnitine acétyl-transférase (E.C.2.3.1.7) était élevée dans la thyrotoxicose. On n'a pas observé une augmentation du  $\alpha$ -glycérol-phosphate DH mitochondrial (E.C.1.1.99.5) chez les malades thyrotoxiques. Ce résultat était inattendu étant donné l'induction connue de cet enzyme dans le foie de rats traités par des hormones thyroïdiennes.

Ces résultats sont discutés en ce qui concerne les différences d'espèce et le mécanisme d'action des hormones thyroïdiennes.

### Auszug

Die Diagnose von Schilddrüsenstörungen stützte sich auf den klinischen Status, Radiojoduntersuchungen,  $PB^{127}J$  und den  $T_3$ -*in-vitro*-Test. Nadelbiopsien aus Lebern wurden sofort eingefroren (flüssiger  $N_2$ ) und die Enzymaktivitäten im Überstand und den Sedimentfraktionen aller Homogenate jeder biopsischen Probe bestimmt.

Die Phosphoglucomutase (E.C.2.7.5.1) ist bei Thyreotoxikose verringert ( $N = 10$ ), im Vergleich zu den Kontrollen ( $N = 19$ ), was auf eine niedrige Glykogenolyse hinweist. Glyceraldehydphosphat DH (E.C.1.2.1.12) war erhöht (erhöhte Glykolyserate). Hexokinase (E.C.2.7.1.1) und Glukose-6-Phosphat DH (E.C.1.1.1.49) blieben bei Thyreotoxikose oder Hypothyreose unverändert ( $N = 5$ ).

Die Aktivität der PEP-Carboxykinase (4.1.1.31) ist bei Thyreotoxikose erhöht und bei Hypothyreose verringert, vermutlich mit äquivalenten Veränderungen der Gluconeogenese. Das letztgenannte Ergebnis muß im Zusammenhang mit dem Nachweis eines erhöhten Titers von freiem Cortisol im Serum bzw. einer verminderten Insulinwirksamkeit bei unseren thyreotoxischen Patienten erörtert werden. Enzyme des Citronensäurezyklus und dessen Nebenwege — Citrat Synthase (E.C.4.1.3.7), NADP Isocitrat DH (E.C.1.1.1.42), Glutamat DH (E.C.1.4.1.2), Aspartataminotransferase (E.C.2.6.1.1), Malat DH (E.C.1.1.1.37) — blieben durch Thyreotoxikose oder Hypothyreose unverändert.

Das Malat Enzym (E.C.1.1.1.40) war in Leberbiopsien von thyreotoxischen Patienten vermehrt. 3-Hydroxyacyl-CoA DH (E.C.1.1.1.35) war nicht verändert, während Carnitin-Acetyltransferase (E.C.2.3.1.7) bei Thyreotoxikose erhöht war. Mitochondriale  $\alpha$ -Glycerolphosphat DH (E.C.1.1.99.5) zeigte keine Erhöhung bei thyreotoxischen Patienten. Dies war im Hinblick auf die bekannte Induktion dieses Enzyms in der Leber von mit Schilddrüsenhormonen behandelten Ratten (LARDY) unerwartet.

Die Ergebnisse werden unter Berücksichtigung der Species-Unterschiede und dem Wirkungsmechanismus der Schilddrüsenhormone besprochen.

The diagnosis of the thyroid status of the patients under study was based on results of tests summarized in Table 1. Only unequivocal cases of thyrotoxicosis or hypothyroidism were selected; however the most severe cases of thyrotoxicosis had to be excluded owing to the need for immediate treatment (SCRIBA et al., 1970). In addition to "typical" values of  $PB^{127}I$  and  $T_3$ -*in-vitro*-test, in thyrotoxicosis elevated concentrations of "free" serum cortisol respectively in thyrotoxicosis and in hypothyroidism

**Table 1**

Clinical data of patients under study (mean + SD).

The techniques used for  $PB^{127}I$  (Autoanalyzer<sup>R</sup>), fluorimetric determination of serum cortisol, assay of "free"  $T_3$ - $^{125}I$  and  $^3H$ -cortisol by dextran gel filtration, i. v. glucose tolerance test and insulin efficiency coefficient have been published (SCRIBA et al., 1970). Age, body weight, clinical thyroid diagnostic index, data of radioiodine studies and various clinical chemical values of patients, have been reported (SCRIBA et al., 1970). Control patients submitted to liver biopsy were suspect for mild liver disease; the normal ranges reported were derived from healthy control persons

	hypo- thyroidism n = 9	control patients n = 20	thyro- toxicosis n = 18	normal range (mean ± 2 S.D.)
$PB^{127}I \times \% \text{ "free" } T_3\text{-}^{125}I$ ( $\mu g/100 \text{ ml}$ )	0.13 ± 0.06	0.98 ± 0.49	3.92 ± 1.95	0.44–8
serum cortisol × % "free" $^3H$ -cortisol ( $\mu g/100 \text{ ml}$ )	2.19 ± 0.71	1.13 ± 0.49	2.34 ± 1.53	—
i. v. glucose tolerance test (kG)	1.04 ± 0.24	1.55 ± 0.39	1.13 ± 0.35	1.2 – 2.2
insulin efficiency coefficient (mg glucose/mE IMI)	14.8 ± 13.7	20.6 ± 6.4	11.8 ± 6.9	20 – 70

decreased glucose tolerance and diminished insulin sensitivity were observed (Table 1). In thyrotoxicosis the histologic examination of liver biopsy samples revealed an increased width of sinusoids throughout the entire hepatic lobule, well distinguishable from congestive failure (SCRIBA et al., 1970).

### Biochemical methods

Biopsy samples from muscles (M. quadriceps, M. erector trunci) and needle biopsy samples from liver were frozen in liquid nitrogen immediately after removal. Extraction of the samples was performed according to (BASS et al., 1969). Soluble and structure-bound enzyme activities representative of glycogen metabolism, glucose oxidation, glycolysis, gluconeogenesis, citric acid cycle and connected pathways, fatty acid oxidation, acetyl transfer and glycerolphosphate oxidation were determined in standardized optical tests (BASS et al., 1969; BÜCHER et al., 1964; BRDIZKA et al., 1969). PEP-carboxikinase (carboxylation reaction) was measured according to CHANG and LANE 1966).

### Results

*Liver.* In thyrotoxicosis (n = 10) (Fig. 1), a distinct decrease ( $p < 0.01$ ) is found in the activity of phosphoglucomutase (PGM, E.C. 2.7.5.1). Hexokinase activity (HK, E.C. 2.7.1.1) is unaffected in thyrotoxicosis (measuring was performed with 2 mM glucose), although it is decreased

significantly in hypothyroidism ( $n = 5$ ) ( $p < 0.02$ ). No changes are found in the activity level of glucose-6-phosphate dehydrogenase (G6PDH, E.C. 1.1.1.49). Glyceraldehyde-phosphate dehydrogenase activity (GAPDH, E.C. 1.2.1.12) is lower in hypothyroidism and increases in thyrotoxicosis ( $p < 0.005$ ). With regard to enzymes of gluconeogenesis, no changes are observed in the activity of fructosediphosphatase (FDP-ase, E.C. 3.1.3.11); however, a marked increase occurs in the activity level of PEP-carboxykinase (PEP-CK, E.C. 4.1.1.32) in thyrotoxicosis. In hypothyroidism the activity of this enzyme decreases. Also, malic enzyme (ME, E.C. 1.1.1.40), is found at significantly higher activity levels in thyrotoxicosis.

The investigated enzymes of the tricarboxylic acid cycle and connected pathways such as condensing enzyme (CE, E.C. 4.1.3.7), isocitrate de-




Fig. 1. Enzyme activity pattern of human livers. Vertical bars indicate the standard error of the mean of 5 hypothyroid ( $\Delta$ ), 15 control ( $O$ ), 10 thyrotoxic ( $\bullet$ ) patients. \*PEP-CK activity was determined at  $37^\circ\text{C}$ . All other enzymes were measured at  $25^\circ\text{C}$ .

hydrogenase (IDH, NADP specific, E.C. 1.1.1.42), succinate dehydrogenase (SDH, E.C. 1.3.99.1), malate dehydrogenase (MDH, E.C. 1.1.1.37), glutamate dehydrogenase (GLUDH, E.C. 1.4.1.2), and glutamate oxaloacetate transaminase (GOT, E.C. 2.6.1.1), reveal little or no significant changes of their activity levels. This is also true for 3-hydroxyacyl-CoA dehydrogenase (HAD, E.C. 1.1.1.35). Carnitine acetyltransferase (CAT, E.C. 2.3.1.7) is markedly increased in thyrotoxicosis, whereas it is decreased in hypothyroidism. In patients with thyrotoxicosis or hypothyroidism the activities of glycerol-3-phosphate dehydrogenase (GDH, E.C. 1.1.1.8) and glycerolphosphate dehydrogenase (GP-OX, E.C. 1.1.99.5) remain normal.

These results are valid in terms of absolute as well as of specific activities. The amount of soluble protein shows only minor changes. The following values (mean  $\pm$  S.D.) were determined in hypothyroid, control and thyrotoxic liver samples:  $139 \pm 57$ ,  $106 \pm 41$ ,  $118 \pm 23$  mg of soluble protein/g of fresh weight.

*Muscle.* The data given in Table 2, refer to results obtained from muscle

**Table 2**

Activities and activity ratios of enzymes representative of different metabolic systems in normal and thyrotoxic human skeletal muscle

	M. erector trunci control <sup>1)</sup>	M. quadriceps thyrotoxic	
PH	514	436	292
HK	34	98	66
GAPDH	12 150	8 748	7 056
LDH	5 750	8 850	5 510
GP-OX	21	21	14
HAD	364	167	212
CE	230	127	92
HK/GAPDH	$2.8 \times 10^{-3}$	$11.2 \times 10^{-3}$	$9 \times 10^{-3}$
HK/CE	$1.4 \times 10^{-1}$	$7.7 \times 10^{-1}$	$7.2 \times 10^{-1}$
PH/GAPDH	$4.2 \times 10^{-2}$	$5 \times 10^{-2}$	$4.1 \times 10^{-2}$
GAPDH/GP-OX	$5.8 \times 10^2$	$4.2 \times 10^2$	$5 \times 10^2$
HAD/CE	1.5	1.3	2.3

<sup>1)</sup> Control values according to SCHIMRIGK et al., 1967.

biopsies of the two thyrotoxic patients examined so far. As is evident, thyrotoxicosis causes a marked increase in the activity level of HK. Corresponding increases in muscle HK activity have been observed in experimental thyrotoxicosis of rats and guinea pigs (SMITH and WILLIAMS-ASHMAN, 1951; BARGONI et al., 1967; KUBISTA et al., 1971). The increased activity level of HK is also obvious from the activity ratios listed at the bottom of Table 2. As has been pointed out elsewhere, the activity ratio

HK/CE is a more or less constant numerical value when different types of vertebrate muscles are compared (PETTE, 1966; BASS et al., 1969). This ratio, however, is elevated in thyrotoxicosis (KUBISTA et al., 1971), and consequently the activity ratio HK/GAPDH is also found to be increased.

## Discussion

Thyrotoxicosis causes certain changes in the enzyme patterns of liver and muscle. Obviously the changes observed in human liver differ qualitatively and quantitatively from those observed in experimental thyrotoxicosis, especially of the rat (LEE and LARDY, 1965; KADENBACH, 1966; NIKKILÄ and PITKÄNEN, 1959). Quantitative differences are probably due to differences existing between the levels of thyroid hormones in "physiological" and experimental thyrotoxicosis. Nevertheless, the decrease observed in the activity level of PGM as well as the increase in the activity levels of GAPDH, PEP-CK, ME and CAT correspond to similar changes found in experimental thyrotoxicosis.

The behaviour of GP-OX is of special interest. This enzyme is involved in extra-intramitochondrial hydrogen transfer by means of the glycerin-1-phosphate cycle. As first shown by LEE and LARDY (1965), experimental thyrotoxicosis causes an increase of GP-OX in rat liver by the factor of 10. As demonstrated by our findings, the human liver enzyme remains unaffected. This result is important with regard to the mechanism of action of thyroid hormones. It is generally believed that the increase of GP-OX assumes a predominant role in thyrotoxicosis. This increase holds for rat liver but not for human liver.

The differences are probably due to a species-specific response to thyroid hormones. This suggestion is also supported by the fact that differences exist between the changes observed in livers of thyrotoxic rats and guinea pigs. Thus, thyrotoxicosis does not cause an increase in activity levels of G6PDH, GP-OX and ME in the liver of the guinea pig, as this is typical for the rat liver. Species-specific differences may also be concluded from the data represented in Table 3. Table 3 lists activities of GAPDH and GP-OX as well as the activity ratios GAPDH/GP-OX (glycolysis/glycerolphosphate oxidation) in livers and skeletal muscles of different species. With regard to absolute activities of the two enzymes as well as their activity ratios, great differences exist in livers of different species. In the case of skeletal muscle, the activity ratio GAPDH/GP-OX varies to a much smaller degree (BASS et al., 1969; PETTE, 1966) than in the liver. In thyrotoxicosis, no changes are found in human muscle. In the rat and the guinea pig, the ratio GAPDH/GP-OX is shifted. These changes, however, occur only in red muscles (e.g. m. soleus), and are due to the fact that thyrotoxicosis induces an increase of GP-OX in this type of muscle only (KUBISTA et al., 1971). The most important change in the enzyme activity pattern of thyrotoxic human muscle is the high elevation of the HK activity. This change appears to be one of the few common and constant changes found in "physiological" and experimental thyrotoxicosis.



Table 3

Activities and activity ratios of glyceraldehydephosphate dehydrogenase and glycerol-phosphate oxidase in normal and thyrotoxic livers and muscles of various species. Activities are given as  $\mu\text{M}/\text{h}\times\text{g}$  w.w. Thyrotoxic animals received  $15\mu\text{g}$  3,3',5'-triiodo-thyronine per 100 g body weight daily for 6 days.

liver

	rat		guinea pig		pig	man	
	control	thyrotoxic	control	thyrotoxic	control	control	thyrotoxic
G A P D H	7 139	10 000	5 980	11 272	5 380	4 851	6 080
G P - O X	56	368	27	27	28	22	18
G A P D H							
G P - O X	130	25	218	417	270	220	340

muscle

	rat				guinea pig				man	
	M. rectus f.		M. soleus		M. rectus f.		M. soleus		M. erector trunci	M. quadriceps
	control	thyro-toxic	control	thyro-toxic	control	thyro-toxic	control	thyro-toxic	control	thyro-toxic
G A P D H										
G P - O X	200	210	300	140	280	166	234	155	580	455

## References

- BARGONI, N., M. A. GRILLO, M. T. RINAUDO and T. FOSSA: Über den Kohlenhydrat-Stoffwechsel in der Niere und im Skelettmuskel von mit Schilddrüsen oder Propylthiouracil gefütterten Ratten. Hoppe-Seyler's Z. Physiol. Chem. 348 (1967) 303.
- BASS, A., D. BRDICZKA, P. EYER, S. HOFER and D. PETTE: Metabolic differentiation of distinct muscle types at the level of enzymatic organization. Eur. J. Biochem. 10 (1969) 198.
- BRDICZKA, D., K. GERBITZ and D. PETTE: Localization and function of external and internal Carnitine acetyltransferases in mitochondria of rat liver and pig kidney. Eur. J. Biochem. 11 (1969) 234.
- BÜCHER, T., W. LUH and D. PETTE: Einfache und zusammengesetzte optische Tests mit Pyridinukleotiden, in: Hoppe-Seyler/Thierfelder, Handb. physiol.- u. pathol.-chem. Analyse, 10. Aufl., Bd. VI/A, S. 292. Springer-Verlag Berlin, Göttingen, Heidelberg, New York (1964).
- CHANG, H.-C. and M. D. LANE: The enzymatic carboxylation of phosphoenolpyruvate. J. Biol. Chem. 241 (1966) 2413.
- KADENBACH, B.: Der Einfluß von Thyroidhormonen *in vivo* auf die oxidative Phosphorylierung und Enzymaktivitäten in Mitochondrien. Biochem. Zschr. 344 (1966) 49.

- KUBISTA, V., J. KUBISTOVA and D. PETTE: Thyroid hormone induced changes in the enzyme activity pattern of energy-supplying metabolism of fast (white), slow (red), and heart muscle of the rat. *Eur. J. Biochem.* 18 (1971) 553.
- LEE, Y. P. and H. A. LARDY: Influence of thyroid hormones on L- $\alpha$ -Glycerophosphate dehydrogenases and other dehydrogenases in various organs of the rat. *J. Biol. Chem.* 240 (1965) 1427.
- NIKKILÄ, E. A. and E. PITKÄNEN: Liver enzyme pattern in thyrotoxicosis. *Acta Endocr. (Kph.)* 31 (1959) 573.
- PETTE, D.: Energieliefernder Stoffwechsel des Muskels unter zellphysiologischem Aspekt. Symposium über progressive Muskeldystrophie. Springer-Verlag Berlin, Göttingen, Heidelberg, New York (1966) S. 492.
- SCHIMRIGK, K., H. G. MERTENS, K. RICKER, J. FÜHR, P. EYER and D. PETTE: McArdle Syndrom. *Klin. Wschr.* 45 (1967) 1.
- SCRIBA, P. C., B. BACHMAIER, B. BAUER, N. BOSS, P. DIETERLE, K. GERBITZ, J. HENNER, K. HORN, P. KIEFHABER, J. NOLTE, D. PETTE, H. SCHNEIDER and A. SOUVATZOGLOU: Der Einfluß von Hyper- und Hypothyreose auf Enzymaktivitäten der menschlichen Leber. *Verh. Dtsch. Ges. Inn. Med.* 76 (1970) 483.
- SMITH, R. H. and H. G. WILLIAMS-ASHMAN: The influence of thyroxine on the enzyme activity of rat tissues. *Biochem. Biophys. Acta* 7 (1951) 295.

The authors are indebted to Dr. K. GERBITZ for measuring the carnitine acetyltransferase activities, to Dr. F. MITTELBACH and Dr. D. PONGRATZ for muscle biopsies and to Mrs. B. BAUER, S. TRIMBORN, I. SICHART and R. FRÖHLICH for conscientious technical assistance.

## List of authors and co-authors with page

- Adams, D. D. 1049  
 Adiga, P. R. 281  
 Ahn, C. S. 825  
 Alexander, N. M. 423  
 Alexander, W. D. 143  
 Almqvist, S. 181  
 Alper, C. A. 1089  
 Al-Saadi, A. A. 807  
 Andreoli, M. 105, 457  
 Appleman jr., M. D. 1173  
 Arima, T. 363  
  
 Bachmaier, B. 1005  
 Bastomsky, C. H. 263, 863  
 Bayse, G. S. 741  
 Beall, G. N. 557, 565  
 Becker, D. V. 603  
 Beckers, C. 255  
 Bénard, B. 771  
 Benotti, J. 1121  
 Ben-Porath, M. 643  
 Bertrand, A. 879  
 Björkman, U. 381  
 Bornet, H. 97  
 Bouchilloux, Simone 371  
 Bouhnik, J. 731  
 Bowers, C. Y. 1029  
 Brault, J. 771  
 Braverman, L. E. 243  
 Bray, G. A. 533  
 Brown, B. L. 1107  
 Brown, M. 663  
 Brownlie, B. E. W. 143  
 Burns, A. H. 943  
 Burrows, B. A. 993  
 Bürgi, H. 225, 519  
 Buttfeld, I. H. 53  
  
 Cahnmann, H. J. 441  
 Camus, M. 85  
 Camuzzini, G. F. 511  
 Cash, A. G. 839  
 Cassano, C. 105, 457  
 Cassidy, C. E. 989  
 Cavaliere, R. R. 327  
 Chabaud, Odile 371  
 Chambers jr., J. B. 233  
 Chang, R. F. 351  
 Chiariello, M. 1205  
 Chiotaki, L. 723  
 Chopra, I. J. 557, 565  
 Clark, M. B. 1233  
  
 Clark, R. L. 1245  
 Comar, D. 511  
 Cook, D. M. 1165  
 Cortese, F. 201  
 Coutsoftides, Th. 1085  
 Covelli, I. 201  
 Cramarossa, L. 105  
  
 D'Angelo, S. A. 1183  
 Danner, D. J. 741  
 D'Armiento, M. 105  
 Degrossi, O. 343  
 Degroot, L. J. 297, 413  
 Dekker, A. 817, 1197  
 Delange, F. 35  
 Dell, E. S. 993  
 De Rubertis, F. 1197  
 Di Stefano, J. J. 351  
 Dobyns, B. M. 603  
 Docter, R. 255  
 Dolman, A. 255  
 Dorrington, K. J. 571  
 Downs, P. 243  
 Dumont, J. E. 291, 307  
 Dussault, J. H. 237, 273  
  
 Edelhoch, H. 97, 113, 201  
 Ekholm, R. 181, 381  
 Ekins, R. P. 1107  
 Ellis, S. M. 1107  
 Ericson, L. E. 779  
 Ermans, A. M. 35, 85  
 Eshchenko, N. D. 701  
 Estrella, E. 61  
  
 Falconer, J. R. 481  
 Feige, Y. 643  
 Fellingner, K. 13  
 Field, J. B. 817, 1197  
 Fierro-Benitez, R. 61  
 Fisher, D. A. 237, 273  
 Florsheim, W. H. 549  
 Folkers, K. 1029  
 Fontana, S. 457  
 Foster, A. E. 243  
 Foster, G. V. 1233  
 Fregly, M. J. 943  
 Fritzsche, H. 907  
 Furnival, B. E. 1057  
  
 Galante, L. 1233  
 Garcia, M. D. 1137

- Ghayasuddin, M. 493  
 Gillespie, F. C. 619  
 Girault, M. 511  
 Gluckman, P. D. 653  
 Goolden, A. W. G. 663  
 Gordon, A. 407, 1085  
 Grahame-Smith, D. 1233  
 Gray, H. W. 619  
 Green, W. L. 965  
 Greer, M. A. 399, 973  
 Greig, W. R. 619  
 Griessen, M. 1069  
 Grimaldi, R. 1121  
 Gross, H. A. 1173  
 Gross, J. 407  
 Gschwendtova, K. 705  
 Gudmundsson, T. V. 1233  
 Guichard, C. 919
- Haddad, A. 387  
 Hadden, D. R. 983  
 Haibach, D. R. 399  
 Harland, W. A. 1127  
 Hart, I. R. 595  
 Hartemann, P. 879  
 Heimann, P. 787  
 Hennemann, G. 255  
 Herrmann, J. 1021  
 Herscovics, Annette 387  
 Hershman, J. M. 35, 1155  
 Hetzel, B. S. 53  
 Hidaka, H. 765  
 Higgins, H. P. 1155  
 Hill Jr., C. S. 1245  
 Hill, J. B. 839  
 Hjort, T. 163  
 Hobel, C. J. 273  
 Höfer, R. 137, 907  
 Horton, R. 1233  
 Hunton, R. B. 653
- Ibbertson, H. K. 653  
 Ibanez, M. L. 1245  
 Ingbar, S. H. 121, 243, 683  
 Inoue, K. 973
- Jaarsveld, P. P. van 465  
 Jereb, Berta 1221  
 Jost, A. 189  
 Jung, A. 225
- Kaneko, T. 817  
 Kapitola, J. 913  
 Kellershohn, C. 511  
 Kendall, J. W. 1165  
 Kennedy, T. H. 1049  
 Keyhani, E. M. 307  
 Kiefhaber, P. 1005  
 Knopp, J. 311  
 Kobayashi, I. 399  
 Kohler, H. 225, 519  
 Koutras, D. A. 79
- Kriss, J. P. 577  
 Krusius, F. E. 149  
 Krüskemper, H. L. 1021  
 Kyncl, F. 797
- Labrique, M. 173  
 Lamas, L. 201  
 Lambert, P. H. 583  
 Lamy, F. M. 307  
 Langer, P. 155, 705  
 Laor, J. 643  
 Lauridsen, U. B. 163  
 Lauringer, P. 137  
 Leclère, J. 879  
 Lecocq, R. E. 307  
 Lemarchand-Béraud, T. 1069  
 Lewitus, Z. 503, 643  
 Lindsay, R. H. 839  
 Lissitzky, S. 189  
 Ljunggren, J. G. 1221  
 Lowry, R. C. 983  
 Lubin, E. 643  
 Lutherer, B. C. 943  
 Lutherer, L. O. 943
- Maayan, M. L. 683  
 Macchia, V. 1205  
 Malamos, B. 79  
 Malan, P. G. 113  
 Maloof, F. 1121  
 Mantzos, J. 79, 723  
 Marchant, B. 143  
 McConahey, W. M. 603  
 McConnon, J. 335  
 McDougall, I. R. 619  
 McGirr, E. M. 619  
 McKenzie, J. M. 281  
 Melander, A. 779  
 Meldolesi, M. F. 1205  
 Michajlovskij, N. 155  
 Michel, O. 731  
 Michel, R. 731  
 Michel-Bechet, M. 371, 797  
 Middlesworth, L. van 525, 891  
 Monaco, F. 457  
 Montgomery, D. A. D. 983  
 Mori, T. 577  
 Morreale de Escobar, G. 1137  
 Morrison, M. 741  
 Mothon, Stella 533  
 Munro, D. S. 571, 611  
 Murthy, P. V. N. 281
- Nagasaka, A. 413, 765  
 Nathanson, B. M. 1233  
 Nayer, P. de 173, 255  
 Nève, P. 291  
 Nichols, E. M. 1165  
 Nicoloff, J. T. 1173  
 Nolte, J. 1005  
 Nunez, J. 433, 693

- Ogawara, H. 441  
 Ohtaki, S. 749  
 Olin, P. 181, 1221  
 Orr, J. S. 1127  
 Ossoinig, K. 871  
 Otten, J. 291  
 Owman, Ch. 779
- Paice, J. C. 1057  
 Palacios, P. 45  
 Paumgartner, G. 137  
 Pavlovic-Hournac, M. 693  
 Pazmanova, L. 705  
 Pecorini, V. 343  
 Peltola, P. 149  
 Persson, I. B. 163  
 Pette, D. 1005  
 Pharmakiotis, A. D. 79  
 Pharoah, P. O. D. 53  
 Pino, S. 1121  
 Pisarev, M. A. 297  
 Pittman, C. S. 233  
 Pitt-Rivers, Rosalind 15  
 Pommier, J. 433  
 Premachandra, B. N. 1097  
 Pretell, E. A. 45  
 Putilina, F. E. 701
- Querido, A. 61
- Rachev, R. R. 701  
 Ramirez, I. 61  
 Rappaport, L. 693  
 Read, V. H. 233  
 Rechnic, J. 643  
 Refetoff, S. 1089  
 Reynolds, G. A. 1029  
 Rigopoulos, G. A. 79  
 Rivièrè, R. 511  
 Robbins, J. 451  
 Robert, J. 879  
 Robin, N. I. 1089  
 Robison, W. L. 891  
 Rocmans, P. 291  
 Rodesch, F. 291  
 Roques, M. 189  
 Rosenberg, I. N. 749, 825  
 Rosenberg, L. L. 957  
 Row, V. V. 335
- Saaman, N. A. 1245  
 Saito, T. 1013  
 Salabé, G. B. 457  
 Salmon, J. 583  
 Salvatore, G. 201  
 Scazziga, B. R. 1069  
 Schally, A. V. 1029  
 Schatz, H. 907  
 Schneider, A. 97  
 Schneider, H. 1005  
 Schreiberova, O. 913
- Schüllerova, M. 913  
 Schussler, G. C. 319  
 Scriba, P. C. 1005  
 Scuncio, G. 105  
 Searle, G. L. 327  
 Sena, L. 465  
 Sfontouris, J. 79  
 Shaham, Y. 503  
 Shapiro, R. J. 683  
 Sheline, G. E. 603  
 Shibasaki, K. 711  
 Shimaoka, K. 1215  
 Shimizu, T. 1013  
 Shishiba, Y. 1013  
 Shizume, K. 1013  
 Shulman, S. 493  
 Slingerland, D. W. 993  
 Smith, B. R. 571  
 Smith, J. F. B. 619  
 Smith, Meredith 387  
 Smith, R. N. 611  
 Sokal, J. E. 1215  
 Sokoloff, L. 433  
 Solomon, D. H. 557, 565  
 Spira, O. 407  
 Spiro, Mary J. 363  
 Spiro, R. G. 363  
 Stanbury, J. B. 61  
 Starnes, W. R. 1155  
 Steinberg, M. 327  
 Sterling, K. 243  
 Stolc, V. 311  
 Studer, H. 225, 519  
 Sunder, F. 779  
 Suzuki, M. 711
- Tashjian Jr., A. H. 1245  
 Taurog, A. 211, 973  
 Tholin, B. 787  
 Thomson, J. A. 619  
 Tompkins, Edythalena 603  
 Tong, W. 311  
 Torresani, J. 189  
 Triantaphyllidis, H. E. 919
- Ui, Nobuo 91  
 Utiger, R. D. 1049
- Vagenakis, A. 243  
 Valenta, L. 797  
 Vassart, G. 307  
 Visscher, M. de 173  
 Vlassis, G. 79  
 Volpé, R. 335  
 Vormittag, E. 137  
 Vormittag, W. 137
- Wall, N. R. 1183  
 Walt, B. van der 465  
 Wan, M. 45  
 Wartofsky, L. 121

Weaver, J. A. 983  
Webster, B. R. 1057  
Weil, A. 1029  
White, B. M. L. 653  
Whur, P. 387  
Wilber, J. F. 297, 1041  
Willems, C. 291  
Williams, E. D. 663  
Williams, E. S. 1107  
Williams, J. A. 677  
Wilson, B. D. 849  
Wilson, G. M. 611

Winand, R. J. 583  
Winship, Th. 25  
Wolff, J. 669, 677  
Workman, J. B. 603  
Wright, R. L. 849

Yamashita, K. 817, 1197

Zakarija, M. 863  
Zaninovich, A. A. 343  
Zor, U. 817  
Zyl, A. van 465