

# $\text{Na}_3(\text{PO}_2\text{NH})_3 \cdot 4\text{H}_2\text{O}$ , $[\text{C}(\text{NH}_2)_3]_3(\text{PO}_2\text{NH})_3 \cdot \text{H}_2\text{O}$ and $(\text{NH}_4)_4(\text{PO}_2\text{NH})_4 \cdot 4\text{H}_2\text{O}$

Annamma John <sup>a</sup>, Daizy Philip <sup>b</sup>, Norbert Stock <sup>c</sup>, Wolfgang Schnick <sup>c</sup>,  
S. Devanarayanan <sup>d,\*</sup>

<sup>a</sup> Department of Physics, St. John's College, Anchal, Kollam 691 306, India

<sup>b</sup> Department of Physics, Mar Ivanios College, Thiruvananthapuram 695 015, India

<sup>c</sup> Laboratorium für Anorganische Chemie der Universität, D-95440 Bayreuth, Germany

<sup>d</sup> Department of Physics, University of Kerala, Kariavattom, Thiruvananthapuram 695 581, India

Received 31 July 2000; accepted 9 August 2000

## Abstract

FT IR and FT Raman spectra of  $\text{Ag}_3(\text{PO}_2\text{NH})_3$  (Compound I),  $\text{Na}_3(\text{PO}_2\text{NH})_3 \cdot \text{H}_2\text{O}$  (Compound II),  $\text{Na}_3(\text{PO}_2\text{NH})_3 \cdot 4\text{H}_2\text{O}$  (Compound III),  $[\text{C}(\text{NH}_2)_3]_3(\text{PO}_2\text{NH})_3 \cdot \text{H}_2\text{O}$  (Compound IV) and  $(\text{NH}_4)_4(\text{PO}_2\text{NH})_4 \cdot 4\text{H}_2\text{O}$  (Compound V) are recorded and analyzed on the basis of the anions, cations and water molecules present in each of them. The  $\text{PO}_2\text{NH}^-$  anion ring in compound I is distorted due to the influence of  $\text{Ag}^+$  cation. Wide variation in the hydrogen bond lengths in compound III is indicated by the splitting of the  $\nu_2$  and  $\nu_3$  modes of vibration of water molecules. The  $\text{NH}_4$  ion in compound V occupies lower site symmetry and exhibits hindered rotation in the lattice. The correlations between the symmetric and asymmetric stretching vibrations of P–N–P bridge and the P–N–P bond angle have also been discussed. © 2001 Elsevier Science B.V. All rights reserved.

**Keywords:** Infrared spectrum; Raman spectrum; Trimetaphosphimate; Tetrametaphosphimate; Hydrogen bonding

## 1. Introduction

Many salts of trimetaphosphimic acid and tetrametaphosphimic acid were known more than 100 years ago [1,2]. These compounds have been mainly characterized by IR spectroscopy [3,4] and

X-ray powder diffraction [5]. Extensive amount of work has been done on the study of the characteristic features of the coordination of metaphosphimate anions by vibrational spectroscopy, involving the calculation of wavenumbers and forms of the normal vibrations [6–16]. The trimetaphosphimate anion is a cyclic system formed by three regularly alternating slightly distorted  $\text{PO}_2\text{NH}$  tetrahedra. The confirmation of the ring in the chair form with symmetry close to

\* Corresponding author. Tel.: +91-471-418920; fax: +91-471-307158.

E-mail address: phy@univker.ernet.in (S. Devanarayanan).

$C_{3v}$  has been established by X-ray structure analysis [17] and from the vibrational spectra of alkali metal trimetaphosphimates [9]. From a study of the vibrational spectra it is known that the tetrametaphosphimate anion in solution has  $S_4$  symmetry and that in salts it has two kinds of structures, viz.,  $C_{2h}$  symmetry and  $S_4$  symmetry [14].

The present authors are interested in the study of the Raman and IR spectra of some newly synthesized phosphate compounds [18–20]. In this chapter a report of the analysis of both the IR and Raman spectra of  $Ag_3(PO_2NH)_3$  (hereafter referred as compound **I**),  $Na_3(PO_2NH)_3 \cdot 3H_2O$  (compound **II**),  $Na_3(PO_2NH)_3 \cdot 4H_2O$  (compound **III**),  $[C(NH_2)_3]_3(PO_2NH)_3 \cdot H_2O$  (compound **IV**) and  $(NH_4)_4(PO_2NH)_4 \cdot 4H_2O$  (compound **V**) is made. The Raman spectra of these samples in aqueous solution are also recorded to clear the ambiguity regarding certain vibrations.

## 2. Experimental

The starting material  $Na_2(PO_2NH)_3 \cdot 4H_2O$  (compound **III**) was prepared by the method reported by Olthof et al. [17]. To an acidified solution of this compound, an aqueous solution of  $AgNO_3$  is added to get  $Ag_3(PO_2NH)_3$  (compound **I**) [21].  $Na_3(PO_2NH)_3 \cdot H_2O$  (compound **II**) was obtained by vapour diffusion of  $CH_3CH_2OH$  at 363 K into a  $3 \times 10^{-4}$  M aqueous solution of the starting material [22].  $[C(NH_2)_3]_3(PO_2NH)_3 \cdot H_2O$  (compound **IV**) was obtained by the addition of  $[C(NH_2)_3]_2CO_3$  to a freshly prepared solution of  $H_3(PO_2NH)_3$  [23]. For the preparation of

$(NH_4)_4(PO_2NH)_4 \cdot 4H_2O$  (compound **V**), the starting material  $H_4(PO_2NH)_4 \cdot 2H_2O$  was obtained according to the literature [6] by hydrolysis (333 K, 48 h) of  $(PNCl_2)_3$  in dioxane. Then an aqueous solution of ammonia (10%) is added to a stoichiometric amount of the acid followed by precipitation with acetone [24].

The IR and Raman spectra were recorded using a Bruker IFS 66V FT-IR/FT-Raman Spectrometer. The Raman spectra were recorded with a resolution of  $4 \text{ cm}^{-1}$ , using an Nd:YAG laser at wavelength  $1.064 \mu\text{m}$  and power 200 mW, over the range  $50\text{--}3500 \text{ cm}^{-1}$ . The IR spectra are obtained in the range  $400\text{--}4000 \text{ cm}^{-1}$  with the samples as KBr pellets using a globar source.

## 3. Factor group analysis

The compounds **I**, **II**, **III** and **V** crystallizes in the monoclinic system and compound **IV** is orthorhombic. The crystal symmetry data of all the five compounds used in the present work are tabulated in Table 1.

The cations, anions and water molecules in all the compounds occupy general position. The trimetaphosphimate anion  $(PO_2NH)_3^{3-}$  and tetrametaphosphimate anion  $(PO_2NH)_4^{4-}$  exhibit chair conformation, but in compound **I** the anion is markedly distorted because of the interaction between  $Ag^+$  and one nitrogen atom of the anion [21]. The  $(PO_2NH)_3^{3-}$  ring has  $C_{3v}$  symmetry [13] and the  $(PO_2NH)_4^{4-}$  ring has  $C_{2h}$  symmetry [15]. The cyclic anion rings in trimetaphosphimates are linked to one another by N–H---O hydrogen bonds to form pairs. These units are interconnected by O–H---O hydrogen bonds through water molecules forming columns. In compound **V** the tetrametaphosphimate rings are linked by N–H---O bonds forming column, along [100]. These columns are interconnected by O–H---O and N–H---O hydrogen bonds through water molecules and ammonium ions [21–24]. All H atoms are involved in hydrogen bonding. In **II** and **III** the sodium ions are approximately octahedrally coordinated by six oxygen atoms, either from the phosphonate groups or from the water

Table 1  
Crystallographic datas of the five compounds

Compounds	Crystallography	Space group	Z <sup>B</sup>	Refs.
<b>I</b>	Monoclinic	$P2_1/c(C_{2h}^5)$	4	[21]
<b>II</b>	Monoclinic	$C2(C_2^3)$	2	[22]
<b>III</b>	Monoclinic	$P2_1/n(C_{2h}^5)$	4	[17]
<b>IV</b>	Orthorhombic	$Pbca(D_{2h}^{15})$	8	[23]
<b>V</b>	Monoclinic	$P2_1/n(C_{2h}^5)$	2	[24]

molecules. The factor group analysis using the standard correlation method [25], for each compound gives the irreducible representation at  $k = 0$  as

Compound I	$54A_g(\text{R}) + 54B_g(\text{R}) + 53A_u(\text{IR}) + 52B_u(\text{IR})$
Compound II	$62A(\text{R,IR}) + 61B(\text{R, IR})$
Compound III	$90A_g(\text{R}) + 90B_g(\text{R}) + 89A_u(\text{IR}) + 88B_u(\text{IR})$
Compound IV	$144A_g(\text{R}) + 144B_1g(\text{R}) + 144B_2g(\text{R})$ $+ 144B_3g(\text{R}) + 144A_u(\text{O}) + 143B_{1u}(\text{IR})$ $+ 143B_{2u}(\text{IR}) + 143B_{3u}(\text{IR})$
Compound V	$78A_g(\text{R}) + 78B_g(\text{R}) + 77A_u(\text{IR}) + 76B_u(\text{IR})$

#### 4. Results and discussions

The IR and Raman spectra recorded for all the five compounds under investigation are shown in Figs. 1–3. The observed spectral lines and their assignments are given in Table 2.

##### 4.1. Vibrations of $\text{PO}_2\text{NH}^-$ anion

Analysis of the spectra obtained shows that  $\nu\text{NH}$  modes are highly characteristic. These are obtained as weak bands in Raman and medium to strong in IR around  $3170\text{ cm}^{-1}$  in all the five compounds. The multiplicity of bands observed in compound I may be due to the participation of the NH groups in intramolecular and intermolecular hydrogen bonding [13]. The frequency shift observed in compound IV may be attributed to the overlapping of the  $\nu\text{NH}$  vibrations with that of  $\nu\text{NH}_2$  vibrations. The NH deformations are observed in the region  $1390$ – $1322$  and  $781$ – $723\text{ cm}^{-1}$ .

As expected the asymmetric and symmetric stretching vibrations of  $\text{PO}_2$  group are observed in the region  $1241$ – $1141$  and  $1118$ – $1039\text{ cm}^{-1}$  respectively [26]. The asymmetric modes are very strong in the IR spectrum and the symmetric modes are very strong in the Raman spectrum. Both the asymmetric and symmetric stretching modes of  $\text{PO}_2$  in compound V appear a little lower in frequency when compared to all the

other four compounds. This is due to the greater value of the average P–O bond length in compound V than that in all the other compounds [24]. The bending modes of  $\text{PO}_2$  group are observed around  $600\text{ cm}^{-1}$ . The wagging, rocking and twisting modes of  $\text{PO}_2$  group are also assigned (Table 2).

The ring vibrations of the anion — the anti-symmetric and symmetric stretches of P–NH–P bridges are expected in the region  $1000$ – $800\text{ cm}^{-1}$  and the ring breathing vibration around  $400\text{ cm}^{-1}$  [12,15]. In compound I the asymmetric stretching gives very weak Raman bands at  $971$  and  $937\text{ cm}^{-1}$  and the symmetric stretching bands are observed at  $831$  and  $793\text{ cm}^{-1}$ . The corresponding IR bands are singlets observed at  $933$  and  $813\text{ cm}^{-1}$  respectively. The large splitting ( $\approx 138\text{ cm}^{-1}$ ) of the symmetric mode observed in the Raman spectrum confirms the distortion of the  $\text{PO}_2\text{NH}^-$  anion ring due to the influence of the stronger  $\text{Ag}^+$  cation on the anion structure [21]. In the Raman spectrum of compound V, a weak band is observed at  $1008\text{ cm}^{-1}$ . Sukova et al. [15], in their studies on lanthanum and lanthanide tetrametaphosphimate had assigned the strong IR band around  $\approx 1005\text{ cm}^{-1}$  to  $\nu_{\text{as}}\text{P-NH-P}$  along with its components in the region  $980$ – $900\text{ cm}^{-1}$ . Accordingly the Raman band at  $1008\text{ cm}^{-1}$ , in the present study, is assigned to  $\nu_{\text{as}}\text{P-NH-P}$ . Its components are the very weak bands at  $980$ – $910\text{ cm}^{-1}$ .

The corresponding IR bands are very strong at  $992$  and  $955\text{ cm}^{-1}$ . The weak to medium intensity Raman bands around  $493\text{ cm}^{-1}$  are due to the ring breathing vibrations. The IR spectra show medium to strong intensity bands in this region. The external deformation mode is assigned to the weak bands around  $225\text{ cm}^{-1}$  [9,12].

##### 4.2. Vibrations of $\text{NH}_4^+$ cation

The normal modes of vibration of free  $\text{NH}_4^+$  ion under perfect  $T_d$  symmetry have frequencies of  $3033$ ,  $1685$ ,  $3134$  and  $1397\text{ cm}^{-1}$  for  $\nu_1(A_1)$ ,  $\nu_2(E)$ ,  $\nu_3(F_2)$  and  $\nu_4(F_2)$  modes, respectively [26].

Table 2  
Spectral data ( $\text{cm}^{-1}$ ) and band assignments of  $\text{Ag}_3(\text{PO}_2\text{NH})_3$ ,  $\text{Na}_3(\text{PO}_2\text{NH})_3 \cdot \text{H}_2\text{O}$ ,  $\text{Na}_3(\text{PO}_2\text{NH})_3 \cdot 4\text{H}_2\text{O}$ ,  $[\text{C}(\text{NH}_2)_3]_3(\text{PO}_2\text{NH})_3 \cdot \text{H}_2\text{O}$  and  $(\text{NH}_4)_4(\text{PO}_2\text{NH})_4 \cdot 4\text{H}_2\text{O}^a$

Raman	$\text{Na}_3(\text{PO}_2\text{NH})_3 \cdot \text{H}_2\text{O}$		$\text{Na}_3(\text{PO}_2\text{NH})_3 \cdot 4\text{H}_2\text{O}$		$[\text{C}(\text{NH}_2)_3]_3(\text{PO}_2\text{NH})_3$		$(\text{NH}_4)_4(\text{PONH})_4 \cdot 4\text{H}_2\text{O}$		Assignments
	IR	Raman	IR	Raman	IR	Raman	IR	Raman	
	3472wbr		3455m		3586m		3571m		Combination
3273w			3372m		3418s		3353sh		$\nu_3\text{H}_2\text{O}$
3233w			3310m		3350sh		3265sh		$\nu_1\text{H}_2\text{O}$
3086vw	3214w	3234s	3300wbr	3352w	3350sh	3301wbr	3213sh		$\nu_3\text{NH}_4$
	3173w	3168s	3173wbr	3173m	3126s		3162vs		$\nu\text{NH}/\nu\text{NH}_2$
	2684w	2678w		2704vww			3018sh		$\nu_1\text{NH}_4$
	2596vw	2659w		2644vww			2853m		Combination
		2575w					2716sh		$\nu\text{NH} \cdots \text{N}(\text{O}^-)$
2208vw		2321vw							Combinations and overtones
2119vw	2124vw	2185w	2085vwbr				2359vw		
		1720w					2145m		
	1643w	1686sh	1670vwbr	1694w	1649sh	1688vw	1699w		$\nu_2\text{NH}_4$
				1661w			1653m		$\nu_2\text{H}_2\text{O}$
				1693w					$\nu_{\text{as}}\text{CN}$
1471w		1545w			1663vs				$\delta\text{NH}_2$
					1442sh				Combinations
					1588vw		1461s		
							1419sh		$\nu_4\text{NH}_4$
							1402s		
1390vww		1355sh		1355m			1385sh		$\delta\text{NH}$
1369vww	1345s	1335vs		1326m			1362sh		$\delta\text{O}-\text{H} \cdots \text{O}$
				1281m			1343s		
				1241s					
				1225s					
	1226vs	1222w	1238m	1234vw					
			1198m	1192m					$\nu_{\text{as}}\text{PO}_2$
							1210s	1192sh	

Table 2 (Continued)

Ag <sub>5</sub> (PO <sub>2</sub> NH) <sub>3</sub>		Na <sub>3</sub> (PO <sub>2</sub> NH) <sub>3</sub> ·H <sub>2</sub> O		Na <sub>3</sub> (PO <sub>2</sub> NH) <sub>3</sub> ·4H <sub>2</sub> O		[C(NH <sub>2</sub> ) <sub>3</sub> ] <sub>3</sub> Δ(PO <sub>2</sub> NH) <sub>3</sub> ·H <sub>2</sub> O		(NH <sub>4</sub> ) <sub>4</sub> (PONH) <sub>4</sub> ·4H <sub>2</sub> O		Assignments
Raman	IR	Raman	IR	Raman	IR	Raman	IR	Raman	IR	
1167w		1185w	1187s	1182m		1174vw		1180w 1148w	1141s	
1094vs	1111m	1107vs	1101s	1107vs	1118w	1106m	1102m	1080vs		<i>v</i> <sub>g</sub> PO <sub>2</sub>
1040w	1054vs	1059w	1060vs	1054m	1066vs	1049vw	1046vs		1063s	
		1039w	1043vs	1047m	1054s	1007vs	1006w	1008w	992s	
971vw								980vw		<i>v</i> <sub>sat</sub> P–NH–P
937vw	933vs	955w	952vs	955vw	960s		946sh	910vw	955vs	
831vw	813m		826s	845w	860vw	853vw	929s	859vw		
793m					821m				818m	<i>v</i> <sub>g</sub> P–NH–P
755s	774sh 731m	803m	781m	806w	802m		779sh			H <sub>2</sub> O liberation δNH
610s	602m 540s	658vw 607m	662m 599m 542s	726m 601m	740m 650sh 608m	723wbr 607vw	608m	596vw 560w	589m 556m	δPO <sub>2</sub>
512w	504s	524m			536m	531m	515w			δCN <sub>3</sub>
470w			513s	517m	504m		493w		506s	Ring breathing
402			480m	478m			492m 424sh	474w 423w	421vw	
		374m		377sh 360s		385sh 351w		368s		ωPO <sub>2</sub>
345w										
334w		318sh		297s		292w		332m		ρPO <sub>2</sub>
297m		295m		247w		277w		288m		
258vw		231vw						228m		Ring deformation
167sh				157vw				209m		
124m		131w		129vw				166m		tPO <sub>2</sub>
		114sh						137m		<i>v</i> N–H---O
84s		84m		84m				105m		<i>v</i> O–H---O
								84m		External mode

<sup>a</sup> Relative intensities: v, very; w, weak; m, medium; s, strong; sh, shoulder; br, broad.




Fig. 1. FT IR spectra of (a)  $\text{Ag}_3(\text{PO}_2\text{NH})_3$ , (b)  $\text{Na}_3(\text{PO}_2\text{NH})_3 \cdot \text{H}_2\text{O}$  (c)  $\text{Na}_3(\text{PO}_2\text{NH})_3 \cdot 4\text{H}_2\text{O}$ , (d)  $[\text{C}(\text{NH})_3]_3(\text{PO}_2\text{NH})_3 \cdot \text{H}_2\text{O}$  and (e)  $(\text{NH}_4)_4(\text{PO}_2\text{NH})_4 \cdot 4\text{H}_2\text{O}$  in the region  $400\text{--}4000\text{ cm}^{-1}$ .

All the four modes are Raman active whereas  $\nu_1$  and  $\nu_2$  are IR inactive. In the crystal, the ion occupies a lower site symmetry, which may lead

to the activation of inactive modes, splitting of degenerate modes and shifting of fundamental modes.

In the Raman spectrum of compound **V**, the weak broad band at  $3058\text{ cm}^{-1}$  is assigned to the symmetric stretching mode  $v_1$ . The corresponding IR band appears as a shoulder at  $3018\text{ cm}^{-1}$ . The broadening ( $\approx 250\text{ cm}^{-1}$ ) of the  $v_1$  mode indicate the participation of the  $\text{NH}_4$  ions in the formation of different types of hydrogen bonds  $\text{N-H}\cdots\text{N}$  and  $\text{N-H}\cdots\text{O}$  of varying strength [27]. The structural investigation of compound **V** [24] reports that the  $\text{NH}_4$  ions take part in extended hydrogen bonding by bridging, via, the lone pairs of  $\text{OW2}$ ,  $\text{N4-H42}\cdots\text{OW2}\cdots\text{H31-N3}$  ( $\text{N}\cdots\text{O}\cdots\text{N}$   $2.801, 2.858\text{ \AA}$ ). The observation of strong IR band extending from about  $3700$  to  $2700\text{ cm}^{-1}$  ( $\text{FWHM} \approx 720\text{ cm}^{-1}$ ) also supports this. The absorption shoulders at

$3265$  and  $3213\text{ cm}^{-1}$  are due to the triply degenerate asymmetric stretching mode  $v_3$ . The  $v_2$  mode is observed as weak absorption band at  $1699\text{ cm}^{-1}$ . The very weak Raman band at  $1688\text{ cm}^{-1}$  is the counterpart of this mode. The absorption triplet at  $1461, 1419$  and  $1402\text{ cm}^{-1}$  are due to the triply degenerate  $v_4$  mode of the ammonium ion.

The splitting of the  $v_3$  and  $v_4$  mode and the activation of the IR inactive modes  $v_1$  and  $v_2$  supports the lowering of site symmetry ( $C_1$ ) of the ion. All the four modes of the  $\text{NH}_4$  ion are found to be shifted from the free state values. This observation, in agreement with the structural reports, shows that the ion is distorted from the ideal tetrahedral geometry [24].




Fig. 2. FT Raman spectra of (a)  $\text{Ag}_3(\text{PO}_2\text{NH})_3$ , (b)  $\text{Na}_3(\text{PO}_2\text{NH})_3 \cdot \text{H}_2\text{O}$  (c)  $\text{Na}_3(\text{PO}_2\text{NH})_3 \cdot 4\text{H}_2\text{O}$ , (d)  $[\text{C}(\text{NH}_2)_3]_3(\text{PO}_2\text{NH})_3 \cdot \text{H}_2\text{O}$  and (e)  $(\text{NH}_4)_4(\text{PO}_2\text{NH})_3 \cdot 4\text{H}_2\text{O}$  in the region  $50\text{--}3500\text{ cm}^{-1}$ .




Fig. 3. FT Raman spectra of the aqueous solution of (a)  $\text{Na}_3(\text{PO}_3\text{NH})_3 \cdot \text{H}_2\text{O}$  (b)  $[\text{C}(\text{NH}_2)_3]_3(\text{PO}_2\text{NH})_3 \cdot \text{H}_2\text{O}$  and (c)  $(\text{NH}_4)_4(\text{PO}_2\text{NH})_4 \cdot 4\text{H}_2\text{O}$  in the region  $50\text{--}2000\text{ cm}^{-1}$ .

In a freely rotating  $\text{NH}_4$  ion, the  $v_3$  and  $v_4$  modes are triply degenerate. However if it is in the locked-in position in the crystal lattice, the degeneracy is expected to be lifted [28]. In the present study, the removal of degeneracy of the  $v_3$  and  $v_4$  modes suggests that the ion is probably in a locked in position in the crystal lattice. The fundamental  $v_6$  could not be identified independently as it merges in the region of the bending mode of  $\text{PO}_2$ . However, its combinations  $v_2 + v_6$  and  $v_4 + v_6$  are not observed. Hence it may be inferred that the  $\text{NH}_4$  ion is executing hindered rotation in the lattice [29–32].

#### 4.3. Vibrations of $[\text{C}(\text{NH}_2)_3]^+$ cation

In the structural studies of triguanidinium tri- $\mu$ -imido-cyclotriphosphate monohydrate (compound IV) Stock et al. have assigned highly intense IR band at  $1662\text{ cm}^{-1}$  to the asymmetric CN stretching vibrations. The IR spectrum of

tetraguanidinium tetra- $\mu$ -imido-cyclotetraphosphate tetrahydrate  $[\text{C}(\text{NH}_2)_3]_4(\text{PO}_2\text{NH})_4 \cdot 4\text{H}_2\text{O}$  also shows a highly intense IR band at  $1659\text{ cm}^{-1}$  due to  $v_{\text{as}}\text{CN}$  [28]. In the present investigation, the very strong absorption band at  $1663\text{ cm}^{-1}$  is assigned to the asymmetric CN stretching vibrations. The very high intensity of this band differentiate it from the  $v_2$  mode of water molecules, which are usually weak to moderate in IR. The symmetric CN stretching is very strong in Raman at  $1007\text{ cm}^{-1}$ . Its corresponding IR band is weak at  $1006\text{ cm}^{-1}$ . The high intensity of this band in the Raman spectra and the low intensity in the IR spectra, differentiate it from the  $v_{\text{as}}\text{P-NH-P}$  mode, which is usually weak in Raman and intense in IR. The Raman spectrum of the aqueous solution also shows, this intense band without much shift ( $1010\text{ cm}^{-1}$ ), which support our assignment of  $v_s\text{CN}$ . The deformation of  $\text{CN}_3$  gives a weak band at  $515\text{ cm}^{-1}$ . The  $v\text{NH}_2$  mode is masked in the broad bands of  $v\text{NH}$ . However, the bands of  $\text{NH}_2$  deformation are clearly identifiable. The weak Raman band at  $1588\text{ cm}^{-1}$  and the shoulder at  $1542\text{ cm}^{-1}$  in the IR spectrum are due to the  $\delta\text{NH}_2$  vibration.

#### 4.4. Vibrations of water molecules

The  $v_1$  and  $v_3$  modes of water molecules are observed in the region  $3472\text{--}3214\text{ cm}^{-1}$  and the bending mode  $v_2$  around  $1661\text{ cm}^{-1}$ . The existence of broad overlapping bands in the stretching region shows the presence of both weakly and strongly bonded molecules of water of crystallization and of hydrogen bridges with different bond energies. The absorption band at  $3472$  and  $1643\text{ cm}^{-1}$  in the unhydrated compound I are due to the water molecules adsorbed during pelletization. In the IR spectrum of compound III the  $v_3$  mode of  $\text{H}_2\text{O}$  appear as a doublet at  $3455$  and  $3372\text{ cm}^{-1}$  and the  $v_2$  mode as triplet at  $1694$ ,  $1661$  and  $1639\text{ cm}^{-1}$ . This splitting of the fundamental vibrations are due to the wide variation in the hydrogen bond lengths, where the  $\text{O}\cdots\text{O}$  ranges from  $2.821$  to  $4.144\text{ \AA}$  [17]. The librational modes of  $\text{H}_2\text{O}$  are observed as weak to medium intense bands in the region  $806\text{--}802\text{ cm}^{-1}$ . This assign-



ment is confirmed by the broad bands observed about this region in their solution spectra.

#### 4.5. Hydrogen bonding

Structural investigations report that, in all the five compounds investigated, two anion rings are linked to each other by N–H---O hydrogen bonds forming pairs [17,21–24]. These units are interconnected by O–H---O hydrogen bonds through water molecules.

As a result of the comparison of the spectra of tetrametaphosphimates with NH group in the ring, with the spectrum of octa silver salt  $\text{Ag}_8(\text{PO}_2\text{N})_4$ , Lunkwitz and Steger [7] were able to distinguish two bands at 3075 and 2675  $\text{cm}^{-1}$  corresponding only to NH groups and bridging hydrogen bonds NH---N(O<sup>-</sup>), respectively. The weak absorption bands in the region 2716–2575  $\text{cm}^{-1}$  are accordingly assigned to NH...N(O<sup>-</sup>) vibrations.

#### 4.6. Correlation between the P–N–P stretching vibrations and the P–N–P angle

The correlation between the P–O–P angle and the asymmetric and symmetric stretching frequencies of the P–O–P bridge are well known [33–37]. The correlation exhibits a definite pattern of relationship. Several authors have used these correlations to calculate the unknown P–O–P angles in pyrophosphates [36,37] and cyclohexaphosphates [19]. Lazarev, [37] has used similar type of correlations in pyrosilicates and determined the Si–O–Si angles.

In the present investigation, a similar type of correlation is attempted for the P–N–P group. The structural and spectral data used are listed in Table 3. The curves (Fig. 4) are plotted for the two relationships

1.  $v_{\text{asPNP}} - v_{\text{sPNP}} = f_1(\angle \text{PNP})$
2.  $(v_{\text{asPNP}} - v_{\text{sPNP}})/(v_{\text{asPNP}} + v_{\text{sPNP}}) = f_2(\angle \text{PNP})$

The curves show a definite pattern as in the case of POP bridge.

Table 3  
Correlation between the PNP bridge angle and its stretching vibrations

Compound	$v_{\text{asPNP}}$ ( $\text{cm}^{-1}$ )	$v_{\text{sPNP}}$ ( $\text{cm}^{-1}$ )	$v_{\text{as}} - v_{\text{s}}$ ( $\text{cm}^{-1}$ )	$(v_{\text{as}} - v_{\text{s}})/$ $(v_{\text{as}} + v_{\text{s}})$	$\angle \text{PNP}$ ( $^\circ$ )	Refs.
$\text{Ag}_3(\text{PO}_2\text{NH})_3$	933	813	120	0.0687	125.5	[21]; present work
$\text{Na}_3(\text{PO}_2\text{NH})_3 \cdot \text{H}_2\text{O}$	952	826	126	0.0709	124.3	[22]; present work
$\text{Na}_3(\text{PO}_2\text{NH})_3 \cdot 4\text{H}_2\text{O}$	960	860	100	0.0549	121.0	[17]; present work
	931	821	110	0.0628	122.6	
		802	129	0.0744	124.4	
$[\text{C}(\text{NH}_2)_3]_3(\text{PO}_2\text{NH})_3 \cdot \text{H}_2\text{O}$	921	853	68	0.0373	122.1	[23]; present work
$(\text{NH}_4)_4(\text{PO}_2\text{NH})_4 \cdot 4\text{H}_2\text{O}$	992	818	137	0.0773	128.61	[24]; present work
	955		174	0.0961	130.1	
$\text{Rh}(\text{PO}_2\text{NH})_3(\text{NH}_3)_3 \cdot \text{H}_2\text{O}$	935	870	65	0.03601	121	[16]
$\text{Na}_4\{\text{Cu}[(\text{PO}_2\text{NH})_3]_2\} \cdot 10\text{H}_2\text{O}$	960	820	110	0.0628	122	[15]
	930	800	160	0.0909	133	
$[\text{C}(\text{NH}_2)_3]_4(\text{PO}_2\text{NH})_4 \cdot 4\text{H}_2\text{O}$	976	820	121	0.0687	123	[23]
	941	773	156	0.0869	130	




Fig. 4. Relationship between (a) the angle PNP and  $\nu_{as\text{PNP}} - \nu_{s\text{PNP}}$  and (b) the angle PNP and  $(\nu_{as\text{PNP}} - \nu_{s\text{PNP}}) / (\nu_{as\text{PNP}} + \nu_{s\text{PNP}})$ .

## 5. Conclusions

The large splitting of  $\nu_s$  P–NH–P modes in compound **I** indicates distortion of the  $\text{PO}_2\text{NH}^-$  anion ring due to the influence of  $\text{Ag}^+$  cation.  $\text{NH}_4^+$  ion exhibits hindered rotation in the lattice. The lifting up of the degeneracy of  $\nu_3$  and  $\nu_4$  mode and the activation of the IR inactive modes indicates lower site symmetry of the NH<sub>4</sub><sup>+</sup> ion. Splitting of the  $\nu_2$  and  $\nu_3$  modes of vibration of water molecules in compound **III** indicates the wide variation in the hydrogen bond lengths. The correlation between angle P–N–P and the stretching vibrations of the P–N–P bridge exhibit a definite pattern.

## References

- [1] H.N. Stokes, Ber. Dtsch. Chem. Ges. 28 (1895) 437.
- [2] H.N. Stokes, Am. Chem. J. 18 (1896) 629.
- [3] D.E.C. Corbridge, E.J. Lowe, J. Chem. Soc. (1954) 4555.
- [4] J.V. Pustinger, W.T. Cave, M.L. Nielsen, Spectrochim. Acta 11 (1959) 909.
- [5] A.H. Herzog, M.L. Nielsen, Anal. Chem. 30 (1958) 1490.
- [6] E. Steger, K. Lunkwitz, Z. Anorg. Allgem. Chem. 313 (1961) 262.
- [7] E. Steger, K. Lunkwitz, Z. Anorg. Allgem. Chem. 313 (1961) 271.
- [8] E. Steger, K. Lunkwitz, Z. Anorg. Allgem. Chem. 316 (1962) 293.
- [9] K. Lunkwitz, E. Steger, Spectrochim. Acta 23A (1967) 2593.
- [10] K. Lunkwitz, E. Steger, Wiss. Z. Techn. Univ. Dresden 17 (1968) 1459.
- [11] K. Lunkwitz, E. Steger, Z. Anorg. Allgem. Chem. 358 (1968) 111.
- [12] E. Steger, K. Lunkwitz, J. Mol. Struct. 3 (1969) 67.
- [13] L.M. Sukova, O.I. Kondratov, K.I. Petrova, I.A. Rozanov, Sov. J. Coord. Chem. 3 (1977) 850.
- [14] L.M. Sukova, K.I. Petrova, I.A. Rozanov, D.A. Murashov, Russ. J. Inorg. Chem. 24 (1979) 1328.
- [15] L.M. Sukova, O.I. Kondratov, K.I. Petrova, I.A. Rozanov, G.G. Novitskii, Russ. J. Inorg. Chem. 24 (1979) 1354.
- [16] L.M. Sukova, O.I. Kondratov, K.I. Petrova, I.A. Rozanov, G.G. Novitskii, Russ. J. Inorg. Chem. 24 (1979) 1509.
- [17] R. Olthof, T. Mighelsen, A. Vos, Acta Cryst. 19 (1965) 596.
- [18] A. John, S. Devanarayanan, Z. Zdravkova, Indian J. Phys 73B (1999) 679.
- [19] A. John, S. Devanarayanan, M. Watanabe, Spectrochim. Acta 56A (2000) 877.
- [20] A. John, D. Philip, A.K. Idrissi, G. Keresztury, S. Devanarayanan, J Raman Spectrosc., in press.
- [21] N. Stock, W. Schnick, Z. Naturforsch 52B (1997) 251.
- [22] N. Stock, W. Schnick, Acta Cryst. C53 (1997) 532.
- [23] N. Stock, B. Jurgens, W. Schnick, Laboratorium for Anorganische Chemie Universitat, D-95440 Bayreuth, Germany, personal communication.

- [24] N. Stock, W. Schnick, *Acta Cryst.* C54 (1998) 171.
- [25] W.G. Fateley, F.R. Dollish, N.T. McDevitt, F.F. Bentley, *Infrared and Raman Selection Rules for Molecules and Lattice Vibrations: The Correlation Method*, Wiley Interscience, New York, 1972.
- [26] K. Nakamoto, *Infrared and Raman Spectra of Inorganic and Coordination Compounds*, fifth ed., Wiley, New York, 1997 Part A.
- [27] R.Ya Melnikova, V.N. Korzhuev, E.A. Prodan, I.E. Malashonok, *J. Appl. Spectrosc.* 41 (1984) 1147.
- [28] T.C. Waddington, *J. Chem. Soc.* (1958) 4340.
- [29] P.K. Acharya, P.S. Narayanan, *Indian J. Pure Appl. Phys.* 11 (1973) 574.
- [30] I.A. Oxtan, O. Knop, M. Falk, *Can. J. Chem.* 53 (1975) 3394.
- [31] G.J. Kearly, I.A. Oxtan, *Adv. Infrared Raman Spectrosc.* 10 (1983) 111.
- [32] R. Hareesh, P. Rajagopal, G. Aruldas, *Spectrochim. Acta* 48A (1992) 1453.
- [33] A. Rulmont, R. Cahay, M. Liegeois-Duyckaerts, P. Tarte, *Eur. J. Solid State Inorg. Chem.* 28 (1991) 207.
- [34] I.H. Joe, G. Aruldas, G. Kersztury, *J. Raman Spectrosc.* 22 (1991) 537.
- [35] V.S. Jayakumar, Ph.D Thesis, University of Kerala, India, 1993.
- [36] M. Harcharras, A. Ennaciri, A. Rulmont, B. Gilbert, *Spectrochim. Acta* 53A (1997) 345.
- [37] A.N. Lazarev, *Vibrational Spectra and Structure of Silicates*, English Translation, Consultant Bureau, New York, 1972.