

Follow-up of vestibular function in bilateral vestibulopathy

V C Zingler,¹ E Weintz,¹ K Jahn,¹ A Mike,² D Huppert,¹ N Rettinger,¹ T Brandt,¹ M Strupp¹

¹Department of Neurology, Ludwig-Maximilians University, Klinikum Grosshadern, Munich, Germany; ²Department of Neurology, University of Pécs, Hungary

Correspondence to:
Dr V C Zingler, Department of Neurology, Ludwig-Maximilians University, Klinikum Grosshadern, Marchioninistraße 15, D-81377 Munich, Germany; vera-carina.zingler@med.uni-muenchen.de

Received 17 April 2007
Revised 8 June 2007
Accepted 11 June 2007
Published Online First
17 July 2007

ABSTRACT

Objective: Bilateral vestibulopathy (BV) leads to a bilateral deficit of the vestibulo-ocular reflex and has various aetiologies. The main goal of this study was to determine the frequency and degree of recovery or worsening of vestibular function over time.

Methods: 82 patients (59 males, 23 females; mean age at the time of diagnosis 56.3 (SD 17.6) years) were re-examined 51 (36) months after the first examination. All patients underwent a standardised neuro-ophthalmological and neuro-otological examination. Electronystagmography with bithermal caloric irrigation was analysed by measurement of the mean peak slow phase velocity (SPV) of the induced nystagmus. Patients evaluated the course of their disease in terms of balance, gait unsteadiness and health related quality of life.

Results: Statistical analysis of the mean peak SPV of caloric induced nystagmus revealed a non-significant worsening over time (initial mean peak SPV 3.0 (3.5)°/s vs 2.1 (2.8)°/s). With respect to subgroups of aetiology, only patients with BV due to meningitis exhibited an increasing, but non-significant SPV (1.0 (1.4)°/s vs 1.9 (1.6)°/s). Vestibular outcome was independent of age, gender, time course of manifestation and severity of BV. Single analysis of all patients showed that a substantial improvement $\geq 5^\circ/\text{s}$ occurred in two patients on both sides (idiopathic $n = 1$, Sjögren's syndrome $n = 1$) and in eight patients on one side (idiopathic $n = 6$, meningitis $n = 1$, Menière's disease $n = 1$). In 84% of patients there was impairment of their health related quality of life (42% slight, 24% moderate, 18% severe). Forty-three per cent of patients rated the course of their disease as stable, 28% as worsened and 29% as improved.

Conclusions: Our data support the view that more than 80% of patients with BV do not improve. Thus the prognosis of BV is less favourable than assumed.

Bilateral vestibulopathy (BV) is characterised by impaired or lost function of both peripheral labyrinths or of the eighth nerves. The most frequent complaint of patients with BV is unsteadiness of gait with worsening in darkness and on uneven ground due to deficient vestibulospinal function. Some patients complain of oscillopsia (ie, apparent motion of the visual scene during head movements or when walking). Oscillopsia is caused by involuntary retinal slip as a result of an insufficient vestibulo-ocular reflex (VOR). Furthermore, BV can also lead to impaired spatial memory and navigation associated with hippocampal atrophy.¹ The diagnosis of BV is based on the head thrust test,^{2,3} a simple bedside test for high frequency VOR function, and bithermal caloric testing with oculographic recordings to test

for a low frequency VOR deficit. BV is a rare vestibular disorder of various aetiologies (eg, ototoxic drugs, meningitis or Menière's disease).⁴⁻⁶ Despite intensive examinations, the aetiology remains unclear in approximately 50%.⁶

Little is known about the course of the disease, especially the recovery. A few small case series have reported contradictory recovery rates.^{5,7-9} In a report on 11 patients with idiopathic BV (eight patients with simultaneous and three patients with sequential onset of BV), partial recovery was detected in four patients with simultaneous and in all three patients with sequential BV after a follow-up of 1-7 years. Complete recovery did not occur in any of the patients.⁵ Improvement of vestibular function was also described in single cases of different aetiologies,⁹ in particular when BV was caused by serous rather than suppurative destructive labyrinthitis.^{4,10,11} In contrast, in a 5 year follow-up study⁷ of seven patients with BV, Baloh *et al* did not find significant improvement in vestibular excitability. Furthermore, in a recent study, the effectiveness of immunosuppressive treatment of patients with BV and inner ear antibodies was moderate in four of 12 patients and only transient in two of them.⁸ Based on our clinical experience, the prognosis of BV is less favourable than hitherto reported. Therefore, we examined 82 patients with BV of various aetiologies and clinically re-evaluated them after a mean follow-up time of 51 (36) months (range 3 months to 13 years). The initial diagnosis of BV and clinical re-evaluation were made according to a standardised protocol, including a neuro-ophthalmological and neuro-otological workup. We not only determined the frequency and extent of recovery or worsening of vestibular function and their dependence on the aetiology of BV, but also the impairment of quality of life.

PATIENTS AND METHODS

The diagnosis of BV was based on the following criteria: (1) a pathological head thrust test,^{2,3} which indicates a high frequency deficit of the VOR and/or (2) bilaterally diminished or absent caloric responses on an electronystagmogram (ENG), indicating a low frequency deficit.^{4,12}

Complete BV was characterised by pathological head thrust tests and the absence of caloric responses on both sides.

Incomplete BV was characterized by one of the following responses:

- ▶ bilaterally pathological head thrust tests and a calorically elicited nystagmus with a mean

peak slow phase velocity (SPV) of $<5^{\circ}/s$ for each caloric irrigation on both sides;

- ▶ bilateral pathological head thrust tests and caloric responses $>5^{\circ}/s$ on one or both sides; or
- ▶ normal head thrust tests and loss of bilateral responses or reduced responses ($<5^{\circ}/s$) to caloric irrigation on both sides.

Definition of aetiology

Patients were classified as having either a “definite” or “probable” cause of BV. This classification was based on the previously published literature on BV.^{4,5} The aetiologies were carefully reassessed by the senior authors.

A total of 163 patients who had been diagnosed with BV in the Department of Neurology, University of Munich, between 1993 and 2005 were contacted. Ninety-five patients responded and 82 finally came to the follow-up examinations. The current address could not be determined for 37 patients, 28 patients did not respond and three had died in the meantime.

A standardised detailed history was obtained from all patients at the first and follow-up examinations. The following parameters were assessed: (1) recurrent (preceding or current) attacks of vertigo or dizziness: onset, duration, time course, frequency and associated symptoms (oscillopsia, double vision, blurred vision, loss of sight, phonophobia, photophobia, headache, nausea, vomiting); (2) unsteadiness of gait, in particular in the dark or on uneven ground; (3) oscillopsia during head movements or during locomotion; (4) recurrent or sustained ear symptoms (tinnitus, hearing loss, fullness of the ear); (5) past medical history with regard to migraine, polyneuropathy, exposure to toxins, antibiotics or diuretics, cardiovascular risk factors (eg, arterial hypertension, diabetes mellitus) or other neurological, ophthalmological, ENT or internal diseases (eg, autoimmune disorders, cancer); (6) family history of impaired audiovestibular function.

All patients underwent a complete neurological, neuro-ophthalmological and neuro-otological examination. The neuro-ophthalmological and neuro-otological examination included the head thrust test, evaluation for spontaneous nystagmus with Frenzel’s goggles, gaze evoked nystagmus, smooth pursuit, saccades, optokinetic nystagmus, visual fixation suppression of the VOR, rebound nystagmus and head

shaking nystagmus (for general neuro-ophthalmological and neuro-otological testing, see Brandt and Strupp¹³). A cerebellar ocular motor disorder was characterised by either one or a combination of the following ocular motor dysfunctions: rebound nystagmus, downbeat nystagmus, pathological saccades and pathological visual fixation suppression of the VOR.

ENG with bithermal caloric testing (30°C and 44°C) was performed in all patients to determine mean peak SPV using Igor Pro Wave Metric software (V.3.13). We determined the mean peak SPV of the 10 fastest slow phases for each side and temperature. For technical reasons, three ENGs performed at follow-up could not be evaluated; these patients were therefore excluded. None of the patients was on medications with a relevant influence on vestibular testing, such as benzodiazepines, analgesics or anticonvulsants.

A cranial MRI was performed in 78 patients. Specific laboratory tests were initiated according to the suspected specific aetiologies (autoantibodies against inner ear structures, vitamin B12 and folate, antinuclear antibodies, complement factors C3 and C4 and CSF test). Autoantibodies against inner ear structures were determined as previously reported by our group.¹⁴ All patients performed vestibular exercises and balance training using a video or DVD for at least 3 months.¹⁵

Subjective parameters

The patients were asked: (1) to evaluate the impairment of their health related quality of life measured at the time of the second evaluation with a self-classification system (with respect to mobility, self-care and daily activities) consisting of four degrees of severity in comparison with their situation before manifestation of BV (not impaired and slightly, moderately or severely impaired); (2) to evaluate the course of their disease by comparing the change in balance and gait unsteadiness (unchanged, worse, improved); (3) if they had noticed changes or difficulties in spatial navigation.

Statistical analysis

The mean peak SPV of each patient was calculated as follows: peak SPV ($^{\circ}/s$) after caloric testing at 30°C and 44°C for the right and left sides were added up, and the sum was divided by 4. For statistical analysis, a main effects ANOVA (Statistica 6.0,

Table 1 Number of patients (%) in subgroups with different aetiologies and types of manifestation of bilateral vestibulopathy* and number of patients with bilateral pathological head thrust tests and electronystagmogram at the initial and follow-up examinations

	Patients (n (%))	ENG (mean (SD) peak SPV)		Bilateral pathological head thrust tests	
		Initial	Follow-up	Initial (n)	Follow-up (n)
Aetiology of BV	79 (100)	3.0 (3.5)	2.1 (2.8)	72	72
Antibiotics	10 (13)	2.3 (2.2)	0.8 (1.1)	8	9
Meningitis	7 (9)	1.0 (1.4)	1.9 (1.6)	7	6
Pathological inner ear antibodies	7 (9)	1.8 (2.0)	1.6 (2.0)	7	7
Menière’s disease	6 (8)	2.5 (3.0)	1.9 (3.2)	6	5
Miscellaneous‡	10 (13)	6.6 (4.8)‡	4.4 (5.6)	9	9
Idiopathic‡	39 (49)	4.9 (6.1)‡	3.9 (5.4)	35	36
Sequential BV	30 (38)	3.0 (3.5)	2.1 (2.8)	28	30
Slowly progressive BV	49 (62)	4.6 (5.8)	3.6 (5.4)	47	45

*Some of the percentages have been rounded up, others rounded down.

†The mean peak SPV of each patient was calculated as follows: all peak SPVs after caloric testing at 30°C and 44°C for the right and left sides were added, and the sum was divided by 4.

‡Only those patients were included who showed either a mean peak SPV $\leq 5^{\circ}/s$ and/or had a bilateral pathological head thrust test.

The total number of all patients was 79.

BV, bilateral vestibulopathy; ENG, electronystagmogram; SPV, slow phase velocity.

Research paper

Statsoft) was performed with mean peak slow phase velocities of caloric responses at the two time points as dependent variables and the following categorical predictors: "aetiology" (see table 1, six levels), "course" (progressive vs sequential, two levels), "gender" (two levels), "age" (patients grouped by decades, eight levels) and "completeness" (complete vs incomplete BV, two levels). Significance was defined as an alpha level of 0.05. Data in the figure are given as mean (95% confidence intervals).

RESULTS

Study population

Eighty-two patients with a mean age of 56.3 (SD 17.6) years at the time of diagnosis were included. The diagnosis was made, on average, 71 (119) months (median 36) after the onset of the first symptoms. The majority were male (72%, n = 59, mean age 55.8 (15.2) years; females 28%, n = 23, mean age 62.9 (15.0) years). Thirty-two per cent of all patients fulfilled the




Figure 1 (A, B) Response to caloric irrigation in the initial and follow-up electronystagmogram (mean (SD) peak slow phase velocity (SPV)) in patients with sequential and slowly progressive bilateral vestibulopathy (BV) (A) and for subgroups with BV of various aetiologies (B). Filled column, first examination; open column, follow-up examination. Misc, miscellaneous; antibodies, pathological inner ear antibodies. Symbols in the columns (square, diamond) = median; horizontal lines of the boxes = 25th and 75th percentiles; whiskers above and below the boxes = minimum and maximum values.

above listed criteria for complete BV (n = 26) and the remaining (n = 56) for incomplete BV. In the latter, a subgroup (n = 5) with bilaterally normal head thrust tests, but pathological caloric responses, was identified.

The ENG of three patients could not be evaluated for technical reasons and thus the following statistical analyses refer to the remaining 79 patients.

Time course of the disease manifestation

Episodes of vertigo or dizziness preceded the manifestation of BV in 38% of patients (n = 30) who had a so-called sequential course. BV developed simultaneously, most often slowly progressively in 62% of all patients (n = 49), as indicated by a patient history of progressive unsteadiness of gait and oscillopsia.

Aetiology of BV

The definite or probable cause was defined in 51% of all patients (n = 40). The most frequent causes were adverse effects of aminoglycosides (13% of total, n = 10): gentamycin only in six (of 10) patients, gentamycin plus other antibiotics (streptomycin, vancomycin, penicillin) in three (of 10) patients and gentamycin plus furosemide in one (of 10) patients. Nine per cent of patients (n = 7) had suffered from meningoencephalitis and another 9% (n = 7) had autoantibodies against inner ear structures. Another group (8% of the total, n = 6) consisted of patients with Menière's disease. There was also a group (13% of the total, n = 10) with "miscellaneous" causes: side effects caused by high dose therapy with furosemide (n = 1), otosclerosis (n = 1), vitamin B12 deficiency (n = 2), alcoholism (n = 2), positive family history for inner ear disease (n = 1), an association of BV with spinocerebellar ataxia type 6 (n = 1) or systemic autoimmune diseases (Sjögren's syndrome and ulcerative colitis) (n = 2). The underlying aetiology of BV remained unclear in 49% (so-called idiopathic BV; n = 39).

BV and its association with cerebellar symptoms

Ten of all included patients with BV (nine males, one female) were also diagnosed as having a cerebellar syndrome at the first examination. The following cerebellar signs were found: cerebellar ocular motor disorders in all 10 patients, ataxia of the limbs in nine patients and dysarthrophonia in four patients. None of the other included patients with BV developed a cerebellar syndrome in the period between the first and the second examinations. BV was caused by spinocerebellar ataxia

Table 2 Subjective evaluation of the course of the disease, health related quality of life and spatial navigation at the time of follow-up

Subjective evaluation	Patients (n (%))
Quality of life	
Not impaired	13 (16)
Slightly impaired	33 (42)
Moderately impaired	19 (24)
Severely impaired	14 (18)
Spatial orientation	
Not impaired	69 (87)
Impaired	10 (13)
Disease course	
Unchanged	34 (43)
Worse	22 (28)
Improved	23 (29)

Table 3 Subjective evaluation of the course of the disease

Aetiology	Worse (n (%))	Unchanged (n (%))	Improved (n (%))
Idiopathic	14 (36)	17 (44)	8 (21)
Antibiotics	3 (30)	4 (40)	3 (30)
Pathological inner ear antibodies	1 (14)	4 (57)	2 (29)
Meningitis	3 (43)	3 (43)	1 (14)
Menière's disease	1 (17)	1 (17)	4 (67)
Miscellaneous	1 (10)	5 (50)	4 (40)

The results are given for different subgroups (some of the percentages have been rounded up, others rounded down).

(n = 1) and alcoholism (n = 2). The other patients had idiopathic BV.

Cranial imaging

MRI studies were performed in 78 of all included 82 patients. The following abnormalities were found: cerebellar atrophy (n = 6), other cerebellar lesions (eg, cerebellar infarction, n = 2), unilateral acoustic neuroma (n = 1), generalised cerebral atrophy (n = 4) and supra- or infratentorial vascular lesions (n = 10).

Follow-up with re-evaluation of vestibular function

The time between the initial diagnosis and follow-up examination was 51 (36) months (range 3 months to 13 years; median 41 months). The initial ENG showed no significant differences in response to caloric irrigation with respect to the aetiology subgroups or time course of manifestation. Table 1 shows the mean peak SPV after caloric irrigation and the number of patients with bilateral pathological head thrust tests at the first and follow-up examinations. The results are also given for different aetiologies and time courses of manifestation. In general, the mean peak SPV after caloric irrigation tended to decrease at follow-up, but not significantly.

This was true for the total study population as well as for patients with different time courses of manifestation of BV (fig 1A) and aetiologies (fig 1B). Only patients with BV due to meningitis had slightly increased mean peak SPVs at the follow-up investigation (n = 7). Single analysis of all patients showed that a substantial improvement in caloric responses $\geq 5^\circ/s$ occurred in two patients on both sides (idiopathic n = 1, Sjögren's syndrome n = 1) and in eight patients on one side (idiopathic n = 6, meningitis n = 1, Menière's disease n = 1). The patient with Sjögren's syndrome had received immunosuppressive treatment with cortisone and azathioprine. The statistical analysis of vestibular outcome with regard to age, gender and complete versus incomplete BV also revealed no significant differences. Oscillopsia during head and body movements were reported by 56% of all patients (n = 44) at the first examination and by 53% (n = 42) at follow-up. Gait unsteadiness was reported by all patients at both examinations.

Subjective evaluation of the course of the disease, quality of life and spatial navigation

Table 2 gives an overview of how patients subjectively evaluated the course of their disease, their health related quality of life and their spatial navigation. Most of the patients (43%) considered the course of their disease stable. The rest estimated their clinical condition as either worsened (29%) or improved (28%).

As regards the different aetiology subgroups (table 3), only four of six patients with Menière's disease felt that they had improved at follow-up. These patients had been treated with

high doses of betahistine dihydrochloride (48 mg three times daily). At follow-up, attacks of vertigo had completely stopped in three of these patients, and the duration and frequency of attacks were markedly decreased in one patient. The majority of patients (42%) stated that BV caused a slight impairment of their health related quality of life (24% moderate and 18% severe impairment). Only a minority of patients (16%) felt that BV had not impaired their quality of life. Thirteen per cent of patients had noticed an impaired spatial navigation.

DISCUSSION

The major finding of this follow-up study in patients with BV was that, in general, vestibular function remained unchanged with a non-significant tendency of slow worsening. The vestibular outcome was independent of the underlying causative factors, type of manifestation, severity of vestibular loss, age and gender. This supports the findings of a longitudinal study⁷ in a small group of seven patients with BV of different aetiologies and 57 healthy subjects who returned for yearly examinations of vestibular function over a period of 5 years. In this study, no improvement was found in BV patients; the normal older subjects (all over the age of 75 years) exhibited an age related decrement in their vestibular responses without any associated symptoms or signs of disequilibrium. It is known from various studies that all structures relevant for vestibular function (ie, the vestibular end organs and central vestibular pathways) degenerate continuously with age. Hair cell loss precedes those seen in the vestibular nerve and Scarpa's ganglion.¹⁶⁻¹⁹ Thus it can be assumed that the mild decrement in vestibular responses in our study can be attributed to progression of the disease as well as vestibular aging. Because of the study design of our and previous investigations,^{5, 7-9} most of the patients presented with a chronic condition. This implies that those patients with acute onset of BV and early recovery (such as patients in the intensive care unit with BV caused by meningitis or treatment with ototoxic antibiotics) might remain undetected by this approach. The latter argument is supported by the improvement of patients with BV due to meningitis in our study. This is supported by previous case reports. It has been stated that patients with serous rather than suppurative meningitis can exhibit partial recovery.^{10, 11}

In our study, none of the patients with BV caused by ototoxic antibiotics showed a substantial improvement. However, regeneration of receptor cells and subsequent functional recovery after (gentamycin induced) damage of the auditory and vestibular systems in many vertebrates is well known.^{20, 21}

The subjective evaluation of the course of the disease was similar to the results of our objectively achieved findings for vestibular testing. Most of the patients estimated their clinical condition as being either unchanged or worse. In particular, only a minority of patients reported an improvement in gait unsteadiness. Part of this improvement might be due to vestibular exercises and balance training, which all of our patients had for at least 3 months. The usefulness of vestibular rehabilitation in unilateral vestibular failure¹⁵ and BV^{22, 23} has been reported previously. The improvement after vestibular rehabilitation with clear locomotor gains has been described in approximately 50% of BV patients who received intensive training over 3-12 months.²² Only patients with Menière's disease frequently assessed their course of disease as being favourable. This might be explained by spontaneous or drug induced reduction or cessation of vertigo attacks in the later stage of the disease.

Research paper

About two-thirds of our patients rated their health related quality of life as slightly to moderately impaired. Nearly 90% of patients estimated their spatial navigation as unchanged since the first manifestation of BV. A previous study had shown that complete BV can lead to impaired spatial memory and navigation associated with hippocampal atrophy.¹

This study has some limitations as to completeness of vestibular testing with respect to otolith (vestibular evoked myogenic potentials) and canal function, dynamic visual acuity, evaluation of hearing impairment, posturography and gait analysis, or the application of a health self-classification system such as the EQ-5D²⁴ or specific dizziness rating scales. None of these tests was included because they were not routinely performed at the first examination. We chose the head thrust test and bithermal caloric testing with 30°C and 44°C (instead of rotational testing) because it is the gold standard for documenting BV¹³ and allows side specific assessment of (residual) vestibular function. One may argue that in patients who already have minimal to no caloric responses to start with, it is not surprising that no statistical difference was seen at follow-up. Caloric testing with ice water would have been an additional tool to detect slight improvement. However, the main goal of this study was to investigate a relevant improvement or worsening of vestibular function over time. In addition, according to our experience with many patients, caloric testing with ice water is not well tolerated and was therefore not routinely used in our patients.

In conclusion, our data support the view that more than 80% of patients with BV do not improve. Thus the prognosis of BV is less favourable than assumed. This emphasises the need for intensive vestibular rehabilitation²⁵ to reduce the burden of disease.

Acknowledgements: We thank Ms Judy Benson for copyediting the manuscript.

Competing interests: None.

Ethics approval: Ethics approval was obtained.

REFERENCES

1. Brandt T, Schautzer F, Hamilton DA, *et al.* Vestibular loss causes hippocampal atrophy and impaired spatial memory in humans. *Brain* 2005;**128**:2732–41.
2. Halmagyi GM, Curthoys IS. A clinical sign of canal paresis. *Arch Neurol* 1988;**45**:737–9.
3. Jorns-Häderli M, Straumann D, Palla A. Accuracy of the bedside head impulse test in detecting vestibular hypofunction. *J Neural Neurosurg Psychiatry* 2007;**78**:1113–18.
4. Rinne T, Bronstein AM, Rudge P, *et al.* Bilateral loss of vestibular function: clinical findings in 53 patients. *J Neurol* 1998;**245**:314–21.
5. Vibert D, Liard P, Häusler R. Bilateral idiopathic loss of peripheral vestibular function with normal hearing. *Acta Otolaryngol* 1995;**115**:611–15.
6. Zingler VC, Cnyrim C, Jahn K, *et al.* Causative factors and epidemiology of bilateral vestibulopathy in 255 patients. *Ann Neurol* 2007;**61**:524–32.
7. Baloh RW, Enrietto J, Jacobson KM, *et al.* Age-related changes in vestibular function: a longitudinal study. *Ann N Y Acad Sci* 2001;**942**:210–19.
8. Deutschlander A, Glaser M, Strupp M, *et al.* Immunosuppressive treatment in bilateral vestibulopathy with inner ear antibodies. *Acta Otolaryngol* 2005;**125**:848–51.
9. Frese KA, Reker U, Maune S. Der beidseitige Vestibularisausfall. *HNO* 2003;**51**:221–5.
10. Fortnum HM. Hearing impairment after bacterial meningitis. *Arch Dis Child* 1982;**67**:1128–33.
11. Bronstein AM, Morland AB, Ruddock KH, *et al.* Recovery from bilateral vestibular failure: implications for visual and cervico-ocular function. *Acta Otolaryngol Suppl* 1995;**520**:405–7.
12. Baloh RW, Honrubia V, Yee RD, *et al.* Changes in the human vestibulo-ocular reflex after loss of peripheral sensitivity. *Ann Neurol* 1984;**16**:222–8.
13. Brandt T, Strupp M. General vestibular testing. *Clin Neurophysiol* 2005;**116**:406–26.
14. Arbusow V, Strupp M, Dieterich M, *et al.* Serum antibodies against membranous labyrinth in patients with “idiopathic” bilateral vestibulopathy. *J Neurol* 1998;**245**:132–6.
15. Strupp M, Arbusow V, Maag KP, *et al.* Vestibular exercises improve central vestibulospinal compensation after vestibular neuritis. *Neurology* 1998;**51**:838–44.
16. Bergstrom B. Morphology of the vestibular nerve. II The number of myelinated vestibular nerve fibers in man at various stages. *Acta Otolaryngol* 1973;**76**:173–9.
17. Rosenhall U. Degenerative patterns in the ageing human vestibular neuroepithelia. *Acta Otolaryngol* 1973;**76**:208–28.
18. Lopez I, Honrubia V, Baloh RW. Ageing and the human vestibular nucleus. *J Vestib Res* 1997;**7**:77–85.
19. Jahn K, Naeßl A, Schneider E, *et al.* Inverse U-shaped curve for age dependency of torsional eye movement responses to galvanic vestibular stimulation. *Brain* 2003;**126**:1579–89.
20. Zakir M, Dickmann JD. Regeneration of vestibular otolith afferents after ototoxic damage. *J Neurosci* 2006;**26**:2881–93.
21. Taura A, Kojima K, Ito J, *et al.* Recovery of hair cell function after damage induced by gentamicin in organ culture of rat vestibular maculae. *Brain Res* 2006;**1098**:33–48.
22. Krebs DE, Gill-Body KM, Parker SW, *et al.* Vestibular rehabilitation: usefull but not universally so. *Otolaryngol Head Neck Surg* 2003;**128**:240–50.
23. Gillespie MB, Minor LB. Prognosis in bilateral vestibular hypofunction. *Laryngoscope* 1999;**109**:35–41.
24. EuroQuol Group. Euroquol: a new facility for the measurement of health-related quality of life. *Health Policy* 1990;**16**:199–208.
25. Herdmann SJ. Role of vestibular adaptation in vestibular rehabilitation. *Otolaryngol Head Neck Surg* 1998;**119**:49–54.



Follow-up of vestibular function in bilateral vestibulopathy

V C Zingler, E Weintz, K Jahn, et al.

J Neurol Neurosurg Psychiatry 2008 79: 284-288 originally published online July 17, 2007

doi: 10.1136/jnp.2007.122952

Updated information and services can be found at:

<http://jnp.bmj.com/content/79/3/284.full.html>

These include:

References

This article cites 25 articles, 6 of which can be accessed free at:
<http://jnp.bmj.com/content/79/3/284.full.html#ref-list-1>

Article cited in:

<http://jnp.bmj.com/content/79/3/284.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Infection \(neurology\)](#) (392 articles)
[Ophthalmology](#) (669 articles)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>