

The Hepatitis B Virus Ribonuclease H Is Sensitive to Inhibitors of the Human Immunodeficiency Virus Ribonuclease H and Integrase Enzymes

John E. Tavis^{1,2*}, Xiaohong Cheng^{1,3}, Yuan Hu^{1,3,9}, Michael Totten¹, Feng Cao^{1,2}, Eleftherios Michailidis⁴, Rajeev Aurora¹, Marvin J. Meyers⁵, E. Jon Jacobsen⁵, Michael A. Parniak⁴, Stefan G. Sarafianos⁶

1 Department of Molecular Microbiology and Immunology, Saint Louis University School of Medicine, Saint Louis, Missouri, United States of America, **2** Saint Louis University Liver Center, Saint Louis University School of Medicine, Saint Louis, Missouri, United States of America, **3** Key Laboratory of Molecular Infectious Diseases, Ministry of Education, Chongqing Medical University, Chongqing, People's Republic of China, **4** Department of Infectious Diseases and Microbiology, University of Pittsburgh School of Public Health, Pittsburgh, Pennsylvania, United States of America, **5** Center for World Health and Medicine, Saint Louis University School of Medicine, Saint Louis, Missouri, United States of America, **6** Department of Molecular Microbiology and Immunology and Department of Biochemistry, University of Missouri School of Medicine, Christopher S. Bond Life Sciences Center, Columbia, Missouri, United States of America

Abstract

Nucleos(t)ide analog therapy blocks DNA synthesis by the hepatitis B virus (HBV) reverse transcriptase and can control the infection, but treatment is life-long and has high costs and unpredictable long-term side effects. The profound suppression of HBV by the nucleos(t)ide analogs and their ability to cure some patients indicates that they can push HBV to the brink of extinction. Consequently, more patients could be cured by suppressing HBV replication further using a new drug in combination with the nucleos(t)ide analogs. The HBV ribonuclease H (RNaseH) is a logical drug target because it is the second of only two viral enzymes that are essential for viral replication, but it has not been exploited, primarily because it is very difficult to produce active enzyme. To address this difficulty, we expressed HBV genotype D and H RNaseHs in *E. coli* and enriched the enzymes by nickel-affinity chromatography. HBV RNaseH activity in the enriched lysates was characterized in preparation for drug screening. Twenty-one candidate HBV RNaseH inhibitors were identified using chemical structure-activity analyses based on inhibitors of the HIV RNaseH and integrase. Twelve anti-RNaseH and anti-integrase compounds inhibited the HBV RNaseH at 10 μ M, the best compounds had low micromolar IC₅₀ values against the RNaseH, and one compound inhibited HBV replication in tissue culture at 10 μ M. Recombinant HBV genotype D RNaseH was more sensitive to inhibition than genotype H. This study demonstrates that recombinant HBV RNaseH suitable for low-throughput antiviral drug screening has been produced. The high percentage of compounds developed against the HIV RNaseH and integrase that were active against the HBV RNaseH indicates that the extensive drug design efforts against these HIV enzymes can guide anti-HBV RNaseH drug discovery. Finally, differential inhibition of HBV genotype D and H RNaseHs indicates that viral genetic variability will be a factor during drug development.

Citation: Tavis JE, Cheng X, Hu Y, Totten M, Cao F, et al. (2013) The Hepatitis B Virus Ribonuclease H Is Sensitive to Inhibitors of the Human Immunodeficiency Virus Ribonuclease H and Integrase Enzymes. *PLoS Pathog* 9(1): e1003125. doi:10.1371/journal.ppat.1003125

Editor: Aleem Siddiqui, University of California, San Diego, United States of America

Received: September 11, 2012; **Accepted:** November 27, 2012; **Published:** January 22, 2013

Copyright: © 2013 Tavis et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by a Saint Louis University School of Medicine President's Research Fund award, a Saint Louis University Molecular Microbiology and Immunology seed grant to J.E.T. H.Y.'s efforts were supported by a Chongqing Medical University International Scholar Award and National Sciences Foundation of China Award NSFC 81101310. S.G.S. acknowledges the Ministry of Knowledge and Economy, Bilateral International Collaborative Research and Development Program, Republic of Korea, and National Institutes of Health Grants MPI-R01AI100890, R01AI099284, R01AI076119, and R33AI079801. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. Naphthyridinone is covered by US patent application 20100056516.

Competing Interests: I have read the journal's policy and have the following conflicts: JET will be an inventor on a pending patent application covering this RNaseH assay and the compounds identified with it. SGS and MAP will be inventors on patent application(s) covering compounds #12-15. This does not alter our adherence to all PLoS Pathogens policies on sharing data and materials.

* E-mail: tavisje@slu.edu

 These authors contributed equally to this work.

Introduction

Hepatitis B virus (HBV) is a hepatotropic DNA virus that replicates by reverse transcription [1]. It chronically infects >350 million people world-wide and kills up to 1.2 million patients annually by inducing liver failure and liver cancer [2–4]. Reverse transcription is catalyzed by a virally-encoded polymerase that has two enzymatic activities: a DNA polymerase that synthesizes new DNA and a ribonuclease H (RNaseH) that destroys the viral RNA after it has been copied into DNA [1,5]. Both activities are essential for viral replication.

HBV infections are treated with interferon α or one of five nucleos(t)ide analogs [6–8]. Interferon α leads to sustained clinical improvement in 20–30% of patients, but the infection is very rarely cleared [1,3,9]. The nucleos(t)ide analogs are used more frequently than interferon. They inhibit DNA synthesis and suppress viral replication by 4–5 log₁₀ in up to 70–90% patients, often to below the standard clinical detection limit of 300–400 copies/ml [10–12]. However, treatment eradicates the infection as measured by loss of the viral surface antigen (HBsAg) from the serum in only 3–6% of patients even after years of therapy [10–

Author Summary

Current therapy for HBV blocks DNA synthesis by the viral reverse transcriptase and can control the infection indefinitely, but treatment rarely cures patients. More patients could be cured by suppressing HBV replication further using a new drug in combination with the existing ones. The HBV RNaseH is a logical drug target because it is the second of only two viral enzymes that are essential for viral replication, but it has not been exploited, primarily because it is very difficult to produce active enzyme. We expressed active recombinant HBV RNaseHs and demonstrated that it was suitable for antiviral drug screening. Twenty-one candidate HBV RNaseH inhibitors were identified based on antagonists of the HIV RNaseH and integrase enzymes. Twelve of these compounds inhibited the HBV RNaseH in enzymatic assays, and one inhibited HBV replication in cell-based assays. The high percentage of compounds developed against the HIV RNaseH and integrase that were also active against the HBV RNaseH indicates that the extensive drug design efforts against these HIV enzymes can be used to guide anti-HBV RNaseH drug discovery.




Figure 1. The HBV replication cycle. HBV replicates by reverse transcription in the cytoplasm of infected hepatocytes. After completion of reverse transcription, intracellular capsids can either be transported into the nucleus to maintain the cccDNA pool (*Recycling*), or they can be enveloped and secreted from the cells as mature virions (*Secretion*). Inhibiting RNaseH activity blocks plus-strand DNA synthesis during reverse transcription; this would prevent both recycling and secretion of virions. The hepatocyte is represented as a rectangle, the nucleus as an oval, HBV capsids as the hexagon, and the viral lipid envelop as a circle surrounding the extracellular capsids. HBV proteins are green or black, RNAs are red, and DNAs are blue.
doi:10.1371/journal.ppat.1003125.g001

13]. Antiviral resistance was a major problem with the earlier nucleos(t)ide analogs, but resistance to the newer drugs entecavir and tenofovir is very low [6,14,15]. This has converted hepatitis B from a steadily worsening disease into a controllable condition for most individuals [16]. The cost of this control is indefinite administration of the drugs (probably life-long; [7]), with ongoing expenses of \$400–600/month [17,18] and unpredictable adverse effects associated with decades-long exposure to the drugs.

The key form of the HBV genome in cells that must be eliminated to clear the infection is the nuclear episomal covalently-closed circular DNA (cccDNA) that is the template for transcription of all HBV RNAs [19]. Following reverse transcription in the cytoplasm, newly synthesized genomes can either be enveloped and secreted from the cell as virions, or they can be transported into the nucleus to replenish the cccDNA pool (Fig. 1) [19,20]. Transfer of newly synthesized viral genomes into the nucleus via “recycling” is the default pathway, and virion secretion occurs only if the cccDNA pool is large enough to support adequate synthesis of the HBsAgs.

The cccDNA pool is very stable, but nucleos(t)ide therapy can suppress cccDNA levels in the liver by $\sim 1 \log_{10}$ after 1–2 years [21–23]. The indefinite persistence of the cccDNA even in patients whose HBV titres in serum have been suppressed below the limit of clinical detection by the nucleos(t)ide analogs is due to residual viral replication, leading to replenishment of the cccDNA pool by a combination of intracellular recycling and low-level infection of new cells [24,25]. The sequential accumulation of resistance mutations during nucleos(t)ide therapy confirms that cccDNA maintenance by residual viral replication occurs in the absence of clinically detectable viremia [15,26,27]. A recent genetic analysis of HBV DNA in the liver explicitly demonstrated that low levels of cccDNA replenishment occurs even when nucleos(t)ide analog therapy has reduced viral titres below the clinical detection limit [24].

RNaseH enzymes hydrolyze RNA in an RNA:DNA heteroduplex [28]. They belong to the nucleotidyl transferase superfamily whose members share a similar protein fold and presumably have similar enzymatic mechanisms [29]. This family includes *E. coli* RNaseH I and II [30], DNA transposases including the Tn5 transposase [31], retroviral integrases including the HIV integrase

[32], the RuvC Holliday junction resolvase [33], the Argonaute RNase [34], and human RNaseH 1 and 2 [35,36]. The canonical RNaseH structure contains about 100 aa including four conserved carboxylates (the “DEDD” motif) that coordinate two divalent cations [37]. The RNaseH mechanism is believed to involve both divalent cations [29,38,39], although a one-ion mechanism has also been proposed [40,41]. The HBV RNaseH domain shares low but recognizable ($\sim 20\%$) sequence identity with the RNaseH domains of reverse transcriptases and other retro-elements [42–44]. Manually optimizing alignment of the HBV RNaseH and the HIV-1 RNaseH yielded 23% identity and 33% similarity (Fig. 2). A similar alignment between the HBV RNaseH and the HIV integrase revealed 19% identity and 33% similarity.

The HBV RNaseH is encoded at the carboxy-terminus of the viral polymerase protein that also encodes the viral DNA polymerase activity (reverse transcriptase). The high hydrophobicity of the HBV polymerase and its existence as a complex with host chaperones [45] have severely restricted study of the HBV RNaseH. Furthermore, we demonstrated that the RNaseH in its native context within the polymerase protein is unable to accept exogenous heteroduplex substrates [46], analogous to the inability of the DNA polymerase active site to engage exogenous primer-templates [47]. Consequently, most of our limited knowledge of the RNaseH comes from mutational studies of the viral genome in the context of viral replication conducted by us and others [48–53]. These restrictions have prevented biochemical characterization of the RNaseH and blocked biochemical screens for anti-HBV RNaseH drugs to date.

A few reports of recombinant forms of the hepadnaviral RNaseH exist. Wei and co-workers [54] expressed the HBV RNaseH domain in *E. coli* and purified it by denaturing nickel-affinity chromatography. Following refolding, they found an RNase activity. Lee et al. [55] expressed the HBV RNaseH domain in *E. coli* as a dual maltose-binding protein/hexahistidine fusion and purified soluble protein by two-step affinity chromatography; this enzyme had RNaseH activity. Choi and co-workers [56] expressed the intact duck hepatitis B virus polymerase in yeast and reported that it had a weak RNase activity. Finally, Potenza

intracellular viral capsids were purified, and then HBV DNAs within the particles were detected by Southern analysis. All four mutants supported DNA synthesis and hence could be analyzed by this approach. The signature of an RNaseH-deficient enzyme is production of RNA:DNA heteroduplexes that migrate like double-stranded DNAs on native gels but as faster-migrating single-stranded DNAs of multiple lengths following digestion of the capsid-derived nucleic acids with exogenous RNaseH. DNAs produced by the wild-type genome were unaffected by treatment with RNaseH prior to electrophoresis (Fig. 3). Mutating each of the four predicted RNaseH DEDD residues blocked production of the slowest-migrating double stranded forms (mature relaxed-circular DNAs) and led to accumulation of smaller forms that migrated similar to the less-mature relaxed-circular DNAs produced by the wild-type genome. Treatment of the nucleic acids from the mutant genomes with exogenous RNaseH collapsed the double-stranded forms to single-stranded forms (Fig. 3). Therefore, all four mutants were RNaseH deficient.




Figure 3. Identification of the DEDD motif in the HBV RNaseH active site. Wild-type and mutant HBV genotype A genomic expression vectors were transfected into cells, intracellular capsids were isolated five days later, and viral nucleic acids were purified from the capsids. The nucleic acids were divided into two aliquots; one aliquot was treated with DNase-free *E. coli* RNaseH to destroy RNA:DNA heteroduplexes and the other was mock treated. The nucleic acids were resolved by agarose electrophoresis and HBV DNAs were detected by Southern analysis. The signature of an RNaseH-deficient genome is production of RNA:DNA heteroduplexes in which the DNA migrates as double-stranded species without treatment with exogenous RNaseH treatment but as single-stranded species following degradation of the RNA. The positions of the duplex linear (DL) and full-length single-stranded linear (SL) HBV DNA markers are shown. DS indicates the spectrum of double-stranded nucleic acids produced by reverse transcription, and SS indicates the spectrum of single-stranded nucleic acids.

doi:10.1371/journal.ppat.1003125.g003

Production of enzymatically active recombinant HBV RNaseH

We expressed HBV RNaseH sequences from the HBV isolate employed by Potenza et al. [57] in *E. coli* as a carboxy-terminally hexahistidine tagged recombinant protein, but we moved the amino terminus nine residues upstream to residue 684 of the HBV polymerase because we felt this site was more probable to yield soluble protein (HRHPL; Fig. 4A). As a negative control, we mutated two of the DEDD active site residues (D702A and E731A). These constructs were expressed in *E. coli*, soluble lysates were prepared, and the lysates were subjected to nickel-affinity chromatography.

Five proteins of approximately 80, 70, 26, 14, and 11 kDa detectable by Coomassie staining were recovered following chromatography, none of which correlated with the predicted mass of 18.9 kDa for HRHPL (Fig. 4B). Mass spectrometry identified the dominant 26 kDa band as the *E. coli* prolyl isomerase SlyD. Concentrating the samples seven-fold did not increase the RNaseH to levels detectable by Coomassie staining. Western analysis with anti-polyhistidine antibodies revealed a large number of cellular bands but failed to unambiguously identify HRHPL. This was presumably due to the presence of histidine-rich regions in the bacterial proteins that promoted their binding to the nickel-affinity resin (e.g., SlyD). However, western analysis with the anti-HBV RNaseH domain antibody 9F9 ([80]; Santa Cruz Biotechnology) revealed a small amount of recombinant HBV RNaseH that migrated close to its predicted mass plus a larger amount of the protein that migrated as a doublet near 15 kDa (Fig. 4B). The doublet is presumably due to proteolysis near the protein's N-terminus because the antibody epitope and hexahistidine tag are at the C-terminus. The sizes of the truncation products imply that they were cleaved near HRHPL residue 36, which would remove the essential D702 carboxylate (HRHPL residue 20) and inactivate the protein. These experiments indicate we could express and enrich small but detectable amounts of soluble recombinant HBV RNaseH.

We tested activity of the recombinant HBV RNaseHs in a DNA oligonucleotide-directed RNA cleavage assay. In this assay, a DNA oligonucleotide is annealed to a uniformly-labeled RNA to create an RNA:DNA heteroduplex. Cleavage of the RNA in the heteroduplex yields two RNA fragments of predictable size that are resolved by electrophoresis and detected by autoradiography (Fig. 5A). We employed the 264 nt RNA (DRF+) used in our previous RNaseH assays [46] in combination with two DNA oligonucleotide pairs. One oligonucleotide in each pair was the correct polarity to anneal to the DRF+ RNA and the other was its inverse complement as a negative control.

Oligonucleotide-directed RNaseH assays were conducted with wild-type HRHPL enzyme and the RNaseH-deficient D702A mutant. The RNA was not cleaved when the non-complementary oligonucleotides were employed in the reactions (Fig. 5B), demonstrating that the enzyme preparations did not contain non-specific RNase activity. Use of complementary oligonucleotide #1 (D2507-) led to complete cleavage of the DRF+ RNA by *E. coli* RNaseH into products of 154 and 94 nt, and to partial cleavage of the RNA at the same site by wild-type HRHPL (Fig. 5B). The large majority of this RNaseH activity was due to the HBV enzyme because mutating DEDD residues D702A and/or E731A sharply reduced cleavage of the RNA. Note that although the relative yield of full-length mutant RNaseH was less than the wild-type enzyme in Fig. 4, in other preparations the amount of mutant RNaseH exceeded the amount of wild-type enzyme (e.g., Fig. 6). In all cases, the enzymatic activity associated with the mutant RNaseH preparations was far lower than in the




Figure 4. Recombinant HBV RNaseH proteins. **A.** Structure of the recombinant RNaseHs. The HBV polymerase with its major domains labeled is at top. The recombinant RNaseH derivatives are shown below with the C-terminal hexahistidine tag in brown. TP, terminal protein domain; RT, reverse transcriptase domain; *, mutations D702A or E731A to RNaseH active site residues. **B.** Proteins in the enriched lysates. The left panel is a Coomassie-blue stained SDS-PAGE gel of enriched RNaseH extracts as employed in the RNaseH assays. The right panel is a western blot of the extracts employing monoclonal antibody 9F9 which recognizes an epitope near the C-terminus of the HBV polymerase.
doi:10.1371/journal.ppat.1003125.g004

wild-type preparations. The residual cleavage products in reactions with the mutant enzymes appear to be non-specific breakdown products from the RNA substrate and/or digestion products from trace contamination with bacterial RNaseH. The RNA products shifted sizes as expected when complementary oligonucleotide #2 (D2543M-Sal, which anneals 33 nt closer to the 3' end of the RNA) was employed in the RNaseH assays (Fig. 5B): the larger fragment became larger (187 nt) and the smaller fragment became smaller (61 nt). These data demonstrate that the RNase activity in HRHP is specific for RNA annealed to the DNA oligonucleotides, and hence confirm that it is an RNaseH activity.

Finally, we synthesized a quenched fluorescent RNA:DNA chimeric hairpin oligonucleotide substrate (RHF1) to confirm RNaseH activity with a different assay. RHF1 has fluorescein at its 5' end, 20 nt of RNA, a 4 nt DNA hairpin, 20 nt of DNA complementary to the RNA, and an Iowa Black FQ quencher at the 3' terminus. The hairpin brings the fluorescein and quencher into close proximity, and digesting the RNA frees the fluorescein and increases its fluorescence (Fig. 5C). RHF1 was terminally digested with *E. coli* RNaseH, the reactions were terminated with 10 mM EDTA, and fluorescence was measured. This digestion amplified the fluorescence of RHF1 22-fold, indicating a 95% quenching efficiency. RHF1 was then employed in an RNaseH assay with buffer alone, wild type HBV RNaseH (HRHPL), and HRHPL-D702A/E731A. RNaseH activity for HRHPL was about 2-fold higher than the no-enzyme control, and mutating the RNaseH active site eliminated this activity (Fig. 5D). This weak signal (7% of the maximal signal in this assay) appears to be

due to poor binding between the small substrate and the RNaseH in the relatively high ionic strength of the reactions because detection of RNaseH activity required reducing the NaCl concentration from 190 to 130 mM.

These data indicate that we can readily detect HBV RNaseH activity in the enriched bacterial extracts despite the fact that the HBV RNaseH is a minor component of the mixture.

Optimization of reaction conditions

The optimal enzymatic conditions for the HRHPL HBV RNaseH were determined by systematically varying the reaction components in the oligonucleotide-directed RNaseH assay (Table 1). Recombinant HBV RNaseH was active over a wide range of pH values but was most active near 8.0. Its activity maximum was at 190 mM NaCl and it became able to digest single-stranded RNA below ~100 mM NaCl. The RNaseH required ~5 mM Mg⁺⁺ for maximal activity; increasing Mg⁺⁺ beyond ~7 mM suppressed RNaseH activity, and inclusion of Mn⁺⁺ in the reactions led to nonspecific degradation of single-stranded RNA. The enzyme became inactive at low reductant concentrations, but it could tolerate up to 2% DMSO. It was stable upon storage in liquid nitrogen, and only marginal loss of activity was observed following five sequential freeze-thaw cycles.

Recombinant RNaseH enzymes from other HBV genotypes

HBV has eight genotypes (A–H, plus provisional identification of genotypes I and J) that differ by >8% at the sequence level [81]. We cloned HBV RNaseH domains for genotype A, B, C, and H




Figure 5. Recombinant HBV RNaseH is enzymatically active. **A.** Oligonucleotide-directed RNaseH assay. Uniformly ^{32}P -labeled RNA (blue or red) is annealed to a complementary DNA oligonucleotide (black). RNaseH activity cleaves the RNA in the heteroduplex formed where the oligonucleotide anneals to the RNA and yields two products (P1 and P2). **B.** Recombinant HBV RNaseH is active. An oligonucleotide-directed RNaseH assay was conducted with *E. coli* RNaseH, wild-type HBV RNaseH (HRHPL), or RNaseH-deficient HRHPL (D702A). A complementary oligonucleotide (+) or non-complementary oligonucleotide (-) was mixed with labeled DRF+ RNA and the reactions were incubated to allow RNaseH activity. The products were resolved by SDS-PAGE and the RNAs were detected by autoradiography. Oligonucleotide set 1 was D2507- and D2526+ and oligonucleotide set #2 was D2543M-Sal and D2453+. The positions of the cleavage products (P1 and P2) are indicated in blue for reactions containing oligonucleotide D2507- and in red for reactions containing oligonucleotide D2543M-Sal. **C.** FRET-based RNaseH assay. A self-complementary chimeric RNA:DNA synthetic oligonucleotide (RHF1) forms a stem-loop in which the stem is an RNA:DNA heteroduplex. The stem brings the fluorescein (F) and quencher (Q) at the 5' and 3' ends of the oligonucleotide into close proximity. Cleavage of the RNA releases the fluorescein and increases its fluorescence. **D.** Detection of HBV RNaseH activity employing the fluorescent assay. The substrate in panel C was employed in an RNaseH assay employing buffer alone, wild-type HBV RNaseH (HRHPL), or RNaseH-deficient HRHPL (D702A/E731A). *, $P < 0.05$. doi:10.1371/journal.ppat.1003125.g005

isolates using the same structure as the HRHPL construct (genotype D) to determine whether HBV's genetic diversity leads to variable sensitivity to inhibitors that must be taken into account during drug development (Fig. 6A). The protein profile detectable by Coomassie staining following expression and nickel-affinity enrichment for all additional constructs was the same as for HRHPL. Western blotting with antibody 9F9 detected the genotype B, C, and D RNaseHs, with the genotype C enzyme appearing primarily as the full-length protein (Fig. 6B). The failure to detect the genotype A and H RNaseHs was due either to lack of accumulation of the proteins or to amino acid variations in the C-terminus of the protein where the antibody epitope is located [80].

The genotype A, B, C, D, and H RNaseH extracts were assessed with the oligonucleotide-directed RNaseH assay (Fig. 6C). The genotype A and B enzymes were inactive, activity of the genotype C RNaseH ranged from undetectable to modest in replicate experiments, and activity of the genotype H enzyme was similar to that of the genotype D RNaseH. The $[\text{NaCl}]^-$, $[\text{Mg}^{++}]^-$, temperature-, and pH-profiles of the genotype H RNaseH were very similar to those of the genotype D enzyme (data not shown).

Therefore, we can express recombinant HBV genotype B, C, D, and H RNaseH proteins that are detectable by enzymatic assays and/or western blotting, but only the genotype D and H proteins are consistently active.

Identification of anti-HBV RNaseH compounds

We hypothesized that the HBV RNaseH may be inhibited by antagonists of the HIV RNaseH based on the similarity of the

reactions they catalyze. We identified 10 compounds known to inhibit the HIV RNaseH or that were predicted by chemical structure-activity relationships to do so (Table 2 and Supplementary Fig. S1). We further hypothesized that anti-HIV integrase compounds may inhibit the HBV RNaseH because the integrase and RNaseH are both members of the nucleotidyl transferase superfamily and because some anti-HIV RNaseH and integrase compounds can cross-inhibit their target enzymes [59,68,70,74,79]. Consequently, we also obtained 11 compounds either known to inhibit the HIV integrase or predicted to do so by chemical structure-activity relationships (Table 2 and Supplementary Fig. S1). We first measured the effect of irrelevant compounds (tryptophan, sucrose, and IPTG) on the RNaseH assay. These compounds reduced RNaseH activity of HRHPL to $52 \pm 9\%$ relative to the DMSO vehicle control (Figs. 7 and 8A). This allowed us to define the mean of the residual activity in the presence of the irrelevant compounds minus two standard deviations of the irrelevant controls as a threshold reduction of the RNaseH activity that must be exceeded before we considered inhibition by the test compounds to be relevant. Using this threshold, 12 of the 21 compounds inhibited the HBV genotype D RNaseH at $10 \mu\text{M}$ (Fig. 7, Table 2, and Supplementary Table S1). These 21 compounds were also screened against the HBV genotype H RNaseH using the oligonucleotide-directed RNaseH assay.

The unexpectedly high frequency of inhibition of the genotype D enzyme led us to question the mechanism(s) by which it was inhibited by the compounds. We addressed this in two manners. First, RNaseH inhibitors usually block the HIV enzyme by interfering with

Table 2. Candidate RNAseH inhibitors and data summary.

Compound			RNAseH assays ²		HBV Replication ³	
Number	Name ¹	HIV template	gtD	gtH	gtA	gtD
2	Sigma 74540	RNAseH	-	-	-	-
3	Sigma n8164	RNAseH	-	-	-	-
4	TimTec ST029023	RNAseH	+/-	-	-	-
5	Enamine T0506-3483	RNAseH	+/-	-	-	-
6	Chembridge 7929959	RNAseH	+/-	-	-	-
7	Idofine 02030	Integrase	-	-	-	-
8	Sigma S439274	Integrase	+	-	-	-
9	Sigma 70050	Integrase	+	-	-	-
10	Selleck S2001	Integrase	+	-	-	-
11	Selleck S2005	Integrase	+	-	-	-
12	Naphthyridinone ⁴	RNAseH	+	++	+	+
13	DHBNH ⁴	RNAseH	-	+/-	-	-
14	THBNH ⁴	RNAseH	-	+	+/-	-
15	BHMP07 ⁴	RNAseH	-	+/-	-	-
30	Chembridge 7248520	Integrase	++	+/-	-	-
31	Chembridge 5104346	Integrase	+	-	-	-
34	Indofine D-009	Integrase	+	+	-	-
35	TCI America D1118	Integrase	+	+/-	-	-
38	Vistas M Lab STK317995	RNAseH	++	+/-	-	-
39	Asinex BAS0223612	Integrase	++	+	-	-
40	NIH 118-D-24	Integrase	++	++	+/-	-

¹Structures are in Supplemental Fig. S1.

²Inhibitory activity in the biochemical RNAseH assay at 10 μ M; quantitative data are in Supplemental Table S1. Cutoff values were established relative to the mean and standard deviation of RNAseH activity in the presence of the irrelevant compounds tryptophan, sucrose, and IPTG normalized to the vehicle control. -, Residual activity greater than one standard deviation below the mean irrelevant inhibition ($\geq 43\%$ activity); +/-, Residual activity between one and two standard deviations below the mean irrelevant inhibition (42-34% activity); +, residual activity between two and three standard deviations below the mean irrelevant inhibition (33-25% activity); ++, residual activity less than three standard deviations below the mean irrelevant inhibition ($\leq 24\%$ activity).

³Inhibitory activity against HBV replication in Huh7 cells at 10 μ M. +, clear inhibition; -, no detectable inhibition; +/-, inhibition observed in some but not most assays.

⁴Naphthyridinone [68]; DHBNH, dihydroxy benzoyl naphthyl hydrazone [75,78]; THBNH, trihydroxy benzoyl naphthyl hydrazone [106]; BHMP07 [78].

doi:10.1371/journal.ppat.1003125.t002




Figure 7. Inhibition of the HBV RNAseH by candidate compounds selected for their similarity to antagonists of the HIV RNAseH and integrase. Candidate inhibitors (compounds #2-40) and irrelevant compounds (tryptophan, sucrose, and IPTG) were included at 10 μ M in a standard oligonucleotide-directed RNAseH assay employing wild-type genotype D HBV RNAseH (HRHPL). DMSO, vehicle control. Error bars are \pm one standard deviation from three to seven replicates. The dashed red line indicates the mean residual activity in the irrelevant control reactions (52%) and the solid red line is two standard deviations of the irrelevant control assays below their mean (33%). Compounds that inhibited the RNAseH to 33% or below were considered to be positive ("+" in Table 2). *, $P < 0.05$ by T-test against the pooled data for the irrelevant controls; **, $P < 0.01$. doi:10.1371/journal.ppat.1003125.g007




Figure 8. Specificity of anti-HBV RNaseH compounds. **A.** Inhibition of HBV genotype D RNaseH by irrelevant compounds at 10 μM in the oligonucleotide-directed RNaseH assay. Compound #4 was employed as an example HBV RNaseH inhibitor. **B.** Anti-HBV RNaseH inhibitors do not significantly inhibit the HCV RNA polymerase. The ability of compounds #5, 6 and 8 to inhibit production of poly-G by the HCV RNA-directed RNA polymerase was measured in a primed homopolymeric RNA synthesis assay [82]. The compounds were employed at 10 μM. DMSO, vehicle control. **C.** Dose-responsiveness of HBV RNaseH inhibition. The effects of compounds #6, 8, and 12 on the RNaseH activity of HRHPL (genotype D) were measured at concentrations ranging from 0.5 to 50 μM. The dose-response profile is plotted for compound #12. doi:10.1371/journal.ppat.1003125.g008

Activity of HBV RNaseH inhibitors against human RNaseH1

A likely cause of cellular toxicity for anti-HBV RNaseH drugs would be inhibition of human RNaseH1 because it is responsible for about 80% of the RNaseH activity in human cells [83,84]. Therefore, we cloned the human RNaseH1 with an N-terminal hexahistidine tag, expressed it in *E. coli*, and enriched the protein by nickel affinity chromatography. The same spectrum of contaminating *E. coli* proteins as was observed for the other RNaseH preparations was detectable by Coomassie staining, but RNaseH1 could be detected at its predicted mass of 32 kDa (Fig. 9A). This enzyme was active in the oligonucleotide-directed and fluorescent RNaseH assays (Fig. 9B and data not shown). To determine how inhibition of human RNaseH1 compared to inhibition of the HBV RNaseH, we titrated RNaseH1 to yield similar levels of activity as the HBV enzyme, and then we directly compared the ability of compounds #8-12 to inhibit human RNaseH1 and HRHPL at 10 μM. All five compounds inhibited the HBV RNaseH. Compound #8 inhibited RNaseH1 well, #9 and 12 inhibited it weakly, and #10 and 11 had no effect on RNaseH1. Therefore, it is possible to inhibit the HBV RNaseH without inhibiting human RNaseH1.

Anti-HBV RNaseH compounds can inhibit HBV replication in culture

Finally, we asked whether HBV RNaseH inhibitors could block HBV replication in culture. Huh7 cells were transfected with genomic expression vectors for HBV genotype A or D isolates, the cells were treated with 10 or 50 μM compounds, and viral nucleic acids were isolated from intracellular HBV capsids after four days. Replicate nucleic acid aliquots were mock treated or treated with DNase-free *E. coli* RNaseH to destroy RNA:DNA heteroduplexes, and then HBV DNAs were detected by Southern blotting. The signature of RNaseH inhibition is accumulation of RNA:DNA heteroduplexes that migrate as double-stranded species without exogenous RNaseH treatment but as faster-migrating single-stranded DNAs following RNaseH treatment.

The mobility of the DNAs synthesized in cells containing the wild-type genotype A genome was unaffected by exogenous RNaseH treatment (Fig. 10). Ablation of RNaseH activity by the D702A mutant altered migration of the double-stranded forms, and treatment of these samples with RNaseH collapsed the double-stranded forms to single-stranded DNAs (Fig. 10 left panel). The mobility of HBV DNAs from cells replicating HBV genotype A treated with DMSO was unaffected by RNaseH digestion (Fig. 10 center panel), but treatment of cells with




Figure 9. Activity of HBV RNaseH inhibitors against human RNaseH1. **A.** Proteins in the enriched recombinant human RNaseH1 lysates employed in the RNaseH reactions were detected by Coomassie-blue staining following SDS-PAGE. **B.** An oligonucleotide-directed RNaseH assay was conducted with wild-type HBV RNaseH (genotype D) and recombinant human RNaseH1 under identical reaction conditions. The inhibitory compounds were employed at 10 μ M. The upper and lower panels are from the same experiment and the data were collected on a single sheet of film, so the reactions can be directly compared. DMSO, vehicle control. S, the DRF+ substrate; P1 and P2, RNaseH cleavage products.
doi:10.1371/journal.ppat.1003125.g009

compound #12 at 10 μ M blocked production of the slowest-migrating double-stranded forms and led to accumulation of RNA:DNA heteroduplexes whose mobility increased upon removal of RNA. Treatment of cells with 3 to 50 μ M compound #12 revealed that the degree of inhibition was proportional to the concentration of the compound (data not shown). Plus-strand preferential real-time PCR across the gap in the minus-polarity viral DNA revealed that 10 μ M compound #12 reduced plus-strand DNA accumulation to 7.3% of the DMSO-treated control (data not shown). None of the other compounds reproducibly inhibited HBV genome synthesis (Table 2), but compound #14 (25 μ M) inhibited HBV replication in one experiment and #40 (50 μ M) inhibited replication in another experiment. Overt cellular toxicity was not observed for any of the compounds at 10 μ M. Toxicity was often observed at higher concentrations; this led to the reduced yield of HBV DNA from cultures treated with 50 μ M compounds #5, 6, and 8 in Fig. 10.

The effect of the compounds on replication of a genotype D isolate was tested to evaluate the generality of the results with the genotype A isolate. Treatment of capsid-derived nucleic acids from

the DMSO control cells with exogenous RNaseH led to partial conversion of the double-stranded molecules to single-stranded forms. Therefore, RNA:DNA heteroduplexes accumulated in capsids even in the absence of RNaseH inhibitors. This indicates that the RNaseH activity during reverse transcription was incomplete for this isolate. Very few of the most slowly-migrating double-stranded nucleic acids accumulated in cells treated with 10 μ M compound #12, and many of the duplex DNAs collapsed to single-stranded forms upon treatment with exogenous RNaseH. Therefore, the inefficient HBV RNaseH in this isolate created a high background, but we were able to detect suppression of the HBV RNaseH activity above background by compound #12. None of the other compounds tested against the genotype D isolate detectably inhibited HBV replication (Table 2).

Therefore, compound #12 inhibited replication of HBV genotypes A and D in cells at low μ M concentrations by blocking RNaseH activity, with the anti-RNaseH effect being somewhat less pronounced than complete ablation of the activity by mutating the RNaseH active site.

Discussion

Nucleos(t)ide analog therapy has turned chronic HBV infection into a disease that can be controlled indefinitely, with enormous benefits to patients [6,7,85]. However, the infection is very rarely cleared, so treatment is essentially life-long, very expensive, and may be associated with unpredictable long-term side effects. Despite these limitations, the ability of protracted nucleos(t)ide analog therapy to slowly suppress cccDNA and HBsAg and to cure a small minority of HBV patients [10–13,21–23] indicates that the nucleos(t)ide analogs can push the virus to the brink of elimination. This implies that many more patients could be cured by employing a new drug against a novel HBV target in combination with the nucleos(t)ide analogs to further suppress HBV replication. Here, we report production of recombinant HBV RNaseH suitable for low throughput antiviral drug screening and demonstrate that chemical structure-activity relationships based on HIV RNaseH and integrase inhibitors can guide identification of compounds likely to inhibit the HBV enzyme.

Production of soluble recombinant HBV polymerase or domains of the polymerase is notoriously difficult, and our experience with the HBV RNaseH domain was no exception. Soluble HBV RNaseH accumulated to low levels in *E. coli* and was a minor component of the extracts even after nickel-affinity enrichment (Fig. 4). Much of the RNaseH was apparently cleaved near its N-terminus, and these cleavage products are unlikely to be active because their sizes imply that they lack D702. Although the concentration of the intact enzyme was very low, its specific activity was high enough to yield readily detectable signals in both radioactive and fluorescent RNaseH assays (Fig. 5). Potenza et al. [57] previously expressed recombinant HBV RNaseH that was very similar to HRHPL (genotype D), but their expression conditions led to accumulation of the enzyme in inclusion bodies, necessitating refolding following purification under denaturing conditions. The refolded enzyme possessed RNaseH activity, but this activity was not demonstrated to be an RNaseH. Differences between the assays employed here and in Potenza's study prevent comparison of the specificity and specific activity of the enzyme prepared under native and denaturing conditions.

The optimal reaction conditions for the recombinant HBV RNaseH (Table 1) were typical for nucleic-acid modifying enzymes and were similar to conditions in which recombinant hepadnaviral reverse transcriptase is active [86–88]. Its activity was dependent upon a divalent cation, but it became active against




Figure 10. Inhibition of HBV replication in culture by RNaseH inhibitors. Genotype A or D HBV genomic expression vectors were transfected into cells, intracellular capsids were isolated four days later, and viral nucleic acids were purified from the capsids. The nucleic acids were divided into two aliquots; one aliquot was treated with DNase-free *E. coli* RNaseH to destroy RNA:DNA heteroduplexes and the other was mock treated. The nucleic acids were resolved by agarose electrophoresis and HBV DNAs were detected by Southern analysis. Inhibition of RNaseH activity leads to accumulation of RNA:DNA heteroduplexes in which the DNA migrates as double-stranded species in the mock-treated sample but as faster-migrating single-stranded species following RNaseH treatment. The left panel is a control in which wild-type and RNaseH-deficient D702A HBV genomes were compared. The right two panels employed wild-type HBV in the presence of the test compounds. Compounds were employed at 10 or 50 μM as indicated. DMSO, vehicle control. The positions of the duplex linear (DL) and full-length single-stranded linear (SL) HBV DNA markers are shown. DS indicates the spectrum of double-stranded nucleic acids produced by reverse transcription, and SS indicates the spectrum of single-stranded nucleic acids.

doi:10.1371/journal.ppat.1003125.g010

single-stranded RNA in addition to RNA in a heteroduplex when Mn^{++} was substituted for Mg^{++} (data not shown). This is similar to the reduced fidelity of nucleic acid polymerases (including the duck HBV polymerase) in the presence of Mn^{++} [89–91]. The RNaseH had a relatively high NaCl optimum of 190 mM and it lost specificity for heteroduplex RNA at low ionic strength (data not shown). Importantly given that a primary goal of this study was to produce enzyme suitable for antiviral drug screening, recombinant HBV RNaseH was stable upon storage in liquid nitrogen, could be repeatedly frozen and thawed, and was fully active in up to 2% DMSO. Therefore, enzyme suitable for low-throughput anti-HBV RNaseH drug screening has been produced.

The HIV RNaseH is a very active target of ongoing antiviral drug discovery [38,59–78], but to our knowledge none of the anti-HIV RNaseH compounds have entered clinical trials yet. This is primarily due to the relatively low therapeutic indexes of most known anti-HIV RNaseH compounds. Similar challenges were faced by the HIV integrase field in the early stages of development of anti-integrase drugs. Many inhibitors were discovered, but clinical development did not begin until strand transfer inhibitors, active site metal binders, etc. were discovered. The failure to advance to HIV RNaseH inhibitors to clinical trials may also be partially due to the large number, high potency, and diverse profile of existing anti-HIV drugs. In contrast, current anti-HBV therapies are primarily based on a single class of inhibitors, nucleos(t)ide analogs. Hence, inhibitors of a new HBV enzymatic function would address the current challenges of limited efficacy and cross-resistance among the nucleos(t)ide analogs, and this would allow meaningful combination therapies for HBV similar to HAART that dramatically changed the landscape of anti-HIV therapy.

The ability to template HBV RNaseH drug discovery on the HIV experience would greatly accelerate anti-HBV efforts. The HIV data could narrow the chemical space to be assessed during screening, compounds synthesized during anti-HIV RNaseH screening would be available for immediate screening against HBV, and the toxicity profile of some of these compounds is known. Templating anti-HBV RNaseH drug development on HIV efforts would be analogous to the development of the anti-HBV nucleos(t)ide analogs, which was greatly facilitated by the parallel development of anti-HIV nucleoside analogs [92].

Twenty-one candidate RNaseH inhibitors were selected due to their similarity to known inhibitors of the HIV RNaseH or integrase. Twelve of these compounds (57%) inhibited the HBV RNaseH at 10 μM to below the threshold defined by control reactions with irrelevant compounds (Fig. 7 and Table 2). Importantly, 10 of 11 compounds analogous to anti-HIV integrase compounds inhibited the HBV RNaseH, including both approved anti-HIV integrase drugs, raltegravir (compound #11) and elvitegravir (#10). This is consistent with the membership of both the RNaseH and integrase in the nucleotidyl transferase superfamily of enzymes. Therefore, there is enough similarity between the HBV RNaseH and the HIV RNaseH and integrase active sites to guide screening for anti-HBV RNaseH compounds.

Most anti-HIV RNaseH inhibitors bind to the enzyme and chelate the divalent cations in the active site [64,65,69,70,73,76]. Similarly, anti-HIV integrase compounds that target the active site typically do so by binding to the enzyme or the enzyme plus DNA and chelating the active site divalent cations [93]. The compounds tested here were selected for the ability to bind to Mg^{++} ions oriented as they are in the HIV RNaseH or integrase active sites, and hence inhibition of the HBV enzyme is predicted to be through binding to the active site and interfering with the Mg^{++}

ions. The mechanisms by which the HBV RNaseH inhibitors function have not been determined, but IC_{50} curves reveal at least two patterns. The profiles for compounds #12, 39, and 40 were consistent with the predicted competitive inhibition mechanism (Fig. 8C and data not shown). In these cases, inhibition appears to be specific. Other compounds, such as #6 and #8, had inhibition profiles with one or more broad plateaus that were inconsistent with simple competitive binding to the active site. In addition, the electrophoretic mobility of the RNA was retarded at high concentrations of compound #8 (Fig. 8C), implying that this compound may react with the RNA substrate.

The compounds employed here were selected by structure-activity relationships with the goal of testing whether these relationships could predict biochemical inhibition of the HBV RNaseH. The compounds were not selected to have other properties necessary for a drug, such as the ability to enter cells. Nevertheless, compound #12 inhibited HBV replication in cell culture at 10 μ M without extensive cellular toxicity (Fig. 10). The reduction in mobility following treatment of capsid-derived nucleic acids with *E. coli* RNaseH demonstrates that RNA:DNA heteroduplexes accumulated in the viral capsid in the presence of compound #12, confirming that these compounds blocked HBV RNaseH activity in culture. Therefore, it is possible to pharmacologically inhibit the HBV RNaseH in cells, and identification of anti-HBV compounds that are active in cells can be achieved employing structure-activity relationships based on anti-HIV compounds. Furthermore, the ability of compounds identified by screening against recombinant genotype D and H enzymes to inhibit both genotype A and D isolates in culture demonstrates that it is possible to identify RNaseH inhibitors that are active against a range of HBV isolates.

The sensitivity profile of the HBV genotype D and H RNaseHs to the inhibitors was not the same (Table 2). This has two implications. First, the genotype H RNaseH may be a better candidate for primary drug screening than the genotype D enzyme because its inhibition profile more accurately predicted inhibition of HBV replication in culture. Second, the variable sensitivity of the genotype D and H enzymes to the compounds indicates that HBV's high genetic diversity is likely to be an important issue during development of anti-HBV RNaseH drugs.

The key HBV molecule that must be eradicated to cure patients is the viral cccDNA (Fig. 1) [25]. Ideally, clearing the cccDNA would be achieved by simultaneously suppressing its synthesis rate with the existing nucleos(t)ide inhibitors and increasing its degradation rate with a new drug. The problem with this approach is that we do not know how to safely destabilize the cccDNA, so the approach that has the most realistic chance of clearing HBV in the foreseeable future is to further suppress its synthesis rate. Importantly, pharmacological suppression of viral genomic synthesis may not need to completely eradicate the cccDNA by itself because the latter stages of viral clearance may be assisted by the immune system. HBV's proteins, including HBsAg [94–99], HBeAg [100,101], and the polymerase [102–104], have immunosuppressive activities. Consequently, if viral genomic replication can be suppressed far enough to inhibit cccDNA synthesis rather than just virion secretion (Fig. 1) as is usually achieved with the nucleos(t)ide analogs, levels of the cccDNA would drop. This reduction in the transcriptional template would reduce production of HBV's proteins, presumably weakening HBV's immunosuppression and promoting immune-mediated viral clearance.

Three challenges remain prior to beginning full-scale antiviral drug screening against the HBV RNaseH. First, the majority of HBV's disease burden is caused by genotypes B and C, and we

have been unsuccessful to date in generating consistently active recombinant RNaseH from these genotypes. This challenge is likely to be surmountable because only a few isolates of these genotypes have been tested for activity and because compound #12 identified by screening against genotypes D and H inhibited replication of HBV genotype A in culture, confirming that cross-genotype inhibition is possible. Second, the existing tissue culture and biochemical assays are sufficient for low throughput drug screening, but anti-HBV RNaseH drug development is anticipated to require screening many thousands of compounds even when the chemical search space is constrained by prior studies with HIV. Therefore, full-scale drug screening and subsequent mechanistic assessment of hit compounds will require improving the yield and purity of the biochemical RNaseH assay. This challenge should be met by further optimizing the induction and extraction conditions, expanding the bacterial induction cultures beyond the 100 ml scale used in this study, adding a second purification step such as ion-exchange chromatography, and expanding efforts to control proteolysis of the enzyme. We are optimistic this goal can be achieved because recent improvements to the induction and extraction conditions have increased the specific activity of the enzyme approximately four-fold, and initial scale-up experiments have not met with difficulty. Finally, the HBV RNaseH assay must be adapted to a format suitable for high throughput screening. This challenge should also be surmountable because fluorescent RNaseH assays have been widely employed to screen for anti-HIV RNaseH inhibitors and because the signal:background ratio for the first-generation HBV RNaseH fluorescent assay in Fig. 5 should be improved by increasing the concentration of the RNaseH and/or by optimizing the substrate structure.

Materials and Methods

Plasmids and viral strains employed

pCMV-HBV-LE- (CMV-HBV) is an HBV over-length genomic expression vector containing 1.2 copies of the HBV(adw2) genome (Genbank X02763.1) downstream of the CMV promoter cloned into pBS (Promega). Surface protein expression from this vector is ablated by mutating the preS and S open reading frames. pCMV-HBV(genotype D) is an analogous HBV genomic expression construct and was a gift from Dr. Shuping Tong. For bacterial expression, codon-optimized cDNA sequences for HRHPL (genotypes A, B, C, D, and H) were cloned by gene synthesis (Genscript) between the NcoI and EcoRI sites into pTrcHis2B (Invitrogen) with a C-terminal hexahistidine tag. HRHPL contains HBV genotype D (Genbank V01460) polymerase residues 684–845. The human RNaseH1 gene (NP_002927.2) was cloned with an N-terminal hexahistidine-tag between the BamHI and XhoI sites of pRsetB (Invitrogen) by gene synthesis.

RNaseH expression and enrichment

HRHPL and human RNaseH1 were expressed in *E. coli* BL21 codon+ cells (Invitrogen). Saturated overnight bacterial cultures were diluted 4-fold into 100 ml fresh medium and protein expression was induced with 0.5 mM IPTG at 30°C for six hours. The cells were lysed by sonication in lysis buffer [50 mM HEPES pH 8.0, 800 mM NaCl, 0.1% NP40, 27.5% glycerol, 2 mM DTT, 20 mM imidazole, and protease inhibitor cocktail (Sigma)]. RNaseH proteins were enriched by nickel-agarose affinity chromatography, eluted with 350 mM imidazole, dialyzed into 50 mM HEPES pH 7.3, 300 mM NaCl, 20% glycerol, and 5 mM DTT, and stored in liquid nitrogen.

In vitro RNaseH assays

For the oligonucleotide-directed RNaseH cleavage assay [46], 6 μ l protein extract (typical protein concentration 2.8 mg/ml) was mixed with 0.5 μ g internally 32 P-labeled DRF+ RNA (nucleotides 2401–2605 of the duck HBV genome plus 60 nucleotides of flanking sequences from pBluescript) and 3 μ g oligonucleotide D2507– or its corresponding negative control D2526+ on ice in 20 μ l under the conditions in Table 1. Some reactions in Fig. 5 employed oligonucleotide D2543M-Sal or its D2453+ negative control as indicated. The reactions were incubated at 42°C for 90 min. and terminated by addition of Laemmli protein loading buffer and boiling. The samples were resolved by 12% SDS-PAGE, the gels were stained with Coomassie blue to monitor protein loading, and labeled RNA was detected by autoradiography. Candidate inhibitors were dissolved in DMSO and added at the indicated concentrations during assembly of the reactions. Control reactions lacking the compounds contained DMSO as a vehicle control. The RNaseH autoradiograms were scanned and quantified with ImageJ. The oligonucleotides were: D2526+ (CCACATAGGCTATGTGGAAC), D2507– (GTTCCACATAGCCTATGTGG), D2453+ (CCGCCTGATTGGACGGCTTTCC), and D2543M-Sal (GCAACTGTGTCGACAGCAGCTCCGAAGGAGA).

For the fluorescent RNaseH assay, the DRF+ RNA and DNA oligonucleotides were omitted from the RNaseH reactions and replaced with 20 μ M of the quenched fluorescent chimeric RNA:DNA oligonucleotide RHF1; the reaction conditions were identical to the oligonucleotide-directed reactions except that the NaCl concentration was reduced to 130 mM. The reactions were incubated in the dark at 42°C for 90 min. prior to termination by addition EDTA to 10 mM and detection of fluorescence at 520 nm on Synergy 4 plate reader (Biotec, Inc.). The sequence of the RHF1 substrate (IDT, Inc.) was: 5'-56-FAM/rCrCrArCrArUrArGrGrCrUrArUrGrUrGrGrArArCTTTTGTTCACATAGCCTATGTGG/3IBkFQ/-3'. The RNA:DNA heteroduplex in the RHF1 substrate was the same as the heteroduplex formed by oligo D2507– annealed to DRF+.

Cell-based HBV replication inhibition assays

Huh7 cells were maintained in Dulbecco's modified Eagle's medium with 10% fetal bovine serum at 37°C in 5% CO₂. Cells were seeded into 60 mm dishes and transfected at 70% confluency with 2.6 μ g of plasmids using TransIT-LT1 (Mirus, Inc.). Test compounds were added the morning following transfection at 10 or 50 μ M, and fresh medium containing the compounds was

provided every 1–2 days. Four or five days post-transfection HBV cores were isolated by lysis of the cells in 10 mM Tris pH 7.5, 1 mM EDTA, 0.25% NP40, 50 mM NaCl, and 8% sucrose followed by sedimentation through a 30% sucrose cushion as described [105]. Viral DNAs were isolated from cytoplasmic core particle preparations by proteinase K digestion followed by phenol/chloroform extraction as described [46]. Duplicate aliquots of the nucleic acids were treated with 2 U *E. coli* DNase-free RNaseH (Invitrogen) at 37°C for 30 min. or were mock treated. The nucleic acid samples were resolved by electrophoresis on 1.2% agarose gels and detected by Southern blotting with 32 P-labeled HBV DNA as a probe.

Supporting Information

Figure S1 Chemical structures of the compounds tested. Compounds are named by the company/product number or their formal names, as appropriate. The approved anti-HIV integrase drugs Elvitegravir (#10) and Raltegravir (#11) are listed by their common names, pharmaceutical developer's codes, and company/product numbers. Compound #40 is listed by its NIH AIDS Research and Reference Reagent Program number. (PDF)

Table S1 Residual activity in RNaseH reactions conducted in the presence of 10 μ M of the test compounds. Values are normalized to vehicle control reactions containing 1% DMSO and the error ranges are \pm the standard deviation from 3 to 7 replicate experiments. (PDF)

Acknowledgments

We thank Nicola Pozzi for conducting the isothermal calorimetry experiments and Dr. Donghui Zhou for technical assistance. We thank Drs. David Wood, David Griggs, Peter Ruminski, Duane Grandgenett, and Enrico Di Cera for many helpful discussions. Compound 118-D-24 was obtained through the NIH AIDS Research and Reference Reagent Program, Division of AIDS, NIAID, NIH, through collaboration with Duane Grandgenett.

Author Contributions

Conceived and designed the experiments: JET YH FC EM RA MJM EJJ MAP SGS. Performed the experiments: JET XC YH MT FC EM. Analyzed the data: JET XC YH FC EM SGS. Contributed reagents/materials/analysis tools: JET RA MJM EJJ MAP SGS. Wrote the paper: JET YH SGS.

References

- Seeger C, Zoulim F, Mason WS (2007) Hepadnaviruses. In: Knipe DM, Howley P, Griffin DE, Lamb RA, Martin MA et al., editors. *Fields Virology*. Philadelphia: Lippincott Williams & Wilkins. pp. 2977–3029.
- Shepard CW, Simard EP, Finelli L, Fiore AE, Bell BP (2006) Hepatitis B virus infection: epidemiology and vaccination. *Epidemiol Rev* 28: 112–125.
- Ganem D, Prince AM (2004) Hepatitis B virus infection—natural history and clinical consequences. *N Engl J Med* 350: 1118–1129.
- Lavanchy D (2004) Hepatitis B virus epidemiology, disease burden, treatment, and current and emerging prevention and control measures. *J Viral Hepat* 11: 97–107.
- Tavis JE, Badtke MP (2009) Hepadnaviral Genomic Replication. In: Cameron CE, Götte M, Raney KD, editors. *Viral Genome Replication*. New York: Springer Science+Business Media, LLC. pp. 129–143.
- Cox N, Tillmann H (2011) Emerging pipeline drugs for hepatitis B infection. *Expert Opin Emerg Drugs* 16: 713–729.
- Kwon H, Lok AS (2011) Hepatitis B therapy. *Nat Rev Gastroenterol Hepatol* 8: 275–284.
- Michailidis E, Kirby KA, Hachiya A, Yoo W, Hong SP, et al. (2012) Antiviral therapies: focus on hepatitis B reverse transcriptase. *Int J Biochem Cell Biol* 44: 1060–1071.
- Zoulim F (2006) Antiviral therapy of chronic hepatitis B. *Antiviral Res* 71: 206–215.
- van Bommel F, De Man RA, Wedemeyer H, Deterding K, Petersen J, et al. (2010) Long-term efficacy of tenofovir monotherapy for hepatitis B virus-monoinfected patients after failure of nucleoside/nucleotide analogues. *Hepatology* 51: 73–80.
- Woo G, Tomlinson G, Nishikawa Y, Kowgier M, Sherman M, et al. (2010) Tenofovir and entecavir are the most effective antiviral agents for chronic hepatitis B: a systematic review and Bayesian meta-analyses. *Gastroenterology* 139: 1218–1229.
- Marcellin P, Heathcote EJ, Buti M, Gane E, De Man RA, et al. (2008) Tenofovir disoproxil fumarate versus adefovir dipivoxil for chronic hepatitis B. *N Engl J Med* 359: 2442–2455.
- Wursthorn K, Jung M, Riva A, Goodman ZD, Lopez P, et al. (2010) Kinetics of hepatitis B surface antigen decline during 3 years of telbivudine treatment in hepatitis B e antigen-positive patients. *Hepatology* 52: 1611–1620.
- Perrillo RP, Marcellin P (2010) Effect of newer oral antiviral agents on future therapy of chronic hepatitis B. *Antivir Ther* 15: 13–22.
- Zoulim F, Locarnini S (2009) Hepatitis B virus resistance to nucleos(t)ide analogues. *Gastroenterology* 137: 1593–1608.

16. Lau GK (2009) A new magic bullet for chronic hepatitis B infection: Is this the end of the story? *Gastroenterology* 136: 1830–1832.
17. Buti M, Brosa M, Casado MA, Rueda M, Esteban R (2009) Modeling the cost-effectiveness of different oral antiviral therapies in patients with chronic hepatitis B. *J Hepatol* 51: 640–646.
18. Ruggeri M, Cicchetti A, Gasbarrini A (2011) The cost-effectiveness of alternative strategies against HBV in Italy. *Health Policy* 102: 72–80.
19. Levrero M, Pollicino T, Petersen J, Belloni L, Raimondo G, et al. (2009) Control of cccDNA function in hepatitis B virus infection. *J Hepatol* 51: 581–592.
20. Tuttleman JS, POURCEL C, Summers J (1986) Formation of the pool of covalently closed circular viral DNA in hepadnavirus-infected cells. *Cell* 47: 451–460.
21. Werle-Lapostolle B, Bowden S, Locarnini S, Wursthorn K, Petersen J, et al. (2004) Persistence of cccDNA during the natural history of chronic hepatitis B and decline during adefovir dipivoxil therapy. *Gastroenterology* 126: 1750–1758.
22. Cheng PN, Liu WC, Tsai HW, Wu IC, Chang TT, et al. (2011) Association of intrahepatic cccDNA reduction with the improvement of liver histology in chronic hepatitis B patients receiving oral antiviral agents. *J Med Virol* 83: 602–607.
23. Wong DK, Yuen MF, Ngai VW, Fung J, Lai CL (2006) One-year entecavir or lamivudine therapy results in reduction of hepatitis B virus intrahepatic covalently closed circular DNA levels. *Antivir Ther* 11: 909–916.
24. Coffin CS, Mulrooney-Cousins PM, Peters MG, van MG, Roberts JP, et al. (2011) Molecular characterization of intrahepatic and extrahepatic hepatitis B virus (HBV) reservoirs in patients on suppressive antiviral therapy. *J Viral Hepat* 18: 415–423.
25. Zoulim F (2004) Antiviral therapy of chronic hepatitis B: can we clear the virus and prevent drug resistance? *Antivir Chem Chemother* 15: 299–305.
26. Monto A, Schooley RT, Lai JC, Sulkowski MS, Chung RT, et al. (2010) Lessons from HIV therapy applied to viral hepatitis therapy: summary of a workshop. *Am J Gastroenterol* 105: 989–1004.
27. Ghany M, Liang TJ (2007) Drug targets and molecular mechanisms of drug resistance in chronic hepatitis B. *Gastroenterology* 132: 1574–1585.
28. Hostomsky Z, Hostomska Z, Matthews DA (1993) Ribonuclease H. In: Linn SM, Lloyd RS, Roberts RJ, editors. *Nucleases*. Plainview, NY: Cold Spring Harbor Laboratory Press. pp. 341–376.
29. Yang W, Steitz TA (1995) Recombining the structures of HIV integrase, RuvC and RNase H. *Structure* 3: 131–134.
30. Katayanagi K, Miyagawa M, Matsushima M, Ishikawa M, Kanaya S, et al. (1990) Three-dimensional structure of ribonuclease H from *E. coli*. *Nature* 347: 306–309.
31. Rice PA, Baker TA (2001) Comparative architecture of transposase and integrase complexes. *Nat Struct Biol* 8: 302–307.
32. Dyda F, Hickman AB, Jenkins TM, Engelman A, Craigie R, et al. (1994) Crystal structure of the catalytic domain of HIV-1 integrase: similarity to other polynucleotidyl transferases. *Science* 266: 1981–1986.
33. Ariyoshi M, Vassilyev DG, Iwasaki H, Nakamura H, Shinagawa H, et al. (1994) Atomic structure of the RuvC resolvase: a holliday junction-specific endonuclease from *E. coli*. *Cell* 78: 1063–1072.
34. Song JJ, Smith SK, Hannon GJ, Joshua-Tor L (2004) Crystal structure of Argonaute and its implications for RISC slicer activity. *Science* 305: 1434–1437.
35. Frank P, Braunschöfer-Reiter C, Wintersberger U, Grimm R, Busen W (1998) Cloning of the cDNA encoding the large subunit of human RNase HI, a homologue of the prokaryotic RNase HII. *Proc Natl Acad Sci USA* 95: 12872–12877.
36. Frank P, Braunschöfer-Reiter C, Polt A, Holzmann K (1998) Cloning, subcellular localization and functional expression of human RNase HII. *Biol Chem* 379: 1407–1412.
37. Nowotny M, Gaidamakov SA, Crouch RJ, Yang W (2005) Crystal structures of RNase H bound to an RNA/DNA hybrid: substrate specificity and metal-dependent catalysis. *Cell* 121: 1005–1016.
38. Klumpp K, Hang JQ, Rajendran S, Yang Y, Derosier A, et al. (2003) Two-metal ion mechanism of RNA cleavage by HIV RNase H and mechanism-based design of selective HIV RNase H inhibitors. *Nucleic Acids Res* 31: 6852–6859.
39. Nowotny M, Yang W (2006) Stepwise analyses of metal ions in RNase H catalysis from substrate destabilization to product release. *EMBO J* 25: 1924–1933.
40. Keck JL, Goedken ER, Marqusee S (1998) Activation/attenuation model for RNase H. A one-metal mechanism with second-metal inhibition. *J Biol Chem* 273: 34128–34133.
41. Goedken ER, Marqusee S (2001) Co-crystal of *Escherichia coli* RNase HI with Mn²⁺ ions reveals two divalent metals bound in the active site. *J Biol Chem* 276: 7266–7271.
42. Li MD, Bronson DL, Lemke TD, Faras AJ (1995) Phylogenetic analyses of 55 retroelements on the basis of the nucleotide and product amino acid sequences of the pol gene. *Mol Biol Evol* 12: 657–670.
43. McClure MA (1993) Evolutionary History of Reverse Transcriptase. In: Skalka AM, editor. *Reverse Transcriptase*. Cold Spring Harbor: Cold Spring Harbor Laboratory Press. pp. 425–444.
44. Poch O, Sauvaget I, Delarue M, Tordo N (1989) Identification of four conserved motifs among the RNA-dependent polymerase encoding elements. *EMBO J* 8: 3867–3874.
45. Hu J, Seeger C (1996) Hsp90 is required for the activity of a hepatitis B virus reverse transcriptase. *Proc Natl Acad Sci USA* 93: 1060–1064.
46. Gong Y, Yao E, Tavis JE (2001) Evidence that the RNaseH activity of the duck hepatitis B virus is unable to act on exogenous substrates. *BMC Microbiology* 1: 12.
47. Radziwill G, Zentgraf H, Schaller H, Bosch V (1988) The duck hepatitis B virus DNA polymerase is tightly associated with the viral core structure and unable to switch to an exogenous template. *Virology* 163: 123–132.
48. Gerelsaikhan T, Tavis JE, Bruss V (1996) Hepatitis B Virus Nucleocapsid Envelopment Does Not Occur without Genomic DNA Synthesis. *J Virol* 70: 4269–4274.
49. Wei Y, Tavis JE, Ganem D (1996) Relationship between Viral DNA Synthesis and Virion Envelopment in Hepatitis B Viruses. *J Virol* 70: 6455–6458.
50. Chen Y, Robinson WS, Marion PL (1994) Selected Mutations of the Duck Hepatitis B Virus P Gene RNase H Domain Affect both RNA Packaging and Priming of Minus-Strand DNA Synthesis. *J Virol* 68: 5232–5238.
51. Chang C, Zhou S, Ganem D, Standring DN (1994) Phenotypic Mixing between Different Hepadnavirus Nucleocapsid Proteins Reveals C Protein Dimerization To Be cis Preferential. *J Virol* 68: 5225–5231.
52. Radziwill G, Tucker W, Schaller H (1990) Mutational analysis of the hepatitis B virus P gene product: domain structure and RNase H activity. *J Virol* 64: 613–620.
53. Chang LJ, Hirsch RC, Ganem D, Varmus HE (1990) Effects of insertional and point mutations on the functions of the duck hepatitis B virus polymerase. *J Virol* 64: 5553–5558.
54. Wei X, Peterson DL (1996) Expression, purification, and characterization of an active RNase H domain of the hepatitis B viral polymerase. *J Biol Chem* 271: 32617–32622.
55. Lee YI, Hong YB, Kim Y, Rho HM, Jung G (1997) RNase H activity of human hepatitis B virus polymerase expressed in *Escherichia coli*. *Biochem Biophys Res Commun* 233: 401–407.
56. Choi J, Kim EE, Park YI, Han YS (2002) Expression of the active human and duck hepatitis B virus polymerases in heterologous system of *Pichia methanolica*. *Antiviral Res* 55: 279–290.
57. Potenza N, Salvatore V, Raimondo D, Falanga D, Nobile V, et al. (2007) Optimized expression from a synthetic gene of an untagged RNase H domain of human hepatitis B virus polymerase which is enzymatically active. *Protein Expr Purif* 55: 93–99.
58. Freed EO, Martin MA (2007) HIVs and their replication. In: Knipe DM, Howley PM, Griffin DE, Lamb RA, Martin MA et al., editors. *Fields Virology*. Philadelphia: Lippincott Williams & Wilkins. pp. 2107–2185.
59. Klarmann GJ, Hawkins ME, Le Grice SF (2002) Uncovering the complexities of retroviral ribonuclease H reveals its potential as a therapeutic target. *AIDS Rev* 4: 183–194.
60. Klumpp K, Mirzadegan T (2006) Recent progress in the design of small molecule inhibitors of HIV RNase H. *Curr Pharm Des* 12: 1909–1922.
61. Takada K, Bermingham A, O'Keefe BR, Wamiru A, Beutler JA, et al. (2007) An HIV RNase H inhibitory 1,3,4,5-tetraalloylapitiol from the African plant *Hydrodendron gabunensis*. *J Nat Prod* 70: 1647–1649.
62. Bokesch HR, Wamiru A, Le Grice SF, Beutler JA, McKee TC, et al. (2008) HIV-1 ribonuclease H inhibitory phenolic glycosides from *Eugenia hymalis*. *J Nat Prod* 71: 1634–1636.
63. Wendeler M, Lee HF, Bermingham A, Miller JT, Chertov O, et al. (2008) Vinyllogous ureas as a novel class of inhibitors of reverse transcriptase-associated ribonuclease H activity. *ACS Chem Biol* 3: 635–644.
64. Fuji H, Urano E, Futahashi Y, Hamatake M, Tatsumi J, et al. (2009) Derivatives of 5-nitro-furan-2-carboxylic acid carbamoylmethyl ester inhibit RNase H activity associated with HIV-1 reverse transcriptase. *J Med Chem* 52: 1380–1387.
65. Su HP, Yan Y, Prasad GS, Smith RF, Daniels CL, et al. (2010) Structural basis for the inhibition of RNase H activity of HIV-1 reverse transcriptase by RNase H active site-directed inhibitors. *J Virol* 84: 7625–7633.
66. Di GM, Olson M, Prasad AS, Beberitz G, Luckay A, et al. (2010) Small molecule inhibitors of HIV RT Ribonuclease H. *Bioorg Med Chem Lett* 20: 398–402.
67. Chung S, Wendeler M, Rausch JW, Beilhartz G, Gotte M, et al. (2010) Structure-activity analysis of vinyllogous urea inhibitors of human immunodeficiency virus-encoded ribonuclease H. *Antimicrob Agents Chemother* 54: 3913–3921.
68. Williams PD, Staas DD, Venkatraman S, Loughran HM, Ruzek RD, et al. (2010) Potent and selective HIV-1 ribonuclease H inhibitors based on a 1-hydroxy-1,8-naphthyridin-2(1H)-one scaffold. *Bioorg Med Chem Lett* 20: 6754–6757.
69. Chung S, Himmel DM, Jiang JK, Wojtak K, Bauman JD, et al. (2011) Synthesis, activity, and structural analysis of novel alpha-hydroxytropolone inhibitors of human immunodeficiency virus reverse transcriptase-associated ribonuclease H. *J Med Chem* 54: 4462–4473.
70. Billamboz M, Bailly F, Lion C, Touati N, Vezin H, et al. (2011) Magnesium chelating 2-hydroxyisoquinoline-1,3(2H,4H)-diones, as inhibitors of HIV-1 integrase and/or the HIV-1 reverse transcriptase ribonuclease H domain:

- discovery of a novel selective inhibitor of the ribonuclease H function. *J Med Chem* 54: 1812–1824.
71. Didierjean J, Isel C, Querre F, Mouscadet JF, Aubertin AM, et al. (2005) Inhibition of human immunodeficiency virus type 1 reverse transcriptase, RNase H, and integrase activities by hydroxytropolones. *Antimicrob Agents Chemother* 49: 4884–4894.
 72. Budihias SR, Gorshkova I, Gaidamakov S, Wamiru A, Bona MK, et al. (2005) Selective inhibition of HIV-1 reverse transcriptase-associated ribonuclease H activity by hydroxylated tropolones. *Nucleic Acids Res* 33: 1249–1256.
 73. Himmel DM, Maegley KA, Pauly TA, Bauman JD, Das K, et al. (2009) Structure of HIV-1 reverse transcriptase with the inhibitor beta-Thujaplicinol bound at the RNase H active site. *Structure* 17: 1625–1635.
 74. Shaw-Reid CA, Munshi V, Graham P, Wolfe A, Witmer M, et al. (2003) Inhibition of HIV-1 ribonuclease H by a novel diketo acid, 4-[5-(benzoylamino)thien-2-yl]-2,4-dioxobutanoic acid. *J Biol Chem* 278: 2777–2780.
 75. Himmel DM, Sarafianos SG, Dharmasena S, Hossain MM, Coy-Simandle K, et al. (2006) HIV-1 reverse transcriptase structure with RNase H inhibitor dihydroxy benzoyl naphthyl hydrazone bound at a novel site. *ACS Chem Biol* 1: 702–712.
 76. Kirschberg TA, Balakrishnan M, Squires NH, Barnes T, Brendza KM, et al. (2009) RNase H active site inhibitors of human immunodeficiency virus type 1 reverse transcriptase: design, biochemical activity, and structural information. *J Med Chem* 52: 5781–5784.
 77. Suchaud V, Bailly F, Lion C, Tramontano E, Esposito F, et al. (2012) Development of a series of 3-hydroxyquinolin-2(1H)-ones as selective inhibitors of HIV-1 reverse transcriptase associated RNase H activity. *Bioorg Med Chem Lett* 22: 3988–3992.
 78. Gong Q, Menon L, Ilima T, Miller LG, Ahn J, et al. (2011) Interaction of HIV-1 reverse transcriptase ribonuclease H with an acylhydrazone inhibitor. *Chem Biol Drug Des* 77: 39–47.
 79. Billamboz M, Bailly F, Barreca ML, De LL, Mouscadet JF, et al. (2008) Design, synthesis, and biological evaluation of a series of 2-hydroxyisoquinoline-1,3(2H,4H)-diones as dual inhibitors of human immunodeficiency virus type 1 integrase and the reverse transcriptase RNase H domain. *J Med Chem* 51: 7717–7730.
 80. zu Putnitz J, Lanford RE, Carlson RI, Notvall L, De la Monte SM, et al. (1999) Properties of monoclonal antibodies directed against hepatitis B virus polymerase protein. *J Virol* 73: 4188–4196.
 81. Kurbanov F, Tanaka Y, Mizokami M (2010) Geographical and genetic diversity of the human hepatitis B virus. *Hepato Res* 40: 14–30.
 82. Cao F, Donlin MJ, Turner K, Cheng X, Tavis JE (2011) Genetic and biochemical diversity in the HCV NS5B RNA polymerase in the context of interferon alpha plus ribavirin therapy. *J Viral Hepat* 18: 349–357.
 83. Lima WF, Rose JB, Nichols JG, Wu H, Migawa MT, et al. (2007) Human RNase H1 discriminates between subtle variations in the structure of the heteroduplex substrate. *Mol Pharmacol* 71: 83–91.
 84. Lima WF, Wu H, Crooke ST (2001) Human RNases H. *Methods Enzymol* 341: 430–440.
 85. Liaw YF (2011) Impact of hepatitis B therapy on the long-term outcome of liver disease. *Liver Int* 31 Suppl 1: 117–121.
 86. Tavis JE, Ganem D (1993) Expression of functional hepatitis B virus polymerase in yeast reveals it to be the sole viral protein required for correct initiation of reverse transcription. *Proc Natl Acad Sci U S A* 90: 4107–4111.
 87. Hu J, Anselmo D (2000) In vitro reconstitution of a functional duck hepatitis B virus reverse transcriptase: posttranslational activation by HSP90. *J Virol* 74: 11447–11455.
 88. Beck J, Nassal M (2003) Efficient Hsp90-independent in vitro activation by Hsc70 and Hsp40 of duck hepatitis B virus reverse transcriptase, an assumed Hsp90 client protein. *J Biol Chem* 278: 36128–36138.
 89. Lin L, Wan F, Hu J (2008) Functional and structural dynamics of hepadnavirus reverse transcriptase during protein-primed initiation of reverse transcription: effects of metal ions. *J Virol* 82: 5703–5714.
 90. Arnold JJ, Ghosh SK, Cameron CE (1999) Poliovirus RNA-dependent RNA polymerase (3D(pol)). Divalent cation modulation of primer, template, and nucleotide selection. *J Biol Chem* 274: 37060–37069.
 91. Pelletier H, Sawaya MR, Wollfe W, Wilson SH, Kraut J (1996) A structural basis for metal ion mutagenicity and nucleotide selectivity in human DNA polymerase beta. *Biochemistry* 35: 12762–12777.
 92. Sturmer M, Doerr HW, Gurtler L (2009) Human immunodeficiency virus: 25 years of diagnostic and therapeutic strategies and their impact on hepatitis B and C virus. *Med Microbiol Immunol* 198: 147–155.
 93. Agrawal A, DeSoto J, Fullagar JL, Maddali K, Rostami S, et al. (2012) Probing chelation motifs in HIV integrase inhibitors. *Proc Natl Acad Sci U S A* 109: 2251–2256.
 94. Op den Brouw ML, Binda RS, van Roosmalen MH, Protzer U, Janssen HL, et al. (2009) Hepatitis B virus surface antigen impairs myeloid dendritic cell function: a possible immune escape mechanism of hepatitis B virus. *Immunology* 126: 280–289.
 95. Vanlandschoot P, Van HF, Roobrouck A, Farhoudi A, Leroux-Roels G (2002) Hepatitis B virus surface antigen suppresses the activation of monocytes through interaction with a serum protein and a monocyte-specific receptor. *J Gen Virol* 83: 1281–1289.
 96. Woltman AM, Op den Brouw ML, Biesta PJ, Shi CC, Janssen HL (2011) Hepatitis B virus lacks immune activating capacity, but actively inhibits plasmacytoid dendritic cell function. *PLoS ONE* 6: e15324.
 97. Wu J, Meng Z, Jiang M, Pei R, Trippler M, et al. (2009) Hepatitis B virus suppresses toll-like receptor-mediated innate immune responses in murine parenchymal and nonparenchymal liver cells. *Hepatology* 49: 1132–1140.
 98. Xu Y, Hu Y, Shi B, Zhang X, Wang J, et al. (2009) HBsAg inhibits TLR9-mediated activation and IFN-alpha production in plasmacytoid dendritic cells. *Mol Immunol* 46: 2640–2646.
 99. Cheng J, Imanishi H, Morisaki H, Liu W, Nakamura H, et al. (2005) Recombinant HBsAg inhibits LPS-induced COX-2 expression and IL-18 production by interfering with the NFkappaB pathway in a human monocytic cell line, THP-1. *J Hepatol* 43: 465–471.
 100. Chen M, Sallberg M, Hughes J, Jones J, Guidotti LG, et al. (2005) Immune tolerance split between hepatitis B virus precore and core proteins. *J Virol* 79: 3016–3027.
 101. Chen MT, Billaud JN, Sallberg M, Guidotti LG, Chisari FV, et al. (2004) A function of the hepatitis B virus precore protein is to regulate the immune response to the core antigen. *Proc Natl Acad Sci U S A* 101: 14913–14918.
 102. Wang H, Ryu WS (2010) Hepatitis B virus polymerase blocks pattern recognition receptor signaling via interaction with DDX3: implications for immune evasion. *PLoS Pathog* 6: e1000986.
 103. Foster GR, Ackrill AM, Goldin RD, Kerr IM, Thomas HC, et al. (1991) Expression of the terminal protein region of hepatitis B virus inhibits cellular responses to interferons alpha and gamma and double-stranded RNA. *Proc Natl Acad Sci U S A* 88: 2888–2892.
 104. Wu M, Xu Y, Lin S, Zhang X, Xiang L, et al. (2007) Hepatitis B virus polymerase inhibits the interferon-inducible MyD88 promoter by blocking nuclear translocation of Stat1. *J Gen Virol* 88: 3260–3269.
 105. Tavis JE, Massey B, Gong Y (1998) The Duck Hepatitis B Virus Polymerase Is Activated by Its RNA Packaging Signal, Epsilon. *J Virol* 72: 5789–5796.
 106. Edward JT, Gauthier M, Chubb FL, Ponka P (1988) Synthesis of new acylhydrazones as iron-chelating compounds. *J Chem Eng Data* 33: 538–540.