

Full Length Research Paper

Expression of an expansin carbohydrate-binding module affects xylem and phloem formation

Olawole O. Obembe^{1,2}, Evert Jacobsen¹, Richard Visser¹ and Jean-Paul Vincken^{1*}

¹Graduate School Experimental Plant Sciences, Laboratory of Plant Breeding, Wageningen University P O Box 386, 6700 AJ, Wageningen, The Netherlands.

²Department of Biotechnology, The Bells University of Technology, Ota Nigeria.

Accepted 14 June, 2007

Expansins are believed to be involved in disrupting the non-covalent adhesion of cellulose to matrix polysaccharides, thereby promoting wall creep. We have targeted a putative potato expansin (EXPA) carbohydrate-binding module (CBM) to the cell walls of tobacco plants. Histological examinations and electron microscopy indicated that 30% of the xylem cells of the transgenic stems with high expression of the expansin CBM are wider (radial surface area) than those of the controls. Similarly, 37% of the xylem cells of the stems of the high expressers have thinner cell walls than those of the controls. There were no such phenotypes in the low and none expressers, as well as in the control plants. The transgenic tobacco plants expressing the potato expansin CBM did not exhibit marked change in plant morphology. Analysis of cellulose content in the stem cell walls was similar between the high expresser of the transgene and the control plants. Nonetheless, our results taken together demonstrate that expansin CBM alone can bring about changes in the plant cell walls.

Key words: Expansin, carbohydrate binding modules, cell wall modification, homology modelling, cell expansion, transgenic.

INTRODUCTION

Expansins are plant proteins discovered over a decade ago, which have the capacity to induce extensibility and stress relaxation of plant cell walls *in vitro* and are proposed to mediate cell expansion (McQueen-Mason et al., 1992; McQueen-Mason and Cosgrove, 1995). The proteins are thought to facilitate wall extension by catalyzing the disruption of hydrogen bonds between cellulose microfibrils and matrix polysaccharides, thereby causing turgor-driven slippage between the microfibrils and other components of the wall (McQueen-Mason and

Cosgrove, 1995). Expansins are expressed in growing tissues including hypocotyls, shoot meristems and internodes (Cho and Kende, 1997; Fleming et al., 1997; Reinhardt et al., 1998). They are also expressed in non-growing tissues such as ripening fruits and abscission zones or tissues where cell separation will take place (Brummell et al., 1999; Cosgrove et al., 1998). These findings suggest that the proteins have tissue-specific roles in addition to cell expansion (Chen and Bradford, 2000). Studies with transgenic plants indicate that endogenous expansins are involved in regulating growth and developmental processes in which rearrangements in the cell wall are thought to be important (Cho and Cosgrove, 2000; Choi et al., 2003; Pien et al., 2001; Zenoni et al., 2004).

Based on their phylogenetic relationship, expansins and expansin-like proteins can be classified into four families (website: <http://www.bio.psu.edu/expansins/arabidopsis.htm> and (Kende et al., 2004). The four protein families currently recognized are EXPA, EXPB, EXLA and EXLB. Members of the EXPA and EXPB families

*Corresponding author. E-mail: jean-paul.vincken@wur.nl. Tel: +31317483998 Fax: +31317483457.

Abbreviations: Cryo-SEM, cryo-scanning electron microscopy; CBM, carbohydrate-binding module; CaMV, cauliflower mosaic virus; PCR, polymerase chain reaction; MS, Murashige-Skoog; NAA, α -naphthalene acetic acid; BAP, 6-benzylaminopurine; rubisco, ribulose 1,5-biphosphate carboxylase/oxygenase.

linked to the ExPASy (Expert Protein Analysis System) proteomic server (<http://us.expasy.org/>), designed for detecting protein modular domains. The alignment was manually edited for structurally homologous (according to Risler's matrix) residues occurring in the CBM.

An alignment, based on Clustal W, of the amino acid sequence of the potato expansin CBM and the PhI P2, together with SWISS-MODEL Version 36.0003 program (available from <http://swissmodel.expasy.org/SWISS-MODEL.html>) (Guex and Peitsch 1997; Peitsch, 1995; Schwede et al., 2003) were used for homology modelling, in which the atomic coordinates of PhI P2 (1WHO) served as a template.

Tobacco transformation, regeneration and growth

The expansin CBM construct and the empty vector control construct were used for *Agrobacterium tumefaciens*-mediated transformation of *in vitro* leaf explants of *Nicotiana tabacum* cv. Samsun NN as follows. Cloned binary vector pGreen7k was co-transformed with the helper plasmid, pSoup (Hellens et al., 2000) into *A. tumefaciens* strain LBA4404 by electroporation. This was plated out on LB-agar plates containing kanamycin (100 µg/mL) and rifampicin (30 µg/mL), and incubated for three days at 28°C, to obtain single colonies. The integrity of the binary plasmids was tested by restriction analysis of plasmids isolated from *A. tumefaciens* cultures used for plant transformation. 20 mL LB medium without selection was inoculated with a single colony and incubated overnight at 28°C. 100 µL of the grown culture was added to a petri dish containing 10 mL Murashige-Skoog (MS30). Leaf explants, without major veins and edges, from young seedlings were transferred upside down in to the MS30 medium. *Agrobacterium* infection was done in the dark for 2 days at 24 - 25°C. Two controls were used for the procedure, untransformed wild type control and empty pGreen7k control without an insert. Leaf explants were washed in three changes of liquid washing medium of MS30 with 250 mg/L carbenicillin. The washed explants were transferred upside-up to MS30-phytagel plates containing 0.1 mg/L α -naphthalene acetic acid (NAA), 1 mg/L 6-benzylaminopurine (BAP), 200 mg/L kanamycin and 250 mg/L carbenicillin, and incubated overnight in the dark at 28°C. Plants were transferred to a growth chamber (25°C), where they were gradually adapted to light and incubated for callus induction. After two weeks in culture, calli generated were transferred to shoot-inducing medium, MS20-phytagel, containing 0.2 mg/L NAA, 2 mg/L BAP, 200 mg/L kanamycin and 250 mg/L carbenicillin. Well-formed shoots were harvested and transferred to root-inducing medium, MS15-phytagel plate containing 100 mg/L kanamycin, 250 mg/L carbenicillin and 200 mg/L vancomycin. Twelve transformed plantlets were transferred to the greenhouse to generate mature plants.

Semi-quantitative RT-PCR analysis of transgenic plants

Total RNA was isolated from 3 - 5 g of transformed *in vitro* shoots as described elsewhere (Kuipers et al., 1995). For first strand cDNA synthesis, 1 µg of total RNA was treated with 0.5 µL DNase I RNase free (10 U/µL; Invitrogen) and incubated with 5 µL of 10 x Taqman RT buffer, 11 µL of 25 mM MgCl₂, 10 µL of 10 mM dNTP mix, 2.5 µL of 50 µM random hexamer primers, 1 µL RNase inhibitor (20 U/µL) and H₂O until a final volume of 50 µL for 30 min at 37°C and 5 min at 75°C. The mixture was then incubated for 10 min at 25°C and 30 min at 48°C with 1 µL of MultiScribe reverse transcriptase (50 U/µL; Applied Biosystems). The reaction was then terminated by heating the sample for 5 min at 95°C. 5 µL of cDNA was then used as a template in standard PCR using construct-specific primers. The forward primer, SPFOR 5'-agcttatgggaagtggta-3', was specific for the signal peptide while the reverse primer, HISREV 5'-tccgtgatggtgatggt- 3', was specific for

the hexa-histidine tag. Rubisco-specific primers, RUBFOR 5'-cagaatcatcaggaaaggaaaca - 3' and RUBREV 5'-tctctcacgtttctgctcttg - 3' were used to amplify rubisco cDNA, as internal control.

Light microscopy

Three individual plants from the high expressing and two plants from the low/none expressing class were used for microscopic examination.

The empty pGreen7k transformants were used as the control. Stem samples were taken from the second internode from the top of the plant. 1 mm-thick stem sections were fixed in 3% glutaraldehyde (Merck) and 3% paraformaldehyde (Merck) in 0.1 M phosphate buffer containing 0.1% Triton x100 for 2 h. The samples were then washed and dehydrated in an ethanol series. After dehydration, they were embedded in Technovit 7100 resin (USA) (Kuroiwa et al., 1990). 4 µm-thick sections were stained with 0.1% toluidine blue (Aldrich) and examined under a bright field microscope.

Cryo-scanning electron microscopy

Transgenic line 16 and one individual plant from the empty vector control were used for cryo-scanning electron microscopy (cryo-SEM) examination. Stem sampling was the same as for the light microscopy. 6 mm-thick stem samples were mounted in a brass cylindrical sample holder with TBS (Tissue Freezing Medium EMS, Washington, PA, USA). The frozen samples were placed in a sample holder in a cryo-ultra microtome (Reichert Ultracut E/FC4D) and cut at a specimen temperature of 100°C. These samples were first planed with a glass knife, after which the surface was planed with a diamond knife (Histo no trough, 8 mm 45°C, Drukker International, The Netherlands). After planing, the samples were placed in a dedicated cryo-preparation chamber (CT 1500 HF, Oxford instruments, UK). All the samples in the cryo-preparation chamber were freeze dried for 3 min at -90°C and 8 x 10⁻⁴ Pa, and subsequently sputtered with a layer of 10 nm Pt. The samples were cryo-transferred into the field emission scanning microscope (JEOL 6300F, Japan) on the sample stage at -190°C. All images were recorded digitally (Orion, 6 E.L.I. sprl, Belgium) at a scan rate of 100 s (full frame) at the size of 2528 x 2030, 8 bit. The images were optimized and resized for publication by Adobe Photoshop CS.

Quantification of cell size and cell wall diameter

To quantify the micrographic images, files were opened with Image J software developed at the National Institute of Health, USA (<http://rsb.info.nih.gov/ij/>). The surface area of three hundred xylem cells was measured as shown by the inset in Figure 6A. One hundred cells per field of view were measured in three replicates. Similarly, cell wall thickness of one hundred and fifty cells was quantified as shown by the inset in Figure 6B. Fifty cells per field of view were measured in three replicates. Standard deviation of three replicates was determined.

Isolation of cell wall material and analysis of the cellulose content

Stem samples from transgenic line 16, pGreen7k vector control and a wild type control plant were separately ground to a fine powder in liquid nitrogen. For each isolation, 1 g of this stem material was extracted in a 50 mM Tris[HCl] pH 7.2 solution containing 1% SDS, acetone and air-dried. 10 mg of the different cell wall materials was




Figure 1. Amino acid sequence alignment of At-EXPA1, At-EXPB1, At-EXLA1, At-EXLB1 (all four from *Arabidopsis thaliana*), Phl p2 (pollen allergen from *Phleum pratense*) and the putative potato expansin CBM (EST accession number BG097738). Conserved aromatic residues are shaded and shown in bold letters. Secondary structure elements of Phl P2 and of the putative potato cellulose-binding domain are indicated with letter 'S' and 'c' above the sequences. Asterisk denotes exposed aromatic residues and histidine on the modelled structure. Underlining of secondary structure elements indicates that this region does not have good match with the 3D structure of the template used for modeling.

hydrolyzed in 1 ml 2 M TFA. The TFA-insoluble cellulose was spun down at 13000 rpm for 15 min, and pellet was dissolved in 67% for 3 h at room temperature (RT) with continuous shaking. The cell wall material (CWM) was spun down by centrifugation at 13000 rpm for 15 min.

Subsequently, the residue was washed with water, ethanol and Sulphuric acid. The suspension was heated and diluted appropriately to determine the cellulose content colorimetrically using anthrone as a colouring agent according to Updegraff (1969). The acid hydrolysis was performed in quadruplicate

RESULTS

Characteristics of the putative potato expansin CBM

The size of a complete expansin protein is about 25 - 28 kDa. The potato CBM that was targeted to tobacco cell wall was about 15 kDa (133 amino acids) in size, including the transit peptide and the hexa-histidine epitope tag.

A blast search with the amino acid sequence of the EST clone of the potato expansin CBM revealed that it has high homology with the *Arabidopsis* expansin 18 (e-value, $2e-50$), which belongs to the EXPA family. We were faced with the question of which part of the sequence should be used for engineering a construct expressing a functional domain II of the potato expansin.

Meanwhile, the putative CBM of an EXPB family Lol P2 from rye grass was found to have about 50% sequence identity with a pollen allergen, Phl P2 from timothy grass (Cosgrove, 2000b), whose structure was solved (De Marino et al., 1999). An amino acid alignment with the potato expansin EST clone and Phl P2 was made, in order to deduce the start of the potato expansin CBM more precisely. The 3-D structure of Phl P2 starts with

Val27 (Val1 in the alignment presented in Figure 1), which is five amino acids before the conserved Phe6 (underlined). We deduced an amino acid sequence (96 amino acids) for a potentially functional putative CBM of the potato expansin, starting with Gly1, four positions before the conserved Phe5 (underlined). We particularly singled out Gly1 as the starting point because it is directly preceded by an exposed two-basic residue RR sequence (data not shown), which is a conserved cleavage signal between domain I and domain II of expansins (Barre and Rouge, 2002). An amino acid sequence alignment of the potato expansin CBM, with four *Arabidopsis* expansin CBMs (representing each of the four classes of the expansins family), and a pollen allergen Phl P2 from Timothy grass, was made (Figure 1). The amino acid sequences of the four representatives above were obtained from a web site (<http://www.bio.psu.edu/expansins/arabidopsis.htm>), and correspond to the domain II or CBM as determined by the ExpASy proteomic server (<http://us.expasy.org/>). These four sequences are shorter than the sequence we deduced for potato CBM, which explains the gaps left in the alignment at the beginning of these sequences. We have incorporated the extra GGIRFTINGFR sequence at the N-terminal side to reduce the risk of an incorrectly folding domain, when expressing the construct in tobacco. The alignment information revealed that the potato CBM shared varying homologies with the CBMs of the four *Arabidopsis* expansin families; 64% identity with the CBM of EXPA1, 24% with EXPB1, 28% with EXLA1 and 26% with EXLB1. Furthermore, the alignment result revealed that aromatic amino acids (tryptophan and tyrosine) are mostly conserved in the different sequences




Figure 2. Putative 3-D structure of potato expansin CBM, obtained by homology modelling, using the 3-D structure of pollen allergen Phl P2 as template. Conserved aromatic residues, which are exposed and might be involved in polysaccharide binding, are: Trp39, Trp46, Trp76, Trp83, Tyr12 and His82. Trp50 is mostly buried, hence it is indicated with light colour. Side chains of none conserved residues His82 and Trp76 are also indicated in light colour. Conserved residues are presented in dark colour.

particularly among the four expansins families. Our particular interest in the aromatic residues was premised on their involvement in carbohydrate binding (Toone, 1994) and reviewed in (Boraston et al., 2004).

We have modelled a putative 3D-structure for the potato expansin CBM by homology modelling using the structure of Phl P2 as a template. This was with a view to revealing the location of its conserved aromatic residues. The modelled 3-D structure of the potato expansin CBM consisted of β strands and coils (Figure 2). Accompanied information on the modelled CBM revealed that the putative structure had 85% identity to the template Phl P2. The identical regions include all regions containing the eight β strands (indicated in Figure 1 with letter 'S'). Only four coil-constituting regions of the model are structurally divergent to Phl P2 as indicated in Figure 1 by underlining of the secondary structure elements. All aromatic amino acid residues (tyrosine, tryptophan) of the sequence except one are exposed. The exposed histidine residue was also visualized because of its ability to mediate pH-dependent binding of proteins to carbohydrates (Linder et al., 1999). One side of the modelled CBM revealed a patch of three exposed residues, Tyr12, His82, and Trp83. It should be noted from the alignment result that the His82 is not conserved, which raises the question as to what extent it is involved in binding. At the opposite side of the protein is a straight line of exposed aromatic residues Trp39, Trp46, and Trp76. It should be noted as well that, even though Trp76 is not conserved




Figure 3. Semi quantitative RT-PCR analysis of the selected transformants and the wild type leaf RNA. The upper panel shows the product amplified using the hexa-histidine tag-specific forward primer and pGreen7k construct-specific reverse primer. The lower panel shows the product amplified using specific primers for ribulose 1,5-biphosphate carboxylase/oxygenase (rubisco), as internal control. Plant lines 5, 7, and 17 are classified as low/none expresser, because of low or none visible intensities. The rest are classified as high expressers. EVC = empty vector negative control, PL = CBM-containing plasmid, as positive control.

across the four families, it is quite conserved within the EXPA family (data not shown). The well-conserved Trp50 is mostly buried, and therefore its involvement in binding is not very likely. It may have a role in maintaining protein structure through hydrophobic interactions.

Characterization of tobacco transformants expressing the putative potato expansin CBM

Twelve antibiotic-resistant tobacco transformants carrying the expansin CBM transgene were regenerated and transferred to the green house. RNA was isolated from leaves of the transgenic plants to analyze the expression level of the expansin CBM in each transgenic line by semi-quantitative RT-PCR analysis, using the constitutively expressed rubisco gene as an internal control. Varying amplification intensities of the different transgenic plants revealed differential expression of the potato expansin CBM gene in the individual tobacco transformants, thus making it possible to categorize them into two classes, as high and low/none expressers (Figure 3). Transgenic lines 5, 7 and 17 represent the low/none expressing transformants, while the remaining nine lines represent the high expressers. Preliminary attempts at purifying CBM proteins using affinity purification with the hexa-histidine tag were not successful (data not shown). Similarly, western detection of a hexa-histidine epitope-tagged fungal elicitor protein ECP2 that was infiltrated into the apoplast of leaves did not succeed in the tomato plants but succeeded in *Arabidopsis* (van Esse et al., 2006)

The results indicated that the histidine tag was cleaved off in the tomato plants, but not in *Arabidopsis*. These observations have now led to the idea that the hexa-histidine epitope tags have cleavage sites that are being recognized by proteases, which are specific for the cell walls of the solanaceous species. There was no obvious phenotype in the transgenic plants when compared to the pGreen7k vector control. The average plant height of the transgenic plants expressing the potato expansin CBM




Figure 4. Cross sections of representative transgenic and control stems. Sections of empty Pgreen7k control (A) and transgenic line 16, high expresser of potato expansin CBM (B) were stained with toluidine blue. A and B show the vascular tissue, phloem fibre and xylem cells of the stem sections of the control and the expansin CBM high expresser, respectively. Pf = phloem fibre, xy = xylem.




Figure 5. Scanning electron micrographs of stem vascular tissue of the empty Pgreen7k control (A) and the transgenic tobacco line 16 (B), at x300 magnification. Micrographs are representatives of several fields of view.

gene at maturity was 70 cm, which was comparable to 75 cm average height for the pGreen7k vector control plants. There was no particular trend in plant development with respect to stem elongation and flower development.

Transgenic stems exhibit enlarged xylem cells and thin cell walls

To examine cellular events resulting from the expression of the potato expansin CBM in the tobacco plant, stem sections from three representatives from the high expressers and two from the low/none expressers were stained with toluidine blue and examined by light microscope. Figure 4 reveals relatively larger xylem and phloem fibre cells in a representative high expresser of the potato CBM (Figure 4B) than those of the Pgreen7k control (Figure 4A). Furthermore, the micrographs indicated that the cell walls of the phloem fibres of the transgenic plants were thinner than those of the control. Other cell types of

the transgenic stems were the same as those of the control. No intermediate effect was observed with the two low/none expressers, as they were comparable with the control plants.

In order to quantify the cellular observations made with the light microscopy, cryo-SEM was done with the stem samples. The micrographs confirmed that the xylem cells of the transgenic stem of line 16 (Figure 5B) were larger than those of the control stems (Figure 5A). To validate our SEM observations, we subjected the images to quantitative measurements. Figure 6A shows that 40% of the xylem cells in the pGreen7k control stem were grouped in the small class ($0 - 30 \mu\text{m}^2$), whereas only 10% of the xylem cells of the transgenic stems grouped in this class. It also shows that 25% of the xylem cells in the transgenic stems grouped in the large class ($60 - 90 \mu\text{m}^2$), whereas only 5% of the cells of the pGreen7k control stems grouped in this class. There was no large difference between transgenic and the control plants, with respect to the percentage of xylem cells that were group-




Figure 6. Cell size distribution (A) and cell wall thickness (B) of xylem cells in the stems of transgenic tobacco plants. The surface area of three hundred xylem cells was measured, as shown by the inset in Fig. 6A. One hundred cells per field of view were measured in three replicates. Similarly, cell wall thickness of one hundred and fifty cells was quantified, as shown by the inset in Fig. 6B. Fifty cells per field of view were measured in three replicates. Data were expressed as percentages of xylem cells and cell walls in the different size classes. Open bars represent pGreen7k empty control. Grey bars represent expansin CBM transgenic stems.

ed in the medium class (63 and 54%, respectively).

We thus can infer that the stem xylem cells of the potato expansin CBM-expressing plants were larger than those of the pGreen7k control plants. Similarly, for cell wall quantification, we measured the thickness of the cell walls of the xylem cells of the transgenic and the pGreen7k control plants as presented in Figure 6B. We observed that 32% of the cell walls of the stem xylem cells of the transgenic plants expressing the expansin CBM grouped in the thin class (1.0 – 2.0 μm), whereas 0% of the cell walls of the pGreen7k control stem xylem cells grouped in this class. Figure 6B also shows that only 5% of the cell walls of the stem xylem cells of the transgenic plant grouped in the thick classes (3.0 – 4.0 μm), whereas 32% of the cell walls of the pGreen7k control stem xylem cells grouped in same class. Similarly as for the cell size, there was no difference between the transgenic and the control, with respect to the distribution of cell wall thickness in the medium class (2.0 – 3.0 μm).

We can also infer from this result that the thickness of the cell walls of the stem xylem cells of the transgenic tobacco plants expressing the potato expansin CBM are thinner than those of the pGreen7k control plants.




Figure 7. Cellulose content in transgenic and control tobacco plants. Cellulose content of stem wall material of transgenic line 16 was determined using a colorimetric assay. The cellulose levels are represented as the percentage of cellulose in total cell wall material. All measurements were performed as quadruplicate and standard errors of the means are represented by the error bars

Analysis of cellulose content of transgenic stems

It has been suggested that expansins may play a role in the assembly of the cell wall by affecting cellulose synthesis or deposition (Zenoni et al., 2004). We have determined the cellulose content of the stem cell walls of the expansin CBM-expressing plant and the control plants. Colorimetric assays were performed on four independently hydrolyzed cell wall samples from transgenic tobacco line 16, the empty vector control, and a wild type control (Figure 7). It seems that the cellulose content of the transgenic plant is lower than that of the control plants, but given the large variation in the measurements (due to heterogeneity of the fibrous cell wall material of the stem), the differences are not significant.

DISCUSSION

We have over-expressed a putative CBM of a potato expansin in tobacco, to investigate whether expansin's domain II can bring about changes in the cell wall on its own. Results of the investigation are discussed with respect to its binding specificity and its ability to alter cell wall structure.

Homology modelling of the 3-D structure of the C-terminal domain II of the expansins has been possible with the resolution of the 3-D structure of PhI P2 (De Marino et al., 1999). The modelled three-dimensional structure of the putative potato expansin CBM showed that most of its conserved aromats were located on the surface. The patch of three "aromatic" residues on one side and the linear strip of another three aromatic residues on the opposite side of the protein are consistent with the 3-D structure of CBMIII (CBM of the cellulosome subunit S1 from *Clostridium thermocellum* (1NBC) as resolved by Tormo et al. (1996), which showed a groo-

ve and a planar cleft at opposite sides of the protein. They proposed that the planar cleft binds crystalline cellulose while the groove binds the amorphous single cellulose chain. Shoseyov et al. (2006) speculated further that the parallel arrangement of the binding grooves would enable the protein to slide between two cellulose chains and facilitate the disruption of the hydrogen bonding between adjacent chains in a wedge-like action. The structure of the C-terminal domain II of Lol PI (EXPB) had been modelled by others using the Phl P2 as template (Barre and Rouge, 2002; Cosgrove, 2000a). That model also revealed that most of the conserved aromatic residues were exposed around a linear strip and a groove (Barre and Rouge, 2002).

Even though our *in planta* characterization of the putative CBM is not conclusive, with respect to its binding specificity, it is interesting to note that the effect of over-expression of the protein was mainly observed in the cellulose- and xylan-rich vascular tissue. We observed an abnormal enlargement (radial expansion) of cells of the xylem and phloem fibre cells in the high expressing transgenic tobacco plants. These observations are consistent in the three independent high-expressing lines examined, suggesting that they are related to the expression of the transgene rather than to other factors. Apart from cell enlargement, the cell walls of the vascular tissue were observed to be thinner in the transgenics as compared to the control plants. In contrast to our observations, which were restricted to the vascular tissue, a heterologous expression of high levels of a recombinant cucumber expansin in transgenic tomato was reported to cause altered expansion of the cortical and epidermal cells of the stems (Rochange et al., 2001). These localised cellular events in both cases seem to support the view that the expansin proteins have tissue-specific roles in addition to cell expansion (Brummell et al., 1999; Chen and Bradford, 2000; Cosgrove et al., 1998). It should be noted that the EST clone that was used for this investigation was obtained from an EST library of potato petioles and leaves. These matured organs are rich in secondary walled vascular tissue. The expansin CBM that was targeted to the tobacco cell walls may have preferentially interacted with the secondary wall of the vascular tissue. This interaction might disrupt cellulose-hemicellulose networks, culminating in wall thinning, which in turn might lead to wall weakening and hence abnormal cell enlargement (Kutschera, 1990). The foregoing might explain the more obvious phenotypes in the vascular tissue than in any other tissue of the stem of the transgenic tobacco plants.

In spite of the cellular phenotypes, transgenic tobacco plants exhibit normal morphology and development. However, in contrast to our observation of normal growth phenotype of the transgenic plants, high levels of over-expression or heterologous expression of expansin genes in transgenic plants was shown to lead to impaired growth phenotypes (Choi et al., 2003). A sense expres-

sion of AtEXP10 was also shown to enhance leaf growth in the transgenic *Arabidopsis* plants (Cho and Cosgrove, 2000). Similarly, local expansin expression within the meristem was shown to induce the process of leaf formation (Pien et al., 2001). It is worth noting, however, that all these reports had used the complete expansin protein, while only the CBM has been used in this work. This suggests that both domains of expansins are needed to bring about developmental changes in the plant.

ACKNOWLEDGEMENT

This work was supported by the Netherlands Foundation for the Advancement of Tropical Research (WOTRO) The Netherlands.

REFERENCES

- Barre A, Rouge P (2002). Homology modeling of the cellulose-binding domain of a pollen allergen from rye grass: structural basis for the cellulose recognition and associated allergenic properties. *Biochem. Biophys. Res. Commun.* 296: 1346-1351.
- Boraston AB, Bolam DN, Gilbert HJ, Davies GJ (2004). Carbohydrate-binding modules: fine-tuning polysaccharide recognition. *Biochem. J.* 382: 769-781.
- Brummell DA, Harpster MH, Dunsmuir P (1999). Differential expression of expansin gene family members during growth and ripening of tomato fruit. *Plant Mol. Biol.* 39: 161-169.
- Ceccardi TL, Barthe GA, Derrick KS (1998). A novel protein associated with citrus blight has sequence similarities to expansin. *Plant Mol. Biol.* 38: 775-783.
- Chen F, Bradford KJ (2000). Expression of an expansin is associated with endosperm weakening during tomato seed germination. *Plant Physiol.* 124: 1265-1274.
- Cho HT, Cosgrove DJ (2000). Altered expression of expansin modulates leaf growth and pedicel abscission in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA.* 97: 9783-9788.
- Choi DS, Lee Y, Cho HT, Kende H (2003). Regulation of expansin gene expression affects growth and development in transgenic rice plants. *Plant Cell* 15: 1386-1398.
- Cosgrove DJ (2000a). Loosening of plant cell walls by expansins. *Nature* 407:321-326.
- Cosgrove DJ (2000b). New genes and new biological roles for expansins. *Curr. Opin. Plant Biol.* 3: 73-78.
- Cosgrove DJ, Scherban TY, Durachko DM (1998). Highly specific and distinct expression patterns for two α -expansin genes in *Arabidopsis*. *Plant Physiol. Suppl.* A1(75): 57.
- De Marino S, Morelli MAC, Fraternali F, Tamborini E, Musco G, Vrtala S, Dolecek C, Arosio P, Valenta R, Pastore A (1999). An immunoglobulin-like fold in a major plant allergen: the solution structure of Phl p 2 from timothy grass pollen. *Structure* 7: 943-952.
- Gehring CA, Irving HR (2003). Natriuretic peptides - a class of heterologous molecules in plants. *Int. J. Biochem. Cell Biol.* 35: 1318-1322.
- Gilkes NR, Henrissat B, Kilburn DG, Miller RC, Warren RAJ (1991). Domains in microbial β -1,4-glycanases: sequence conservation, function and enzymes families. *Microbiol. Rev.* 55: 303-315.
- Guex N, Peitsch MC (1997). SWISS-MODEL and the Swiss-PdbViewer: An environment for comparative protein modelling. *Electrophoresis* 18: 2714-2723.
- Hellens RP, Edwards EA, Leyland NR, Bean S, Mullineaux PM (2000). pGreen: a versatile and flexible binary Ti vector for *Agrobacterium*-mediated plant transformation. *Plant Mol. Biol.* 42: 819-832.
- Kende H, Bradford KJ, Brummell DA, Cho HT, Cosgrove DJ, Fleming AJ, Gehring C, Lee Y, McQueen-Mason S, Rose JKC, Voeselek L

- (2004). Nomenclature for members of the expansin superfamily of genes and proteins. *Plant Mol. Biol.* 55: 311-314.
- Kuipers AGJ, Soppe WJJ, Jacobsen E, Visser RGF (1995). Factors affecting the inhibition by antisense RNA of granule-bound starch synthase gene-expression in potato. *Mol. Gen. Genet.* 246: 745-755.
- Kuroiwa T, Kuroiwa H, Mita T, Fujie M (1990). Fluorescence Microscopic Study of the Formation of Giant Mitochondrial Nuclei in the Young Ovules of *Pelargonium-Zonale*. *Protoplasma* 158: 191-194.
- Kutschera U (1990). Cell-wall synthesis and elongation growth in hypocotyls of *Helianthus-annuus* L. *Planta* 181: 316-323.
- Li Y, Jones L, McQueen-Mason S (2003). Expansins and cell growth. *Curr. Opin. Plant Biol.* 6: 603-610.
- Linder M, Nevanen T, Teeri TT (1999). Design of a pH-dependent cellulose-binding domain. *FEBS Lett.* 447: 13-16.
- Ludidi NN, Heazlewood JL, Seoighe C, Irving HR, Gehring CA (2002). Expansin-like molecules: Novel functions derived from common domains. *J. Mol. Evol.* 54: 587-594.
- Maryani MM, Bradley G, Cahill DM, Gehring CA (2001). Natriuretic peptides and immunoreactants modify osmoticum-dependent volume changes in *Solanum tuberosum* L. mesophyll cell protoplasts. *Plant Sci.* 161: 443-452.
- McQueen-Mason SJ, Cosgrove DJ (1994). Disruption of hydrogen-bonding between plant-cell wall polymers by proteins that induce wall extension. *Proc. Natl. Acad. Sci. USA.* 91: 6574-6578.
- McQueen-Mason SJ, Cosgrove DJ (1995). Expansin mode of action on cell-walls- Analysis of wall hydrolysis, stress-relaxation, and binding. *Plant Physiol.* 107: 87-100.
- McQueen-Mason SJ, Durachko DM, Cosgrove DJ (1992). Two endogenous proteins that induce cell-wall extension in plants. *Plant Cell* 4: 1425-1433. Peitsch MC (1995). *Protein Modeling By E-Mail*. *Biotechnology (N. Y.)* 13: 658-660.
- Pharmawati M, Maryani MM, Nikolakopoulos T, Gehring CA, Irving HR (2001). Cyclic GMP modulates stomatal opening induced by natriuretic peptides and immunoreactive analogues. *Plant Physiol. Biochem.* 39: 385-394.
- Pien S, Wyrzykowska J, McQueen-Mason S, Smart C, Fleming A (2001). Local expression of expansin induces the entire process of leaf development and modifies leaf shape. *Proc. Natl. Acad. Sci. USA.* 98: 11812-11817.
- Rochange SF, Wenzel CL, McQueen-Mason SJ (2001). Impaired growth in transgenic plants over-expressing an expansin isoform. *Plant Mol. Biol.* 46: 581-589.
- Schwede T, Kopp J, Guex N, Peitsch MC (2003). SWISS-MODEL: an automated protein homology-modeling server. *Nucleic Acids Res.* 31: 3381-3385.
- Shcherban TY, Shi J, Durachko DM, Gultinan MJ, McQueen-Mason SJ, Shieh M, Cosgrove DJ (1995). Molecular-cloning and sequence-analysis of expansins - a highly conserved, multigene family of Proteins that mediate cell-wall extension in plants. *Proc. Natl. Acad. Sci. USA.* 92: 9245-9249.
- Shoseyov O, Shani Z, Levy I (2006). Carbohydrate binding Modules: Biochemical properties and novel applications. *Microbiol. Mol. Biol. Rev.* 70: 283-295.
- Toone EJ (1994). Structure and energetics of protein carbohydrate complexes. *Curr. Opin. Struct. Biol.* 4: 719-728.
- Tormo J, Lamed R, Chirino AJ, Morag E, Bayer EA, Shoham Y, Steitz TA (1996). Crystal structure of a bacterial family-III cellulose-binding domain: a general mechanism for the attachment to cellulose. *EMBO J.* 15: 5739-5751.
- Updegraff DM (1969). Semi-microdetermination of cellulose in biological materials. *Anal. Biochem.* 32: 420-424.
- Van Esse HP, Thomma BPHJ, Van T, Klooster JW, Wit PJGM (2006). Affinity-tags are removed from *Cladosporium fulvum* effector proteins expressed in the tomato leaf apoplast. *J. Exp. Bot.* 57: 599-608.
- Whitney SEC, Gidley MJ, McQueen-Mason SJ (2000). Probing expansin action using cellulose/hemicellulose composites. *Plant J.* 22: 327-334.
- Wittink FRA, Knuiman B, Derksen J, Capkova V, Twell D, Schrauwen JAM, Wullems GJ (2000). The pollen-specific gene Ntp303 encodes a 69-kDa glycoprotein associated with the vegetative membranes and the cell wall. *Sex Plant Reprod.* 12: 276-284.
- Zenoni S, Reale L, Torielli GB, Lanfaloni L, Porceddu A, Ferrarini A, Moretti C, Zamboni A, Speghini S, Ferranti F, Pezzotti M (2004). Downregulation of the *Petunia hybrida* alpha-expansin gene PhEXP1 reduces the amount of crystalline cellulose in cell walls and leads to phenotypic changes in petal limbs. *Plant Cell* 16: 295-308.