

Acta Medica Okayama

Volume 50, Issue 4

1996

Article 6

AUGUST 1996

Pharmacokinetic Evaluation of Omeprazole Suspension Following Oral Administration in Rats: Effect of Neutralization of Gastric Acid

Kazuhide Watanabe*

Naoyuki Matsuka[†]

Katushi Furuno[‡]

Kohei Eto**

Hiromu Kawasaki^{††}

Yutaka Gomita^{‡‡}

*Okayama University,

[†]Okayama University,

[‡]Okayama University,

**Okayama University,

^{††}Okayama University, kawasaki@pheasant.pharm.okayama-u.ac.jp

^{‡‡}Okayama University,

Pharmacokinetic Evaluation of Omeprazole Suspension Following Oral Administration in Rats: Effect of Neutralization of Gastric Acid*

Kazuhide Watanabe, Naoyuki Matsuka, Katushi Furuno, Kohei Eto, Hiromu Kawasaki, and Yutaka Gomita

Abstract

In order to evaluate a clinical use of omeprazole suspension, we examined the pharmacokinetics of omeprazole after oral administration in rats. Although the administration of omeprazole suspension buffered by NaHCO₃ solution did not produce a significant increase in the area under the concentration-time curve (AUC) value compared with non-buffered group, the administration of NaHCO₃ buffer immediately after dosing of omeprazole suspension buffered by NaHCO₃ caused a significant increase in the AUC value. These results suggest that the NaHCO₃ treatment following the administration of omeprazole buffered suspension effectively decreased the degradation of the compound by gastric acid. Therefore, the successive administration of NaHCO₃ solution after the omeprazole dosing seems to be a simple and useful method for the administration to patients who cannot receive tablets.

KEYWORDS: omeprazole, suspension, pharmacokinetics, rats

Brief Note

Pharmacokinetic Evaluation of Omeprazole Suspension Following Oral Administration in Rats: Effect of Neutralization of Gastric Acid

Kazuhide WATANABE, Naoyuki MATSUKA, Katsushi FURUNO, Kohei ETO^a, Hiromu KAWASAKI and Yutaka GOMITA*

Department of Hospital Pharmacy, Okayama University Medical School and ^aOkayama University Dental School, Okayama 700, Japan

In order to evaluate a clinical use of omeprazole suspension, we examined the pharmacokinetics of omeprazole after oral administration in rats. Although the administration of omeprazole suspension buffered by NaHCO₃ solution did not produce a significant increase in the area under the concentration-time curve (AUC) value compared with non-buffered group, the administration of NaHCO₃ buffer immediately after dosing of omeprazole suspension buffered by NaHCO₃ caused a significant increase in the AUC value. These results suggest that the NaHCO₃ treatment following the administration of omeprazole buffered suspension effectively decreased the degradation of the compound by gastric acid. Therefore, the successive administration of NaHCO₃ solution after the omeprazole dosing seems to be a simple and useful method for the administration to patients who cannot receive tablets.

Key words: omeprazole, suspension, pharmacokinetics, rats

Omeprazole is a substituted benzimidazole that effectively suppresses gastric acid secretion by inhibiting H⁺/K⁺-ATPase in the parietal cell (1, 2). This inhibition is long-lasting, although omeprazole is rapidly eliminated from plasma (3). Thus, the time-dependent antisecretory effect of omeprazole is not correlated with the plasma concentration (3). However, the area under the plasma concentration-time curve (AUC) has been reported to be correlated to the degree of the inhibition (4).

Since omeprazole is unstable in acidic conditions (5), the compound is administered orally to patients as an enteric-coated tablet that is the only formulation on the market in Japan. For patients who have difficulty with tablets, omeprazole is given in the powder or suppository form. Recently, we reported that, in rats, omeprazole administered intrarectally undergoes metabolism by gut flora, and that about 60 % of the dose is degraded before absorption from the intestinal tract can occur (6). Thus, the powder or suspension form of omeprazole is preferred for these patients. However, the pharmacokinetics of the absorption of omeprazole suspension has not been systematically investigated. In the present study, therefore, we examine the pharmacokinetics of omeprazole suspension after oral administration in rats.

Male Wistar rats (Charles River Lab., Atsugi, Japan) each weighing 200–320 g were housed in the experimental animal center of Okayama University Medical School at a controlled ambient temperature of 22 ± 1 °C with approximately 60 % relative humidity and with a 12-h light/12-h dark cycle (lights on at 07:00). Animals were allowed free access to food and water, except for a 12-h fast before experiments. Omeprazole (donated by Fujisawa-Astra Ltd. Osaka, Japan) was suspended in 0.5 % methylcellulose, and administered orally at a dose of 20 mg/kg. The animals were divided into six groups as follows: (a) non-buffered group receiving omeprazole suspension which was not buffered with NaHCO₃; (b) 0.2 % NaHCO₃ group receiving omeprazole suspension

* To whom correspondence should be addressed.

^aPresent address: Pharmaceutical Analysis Center, Okayama Pharmacist Association, Okayama 700, Japan.

Table 1 Method for oral administration of omeprazole suspension in rats

Groups	Omeprazole suspension ^a		Treatment
	Buffering	pH Adjustment ^b	
Non-buffered	Not contained	Not adjusted	—
0.2% NaHCO ₃	0.2% NaHCO ₃	pH 9	—
2.5% NaHCO ₃	2.5% NaHCO ₃	pH 9	—
H ₂ O post-dosed	2.5% NaHCO ₃	pH 9	Oral administration of H ₂ O (3 ml/kg) immediately after omeprazole dosing
NaHCO ₃ pre-dosed	2.5% NaHCO ₃	pH 9	Oral administration of 2.5% NaHCO ₃ (3 ml/kg) 5 min before omeprazole dosing
NaHCO ₃ post-dosed	2.5% NaHCO ₃	pH 9	Oral administration of 2.5% NaHCO ₃ (3 ml/kg) immediately after omeprazole dosing

a: Omeprazole was suspended in 0.5% methylcellulose and administered at 20 mg/kg in a volume of 1 ml/kg.

b: pH of omeprazole suspension was adjusted by 1 N NaOH.




Fig. 1 Time course of plasma omeprazole concentration in rats. Omeprazole suspension was administered orally at a dose of 20 mg/kg. Each point indicates the mean \pm SEM. A: (\circ), non-buffered group; (\bullet), 0.2% NaHCO₃ group; (\square), 2.5% NaHCO₃ group. B: (\blacksquare), H₂O post-dosed group; (\circ), NaHCO₃ pre-dosed group; (\blacktriangle), NaHCO₃ post-dosed group.

containing 0.2% NaHCO₃ (pH 9); (c) 2.5% NaHCO₃ group receiving omeprazole suspension containing 2.5% NaHCO₃ (pH 9); (d) H₂O post-dosed group receiving H₂O (3 ml/kg) immediately after the administration of omeprazole suspension containing 2.5% NaHCO₃ (pH 9); (e) NaHCO₃ pre-dosed group receiving 2.5% NaHCO₃ solution (3 ml/kg) 5 min before the administration of omeprazole suspension containing 2.5% NaHCO₃ (pH 9); (f) NaHCO₃ post-dosed group receiving 2.5% NaHCO₃ solution (3 ml/kg) immediately after the administration of omeprazole suspension containing 2.5% NaHCO₃ (pH 9) (Table 1). At various times after the administration, blood samples of 60 μ l were collected

from the tail vein, and the plasma concentrations of omeprazole were determined by high performance liquid chromatography, as described previously (7).

Pharmacokinetic parameters were obtained from the plasma concentration-time data of omeprazole for each animal, using a personal computer program for nonlinear least squares regression analysis (MULTI, Nankodo, Tokyo, Japan) (8). The maximum plasma concentration (C_{max}) and the time to reach the maximum concentration (T_{max}) were estimated from these pharmacokinetic parameters. The AUC values were calculated for 0–3 h period by the trapezoidal method. The mean residence time (MRT) was calculated by the model-independent moment

Table 2 Pharmacokinetic parameters of omeprazole suspension after oral administration in rats

Parameter	non-buffered (n = 5)	0.2% NaHCO ₃ (n = 5)	2.5% NaHCO ₃ (n = 5)	H ₂ O post-dosed (n = 5)	NaHCO ₃ pre-dosed (n = 5)	NaHCO ₃ post-dosed (n = 5)
C _{max} (μg/ml)	0.40 ± 0.06	0.42 ± 0.11	0.56 ± 0.10	0.61 ± 0.18	1.18 ± 0.20 ^a	1.53 ± 0.26 ^{b,d,e}
T _{max} (h)	0.16 ± 0.02	0.24 ± 0.04	0.19 ± 0.02	0.14 ± 0.02	0.12 ± 0.03 ^c	0.21 ± 0.02
AUC (μg·h/ml)	0.24 ± 0.02	0.40 ± 0.09	0.48 ± 0.08	0.46 ± 0.09	0.70 ± 0.09 ^a	0.87 ± 0.14 ^{b,c,e}
MRT	0.47 ± 0.03	0.75 ± 0.08 ^b	0.86 ± 0.10 ^{a,e}	0.51 ± 0.09	0.68 ± 0.07 ^a	0.52 ± 0.03 ^c

Omeprazole suspension was administered orally at a dose of 20 mg/kg. Each value represents the mean ± SEM.

C_{max}: The maximum plasma concentration; T_{max}: The time to reach the maximum concentration; AUC: The area under the concentration-time curve; MRT: mean residence time.

^aP < 0.05 vs. non-buffered group. ^bP < 0.01 vs. non-buffered group. ^cP < 0.05 vs. 2.5% NaHCO₃ group.

^dP < 0.01 vs. 2.5% NaHCO₃ group. ^eP < 0.05 vs. H₂O post-dosed group. ^fP < 0.01 vs. H₂O post-dosed group.

analysis using a personal computer program (9). The results were statistically evaluated by analysis of variance followed by Dunnett's test.

Figure 1 shows the time-course of plasma omeprazole concentration after oral administration in each group. The plasma concentration of omeprazole in the non-buffered group increased rapidly, reached a maximum at about 10 min, and became undetectable by 1 h after the administration. Similar rapid increases in plasma concentrations were obtained in the other groups with the maximum being reached at about 7-14 min; however, plasma concentrations were detectable for up to 2 h. Table 2 summarizes the pharmacokinetic parameters of omeprazole suspension after oral administration. The AUC values in the 0.2% NaHCO₃ and 2.5% NaHCO₃ groups were higher than in the non-buffered group, but the difference was not statistically significant. The MRT values in the 0.2% NaHCO₃ and 2.5% NaHCO₃ groups were significantly higher than in the non-buffered group ($P < 0.01$ and $P < 0.05$, respectively). The H₂O post-dosed group had an AUC value similar to the 2.5% NaHCO₃ group but a significantly decreased MRT value ($P < 0.05$). In the NaHCO₃ pre-dosed group, C_{max}, AUC and MRT values were significantly higher than in the non-buffered group ($P < 0.05$), whereas T_{max} value in the NaHCO₃ pre-dosed group was significantly lower than in the 2.5% NaHCO₃ group ($P < 0.05$). The NaHCO₃ post-dosed group showed significant increases in C_{max} and AUC values compared with the non-buffered group ($P < 0.01$), H₂O post-dosed group ($P < 0.01$ and 0.05 , respectively) and the 2.5% NaHCO₃ group ($P < 0.01$ and $P < 0.05$, respectively). The MRT value in the NaHCO₃ post-dosed group was significantly lower than in the 2.5%

NaHCO₃ group ($P < 0.05$).

Omeprazole is a potential inhibitor of gastric acid secretion (10), and is effective for intractable peptic ulcer that fails to respond to histamine H₂ receptor antagonists (11). Recently, proton pump inhibitors (PPI) including omeprazole have been used in combination with antibiotics to try to eradicate *Helicobacter pylori* (12, 13), which is involved in the pathogenesis of gastritis and peptic ulcer (14). Thus, PPI will probably be used more frequently, and so it seems that an occasion increases for administering the PPI to patients who cannot receive tablets.

In the present study, we examined the clinical use of omeprazole suspension. In the first place, the effect of NaHCO₃ buffer on the bioavailability of omeprazole suspension was investigated. The AUC values in the 0.2 and 2.5% NaHCO₃ groups were 1.7 and 2 times higher than in the non-buffered group, but the difference was not statistically significant. This suggests that the addition of NaHCO₃ did not sufficiently prevent degradation of omeprazole in the stomach. The MRT values in the NaHCO₃-buffered groups were significantly higher than in the non-buffered group. One explanation is that omeprazole in the non-buffered group was eliminated from plasma within 1 h due to rapid degradation in the stomach, whereas the addition of NaHCO₃ buffer decreased degradation of omeprazole and prolonged the MRT value.

We further examined the effect of administration of NaHCO₃ buffer before or after dosing of omeprazole buffered suspension. The AUC values in the NaHCO₃ pre-dosed and NaHCO₃ post-dosed groups were 1.5 and 1.8 times higher than in the 2.5% NaHCO₃ group, respectively, while the AUC value in the H₂O post-dosed group was similar to that in the 2.5% NaHCO₃ group.

These results indicate that neutralization of gastric acid is essential for improvement of the bioavailability of omeprazole, although it is still possible that the NaHCO_3 treatment increases absorption of omeprazole from the gastrointestinal tract. It appears that the NaHCO_3 pretreatment effectively decreased the degradation of omeprazole in the stomach, although partial degradation still occurs. However, the administration of NaHCO_3 buffer after omeprazole dosing accelerates gastric transit of the compound, thereby effectively reducing the degradation by gastric acid. Thus, post-treatment with the NaHCO_3 buffer is useful when the omeprazole suspension is administered to patients who cannot receive tablets.

In conclusion, the present study showed that the administration of NaHCO_3 solution following the dosing of omeprazole in buffered suspension attenuates effectively degradation of omeprazole in the stomach and causes an increase in systemic availability of the compound. The treatment of NaHCO_3 solution after omeprazole administration is a simple and useful method for administering to patients who cannot receive tablets.

Acknowledgments. The authors thank Mr. M. Okazaki and Y. Hashimoto for their technical assistance.

References

1. Fellenius E, Berglinth T and Sachs G: Substituted benzimidazoles inhibit gastric acid secretion by blocking (H^+ , K^+)-ATPase. *Nature* (1981) **290**, 159-161.
2. Sachs G and Wallmark B: The gastric H^+ , K^+ -ATPase: The site of action of omeprazole. *Scand J Gastroenterol* (1989) **24** (Suppl 166) 3-11.
3. Clissold SP and Campoli-Richards DM: Omeprazole: A preliminary review of its pharmacodynamic and pharmacokinetic properties, and therapeutic potential in peptic ulcer disease and Zollinger-Ellison syndrome. *Drugs* (1986) **32**, 15-47.
4. Cederberg C, Ekenved G, Lind T and Olbe L: Acid inhibitory characteristics of omeprazole in man. *Scand J Gastroenterol* (1985) **20** (Suppl 108), 105-112.
5. Pilbrant Å and Cederberg C: Development of an oral formulation of omeprazole. *Scand J Gastroenterol* (1985) **20** (Suppl 108), 113-120.
6. Watanabe K, Yamashita S, Furuno K, Kawasaki H and Gomita Y: Metabolism of omeprazole by gut flora in rats. *J Pharm Sci* (1995) **84**, 516-517.
7. Watanabe K, Furuno K, Eto K, Oishi R and Gomita Y: First-pass metabolism of omeprazole in rats. *J Pharm Sci* (1994) **83**, 1131-1134.
8. Yamaoka K, Tanigawara T, Nakagawa T and Uno T: A pharmacokinetic analysis program (MULTI) for microcomputer. *J Pharmacobiodyn* (1981) **4**, 879-885.
9. Yamaoka K, Nakagawa T and Uno T: Statistical moments in pharmacokinetics. *J Pharmacokin Biopharm* (1978) **6**, 547-558.
10. Larsson H, Mattson H, Sundell G and Carlsson E: Animal pharmacodynamics of omeprazole. *Scand J Gastroenterol* (1985) **20** (Suppl 108), 23-35.
11. Asaka M and Miyazaki T: Proton pump inhibitors in the treatment of H_2 -blocker resistant ulcer. *Igaku no Ayumi* (1991) **159**, 775-778 (in Japanese).
12. Labenz J, Gyenes E, Rühl GH and Börsch G: Amoxicillin plus omeprazole versus triple therapy for eradication of *Helicobacter pylori* in duodenal ulcer disease: A prospective, randomized, and controlled study. *Gut* (1993) **34**, 1167-1170.
13. Bayerdörffer E, Miehle S, Mannes GA, Sommer A, Höchter W, Weingart J, Heldwein W, Klann H, Simon T, Schmitt W, Bästlein E, Eimiller A, Hatz R, Lehn N, Dirschedl P and Stolte M: Double-blind trial of omeprazole and amoxicillin to cure *Helicobacter pylori* infection in patients with duodenal ulcers. *Gastroenterology* (1995) **108**, 1412-1417.
14. Sipponen P and Hyvarinen H: Role of *Helicobacter pylori* in the pathogenesis of gastritis, peptic ulcer and gastric cancer. *Scand J Gastroenterol* (1993) **28** (Suppl 196) 3-6.

Received February 7, 1996; accepted May 13, 1996.