

Acta Medica Okayama

Volume 42, Issue 4

1988

Article 2

AUGUST 1988

Fluorescence and Electron Microscopic Studies of the Cytoskeletal Organi-zation of Normal, Established and Transformed Chick Embryo Cells

Junzo Sasaki*

Sadahiro Watanabe[†]

Takako Nomura[‡]

Tazuko Fujiwara**

Hajime Ogura^{††}

*Okayama University,

[†]Okayama University,

[‡]Okayama University,

**Okayama University,

^{††}Okayama University,

Fluorescence and Electron Microscopic Studies of the Cytoskeletal Organization of Normal, Established and Transformed Chick Embryo Cells*

Junzo Sasaki, Sadahiro Watanabe, Takako Nomura, Tazuko Fujiwara, and Hajime Ogura

Abstract

The cytoskeletons of two established chick embryo cell (CEC) lines were examined by fluorescence and electron microscopy and compared with those of control cells and cells transformed by Rous sarcoma virus (RSV). In normal CEC, many stress fibers were observed. On the other hand, stress fibers were disorganized in nontransformed spontaneously established CEC, non-tumorigenic CEC partially transformed with a chemical carcinogen, and tumorigenic RSV-transformed CEC. In the normal CEC, actin filaments formed several bundles along the processes of the cell. Stereo-images of the peripheral region revealed bundles of filaments which were located along the attached side to the substrate. A fine well preserved network of filaments was also observed. On the other hand, in spontaneously established, partially transformed and RSV-transformed CEC, a fine network of filaments, but no actin cables, was found. These results support previous evidence that the cytoskeletal changes themselves are not directly related to the transformation or tumorigenicity of cells.

KEYWORDS: chick embryo cells, cytoskeleton, actin, Triton X-100, Rous sarcoma virus.

*PMID: 3177004 [PubMed - indexed for MEDLINE]

Copyright (C) OKAYAMA UNIVERSITY MEDICAL SCHOOL

Fluorescence and Electron Microscopic Studies of the Cytoskeletal Organization of Normal, Established and Transformed Chick Embryo Cells

Junzo Sasaki*, Sadahiro Watanabe, Takako Nomura, Tazuko Fujiwara^a and Hajime Ogura^a

Department of Anatomy, Okayama University Medical School and ^aDepartment of Virology, Cancer Institute, Okayama University Medical School, Okayama 700, Japan

The cytoskeletons of two established chick embryo cell (CEC) lines were examined by fluorescence and electron microscopy and compared with those of control cells and cells transformed by Rous sarcoma virus (RSV). In normal CEC, many stress fibers were observed. On the other hand, stress fibers were disorganized in non-transformed spontaneously established CEC, non-tumorigenic CEC partially transformed with a chemical carcinogen, and tumorigenic RSV-transformed CEC. In the normal CEC, actin filaments formed several bundles along the processes of the cell. Stereo-images of the peripheral region revealed bundles of filaments which were located along the attached side to the substrate. A fine well preserved network of filaments was also observed. On the other hand, in spontaneously established, partially transformed and RSV-transformed CEC, a fine network of filaments, but no actin cables, was found. These results support previous evidence that the cytoskeletal changes themselves are not directly related to the transformation or tumorigenicity of cells.

Key words : chick embryo cells, cytoskeleton, actin, Triton X-100, Rous sarcoma virus.

Stress fibers, a major cytoskeletal component first described by Lewis *et al.* (1), consist of microfilament bundles (2). By the use of immunofluorescence microscopy, they have been shown to contain several contractile proteins such as actin (3), myosin (4), alpha-actinin (5) and tropomyosin (6). The organization of these stress fibers was shown to change following transformation, and disorganized microfilaments were observed following transformation by RSV.

Intensive studies have been made to elucidate the correlation of transformation by RSV and intracellular cytoskeletal changes, especially concerning the organization of stress fibers (7). Cytoskeletal changes are thought to result from the direct interaction of pp60^{src}, a single viral oncogene product of RSV, with microfilaments or microfilament-associated proteins (7-12). On the other hand, Notter and Balduzzi (13) recently reported that the organization of actin and tubulin was unaffected in CEC transformed by certain strains of RSV. More recently,

* To whom correspondence should be addressed.

Nigg *et al.* (14) reported that the morphology of a transformed cell is not determined simply by pp60^{src} but by a combination of several factors. In systems other than RSV, some research groups have observed the loss of actin organization in a benign-to-malignant tumor transition (15) while other groups have shown that the loss of microfilament bundles was not related to tumorigenicity (16-18). Thus, there are discrepancies about the correlation between cytoskeletal changes and transformation or tumorigenicity.

To study systematically the correlation of cytoskeletal structure and cellular transformation, we used two avian fibroblastic cell lines established by Ogura *et al.* (19), their normal counterparts and RSV-transformed cells. Particular attention was paid to the changes of stress fibers in these cells.

Materials and Methods

Cells and virus. Embryonated eggs were obtained from Kanonji Institute, the Research Foundation for Microbial Disease of Osaka University. The CEC culture was prepared by the routine method. The Schmidt-Ruppin strain of RSV was used for transformation of CEC. The non-transformed spontaneously established CEC line (SPCC-OU1) and the CEC line (CHCC-OU1) partially transformed by treatment with N-methyl-N'-nitro-N-nitrosoguanidine were described previously (19). The cultures were maintained in a CO₂-incubator at 41°C.

Fluorescence microscopy. Cells of the last passage were grown on coverslips. The cells were fixed with 3.7% paraformaldehyde, treated with absolute acetone at -20°C for 5 min, and then air dried. The cells on coverslips were stained with a few drops of rhodamine-labeled phalloidin, a cyclic peptide that specifically binds to F-actin (20, 21), in a moist chamber at room temperature for 20 min, washed with a buffer solution and kept in

50% glycerin solution. The samples were observed with an Olympus fluorescence microscope (BH-2). Photographs were taken with Kodak Tri-X film.

Electron microscopy. Gold grids, sandwiched between a thin layer of Formvar and coverslips, were coated with carbon and sterilized by ultraviolet irradiation (12). Chick embryo cells were grown on these carbon- and Formvar-coated EM grids on coverslips. These coverslips were treated with 0.15% Triton X-100 dissolved in a PHEM buffer solution containing 60 mM piperazine-N, N'-bis(2-ethanesulfonic acid), 25 mM N-2-hydroxy ethylpiperazine-N'-2-ethanesulfonic acid, 10 mM ethylene glycol bis-(β -aminoethyl-ether) N, N, N', N'-tetraacetic acid (EGTA) and 2 mM MgCl₂ (22) for 1 min at room temperature. They were fixed with 2.5% glutaraldehyde solution for 10-20 min and postfixed with 0.5% osmium tetroxide in 0.05 M phosphate buffer, pH 6.0, for 1 min at 0°C. They were dehydrated with a graded ethanol series, immersed in isoamyl acetate and dried by critical point drying, and observed without staining with a transmission electron microscope (JEM 100 CX) at an accelerating voltage of 80 kV.

Results

Fig. 1a is a fluorescence micrograph showing the distribution of actin stained with rhodamine-labeled phalloidin in a normal C-EC. Many stress fibers extended along the long axis of normal cells, and some fibers crossed them to give a lattice appearance. These fibers extended into the ruffles and terminated at the edges or a few micrometers from the edges of the ruffles. The sub-membranous region without membrane ruffles was also stained intensely so that the concave of the cell border was delimited. Both spontaneously established SPCC-OU1 cells and partially transformed CHCC-OU1 cells showed similar structures. Stress fibers were disorganized and the background fluorescent intensity decreased. Although stress




Fig. 1 Fluorescence micrographs of chick embryo cells (CEC) stained with rhodamin (rh)-phalloidin. (a) Normal CEC. Many stress fibers run along the long axis of the cells and into the ruffles. The cell cortex without membrane ruffles is also stained intensely. $\times 600$. (b) Spontaneously established CEC. Stress fibers are disorganized although filaments remain in the cytoplasm. Fluorescent intensity is decreased. $\times 600$. (c) Partially transformed CEC. Fluorescent intensity in the cytoplasm is decreased. Intense fluorescence is seen in the center of the cell. $\times 600$. (d) RSV-transformed CEC. The cells are round and have fine filaments in the cytoplasm, but stress fibers are not observed. Actin is concentrated at the ruffle and around the nucleus. Actin aggregates (actin patches) are present. $\times 500$.




Fig. 2 Transmission electron micrographs of normal CEC treated with 0.15% Triton X-100. (a) Many filaments form several bundles identified as stress fibers by fluorescent microscopy. Bar=5 μm . (b) These bundles terminate before the edges of the cell and form focal patches where these cells adhered to the substratum. Bar=2 μm . (c) Stereopair of micrographs showing radiating filament bundles. Granular bodies exist in an area more dorsal than these filaments. Bar=1 μm .




Fig. 3 Transmission electron micrographs of spontaneously established CEC treated with detergent. (a) Flattened cells have several processes whose shapes are the same as observed by fluorescence microscopy. Bar=5 μ m (b) In the processes, thicker filaments are observed. Bar=1 μ m.

Fig. 4 Transmission electron micrographs of partially transformed CEC treated with detergent. (a) The morphology is quite similar to spontaneously established CEC. Bar=5 μ m. (b) Thicker filaments are observed around the nucleus. Bar=2 μ m.




Fig. 5 Transmission electron micrographs of RSV-transformed CEC treated with detergent. (a) Stress fibers are not found in these cells, but fine filaments run radially from the nucleus to the periphery. Bar=5 μ m. (b) Some filaments run along the cell periphery. Bar=1 μ m.

fibers were no more visible, weakly stained fine filaments were present in the cytoplasm (Fig. 1 b). Actin often was concentrated at the periphery of the cell and around the nucleus (Fig. 1 c). RSV-transformed CEC became round with fine filaments in the cytoplasm, but stress fibers were not observed. Actin was concentrated at the ruffle or around the nucleus. Actin aggregates (actin patches) were often present near the ventral surface of the cells (Fig. 1 d).

The treatment of the cells with a detergent solution removed the cytoplasmic materials satisfactorily and revealed the cytoskeletal organization of these cells (Figs. 2-5). Figs. 2 a-c show normal CEC treated with detergent. Many filaments formed several bundles identified as stress fibers by fluorescence microscopy. These bundles terminated before the edges of the cell where

these cells adhered to the substratum, forming focal patches. Fine filaments splayed to the edges and formed a network at the ruffles where cytoplasmic materials were scarce. Stereo-pairs of micrographs showed that radiating filament bundles ran along the attached side to the substratum, and fibers which delimited the cell edges crossed over these filament bundles. Granular bodies existed in an area more dorsal to these filaments. The network of filaments was well preserved (Fig. 2 c). In spontaneously established SPCC-OU1 cells (Figs. 3 a, b) and partially transformed CHCC-OU1 cells (Figs. 4 a, b), the shape of the cell margin was the same as that seen by fluorescence microscopy. Stress fibers were not observed, and thicker filaments were often seen around the nucleus. A network of filaments was present in the cell body. RSV-

transformed CEC (Figs. 5 a,b) were round, and stress fibers were not formed in these cells, but fine filaments radiated from the nucleus region to the periphery. Some filaments ran along the cell periphery.

Discussion

Previous reports about the correlation between cytoskeletal changes and transformation (7, 11) led us to investigate systematically the cytoskeletal structures in CEC. Transformed cells contained far fewer stress fibers than did their normal counterparts. The cells investigated in this study were: a) normal, b) spontaneously established but not transformed (SPCC-OU1), c) partially transformed by a chemical carcinogen but not tumorigenic and d) tumorigenic RSV-transformed CECs. The SPCC-OU1 cells are morphologically normal and form no colony in soft agar, while CHCC-OU1 cells pile up and form colonies in soft agar, but do not produce tumors either in the syngeneic chicken or in nude mice (19).

To study the fine structure of the cytoskeleton, we applied fluorescence microscopy using rhodamine-labeled phalloidin and electron microscopy using a detergent (23, 24). Typical stress fibers were observed by fluorescence microscopy only in normal CEC. In spontaneously established non-transformed SPCC-OU1, partially transformed, non-tumorigenic CHCC-OU1 and tumorigenic RSV-transformed CEC, stress fibers were disorganized. Instead of the bundles of filaments which constitute stress fibers, a fine network of filaments was observed in these cells by electron microscopy. The fluorescence microscopic findings correlated well with the electron microscopic findings.

These observations are in agreement with the reports which state that there is no direct correlation between cytoskeletal

changes and transformation (13, 14), and, accordingly, disagree with the proposal that a correlation exists between stress fibers and transformation (8, 9, 12). The idea that the cytoskeletal change causes a loss of adhesiveness to the substratum and allows cells to deform and invade neighboring tissues (15) is simple and tempting, but not certain (16-18).

The cytoplasmic aggregates described as a flower (7, 11) or patch (25) were observed in CEC transformed by RSV but not in CEC transformed by a chemical carcinogen. At present, the role and meaning of these structures in RSV-transformed cells are unknown.

Acknowledgments. We are grateful to Professor N. Otsuka, Department of Anatomy and Professor Y. Yabe, Department of Virology, Cancer Institute for their continued support throughout this study. We also wish to express our gratitude to Professor Y. Kanemasa, Department of Microbiology for the use of the fluorescence microscope. This work was supported in part by a Grant-in Aid for general scientific research (No. 62570007) from the Ministry of Education, Science, and Culture of Japan.

References

1. Lewis WH and Lewis MR: Behavior of cells in tissue cultures; in *General Cytology*, Cowdry ed, The University of Chicago Press, Chicago (1924) pp385-447.
2. Buckley IK and Porter KR: Cytoplasmic fibrils in living cultured cells: A light and electron microscope study. *Protoplasma* (1967) **64**, 349-380.
3. Lazarides E and Weber K: Actin antibody: The specific visualization of actin filaments in non-muscle cells. *Proc Natl Acad Sci USA* (1974) **71**, 2268-2272.
4. Weber K and Gröschel-Stewart U: Antibody to myosin: The specific visualization of myosin-containing filaments in non-muscle cells. *Proc Natl Acad Sci USA* (1974) **71**, 4561-4564.
5. Lazarides E and Burridge K: Alpha-actinin: Immunofluorescent localization of a muscle structural protein in non-muscle cells. *Cell* (1975) **6**, 289-298.
6. Lazarides E: Tropomyosin antibody: The specific localization of tropomyosin in non muscle cells. *J Cell Biol* (1975) **65**, 549-561.
7. Boschek CB: Organizational changes of cytoskeletal

- proteins during cell transformation: *Adv Viral Oncol* (1982) **1**, 173-187.
8. Edelman GM and Yahara I: Temperature-sensitive changes in surface modulating assemblies of fibroblasts transformed by mutants of Rous sarcoma virus. *Proc Natl Acad Sci USA* (1976) **73**, 2047-2051.
 9. Wang E and Goldberg AR: Changes in microfilament organization and surface topography upon transformation of chick embryo fibroblasts with Rous sarcoma virus. *Proc Natl Acad Sci USA* (1976) **73**, 4065-4069.
 10. Sefton BM, Hunter T, Ball EH and Singer SJ: Vinculin: A cytoskeletal target of the transforming protein of Rous sarcoma virus. *Cell* (1981) **24**, 165-174.
 11. Boschek CB, Jockusch BM, Friis RR, Back R, Grundmann E and Bauer H: Early changes in the distribution and organization of microfilament proteins during cell transformation. *Cell* (1981) **24**, 175-184.
 12. Witt DP, Brown DJ and Gordon JA: Transformation-sensitive isoactin in passaged chick embryo fibroblasts transformed by Rous sarcoma virus. *J Cell Biol* (1983) **96**, 1766-1771.
 13. Notter MFD and Balduzzi PC: Cytoskeletal changes induced by two avian sarcoma viruses: UR2 and Rous sarcoma virus. *Virology* (1984) **136**, 56 - 68.
 14. Nigg EA, Sefton BM, Singer SJ and Vogt PK: Cytoskeletal organization, vinculin-phosphorylation, and fibronectin expression in transformed fibroblasts with different cell morphologies. *Virology* (1986) **151**, 50-65.
 15. Friedman E, Verderame M, Winawer S and Pollack R: Actin cytoskeletal organization loss in the benign-to-malignant tumor transition in cultured human colonic epithelial cells. *Cancer Res* (1984) **44**, 3040-3050.
 16. Fine RE and Taylor L: Decreased actin and tubulin synthesis in 3T3 cells after transformation by SV 40 virus. *Exp Cell Res* (1976) **102**, 162-168.
 17. Willingham MC, Yamada KM, Yamada SS, Pouyssegur J and Pastan I: Microfilament bundles and cell shape are related to adhesiveness to substratum and are dissociable from growth control in cultured fibroblasts. *Cell* (1977) **10**, 375-380.
 18. Celis JE, Small JV, Kaltoft K and Celis A: Microfilament bundles in transformed mouse CLID x transformed CHO cell hybrids: Correlation with tumorigenicity in nude mice. *Exp Cell Res* (1979) **120**, 79-86.
 19. Ogura H, Fujiwara T and Namba M: Establishment of two chick embryo fibroblastic cell lines. *Gann* (1984) **75**, 410-414.
 20. Wulf E, Deboen A, Bauts FA, Faulstich H and Wieland T: Fluorescent phallotoxin, a tool for the visualization of cellular actin. *Proc Natl Acad Sci USA* (1979) **76**, 4498-4502.
 21. Wieland T, Miura T and Seeliger A: Analogs of phalloidin. *Int J Pept Protein Res* (1983) **21**, 3-10.
 22. Schliwa M and van Blerkom J: Structural interaction of cytoskeletal components. *J Cell Biol* (1981) **90**, 222-235.
 23. Watanabe S, Takehara Y, Fujii Y, Okimasu E, Moromizato Y and Sasaki J: Fluorescence and electron microscopic study of intracellular F-actin in concanavalin A-treated or cytochalasin B-treated Ehrlich ascites tumor cells. *Acta Med Okayama* (1986) **40**, 301-311.
 24. Sasaki J, Takehara Y, Fujii Y, Nomura T and Watanabe S: Presence of abundant microfilaments in apical caps of the nonciliated bronchiolar epithelial (Clara) cells. *Am J Anat* (1987) **179**, 1-9.
 25. Carley WW, Barak LS and Webb WW: F-actin aggregates in transformed cells. *J Cell Biol* (1981) **90**, 797-802.

Received April 21, 1988; accepted June 7, 1988