

# *Acta Medica Okayama*

---

*Volume 60, Issue 6*

2006

*Article 6*

DECEMBER 2006

---

## Three Type 6 Hepatitis C Virus Subgroups among Blood Donors in the Yangon Area of Myanmar Are Identified as Subtypes 6m and 6n, and a Novel Subtype by Sequence Analysis of the Core Region.

Toshiyuki Shinji\*

Aye Aye Lwin†

Katsunori Gokan‡

Mikako Obika\*\*

Hiromasa Ryuko††

Myo Khin‡‡

Shigeru Okada§

Norio Koide¶

\*Okayama University,

†Okayama University,

‡Okayama University,

\*\*Okayama University,

††Okayama University,

‡‡Lower Myanmar, Myanmar,

§Okayama University,

¶Okayama University,

# Three Type 6 Hepatitis C Virus Subgroups among Blood Donors in the Yangon Area of Myanmar Are Identified as Subtypes 6m and 6n, and a Novel Subtype by Sequence Analysis of the Core Region.\*

Toshiyuki Shinji, Aye Aye Lwin, Katsunori Gokan, Mikako Obika, Hiromasa Ryuko, Myo Khin, Shigeru Okada, and Norio Koide

## Abstract

Previously, using phylogenetic analysis of NS5b sequences, we found that three type 6 variant subgroups (M6#x9D;1, M6#x9D;2 and M6#x9D;3) exist in Myanmar. According to the new nomenclature of hepatitis C, M6#x9D;1 and M6#x9D;2 belong to subtypes 6m and 6n, respectively, but M6#x9D;3 is unassigned. In this study, we sequenced and phylogenetically analyzed the core region of these type 6 variant subgroups. Serum samples assigned as 6m or 6n by NS5b sequence were also identified as 6m or 6n by core region analysis. The M6#x9D;3 (sample name MYAN-3E-3) remained unassigned to a subgroup based on its core region analysis. The findings of this study suggest that either the core region or the NS5b region can be analyzed for HCV subtype classification.

**KEYWORDS:** HCV genotype, type 6 subgroup, Myanmar, HCV core, phylogenetic analysis

---

\*PMID: 17189978 [PubMed - in process]

*Short Communication*

## Three Type 6 Hepatitis C Virus Subgroups among Blood Donors in the Yangon Area of Myanmar Are Identified as Subtypes 6m and 6n, and a Novel Subtype by Sequence Analysis of the Core Region

Toshiyuki Shinji<sup>a\*</sup>, Aye Aye Lwin<sup>a</sup>, Katsunori Gokan<sup>a</sup>, Mikako Obika<sup>a</sup>,  
Hiromasa Ryuko<sup>a</sup>, Myo Khin<sup>b</sup>, Shigeru Okada<sup>c</sup>, and Norio Koide<sup>a</sup>

*Departments of <sup>a</sup>Laboratory Medicine, and <sup>c</sup>Food and Health Science, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama 700-8558, Japan, and <sup>b</sup>Department of Medical Research (Lower Myanmar), Yangon 11191, Myanmar*

Previously, using phylogenetic analysis of NS5b sequences, we found that three type 6 variant subgroups (M6-1, M6-2 and M6-3) exist in Myanmar. According to the new nomenclature of hepatitis C, M6-1 and M6-2 belong to subtypes 6m and 6n, respectively, but M6-3 is unassigned. In this study, we sequenced and phylogenetically analyzed the core region of these type 6 variant subgroups. Serum samples assigned as 6m or 6n by NS5b sequence were also identified as 6m or 6n by core region analysis. The M6-3 (sample name MYAN-3E-3) remained unassigned to a subgroup based on its core region analysis. The findings of this study suggest that either the core region or the NS5b region can be analyzed for HCV subtype classification.

**Key words:** HCV genotype, type 6 subgroup, Myanmar, HCV core, phylogenetic analysis

A higher prevalence of HCV infection has been found in Southeast Asia than in other geographic locations. Myanmar is located in Southeast Asia and geographically bordered by other Southeast Asian countries, namely China, Thailand, Laos, and South Asian countries such as India or Bangladesh. The HCV genotype distributions of these countries differ [1–5]. Therefore, information on HCV genotypes in Myanmar is epidemiologically significant; yet such information is scarce.

We previously reported HCV genotypes of the Yangon city area of Myanmar [6]. Genotypes 3 and

1 and genotype 6 variants are the main genotypes. Using phylogenetic analysis of NS5b sequences, we found that three type 6 variant subgroups exist in Myanmar. According to the newly proposed nomenclature of hepatitis C, the M6-1 and M6-2 groups belong to subtypes 6m and 6n, respectively [7]. But M6-3 is still unassigned [8].

In this study, we sequenced the core region of these type 6 variant subgroup sera samples, compared the sequences with other reported sequences by phylogenetic analysis, and examined whether these samples are classified the same way by core region as by NS5b sequence analysis.

Received April 13, 2006; accepted May 31, 2006.

\*Corresponding author. Phone: +81-86-235-7342; Fax: +81-86-235-7345  
E-mail: [toshinji@md.okayama-u.ac.jp](mailto:toshinji@md.okayama-u.ac.jp) (T. Shinji)

## Materials and Methods

**Serum samples.** Four HCV antibody positive serum samples representing three type 6 subgroups found for a previous study were used in this study. MYAN-8H-1 and MYAN-9H-1 represent the M6-1 group. MYAN-2I-2 and MYAN-3E-3 represent M6-2 and M6-3, respectively.

**RNA extraction.** Amplicor HCV 2.0 positive sera were HCV-RNA extracted for further experimentation. RNA was extracted from 0.125 ml of serum by the single-step method of acid guanidinium thiocyanate-phenol-chloroform extraction [9] using Isogen-LS reagent (Nippon Gene Co., Toyama, Japan) according to the manufacturer's instructions. Extracted samples were precipitated with isopropanol and washed with 70% ethanol. The resulting pellet was resuspended in 20  $\mu$ l of RNase-free water.

**RT-PCR and nested PCR.** RT-PCR and nested PCR were performed according to the method of Ohno *et al.* [10] with minor modifications. Briefly, extracted RNA was amplified by reverse transcriptase (RT)-PCR using sense primer Sc2 and antisense primer Ac2 at a portion of the core region (Table 1). RT-PCR was performed by the following protocol. We first prepared 20  $\mu$ l of reaction mixture, containing 4  $\mu$ l of 5 x reaction buffer, 0.8  $\mu$ l of RNase inhibitor, 0.8  $\mu$ l of rTTH DNA polymerase (Toyobo, Osaka, Japan), 0.25  $\mu$ M of each primer and 0.3 mM of deoxynucleotide. The thermal cycler was programmed at 60 °C for 30 min and 95 °C for 1 min for the RT step; 95 °C 15 sec, 60 °C 20 sec for 2 cycles; and 90 °C 15 sec, 60 °C 20 sec for 38 cycles, followed by extension at 60 °C for 4 min.

The second PCR was performed using sense primer S7 (Table 1) and antisense primer A5 at

95 °C for 10 min and at 95 °C for 15 sec; and 60 °C for 20 sec for 30 cycles followed by at 60 °C for 4 min.

The resultant PCR product was electrophoresed on 3% agarose gel in 1x TAE, stained in ethidium bromide solution, and evaluated under UV light. To avoid the risk of false-positive results, PCR assays were done with strict precautions against cross-contamination.

**Direct sequencing.** Direct sequencing was performed as follows. Five microliters of crude PCR product was purified by MonoFas DNA kit I (GL Science, Tokyo Japan). 2.5 or 5.0  $\mu$ l of purified PCR product was used as a template, and cycle-sequencing reactions were completed with the addition of 2  $\mu$ l (3.2 pmol) internal sense or antisense sequencing primer using S7 or A5, plus 8  $\mu$ l of the dye terminator included in the Big Dye Terminator 1.1 (Applied Biosystems, Foster City, CA, USA) cycle sequencing reaction kit and RNase-free water (total reaction amount 20  $\mu$ l). Thermal cycling was performed on a Gene Amp 9600 thermal cycler at 96 °C for 10 sec, at 50 °C for 5 sec and at 60 °C 4 min, for a total of 25 cycles. After heat shock (95 °C for 2 min, at -80 °C for 5 min, sequencing was performed using an ABI PRISM 310 Genetic Analyzer (Applied Biosystems).

**Phylogenetic analysis by core region sequences of Myanmar HCV type 6 subgroup samples.** We analyzed a phylogenetic tree with core sequences of the four Myanmar HCV type 6 subgroup samples of this study and with other known genotype sequences obtained from the GenBank database.

ODEN (version 1.1.1) [11] using the six-parameter method [12] was used to determine the number

**Table 1** Oligonucleotide primers used for PCR sequencing of the core region

Primer	Nucleotide Sequence	Nucleotide Position
Sc2	5'-GGGAGGTCTCGTAGACCGTGACCATG-3'	nt.318-344
Ac2	5'-GAGMGGKATRTACCCCATGAGRTCGGC-3'	nt.758-732
S7	5'-AGACCGTGACCATGAGCAC-3'	nt.330-349
A5	5'-TACGCCGGGGTCAKTRGGGCCCA-3'	nt.684-660

R, A.G; M, A.C; K, G.T;

S represents the sense primer; A represents the antisense primer.

Nucleotide numbering is from the 5'end of the HCV genome of HC-J4/83 (accession no. D13558).

of nucleotide substitutions per site (genetic distance) between the isolates. Based on these values, a phylogenetic tree was constructed by the neighbor-joining (N-J) method [13]. The tree was plotted by a program from DDBJ with the mid-point rooting option. To confirm the reliability of the phylogenetic tree, bootstrap resampling tests were performed 1,000

times [14].

**Nucleotide sequence accession numbers.**

The sequences obtained in this study have been submitted to the DDBJ, GeneBank and EMBL nucleotide sequence databases and have been assigned accession numbers AB254860 to AB254863.

```

MYAN-8H-1 1:CAACACCAACCGCCGCCACAGGACGTTAAGTCCCGGGCGGGCCAGATCGTCGGTGG 60
MYAN-9H-1 1:TAACACCAACCGCCGCCACAGGACGTTAAGTCCCGGGCGGGCCAGATCGTCGGTGG 60
MYAN-2I-2 1:AAACACCAACCGTCGTCATGGATGTTAAGTCCCGGGCGGGCCAGATCGTCGGTGG 60
MYAN-3E-3 1:AAACACCAACCGTCGTCATGGATGTTAAGTCCCGGGTGGCGGTGTCAGATCGTGGTGG 60
VN506-6a 1:AAACACCAACCGTCGTCATGGACGTCAGTCCCGGGTGGCGGTGTCAGATCGTGGCGGG 60
VN843-7a 1:AAACACCAACCGCCGCCACAGGACGTCAGTCCCGGGTGGCGGTGTCAGATCGTGGTGG 60
VN530-8a 1:AAACACCAACCGTCGTCATGGACGTCAGTCCCGGGTGGCGGTGTCAGATCGTCGGTGG 60
VN004-9a 1:AAACACTAACCGTCGTCATGGACGTCAGTCCCGGGCGGGCCAGATCGTGGTGG 60
***** ** ** * * ** ***** ** ** ***** ** **

MYAN-8H-1 61:AGTCTACTTGTGCGCGCAGGGGCCACGTTTGGGTGTGCGCGCAGTGAGGAAGACCTC 120
MYAN-9H-1 61:AGTCTACTTGTGCGCGCAGGGGCCACGTTTGGGTGTGCGCGCAGTGAGGAAGACCTC 120
MYAN-2I-2 61:AGTCTACTTATTGCGCGCAGGGGCCACGTTTGGGTGTGCGCGCAGTGAGGAAGACTTC 120
MYAN-3E-3 61:AGTTTACTTGTGCGCGCAGGGGCCCTAGATTGGGTGTGCGCGCAGTTAGGAAGACTTC 120
VN506-6a 61:AGTTTACTTGTGCGCGCAGGGGCCCTAGATTGGGTGTGCGCGCAACGAGAAAGACTTC 120
VN843-7a 61:AGTTTACTTGTGCGCGCAGGGGCCCTAGATTGGGTGTGCGCGCAGGAGAAAGACTTC 120
VN530-8a 61:AGTCTACTTATTGCGCGCAGGGGCCCTAGATTGGGTGTGCGCGCAACGAGGAAGACTC 120
VN004-9a 61:AGTTTACTTGTGCGCGCAGGGGCCCTAGATTGGGTGTGCGCGCAGGAGAAAGACTTC 120
*** ***** ***** * ***** ** ** ** **

MYAN-8H-1 121:TGAACGATCCAGCCTAGAGGCAGGGCCAGCCATACCAAAGGCGCGCCAGCCGACGGG 180
MYAN-9H-1 121:TGAACGATCCAGCCTAGAGGCAGGGCTCAGCCTATACCAAAGGCGCGCCAGCCGACGGG 180
MYAN-2I-2 121:TGAACGATCCAGCCTAGAGGCAGGGCTCAGCCATACCAAAGGCGCGCCAGCCGACGGG 180
MYAN-3E-3 121:TGAACGATCCAGCCTAGAGGCAGGGGAGAGCCAGCCTATACCAAAGCAGGCTCCTCCGACGGG 180
VN506-6a 121:CGAGCGATCCAGCCTAGAGGCAGGGCCAACTATACCAAAGGCGCGCCAGTCCAGGG 180
VN843-7a 121:TGAACGGTCCAGCCAGGGGTAGAGCCAACTATACGAAAAGCGCGTACCAAACGGG 180
VN530-8a 121:CGAGCGGTCCAGCCAGGGGAGGGCTCAGCCATACCAAAGGCGCGCCAGCCAGCCG 180
VN004-9a 121:CGAACGGTCCAGCCTAGAGGCAGGGCCAGCCATACCAAAGGCGCGCCAGCCGATAGG 180
** ** ***** * ** ** ** ** ** ***** * ** * **

MYAN-8H-1 181:CCGGCACTGGGCTCAGCCTGGTTACCCCTGGCCCTTATGGGAATGAGGGCTGCGGTTG 240
MYAN-9H-1 181:CCGGCACTGGGCTCAGCCTGGTTACCCCTGGCCCTTATGGCAACGAGGGCTGCGGTTG 240
MYAN-2I-2 181:CCGGCACTGGGCTCAGCCTGGTTACCCCTGGCCCTTATGGGAATGAGGGCTGCGGATG 240
MYAN-3E-3 181:CCGGCACTGGGCTCAGCCCGTTACCCCTGGCCCTTACGGTAACGAGGGCTGCGGCTG 240
VN506-6a 181:CCGGCACTGGGCTCAGCCCGATACCCCTGGCCCTTATGGAAACGAGGGCTGCGGTTG 240
VN843-7a 181:CCGTACCTGGGCACAGCCTGGGTACCCCTGGCCCTTATGGGAATGAGGGTTGGCTG 240
VN530-8a 181:CCGGCACTGGGCGCAGCCTGGTTACCCCTGGCCCTTATGGTAACGAGGGCTGCGGATG 240
VN004-9a 181:CCGTAGCTGGGTCACCCGGTACCCCTGGCCCTTATGGCAACGAGGGCTGCGGATG 240
*** ***** ** ** ** ***** ***** ** ** ** ***** ** **

MYAN-8H-1 241:GGCAGGGTGGCTCCTGTCCCTCGGGTTCGCGCCTAAT 280
MYAN-9H-1 241:GGCAGGGTGGCTCCTGTCCCTCGGGTTCGCGCCTCAT 280
MYAN-2I-2 241:GGCAGGGTGGCTCCTGTCCCTCGGGTTCGCGCCTAAT 280
MYAN-3E-3 241:GGCAGGGTGGCTCCTGTCCCTCGGGTTCGCGCCTAAT 280
VN506-6a 241:GGCAGGTTGGCTCCTGTCCCTCGGGTTCGCGCCACAT 280
VN843-7a 241:GGCAGGGTGGCTCCTGTCCCTCGGGTTCGCGCCTAAT 280
VN530-8a 241:GGCAGGGTGGCTCCTGTCCCTCGGGTTCGCGCCTCAC 280
VN004-9a 241:GGCAGGGTGGCTCCTGTCCCTCGGGTTCGCGCCTAAT 280
*** ** ***** ** ** ** ***** ** ** ** ***** ** **
    
```


Fig. 1 Comparison of sequences of the core region (nucleotides 380–659) of type 6 subgroups. VN506 represents type 6a. VN843, VN530 and VN004 represent types 7a, 8a and 9a respectively. MYAN-8H-1 and MYAN-9H-1 represent the M6-1 subgroup. MYAN-2I-2 represents the M6-2 subgroup. MYAN-3E-3 represents M6-3. Asterisks indicate identical nucleotides.

## Results

**Sequence results and phylogenetic tree of HCV core region including Myanmar type 6 subgroup samples.** A comparison of sequences of the core region nucleotides 380–659 of type 6 subgroups including representatives of M6-1 to M6-3 is

shown in Fig. 1.

As with the NS5b analysis, phylogenetic analysis of the core region classified the type 6 variants into 3 subgroups (Fig. 2). The first 2 of these subgroups, which we had formerly named M6-1, M6-2 and M6-3, have been assigned according to the new nomenclature, to 6m and 6n [7], but M6-3 (sample name



**Fig. 2** Phylogenetic tree constructed from the core region of 4 HCV type 6 variants in Myanmar and previously reported type 6 group variants, along with representatives of the other 5 genotypes. Bootstrap values are shown along each main branch. The length of the horizontal bar indicates the number of nucleotide substitutions per site. Isolates labeled "MYAN-" were sequenced in the present study.

MYAN-3E-3) has not yet been assigned.

### Discussion

HCV genotype 6 variants are common in the Southeast Asian countries, and also are restricted to this region. They have been classified into many subgroups from 6a to 6q [7]. 5'UTR PCR is often preferred for sensitive diagnostic assays, but sequence analysis of the region is not appropriate to differentiate genotypes 1 and 6 [3, 7]. To propose a new subtype, 3 or more examples of infection and sequences from both the core/ E1 region and the NS5b region are required [7]. Previously we reported our analysis of the NS5b region of type 6 subgroups in Myanmar 6. In this study, we analyzed the core regions of the same samples.

By the new nomenclature system [7], the M6-1 group (MYAN-8H-1, MYAN-9H-1) is classified as subtype 6m, and the M6-2 group (MYAN-2I-2) is classified as subtype 6n. At this point, subtype 6m has been reported only in Myanmar and northern Thailand. On the other hand, subtype 6n has been reported in Myanmar, Thailand and Kunming city in southwestern China [8]. Subtype 6n may more widely spread in this area than 6m.

The M6-3 group (MYAN-3E-3) was an unassigned type 6 subtype by NS5b analysis [6, 8], and remained as an unassigned type 6 subtype by the core sequence analysis in this study. Phylogenetic analysis of both the core region and NS5b region showed the same results. These findings suggest that, for HCV subtype classification, either the core region or the NS5b region can be analyzed.

Currently, we are analyzing sequences of HCV samples from the Myanmar - China border area and from the Myanmar - India border area. As a preliminary result, we found several samples that form a group of MYAN-3E-3. This group could be provisionally classified as a new subtype of HCV. We are performing further study for the assignment of new subtype of HCV from Myanmar.

**Acknowledgements.** This work was supported by a grant from the Okayama University COE Project (2004, 2005), Japan and a grant from Ryobi-Teien Memorial Foundation (2004), Japan.

### References

1. Tokita H, Okamoto H, Luengrojanakul P, Vareesangthip K, Chainuvati T, Iizuka H, Tsuda F, Miyakawa Y and Mayumi M: Hepatitis C virus variants from Thailand classifiable into five novel genotypes in the sixth (6b), seventh (7c, 7d) and ninth (9b, 9c) major genetic groups. *J Gen Virol* (1995) 76: 2329-2335.
2. Tokita H, Okamoto H, Iizuka H, Kishimoto J, Tsuda F, Lesmana LA, Miyakawa Y and Mayumi M: Hepatitis C virus variants from Jakarta, Indonesia classifiable into novel genotypes in the second (2e and 2f), tenth (10a) and eleventh (11a) genetic groups. *J Gen Virol* (1996) 77: 293-301.
3. Mellor J, Walsh EA, Prescott LE, Jarvis LM, Davidson F, Yap PL and Simmonds P: Survey of type 6 group variants of hepatitis C virus in Southeast Asia by using a core-based genotyping assay. *J Clin Microbiol* (1996) 34: 417-423.
4. Hissar SS, Goyal A, Kumar M, Pandey C, Suneetha PV, Sood A, Midha V, Sakhuja P, Malhotra V and Sarin SK: Hepatitis C virus genotype 3 predominates in North and Central India and is associated with significant histopathologic liver disease. *J Med Virol* (2006) 78: 452-458.
5. Khaja MN, Madhavi C, Thippavazzula R, Nafeesa F, Habib AM, Habibullah CM and Guntaka RV: High prevalence of hepatitis C virus infection and genotype distribution among general population, blood donors and risk groups. *Infect Genet Evol* (2006) 6: 198-204.
6. Shinji T, Kyaw YY, Gokan K, Tanaka Y, Ochi K, Kusano N, Mizushima T, Fujioka S, Shiraha H, Lwin AA, Shiratori Y, Mizokami M, Khin M, Miyahara M, Okada S and Koide N: Analysis of HCV genotypes from blood donors shows three new HCV type 6 subgroups exist in Myanmar. *Acta Med Okayama* (2004) 58: 135-142.
7. Simmonds P, Bukh J, Combet C, Deleage G, Enomoto N, Feinstone S, Halfon P, Inchauspe G, Kuiken C, Maertens G, Mizokami M, Murphy DG, Okamoto H, Pawlitsky JM, Penin F, Sablon E, Shin-I T, Stuyver LJ, Thiel HJ, Viazov S, Weiner AJ and Widell A: Consensus Proposals for a Unified System of Nomenclature of Hepatitis C Virus Genotypes. *Hepatology* (2005) 42: 962-973.
8. Lu L, Nakano T, Li C, Fu Y, Miller S, Kuiken C, Robertson BH and Hagedorn CH: Hepatitis C virus complete genome sequences identified from China representing subtypes 6k and 6n and a novel, as yet unassigned subtype within genotype 6. *J Gen Virol* (2006) 87: 629-634.
9. Chomczynski P and Sacchi N: 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* (1987) 162: 156-159.
10. Ohno T, Mizokami M, Wu RR, Saleh MG, Ohba K, Orito E, Mukaide M, Williams R and Lau JY: New Genotyping system that allows for identification of HCV Genotypes 1a, 1b, 2a, 2b, 3a, 3b, 4, 5a, and 6a. *J Clin Microbiol* (1997) 35: 201-207.
11. Ina Y: ODN: A program package for molecular evolutionary analysis and database search of DNA and amino acid sequences. *Comput Appl Biosci* (1994) 10: 11-12.
12. Gojobori T, Ishii T and Nei M: Estimation of average number of nucleotide substitutions when the rate of substitution varies with nucleotide. *J Mol Evol* (1982) 18: 414-423.
13. Saitou N and Nei M: The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* (1987) 4: 406-425.
14. Felsenstein J: Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* (1985) 39: 783-791.