

# *Acta Medica Okayama*

---

*Volume 50, Issue 1*

1996

*Article 8*

FEBRUARY 1996

---

## cDNA cloning of rat major AP endonuclease (APEX nuclease) and analyses of its mRNA expression in rat tissues.

Yunshan Tan\*

Yuko Nakagawa†

Kosuke Akiyama‡

Hajime Wakabayashi\*\*

Altaf H. Sarker††

Shuji Seki‡‡

\*Okayama University,

†Okayama University,

‡Okayama University,

\*\*Okayama University,

††Okayama University,

‡‡Okayama University,

# cDNA cloning of rat major AP endonuclease (APEX nuclease) and analyses of its mRNA expression in rat tissues.\*

Yunshan Tan, Yuko Nakagawa, Kosuke Akiyama, Hajime Wakabayashi, Altaf H. Sarker, and Shuji Seki

## Abstract

APEX nuclease is a mammalian DNA repair enzyme having apurinic/apyrimidinic (AP) endonuclease, 3'-5'-exonuclease, DNA 3' repair diesterase and DNA 3'-phosphatase activities. It is also a redox factor (Ref-1), stimulating DNA binding activity of AP-1 binding proteins such as Fos and Jun. In the present paper, a cDNA for the enzyme was isolated from a rat brain cDNA library using mouse Apex cDNA as a probe and sequenced. The rat Apex cDNA was 1221 nucleotides (nt) long, with a 951-nt coding region. The amino acid sequence of rat APEX nuclease has 98.4% identity with mouse APEX nuclease. Using the rat Apex cDNA as a probe for Northern blot analysis, the size of rat Apex mRNA was shown to be approximately 1.5 kb. Its expression was compared in 9 rat organs on postnatal days 7 and 28. Although Apex mRNA was expressed ubiquitously, the levels varied significantly, suggesting organ- or tissue-specific expression of the Apex gene. The highest level was observed in the testis, relatively high levels in the thymus, spleen, kidney and brain, and the lowest level in the liver. The level of expression at postnatal day 28, with the exception of the testis, was almost the same as or lower in respective organs than that at postnatal day 7. Postnatal developmental changes of Apex mRNA expression in the testis and thymus were further studied. The expression in testis was markedly increased on postnatal days 21 and 28. The expression in thymus increased once at postnatal day 14, and then decreased. The developmental changes of Apex mRNA expression in testis and thymus suggest that APEX nuclease is involved in processes such as recombinational events.

**KEYWORDS:** Apurinic/apyrimidinic endonuclease; APEX nuclease; Repair enzyme; Apex mRNA; Northern blot; development; testis; rat

---

\*PMID: 8701782 [PubMed - indexed for MEDLINE]

Copyright (C) OKAYAMA UNIVERSITY MEDICAL SCHOOL

## cDNA Cloning of Rat Major AP Endonuclease (APEX Nuclease) and Analyses of Its mRNA Expression in Rat Tissues<sup>†</sup>

Yunshan TAN, Yuko NAKAGAWA, Kosuke AKIYAMA, Hajime WAKABAYASHI, Altaf H. SARKER and Shuji SEKI\*

Department of Molecular Biology, Institute of Cellular and Molecular Biology, Okayama University Medical School, Okayama 700, Japan

APEX nuclease is a mammalian DNA repair enzyme having apurinic/aprimidinic (AP) endonuclease, 3'-5'-exonuclease, DNA 3' repair diesterase and DNA 3'-phosphatase activities. It is also a redox factor (Ref-1), stimulating DNA binding activity of AP-1 binding proteins such as Fos and Jun. In the present paper, a cDNA for the enzyme was isolated from a rat brain cDNA library using mouse Apex cDNA as a probe and sequenced. The rat Apex cDNA was 1221 nucleotides (nt) long, with a 951-nt coding region. The amino acid sequence of rat APEX nuclease has 98.4% identity with mouse APEX nuclease. Using the rat Apex cDNA as a probe for Northern blot analysis, the size of rat Apex mRNA was shown to be approximately 1.5 kb. Its expression was compared in 9 rat organs on postnatal days 7 and 28. Although Apex mRNA was expressed ubiquitously, the levels varied significantly, suggesting organ- or tissue-specific expression of the Apex gene. The highest level was observed in the testis, relatively high levels in the thymus, spleen, kidney and brain, and the lowest level in the liver. The level of expression at postnatal day 28, with the exception of the testis, was almost the same as or lower in respective organs than that at postnatal day 7. Postnatal developmental changes of Apex mRNA expression in the testis and thymus were further studied. The expression in testis was markedly increased on postnatal days 21 and 28. The expression in thymus increased once at postnatal day 14, and then decreased. The developmental changes of Apex mRNA expression in testis and thymus suggest that APEX nuclease is involved in processes such as recombinational events.

**Key words:** Apurinic/aprimidinic endonuclease; APEX

nuclease; Repair enzyme; Apex mRNA; Northern blot; development; testis; rat

**A**purinic/aprimidinic (AP) sites resulting from loss of bases are the most frequent DNA lesion in cells, and are generated in DNA by spontaneous hydrolysis, radiation, oxidative damage and action of DNA glycosylases on modified bases (1-4). Single-strand breaks with 3'-blocked termini are also frequently produced by radical-induced DNA damage (3-5). The resulting AP sites and 3'-blocked single-strand breaks, if they can not be repaired, block transcription and replication of DNA, and have cytotoxic and mutagenic effects on cells (2-6). Repair of these lesions is thought to be mostly initiated by AP endonucleases having 5' AP endonuclease and DNA 3' repair diesterase activities (2-5). Although multiple forms of AP endonucleases have been reported in mammalian cells, so far only one form (the major 5' AP endonuclease designated as APEX nuclease, or HAP1, APE or Ref-1 protein) of AP endonuclease/DNA 3' repair diesterase has been clearly identified (7-11). *In vitro* studies have shown that the major AP endonuclease (APEX nuclease) is multifunctional. It shows 5' AP endonuclease, 3'-5' exonuclease, DNA 3' repair diesterase, DNA 3'-phosphatase and RNase H activities as well as reductive activation of AP-1-binding proteins such as Fos and Jun (4, 7-15). However, at present, we have not enough information on actual *in vivo* functions and regulation of APEX nuclease.

Studies on changes of its expression *in vivo* depending on specific DNA damage, cell proliferation, cell differentiation or organ development are thought to be helpful to elucidate *in vivo* functions of the enzyme. We

\* To whom correspondence should be addressed.

<sup>†</sup> The nucleotide sequence data reported in this paper appear in the GSDB, DDBJ, EMBL and NCBI nucleotide sequence databases under accession number D44495.

considered that rats are the well usable laboratory animal with appropriate organ sizes for studying difference of its expression in various conditions, organs or developmental stages. In the present paper, therefore, we cloned and sequenced rat Apex cDNA, and studied Apex mRNA expression in various rat tissues using the cDNA as a probe.

## Materials and Methods

**Materials.** The reagents used in these experiments were obtained from the following sources: restriction enzymes from Toyobo Biochem., Osaka, Japan; T4 DNA ligase from Promega Corp. (Madison, WI, USA); Taq DNA polymerase from Wako Pure Chemical Industries (Osaka, Japan); human glyceraldehyde 3-phosphate dehydrogenase (G3PDH) cDNA and a rat brain cDNA library in phage  $\lambda$ gt10 from Clontech Lab., CA, USA; PRISM Ready Reaction DyeDeoxy Terminator Cycle Sequencing Kit from Applied Biosystems (Mountain View, CA, USA); Megaprime DNA Labeling System, [ $\alpha$ - $^{32}$ P] dCTP (3000 Ci/mmol) and positively charged nylon membrane (Hybond-N+) from Amersham Japan, Tokyo; RNazol B from Biotech Lab., Texas, USA.

**Cloning and DNA sequencing.** The recombinant phages in the rat brain cDNA library (Clontech) were plated with *Escherichia coli* C600Hfl, and three replica nylon membranes per plate were prepared. The coding region of mouse Apex cDNA was amplified by polymerase chain reaction using a mouse cDNA clone (pExE2) (7) and appropriate oligonucleotide primers. A  $^{32}$ P-labeled probe was prepared using the amplified DNA fragments as the template and [ $\alpha$ - $^{32}$ P] dCTP (3000 Ci/mmol) as the labeled nucleotide by the random priming method (16) using the Megaprime DNA Labeling System. About  $7.5 \times 10^5$  independent clones of the rat brain cDNA library were screened with the  $^{32}$ P-labeled probe by the plaque hybridization technique (17).

DNA sequences were determined by the dideoxy-termination method (18) using an ABI 373A DNA Sequencer (Applied Biosystems, Japan).

**Animals and RNA extraction.** Male Donryu rats were purchased from Sizuoka Laboratories Center (Shizuoka, Japan), and maintained on a laboratory diet (RC4, Oriental Yeast Co., Tokyo, Japan) and water *ad libitum*. Rats were sacrificed under ether anesthesia on postnatal days 1, 7, 14, 21, 28, 147 (21 w) or

224 (32 w). At least 3 rats were used for each experimental point.

Rat organs to be used for total RNA isolation were flash-frozen in liquid nitrogen immediately after sacrifice and stored at  $-80^\circ\text{C}$ . The extraction of total RNA was performed using RNazol B as described previously (19, 20). The RNA concentration was determined spectrophotometrically, and the amount and property of RNA electrophoresed on a denaturing 1% agarose gel were monitored after ethidium bromide staining.

**Northern blot hybridization.** The RNA fractions were denatured with 6% formaldehyde, and electrophoresed at  $30\ \mu\text{g}/\text{lane}$  at 20V for 18h on a 1% agarose gel in the presence of 6% formaldehyde. After the electrophoresis, the gel was treated with 0.25M ammonium acetate, and then RNA on the gel was transferred overnight onto a positively charged nylon membrane (Hybond-N+) by the capillary blotting technique (17). To analyse the results of Northern blotting, three or four blotted membranes were prepared for each set of samples. The membranes were prehybridized at  $42^\circ\text{C}$  in a prehybridization buffer consisting of  $5 \times \text{SSPE}$  ( $1 \times \text{SSPE}$ : 150 mM NaCl, 10 mM  $\text{NaH}_2\text{PO}_4$ , 1 mM EDTA, pH 7.6),  $5 \times \text{Denhardt's reagent}$ , 0.5% SDS, 50% formamide and  $100\ \mu\text{g}/\text{ml}$  denatured salmon sperm DNA. Hybridization was performed by incubating the membranes at  $42^\circ\text{C}$  for 18h in the prehybridization buffer supplemented with a denatured  $^{32}$ P-labeled probe prepared using the cloned rat Apex cDNA. After removing the excess probe by washing, the membranes were processed for autoradiography.

The preparation procedure for the  $^{32}$ P-labeled G3PDH probe was the same as that for the Apex probe. After removal of the Apex probe, the RNA-blotting membranes were incubated at  $65^\circ\text{C}$  for 20h for hybridization with the denatured,  $^{32}$ P-labeled G3PDH probe in a buffer containing  $6 \times \text{SSC}$ ,  $5 \times \text{Denhardt's solution}$ , 0.5% SDS and  $100\ \mu\text{g}/\text{ml}$  salmon sperm DNA. The membranes were processed in the same way as that for hybridization with the Apex probe.

**Analysis of autoradiograms.** Quantitative analysis of the autoradiograms was performed using a computerized imaging analysis system using Fujix Bio-Imaging Analyzer BAS2000 (Fuji Photo Film, Tokyo, Japan) as described previously (21). The hybridized membranes were exposed for about 40h at room temperature to phosphor imaging plates. The exposed imaging plates were inserted into an image reading unit and

scanned with a fine laser beam. To compare Apex mRNA expression in various tissues, a mean value (arbitrary units), which was obtained by the analyzer, of Apex mRNA in 7-day-old rat livers was used as the standard for normalization of the data obtained in separate experiments and for comparing the values in different tissues and different developmental stages. Each datum is expressed as a percentage of the mean value in 7-day-old rat livers.

**Results**

**Cloning of rat Apex cDNA.** A rat brain cDNA library was screened with <sup>32</sup>P-labeled mouse Apex cDNA. Seven positive clones were isolated from 7.5 × 10<sup>5</sup> plaques. The insert of the positive clone was subcloned into the EcoRI site of the pUC18 plasmid vector. Among these positive clones, two clones having inserts of roughly the Apex mRNA size were selected and the inserts were sequenced. The inserts of the two clones consisted of an identical region in both clones and a non-identical region. The nucleotide sequence of the identical region was highly homologous to that of mouse Apex cDNA, and was thought to be downstream of the 5' non-coding sequence and the complete sequence of the coding and 3' non-coding regions. However, the upstream sequences of the two inserts were not identical to each other and were not homologous to that of mouse Apex cDNA. The *Sma* I/*Pma*CI and *Eco*RI fragment, which belongs to the identical region, was subcloned into pUC18 plasmid to generate the rat Apex cDNA clone pUC18-RAPEX80. The cDNA sequence of the clone pUC18-RAPEX80 was determined for both strands from overlapping DNA regions. The DNA sequencing identified a coding region of 951 nucleotides (nt) with 115 nt of 5' flanking and 155 nt of 3' flanking regions followed by a poly(A) tail at 17 nt downstream of an authentic polyadenylation signal, AATAAA (22) (Fig. 1). The cDNA for rat APEX nuclease (951 nt) encodes 317 amino acids.

**Analysis of deduced amino acid sequence of rat APEX nuclease.** The amino acid sequence of rat APEX nuclease is highly homologous to that of mouse APEX nuclease (98.4 % identity/317 residues), human APEX nuclease (93.1 % identity/318 residues), and bovine AP endonuclease 1 (92.1 % identity/318 residues). Amino acid variation in these sequences is mostly in the N-terminal 60 amino acids (Fig. 2). Recent-

ly, Wilson *et al.* (23) reported the amino acid sequence of a rat AP endonuclease (rAPEN), deduced from the cDNA cloned by using a rat testis gt11 cDNA library and mouse Apex cDNA. The rAPEN amino acid sequence is 99.1 % identical with that of rat APEX nuclease.

**Expression of Apex mRNA in various rat organs.** The levels of Apex mRNA in various rat

GTGGAAACAGACAGACTCCATTCCTTTGTGCAGTGAGGGCTCCCTGCCTCGTTGGGAGGC	60
AGCGTAGTAAACACTGCTTCGGTGTCTCCAGACGCCTAAGGGCTTCGTTCACAGCGATGCC	120
	M P 2
GAAGCGGGGAAGAGAGCGGCAGCGGAAGACGGGAAGAACCAGTCCGAGCCAGAGAC	180
K R G K R A A A E D G E E P K S E P E T	22
CAAGAAGAGTAAGGGGGCAGCAAAAGAAAACGTGAGAAGGAGGGCCGAGGAGGGCCCTGT	240
K K S K G A A K K T E K E A A G E G P V	42
CCTGTATGAGGACCCCTCCAGATCAGAAAACGTCAGCCAGTGGCAAACTCTGCCACACTCAA	300
L Y E D P P D Q K T S A S G K S A T L K	62
GATATGCTCCTGGAATGTGGATGGGCTTCGAGCCTGGATTAATAAGAAAGGCTTGGATTG	360
I C S W N V D G L R A W I K K K G L D W	82
GGTAAAGAAGAAGCACCAGACATCTTGTGCCTCCAAGAGACCAATGCTCAGAGAACA	420
V K E E A P D I L C L Q E T K C S E N K	102
ACTCCCGGCTGAAGTCAAGAGCTGCCTGGACTCACCCATCAGTACTGGTCAGCTCCATC	480
L P A E L Q E L P G L T H Q Y W S A P S	122
AGACAAGAAGGATATAGTGGTGTGGGCTACTTTCCGCCAATGCCCGCTCAAAAGTCTC	540
D K E G Y S G V G L L S R Q C P L K V S	142
TTATGGCATTGGTAGGAAGAACATGATCAAGAAGCCGGGTGATGTGGCTGAATTTGA	600
Y G I G E E E H D Q E G R V I V A E F E	162
GTCCCTTTATCTGGTAACAGCCTATGTTCCGAACGCAGGAAGGGGCTGGTAAAGACTGGA	660
S F I L V T A Y V P N A G R G L V R L E	182
GTACCGCAGCGATGGGATGAAGCCTTCAGAAAATTCTAAAGGACTTGGCTTCCCGGAA	720
Y R Q R W D E A F R K F L K D L A S R K	202
ACCTCTGTGCTGTGTGGGATCTCAATGTGGCTCATGAAGAAATAGACCITCGTAAACC	780
P L V L C G D L N V A H E E I D L R N P	222
CAAAGGAACAAAAGAATGCTGGTTTTACTCCCGAGGAGCCAAAGGCTTTGGGAAAT	840
K G N K K N A G F T P Q E R Q G F G E M	242
GCTACAGGCTGTACCCTGGCTGACAGCTTCCGGCATCTCTACCCCAACACTGCCTACGC	900
L Q A V P L A D S F R H L Y P N T A Y A	262
TTATACTTTCTGGACTTACATGATGAATGCCCGCTCTAAGAATGTGGTGGCCCTTGA	960
Y T F W T Y M M N A R S K N V G W R L D	282
TTACTTTTGTGCTCCTCCTCTCTTTTACCTGCTTTGTGCGACAGCAAGATCCGGTCCAA	1020
Y F L L S H S L L P A L C D S K I R S K	302
GGCTTTGGCAGTGACCACTGTCCCATCACCTTTACCTAGCACTGTGACACTCCCTCA	1080
A L G S D H C P I T L Y L A L ***	317
AGTAGCTTCATGCTGGGAAATAGCCTCCTCTCTCCAGGAGACCAGTGCCTTATCTCTTC	1140
TTACGGTGTTTACTCCCTCTAAACCAACTTCTGGTTTCCTTAAACAATCCAAGTAA	1200
<u>ATAAAGTCTACTTTTCAAC</u>	1221

**Fig. 1** Nucleotide sequence and the deduced amino acid sequence of the cDNA for rat APEX nuclease. The deduced amino acid sequence in the standard one-letter code is shown beneath the nucleotide sequence. Amino acids are numbered, starting with the first in-frame methionine as number 1. The translation termination codon and polyadenylation signal are indicated with triple asterisks and double underlines, respectively.

rApex	1	MPKRGKRAA-AEDGEEPKESEPETKKSCKGAACKTEKEAAGEGVPVLYEDPPDQKTSASGKSA	
mApex	1	.....K...D.....	.....P.....
BAP1	1	.....VV...A...T...A...AG...N...V...A.....	.....P.....
hApex	1	.....KG·V...D·LRT...A...T...ND.....	.....A.....H...P...P.....
rApex	60	TLKICSWNVVDGLRAWIKKKGLDWVKEEAPDILCLQETKCSENKLPaelQELPGLTHQYWS	
mApex	60	.....	.....V.....S...S.....
BAP1	61	.....	.....S.....
hApex	61	.....	.....S.....
rApex	120	APSDKEGYSGVGLLSRQCPLKVSYGIGEEHDQEGRVIVAEFESFILVTAYVFNAGRGLV	
mApex	120	.....	.....V.....
BAP1	121	.....	.....YDA·V.....
hApex	121	.....	.....D...D...V.....
rApex	180	RLEYRQRWDEAFRFLKDLASRKPLVLCGDLNVAHEEIDLNPKNKKNAGFTPQERQGF	
mApex	180	.....	.....
BAP1	181	.....	.....G.....
hApex	181	.....	.....G.....
rApex	240	GEMLQAVPLADSPFRHLYPNTAYAYTFWYMMNARSKNVGWRLDYFLLSHSLPALCDSKI	
mApex	240	..L.....	.....Q·V.....
BAP1	241	..L.....T.....	.....P.....
hApex	241	..L.....	.....P.....
rApex	300	RSKALGSDHCPITLYLAL	317
mApex	300	.....	317
BAP1	301	.....	318
hApex	301	.....	318

Fig. 2 Comparison of the amino acid sequence of rat APEX nuclease (rApex) with those of mouse APEX nuclease (mApex) (7), bovine AP endonuclease I (BAP1) (27) and human APEX nuclease (hApex) (8). Gaps (indicated by dashes) are introduced at position 10 of the rat and mouse APEX nucleases. Amino acids identical to rat APEX nuclease are shown as a dot.





Fig. 3 (Left) Northern blot analyses of mRNA expression of Apex and G3PDH genes in various organs of 7-day-old rats. Thirty micrograms of total RNA from various rat organs were electrophoresed on a 1% agarose gel containing 6% formaldehyde, blotted onto a nylon membrane, and hybridized first with the  $^{32}\text{P}$ -labeled rat Apex cDNA probe (A), and then with  $^{32}\text{P}$ -labeled G3PDH probe after detaching the Apex probe (B). To monitor the RNA samples, total RNA of each sample was electrophoresed at  $1\ \mu\text{g}/\text{lane}$  and stained with ethidium bromide (C). Numbers to the right in Figs. 3A and 3B indicate sizes of transcripts in thousands of nucleotides.

Fig. 4 (Right) Northern blot analyses of mRNA expression of Apex (A) and G3PDH (B) genes in various organs of 28-day-old rats. The Northern blotting, hybridization and monitoring of the RNA samples (C) were performed as described in Materials and Methods and in the legend to Fig. 3.




**Fig. 5** Levels of Apex mRNA in various organs of 7- and 28-day-old rats. Quantitative analyses of the autoradiograms were performed as described in Materials and Methods. To compare Apex mRNA expression in various organs, its expression level in livers in 7-day-old rats was used as the standard for normalization of the data obtained in separate experiments, and the percentage of Apex mRNA expression in an organ to the expression in livers were calculated. Each point indicates the mean values of 3 independent determinations. Vertical bars indicate the standard errors of the mean values.


organs (skeletal muscle, thymus, testis, spleen, liver, kidney, lung, heart and brain) were examined by Northern blot hybridization using  $^{32}\text{P}$ -labeled rat Apex cDNA as a probe. A 1.5 kb transcript for the Apex gene was detected in all organs examined (Figs. 3 and 4). The amounts of Apex mRNA in various organs relative to the amount in livers of 7-day-old rats were determined as described in Materials and Methods, and the results are shown in Fig. 5. Although the Apex gene is expressed ubiquitously, its expression level varied significantly from tissue to tissue and with the developmental stage of the tissues. In most organs examined, the Apex mRNA level at postnatal day 28 was slightly (skeletal muscle, thymus and brain) or markedly (spleen, liver, kidney, lung and heart) lower than that at postnatal day 7. Among these organs, the liver showed the lowest Apex mRNA expression, and the liver mRNA level in 28-day-old rats was one-sixth of that in 7-day-old rats. Testis showed the highest expression, and the amounts of Apex mRNA in testis on postnatal days 7 and 28 were 4 and 50 times higher, respectively, than those in liver on postnatal days

7 and 28.

The expression patterns of the G3PDH gene, which was used as an internal control for Northern blot hybridization, show that the relative expression levels in these organs at postnatal day 28 are similar to those at postnatal



**Fig. 6** Northern blot analysis of Apex mRNA expression in the developing rat testes. The Northern blotting of total RNA isolated from developing rat testes and hybridization with the  $^{32}\text{P}$ -labeled rat Apex cDNA probe (A), and monitoring of electrophoresis of the RNA samples (B) were performed as described in Materials and Methods and in the legend to Fig. 3. The sample examined was testis at postnatal day 7 in lane 1, 14 in lane 2, 21 in lane 3, 28 in lane 4, 35 in lane 5 and 42 in lane 6. (C) The same experiment was repeated 3 times. The results were analyzed as described in the legend to Fig. 5.



**Fig. 7** Northern blot analysis of Apex mRNA expression in the developing rat thymuses. The Northern blotting of total RNA isolated from developing rat thymuses and hybridization with the  $^{32}\text{P}$ -labeled rat Apex cDNA probe (A), and monitoring of electrophoresis of the RNA samples (B) were performed as described in Materials and Methods and in the legend to Fig. 3. The sample examined was total RNA from the thymus on postnatal day 7 in lane 1, day 14 in lane 2, day 21 in lane 3, day 28 in lane 4, day 147 in lane 5 and day 224 in lane 6. (C) Each experiment was performed in triplicate. The results were analyzed as described in the legend to Fig. 5.

day 7, except that the expression in skeletal muscle is strikingly high in 28-day-old rats (Figs. 3 and 4). Figures 3 and 4 also show that the expression patterns of G3PDH gene in various organs on postnatal days 7 and 28 are quite different from those of the Apex gene.

**Expression patterns of Apex mRNA during testis and thymus development.** The expression pattern of Apex mRNA during the rat testis development is shown in Fig. 6. The Apex mRNA expression in

rat testis was observed at a fairly high and constant level in 7-, 14- and 21-day-old rats. The expression increased sharply from postnatal day 21 to day 28, possibly in association with the increase of pachytene spermatocytes as discussed later, and finally attained the adult level which was almost 3 times and 1.5 times higher than the levels in testes of 7- and 21-day-old rats and of 28-day-old rats, respectively.

Postnatal changes of the Apex gene expression in the thymus are shown in Fig. 7. Although the variation in experimental data are relatively large, the result shows that the Apex gene expression in rat thymuses tends to be the highest in 14-day-old rats, and declines sharply at the beginning and then gradually to half of the highest expression.

## Discussion

APEX nuclease (also designated HAP1 (9), APE (10), or Ref-1 (11) gene product) is a multifunctional DNA repair enzyme which may be involved in DNA repair of AP sites and 3' blocked single-strand breaks. The enzyme is also known to function as a redox factor which stimulates the DNA binding activity of Fos-Jun heterodimers, Jun-Jun homodimers and HeLa cell AP-1 protein as well as that of several other transcription factors including NF- $\kappa$ B, Myb and members of ATP/CREB family (11, 13, 24-26).

Previous studies on the enzyme have been mostly restricted to *in vitro* studies. In the present investigations, cDNA cloning of APEX nuclease from a rat cDNA library and its sequencing were performed for analyzing *in vivo* functions of the enzyme. Apex gene expressions in various rat tissues and in developmental stages of testis and thymus were studied using the cloned rat Apex cDNA.

The amino acid sequence of rat APEX nuclease deduced from the rat Apex cDNA sequence exhibits intensive homology with those of mouse and human APEX nuclease and of bovine AP endonuclease 1 (BAP1), especially with that of mouse APEX nuclease (7-10, 27). During this study, Wilson *et al.* (23) reported cDNA cloning for the multifunctional rat AP endonuclease (rAPEN)/redox factor using mouse Apex cDNA as a probe. The amino acid sequence reported for rAPEN was almost identical (99.1% identity/317 amino acids) to that reported here for rat APEX nuclease. There are some differences in the cDNA and deduced



amino acid sequences between rAPEN and rat APEX nuclease, although both enzymes are thought to be identical. The rat Apex cDNA encodes 317 amino acids, whereas rAPEN encodes 316 amino acids. The alanine residue at position 9 in rat APEX nuclease is missing in rAPEN. The arginine residue at position 236 and the histidine residue at position 288 in rat APEX nuclease are the alanine and glutamine residues, respectively, in rAPEN. At present, we can not say whether the sequence difference between rat APEX nuclease and rAPEN is due to a polymorphism of the same enzyme. Just, we can say that two independent rat Apex cDNA clones tested showed the identical coding sequence as described in the results section.

Studies on *in vivo* expression of the rat Apex gene indicate that the gene is ubiquitously expressed house-keeping gene, and that the level of its expression varies depending on tissues and developmental stages. The expression pattern of the rat Apex gene in various tissues is similar to the patterns of DNA polymerase  $\beta$  gene (28, 39), poly(ADP-ribose) polymerase gene (30, 31) and the Xrcc-1 DNA repair gene (32, 33) in various rat and mouse tissues, although there are some differences among them. The almost parallel expression of the Apex, DNA polymerase  $\beta$ , poly(ADP-ribose) polymerase and Xrcc-1 genes in these tissues suggests that the genes coding for these enzymes may be coexpressed. The mouse RAD51 homolog gene (34) is also expressed highly in testis and thymus, similar to expression of the Apex gene. Wilson *et al.* (23) also reported that expression of the rat major AP endonuclease gene (rAPEN) was highest in the testis and thymus. The possible coexpression of the Apex, DNA polymerase  $\beta$  and Xrcc-1 genes may support the notion that these proteins are involved in X-ray-type short patch repair such as base excision repair and repair of radical-induced, single-strand DNA breaks.

We previously studied the expression of mouse Apex gene during mouse brain development by *in situ* and Northern blot hybridization, and showed that mouse Apex mRNA was expressed at high levels in the proliferative zone of various brain regions, with showing temporal and spatial changes of the expression in the embryonic and postnatal development (35). Its expression decreased in association with brain development to the basal expression level which was observed even in adulthood, with the exception of its high expression in the hippocampal formation.

The present experiments also showed a similar developmental decrease in Apex expression in most tissues. The expression pattern of the Apex gene in most of the rat tissues examined suggested that its expression decrease slightly or markedly in accordance with the development and reached adult levels which are quite different from tissue to tissue. The developmental change of its expression in thymus is phenomenologically associated with the proliferative change of the organ (36).

It is well-known that specific cell types appear in the seminiferous tubules at a well-defined age during postnatal development of the testis (28, 37). In 23- and 26-day-old rats, spermatocytes are seen at all phases of the long meiotic prophase. Increasing numbers of pachytene spermatocytes and spermatids are observed in all seminiferous tubules in 26-day-old rats. The adult cell pattern in seminiferous tubules is seen in 50-day-old rats (37). The Apex mRNA expression in testis increased markedly from postnatal day 21 to day 28 in association with the increase of pachytene spermatocytes, in which the recombination process occurs, and finally reached the adult high level.

The developmental changes and tissue or cellular variations of Apex gene expression suggest that APEX nuclease is multifunctional *in vivo*, as suggested by *in vitro* studies, and that it is involved in cellular processes such as growth, development, transcriptional regulation and recombination as well as base excision repair. To study further the *in vivo* functions of APEX nuclease, the rat Apex cDNA may provide a useful tool.

**Acknowledgments** We thank Dr. K. Tsutsui and Dr. S. Watanabe for their valuable discussion and Mr. T. Nakamura, Mr. T. Tabayashi and Ms. M. Taniyama for their technical assistance. This work was supported in part by a Grant-in-Aid for Scientific Research from the Japan Ministry of Education, Science, Sports and Culture, and in part by a Grant-in-Aid for Scientific Research on Priority Areas from the Ministry of Education, Science, Sports and Culture of Japan (Molecular Mechanism for Maintenance of Genetic Information, 0520101).

## References

1. Lindahl T and Nyberg B: Rate of depurination of native deoxyribonucleic acid. *Biochemistry* (1972) 11, 3610-3618.
2. Sancar A and Sancar GB: DNA repair enzymes. *Annu Rev Biochem* (1988) 57, 29-67.
3. Friedberg EC: DNA Repair. Freeman WH and company, New York (1985)
4. Demple B and Harrison L: Repair of oxidative damage to DNA: Enzymology and biology. *Annu Rev Biochem* (1994) 63, 915-48.
5. Teoule R: Radiation-induced DNA damage and its repair. *Int J Radiat Biol* (1987) 51, 573-589.

6. Loeb LA and Preston BD: Mutagenesis by apurinic/aprimidinic sites. *Annu Rev Genet* (1986) **20**, 201-230.
7. Seki S, Akiyama K, Watanabe S, Hatsushika M, Ikeda S and Tsutsui K: cDNA and deduced amino acid sequence of a mouse DNA repair enzyme (APEX nuclease) with significant homology to *Escherichia coli* exonuclease III. *J Biol Chem* (1991) **266**, 20797-20802.
8. Seki S, Hatsushika M, Watanabe S, Akiyama K, Nagao K and Tsutsui K: cDNA cloning, sequencing, expression and possible domain structure of human APEX nuclease homologous to *Escherichia coli* exonuclease III. *Biochim Biophys Acta* (1992) **1131**, 287-299.
9. Robson CN and Hickson ID: Isolation of cDNA clones encoding a human apurinic/aprimidinic endonuclease that corrects DNA repair and mutagenesis defects in *E. coli xth* (exonuclease III) mutants. *Nucleic Acids Res* (1991) **19**, 5519-5523.
10. Demple B, Herman T and Chen DS: Cloning and expression of APE, the cDNA encoding the major human apurinic endonuclease: Definition of a family of DNA repair enzymes. *Proc Natl Acad Sci USA* (1991) **88**, 11450-11454.
11. Xanthoudakis S, Miao G, Wang F, Pan YCE and Curran T: Redox activation of Fos-Jun DNA binding activity is mediated by a DNA repair enzyme. *EMBO J* (1992) **11**, 3323-3335.
12. Seki S, Ikeda S, Watanabe S, Hatsushika M, Tsutsui K, Akiyama K and Zhang B: A mouse DNA repair enzyme (APEX nuclease) having exonuclease and apurinic/aprimidinic endonuclease activities: Purification and characterization. *Biochim Biophys Acta* (1991) **1079**, 57-64.
13. Walker LJ, Robson CN, Black E, Gillespie D and Hickson ID: Identification of residues in the human DNA repair enzyme HAPI (Ref-1) that are essential for redox regulation of jun DNA binding. *Mol Cell Biol* (1993) **13**, 5370-5376.
14. Barzilay G, Walker LJ, Robson CN and Hickson ID: Site-directed mutagenesis of the human DNA repair enzyme HAPI: Identification of residues important for AP endonuclease and RNase H activity. *Nucleic Acids Res* (1995) **23**, 1544-1550.
15. Wilson DM III, Takeshita M, Grollman AP and Demple B: Incision activity of human apurinic endonuclease (Ape) at abasic site analogs in DNA. *J Biol Chem* (1995) **270**, 16002-16007.
16. Feinberg AP and Vogelstein B: A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal Biochem* (1984) **137**, 266-267.
17. Sambrook J, Fritsch EF and Maniatis T: *Molecular Cloning, A Laboratory Manual*, 2nd Ed, Cold Spring Harbor Laboratory Press, Cold Spring Harbor (1989).
18. Sanger F, Nicklen S and Coulson AR: DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci USA* (1977) **74**, 5463-5467.
19. Chomczynski P and Sacchi N: Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* (1987) **162**, 156-159.
20. Chomczynski P: Solubilization in formamide protects RNA from degradation. *Nucleic Acids Res* (1992) **20**, 3791-3792.
21. Nakagawa Y, Yanai K, Ryu JH, Kiyosawa M, Tamai M and Watanabe T: Marked increase in [<sup>3</sup>H] (R)  $\alpha$ -methylhistamine binding in the superior colliculus of visually deprived rats after unilateral enucleation. *Brain Res* (1994) **643**, 74-80.
22. Proudfoot NJ and Brownlee GG: 3' Non-coding region sequences in eukaryotic messenger RNA. *Nature* (1976) **263**, 211-214.
23. Wilson TM, Carney JP and Kelley MR: Cloning of the multifunctional rat apurinic/aprimidinic endonuclease (rAPEN)/redox factor from an immature T cell line. *Nucleic Acids Res* (1994) **22**, 530-531.
24. Xanthoudakis S and Curran T: Identification and characterization of Ref-1, a nuclear protein that facilitates AP-1 DNA-binding activity. *EMBO J* (1992) **11**, 653-665.
25. Mitomo K, Nakayama K, Fujimoto K, Sun X, Seki S and Yamamoto K: Two different cellular redox systems regulate the DNA-binding activity of the p50 subunit of NF- $\kappa$ B *in vitro*. *Gene* (1994) **145**, 197-203.
26. Okazaki T, Chung U, Nishishita T, Ebisu S, Usuda S, Mishiro S, Xanthoudakis S, Igarashi T and Ogata E: A redox factor protein, ref1, is involved in negative gene regulation by extracellular calcium. *J Biol Chem* (1994) **269**, 27855-27862.
27. Robson CN, Milne AM, Pappin DJC and Hickson ID: Isolation of cDNA clones encoding an enzyme from bovine cells that repairs oxidative DNA damage *in vitro*: homology with bacterial repair enzymes. *Nucleic Acids Res* (1991) **19**, 1087-1092.
28. Nowak R, Woszczyński M and Siedlecki JA: Changes in the DNA polymerase  $\beta$  gene expression during development of lung, brain, and testis suggest an involvement of the enzyme in DNA recombination. *Exp Cell Res* (1990) **191**, 51-56.
29. Nowak R, Siedlecki JA, Kaczmarek L, Zmudzka BZ and Wilson SH: Levels and size complexity of DNA polymerase  $\beta$  mRNA in rat regenerating liver and other organs. *Biochim Biophys Acta* (1989) **1008**, 203-207.
30. Ogura T, Takenouchi N, Yamaguchi M, Matsukage A, Sugimura T and Esumi H: Striking similarity of the distribution patterns of the poly (ADP-ribose) polymerase and polymerase  $\beta$  among various mouse organs. *Biochem Biophys Res Commun* (1990) **172**, 377-384.
31. Menegazzi M, Grassi-Zucconi G, Prati ACD, Ogura T, Poltronieri P, Nyunoya H, Shiratori-Nyunoya Y, Miwa M and Suzuki H: Differential expression of poly (ADP-ribose) polymerase and DNA polymerase  $\beta$  in rat tissues. *Exp Cell Res* (1991) **197**, 66-74.
32. Walter CA, Lu J, Bhakta M, Zhou ZQ, Thompson LH and McCarrey JR: Testis and somatic XRCC-1 DNA repair gene expression. *Somat Cell Mol Genet* (1994) **20**, 451-461.
33. Yoo H, Li L, Sacks PG, Thompson LH, Becker FF and Chan JYH: Alterations in expression and structure of the DNA repair gene XRCC1. *Biochem Biophys Res Commun* (1992) **186**, 900-910.
34. Morita T, Yoshimura Y, Yamamoto A, Murata K, Mori M, Yamamoto H and Matsushiro A: A mouse homolog of the *Escherichia coli* recA and *Saccharomyces cerevisiae* RAD51 genes. *Proc Natl Acad Sci USA* (1993) **90**, 6577-6580.
35. Ono Y, Watanabe M, Inoue Y, Ohmoto T, Akiyama K, Tsutsui K and Seki S: Developmental expression of APEX nuclease, a multifunctional DNA repair enzyme, in mouse brains. *Dev Brain Res* (1995) **86**, 1-6.
36. Fawcett DW: *Histology*. 11th Ed, WB Saunders Company, Philadelphia (1986) pp 436-447.
37. Clermont Y and Perey B: Quantitative study of the cell population of the seminiferous tubules in immature rats. *Am J Anat* (1957) **100**, 241-267.

---

Received December 22, 1995; accepted January 15, 1996.