

Chemistry

Physical & Theoretical Chemistry fields

Okayama University

Year 1992

Studies on the acid-base properties of the
ZnBr₂NaBr molten salt system

Hidetaka Hayashi
Okayama University

Zen-ichiro Takehara
Kyoto University

Kiyofumi Uno
Kyoto University

Akira Katagiri
Kyoto University

This paper is posted at eScholarship@OUDIR : Okayama University Digital Information Repository.

<http://escholarship.lib.okayama-u.ac.jp/physical.and.theoretical.chemistry/23>

8. T. Yoshida, K. Okabayashi, T. Assoka, and K. Abe, Abstract 380, p. 552, The Electrochemical Society Extended Abstracts, Vol. 88-2, Chicago, IL, Meeting, Oct. 9-14, 1988.
9. K. Honda, M. Fujita, H. Ishida, R. Yamamoto, and K. Ohgaki, *This Journal*, **135**, 3151 (1988).
10. T. Kase, T. Miyamoto, T. Yoshimoto, Y. Ohsawa, H. Inaba, and K. Nakasi, in *Large Area Chromogenics: Materials and Devices for Transmittance Control*, C. M. Lampert and C. G. Granqvist, Editors, p. 504, SPIE Optical Engineering Press, Bellingham, WA (1990).
11. H. Tada, Y. Bito, K. Fujino, and H. Kawahara, *Solar Energy Mater.*, **16**, 509 (1987).
12. R. D. Rauh and S. F. Cogan, *Solid State Ionics*, **28-30**, 1707 (1988).
13. P. Baudry, M. A. Aegerter, D. Deroo, and B. Valla, *This Journal*, **138**, 460 (1991).
14. S. F. Cogan, R. D. Rauh, Nguyet M. Nguyen, T. D. Plante, and J. D. Westwood, Submitted to *This Journal*.
15. R. B. Goldner, T. E. Haas, G. Seward, K. K. Wong, P. Norton, G. Foley, G. Berera, G. Wei, S. Schulz, and R. Chapman, *Solid State Ionics*, **28-30**, 1715 (1988).
16. S. F. Cogan, N. Nguyen, and R. D. Rauh, Abstract 733, p. 1045, The Electrochemical Society Extended Abstracts, Vol. 90-2, Seattle, WA, Meeting, Oct. 14-19, 1990.
17. K. H. Cheng and M. S. Whittingham, *Solid State Ionics*, **1**, 151 (1980).
18. P. G. Dickens and S. A. Kay, *ibid.*, **8**, 291 (1983).
19. S. F. Cogan, E. J. Anderson, T. D. Plante, and R. D. Rauh, *Appl. Optics*, **24**, 2282 (1985).
20. S. F. Cogan and R. D. Rauh, in *Large Area Chromogenics: Materials and Devices for Transmittance Control*, C. M. Lampert and C. G. Granqvist, Editors, p. 482, SPIE Optical Engineering Press, Bellingham, WA (1990).
21. S. Hub, A. Tranchant, and R. Messina, *Electrochim. Acta*, **33**, 997 (1988).
22. K. Ho, D. E. Singleton, and C. G. Greenberg, *This Journal*, **137**, 3858 (1990).
23. J. P. Randin, in *Large Area Chromogenics: Materials and Devices for Transmittance Control*, C. M. Lampert and C. G. Granqvist, Editors, p. 539, SPIE Optical Engineering Press, Bellingham, WA (1990).
24. R. B. Goldner and R. D. Rauh, in *Proc. SPIE*, **428**, p. 38 (1983).
25. R. D. Rauh and S. F. Cogan, in *Proceedings of 25th IECEC*, Vol. 4, p. 26 (1990).
26. S. F. Cogan, T. D. Plante, M. A. Parker, and R. D. Rauh, *Solar Energy Mater.*, **14**, 185 (1986).
27. S. F. Cogan, R. D. Rauh, T. D. Plante, N. M. Nguyen, and J. D. Westwood, in *Electrochromic Materials*, M. K. Carpenter and D. A. Corrigan, Editors, PV 90-2, p. 298, The Electrochemical Society Softbound Proceedings Series, Pennington, NJ (1990).
28. W. C. Dautremont-Smith, *Displays*, **3**, 3 (1982).
29. W. C. Dautremont-Smith, *ibid.*, **3**, 67 (1982).
30. J. D. E. McIntyre, W. F. Peck, Jr., and S. Nakahara, *This Journal*, **127**, 1264 (1980).
31. M. B. Armand, in *Proceedings of NATO Symposium on Materials for Advanced Batteries*, D. W. Murphey, J. Broadhead, and B. C. H. Steele, Editors, p. 145, Plenum Press, New York (1980).
32. B. E. Conway, *This Journal*, **138**, 1539 (1991).
33. R. D. Giglia and G. Haake, in *Proc. SID*, **23**, 41 (1982).
34. J. P. Randin, *This Journal*, **129**, 1215 (1982).

Studies on the Acid-Base Properties of the ZnBr₂-NaBr Molten Salt System

Hidetaka Hayashi*

Department of Applied Chemistry, Faculty of Engineering, Okayama University, Okayama 700, Japan

Kiyofumi Uno^a and Zen-ichiro Takehara*



Department of Industrial Chemistry, Faculty of Engineering, Kyoto University, Sakyo-ku, Kyoto 606, Japan

Akira Katagiri*

Faculty of Integrated Human Studies, Kyoto University, Sakyo-ku, Kyoto 606, Japan

ABSTRACT

The acid-base properties of the ZnBr₂-NaBr melts at 623 K were investigated on the basis of the electromotive force measurements of a zinc-zinc concentration cell. The following two chemical equilibria were postulated to describe the acid-base character of the melts



The equilibrium constants K_1 and K_2 were determined to be 5.0×10^2 and 1.0×10^2 , respectively, at 623 K. The acidity of ZnBr₂-NaBr melts is essentially weaker than that of bromoaluminate melts.

It has been pointed out that some kinds of oxyanions and complex ions which have unusual oxidation states are stable in acidic melts,¹ such as sodium chloroaluminate, because the concentration of "free" halide ion is very low.²⁻⁵ It is possible to obtain some metallic elements by the electroreduction of such unusual ions.⁶ Some chalcogen compounds are stable in acidic chloroaluminate melts and have

been tested as an active material for a rechargeable battery.⁷ Chloroaluminate melts are also potential materials for use as solvents for electro-organic synthesis.⁸

Zinc chloride and bromide are also Lewis acids, but their vapor pressure and reactivity with moisture are substantially lower than those of aluminum chloride and aluminum bromide. Many investigations have been performed on the structure and physical properties of zinc chloride-alkali metal chloride systems.⁹⁻²¹ However, to date very few studies^{15,18} have been reported concerning the fundamental properties of bromide systems. From this point of view,

* Electrochemical Society Active Member.



^a Present address: Sankyo Aluminium Industry Co. Ltd., Takaoka, Toyama 933, Japan.

EMF measurements of a zinc-zinc concentration cell were performed with ZnBr_2 -NaBr melts, and the acid-base equilibrium of this system was investigated.

Chemical species in ZnBr_2 -NaBr melts.—Since chlorides and bromides generally have similar properties, it is helpful to consult the literature on ZnCl_2 -MCl [M = alkali metal] systems.⁹⁻²¹

The polymeric nature or network structure of ZnCl_2 -MCl melts has long been recognized from their physical properties, such as density,²⁰ viscosity,¹⁹ and conductivity,¹² and from thermodynamic^{9,13,18} and Raman spectroscopic data.^{10,14,15,21} Thus, pure ZnCl_2 and ZnCl_2 -rich melts contain $(\text{ZnCl}_2)_n$ polymers or aggregates. These species are believed to consist of ZnCl_4 tetrahedra sharing corners as in crystalline ZnCl_2 . After increasing the temperature or after the addition of MCl, the $(\text{ZnCl}_2)_n$ polymers dissociate into smaller clusters. The degree of polymerization (the number n) depends on the temperature and composition of the melt. The $(\text{ZnCl}_2)_n$ polymers are considered to have peripheral groups, such as ZnCl_4^{2-} , ZnCl_3^- , and ZnCl_2 , attached through a chlorine atom. At high MCl/ ZnCl_2 ratios, zinc chloride tends to form a monomeric complex ZnCl_4^{2-} , as evidenced by Raman spectroscopy^{10,11,14,15,21} and EMF data of the Zn/ Cl_2 cell.^{9,13} Other species, such as monomeric ZnCl_2 ,¹⁵ ZnCl_3^- (with a planar configuration),^{10,15} and $\text{Zn}_2\text{Cl}_7^{2-}$,¹⁷ are also suggested. Similar polymers and complex ions are considered in the ZnBr_2 -MBr systems.^{15,18}

Because it is difficult to take into account all of the possible species, a simplification is attempted in order to describe the acid-base properties of ZnBr_2 -NaBr melts. Thus we postulate the following chemical constituents (hypothetical species) and the following equilibria among them




Here, ZnBr_2 represents not only the monomeric ZnBr_2 , but also the ZnBr_2 unit in various $(\text{ZnBr}_2)_n$ polymers. Similarly, ZnBr_3^- and ZnBr_4^{2-} represent both the simple complexes and peripheral groups attached to $(\text{ZnBr}_2)_n$ polymers. The equilibrium constants K_1 and K_2 are defined in terms of mole fractions of these hypothetical species. According to the stoichiometry, the dominant chemical species at various compositions can be estimated as shown in Table I. The border at 50 mole percent (m/o) NaBr corresponds to the equivalent point of Eq. 1, where $[\text{ZnBr}_2] = [\text{Br}^-]$. The border at 66.7 m/o corresponds to the equivalent point of Eq. 2, where $[\text{ZnBr}_3^-] = [\text{Br}^-]$.

Acid-base equilibria in the molten ZnBr_2 -NaBr system.— ZnBr_2 -NaBr melts are expected to exhibit Lewis acidity and to have a concentration dependent acid-base character. In bromide melts, "acid" is defined as a bromide ion acceptor, while "base" is defined as a bromide ion donor



The basicity of this system is indicated by the use of the $p\text{Br}^-$ value. [$p\text{Br}^- = -\log a(\text{Br}^-)$; $a(\text{Br}^-)$ means bromide ion activity.] The bromide ion activity can be determined by EMF measurements of the concentration cell in reference to the activity in a particular melt.

The zinc-zinc concentration cell described below was used for these EMF measurements



where "j" indicates the melt in the measuring electrode compartment, and "ref." means the reference melt which is saturated with NaBr. The activity changes were deduced from the relationship between EMF values and melt composition (j), then the acid-base equilibrium was discussed.

Experimental

Phase diagram of the ZnBr_2 -NaBr system.—No phase diagram data of the ZnBr_2 -NaBr system is available. In order to use this system as a solvent for electrosynthesis, it is necessary to know its liquid range. Therefore, the phase

Table I. Dominant species in ZnBr_2 -NaBr melts.

X(NaBr)	0	0.5	0.67	1
Stoichiometric composition	ZnBr_2	NaZnBr_3	Na_2ZnBr_4	NaBr
Dominant species	ZnBr_2 ZnBr_3^-		ZnBr_3^- ZnBr_4^{2-}	ZnBr_4^{2-} Br^-

diagram was determined by a simple thermal analysis method.

These experiments were performed in a purified argon atmosphere inside a glove box. Certain amounts of sodium bromide and zinc bromide were put in a high-purity alumina crucible. The crucible was placed in an electric furnace, and the temperature of the furnace was increased until the mixture completely melted. After holding the temperature for several hours, cooling was initiated. The temperature was recorded by using a Chromel-Alumel thermocouple. The phase transition points were detected as an inflection of the cooling curves.

EMF measurement.—Figure 1 shows the schematic diagram of the experimental cell. A NaBr saturated melt was put inside of the β -alumina tube which acts as a diaphragm to separate the reference compartment from the main compartment. Zinc wires (1 mm ϕ) of 99.99% purity were used for both electrodes. One was placed in the reference compartment, while the other was placed in the main compartment, as shown in the illustration. The melt composition in the main compartment was adjusted by adding sodium bromide or zinc bromide. Just after the salt addition, argon




Fig. 1. Experimental cell: a, zinc electrode (reference electrode); b, zinc electrode (measuring electrode); c, thermocouple; d, argon gas inlet; e, argon gas outlet; f, β -alumina tube; g, Pyrex container; h, ZnBr_2 -NaBr melt; i, ZnBr_2 -NaBr melt saturated with NaBr.




Fig. 2. Phase diagram for the $\text{ZnBr}_2\text{-NaBr}$ system: (○) this work (●) melting point of pure salt.

gas was bubbled through the melt to ensure complete mixing. All measurements were carried out in the argon atmosphere in a glove box.

Sodium bromide (Wako Chemical) and zinc bromide (Wako Chemical) had been dried at 573 K *in vacuo* in two days, separately. The temperature of the melt was controlled by PID electronic instruments at 623 K within ± 1 K. The EMF values were recorded by using a high impedance digital voltmeter.

Results and Discussion

Phase diagram of the $\text{ZnBr}_2\text{-NaBr}$ system.—Figure 2 shows the liquidus points thus obtained and the melting points of pure sodium bromide and zinc bromide (NaBr , 1020 K; ZnBr_2 , 667 K).²² In this measurement, the inflection at the solidus temperature was not detected, and therefore only the liquidus line is drawn in Fig. 2. In the vicinity of the equimolar point, the liquidus temperature is below 573 K. At 623 K, the liquid ranges expand from 30 to 70 m/o NaBr . Therefore, a temperature of 623 K was chosen for the following experiments.

EMF measurement.—Figure 3 shows the relationship between the observed EMF values and melt compositions. On decreasing the NaBr content in the melt, the EMF values are expected to become constant below 30 m/o, since zinc bromide starts to precipitate at about this composition. Above 70 m/o of NaBr , sodium bromide starts to precipitate, and the observed EMF values approach 0 V.

To simplify the discussion concerning the acid-base equilibria, let us start from A mole of ZnBr_2 and $1-A$ mole of NaBr . On melting the mixture, x moles of ZnBr_3^- and y moles of ZnBr_4^{2-} can be generated. The number of moles (n) of each chemical species in the equilibria can be described as follows

$$n(\text{Na}^+) = 1 - A \quad [4]$$

$$n(\text{ZnBr}_2) = A - x - y \quad [5]$$

$$n(\text{ZnBr}_3^-) = x \quad [6]$$

$$n(\text{ZnBr}_4^{2-}) = y \quad [7]$$

$$n(\text{Br}^-) = (1 - A) - x - 2y \quad [8]$$




Fig. 3. Relationship between the observed EMF and melt composition; (○) observed EMF, dotted line: calculated curve for $K_1 = 5.0 \times 10^2$, $K_2 = 1.0 \times 10^2$.

The total number of moles B is given by the equation below


$$B = 2 - A - x - 2y \quad [9]$$

Assuming that the activity is equal to the mole fraction of the species, the equilibrium constants of Eq. 1 and 2 are described by Eq. 10 and 11, respectively

$$K_1 = \frac{B \cdot n(\text{ZnBr}_3^-)}{n(\text{Br}^-) \cdot n(\text{ZnBr}_2)} = \frac{(2 - A - x - 2y) \cdot x}{(1 - A - x - 2y) \cdot (A - x - y)} \quad [10]$$

$$K_2 = \frac{B \cdot n(\text{ZnBr}_4^{2-})}{n(\text{Br}^-) \cdot n(\text{ZnBr}_3^-)} = \frac{(2 - A - x - 2y) \cdot y}{(1 - A - x - 2y) \cdot x} \quad [11]$$

The potential-determining reaction of zinc electrode and the Nernst equation can be described by Eq. 12 and 13, respectively



$$E = E^\circ - \frac{3RT}{2F} \cdot \ln \frac{n(\text{Br}^-)}{B} + \frac{RT}{2F} \cdot \ln \frac{n(\text{ZnBr}_3^-)}{B} \quad [13]$$

The EMF values of the concentration cell are expressed as

$$\begin{aligned} EMF_1 = & -\frac{3RT}{2F} \cdot \ln \frac{n(\text{Br}^-)_i}{B_i} \cdot \frac{B_{\text{ref}}}{n(\text{Br}^-)_{\text{ref}}} \\ & + \frac{RT}{2F} \cdot \ln \frac{n(\text{ZnBr}_3^-)_i}{B_i} \cdot \frac{B_{\text{ref}}}{n(\text{ZnBr}_3^-)_{\text{ref}}} \end{aligned} \quad [14]$$

where "i" and "ref" indicate the i th composition and reference composition, respectively. Equation 14 implicitly assumes that the membrane potential difference across the β -alumina separator is negligible. This approximation may be justified, as in the case of chloroaluminate melts,^{2,3,5} because the Na^+ ion is the only charge carrier in the β -alumina separator and the ratio of the Na^+ ion concentrations on both sides of the separator does not differ very much from unity.

Equations 10 and 11 give rise to two simultaneous equations in x_i and y_i . Therefore, both x_i and y_i can be described as functions of K_1 , K_2 , A_i

$$x_i = f(K_1, K_2, A_i)$$

$$y_i = g(K_1, K_2, A_i)$$

Substituting appropriate values for K_1 and K_2 yields x_i and y_i for a given value of A_i . The number of moles of each chemical species is calculated by Eq. 4 to 9. Then, the theoretical EMF value is calculated by using Eq. 14 for each melt composition. By changing K_1 and K_2 gradually, the





Fig. 4. Activity profiles in the ZnBr_2 -NaBr melt: (a) ZnBr_2 , (b) ZnBr_3^- , (c) ZnBr_4^{2-} , (d) Br^- .

above procedure is repeated until the difference between the observed and calculated EMF attains a minimum. Thus the most probable values of K_1 and K_2 can be obtained. Using this procedure, the values of K_1 and K_2 were determined to be 5.0×10^2 and 1.0×10^2 , respectively, at 623 K. The dotted line drawn in Fig. 3 indicates the calculated EMF values.

Figure 4 shows the activity profiles of each of the chemical species derived from the K_1 and K_2 values. The activity is indicated in units of mole fraction. It is clear from the illustration that the activity of bromide ion decreases gradually from 70 to 40 m/o NaBr. In the composition ranges near 50 m/o, ZnBr_3^- is the dominant species, whereas ZnBr_4^{2-} becomes dominant near 67 m/o.

The authors also investigated the acid-base properties of the AlBr_3 -NaBr melt and evaluated the equilibrium constant K_{Al} of the following reaction to be 2.5×10^5 at 483 K²³



The $p\text{Br}^-$ value of 40 m/o NaBr in AlBr_3 -NaBr melts at 483 K was 5.6, while the value in ZnBr_2 -NaBr melts at 623 K was 2.9; therefore, it is concluded that the acidity of the ZnBr_2 -NaBr system is essentially weaker than that of the bromoaluminate melts.

Conclusions

The acid-base properties of the ZnBr_2 -NaBr melts were examined on the basis of the EMF measurement of a zinc-zinc concentration cell in these melts. The following conclusions were obtained.

1. The equilibrium constants of the reactions shown in Eq. 1 and 2, were calculated to be

$$K_1 = 5.0 \times 10^2 \quad K_2 = 1.0 \times 10^2$$

2. The ZnBr_2 -NaBr system is less acidic than the bromoaluminate melt.

Acknowledgments

The authors wish to express their gratitude to the Asahi Glass Foundation for Industrial Technology and a Grant-in-Aid for Developmental Scientific Research from the Ministry of Education, Science and Culture of Japan for financial support.

Manuscript submitted April 14, 1992; revised manuscript received Oct. 19, 1992.

Okayama University assisted in meeting the publication costs of this article.

REFERENCES

- G. Mamantov and R. A. Osteryoung, *Characterization of Solute in Nonaqueous Solvents*, G. Mamantov, Editor, Chap. 11, p. 223, Plenum Press, New York (1978).
- B. Tremillon and G. Letisse, *J. Electroanal. Chem.*, **17**, 371 (1968).
- G. Torsi and G. Mamantov, *Inorg. Chem.*, **10**, 1900 (1971).
- A. A. Fannin, L. A. King, and D. W. Seegmiller, *This Journal*, **119**, 801 (1972).
- L. G. Boxall, H. L. Jones, and R. A. Osteryoung, *ibid.*, **120**, 223 (1973).
- H. Hayashi, N. Hayashi, K. Uno, Z. Takehara, and A. Katagiri, in *Proceedings of the Joint International Symposium on Molten Salts*, G. Mamantov, M. Blander, C. Hussey, C. Mamantov, M.-L. Saboungi, and J. Wilkes, Editors, PV 87-7, p. 814, The Electrochemical Society Softbound Proceedings Series, Pennington, NJ (1987).
- G. Mamantov, R. Marassi, M. Matsunaga, Y. Ogata, J. P. Wiaux, and E. J. Frazer, *This Journal*, **127**, 2319 (1980).
- H. L. Jones and R. A. Osteryoung, in *Advances in Molten Salt Chemistry*, Vol. 3, J. Braunstein, G. Mamantov, and G. P. Smith, Editors, Chap. 3, p. 121, Plenum Press, New York (1975).
- M. F. Lantratov and A. F. Alabyshev, *J. Appl. Chem. USSR*, **27**, 685 (1954).
- W. Bues, *Z. Anorg. Allg. Chem.*, **279**, 104 (1955).
- M. A. Bredig and E. R. Van Artsdalen, *J. Chem. Phys.*, **24**, 478 (1956).
- F. R. Duke and R. A. Fleming, *This Journal*, **104**, 251 (1957).
- B. F. Markof and S. V. Volkov, *Ukr. Khim. Zh.*, **30**, 341, 545 (1964).
- J. R. Moyer, J. C. Evans, and G. Y.-S. Lo, *This Journal*, **113**, 158 (1966).
- R. B. Ellis, *ibid.*, **113**, 485 (1966).
- C. Dijkhuis, R. Dijkhuis, and G. J. Janz, *Chem. Rev.*, **68**, 252 (1968).
- W. E. Smith, J. Brynestad, and G. P. Smith, *J. Chem. Phys.*, **52**, 3890 (1970).
- G. N. Papatheodorou and O. J. Kleppa, *Z. Anorg. Allg. Chem.*, **401**, 132 (1973).
- T. Ejima, T. Yoko, and K. Nakashima, *Nippon Kinzoku Gakkaishi*, **41**, 86 (1977).
- T. Yoko, R. Crescent, Y. Tsukagoshi, and T. Ejima, *Nippon Kinzoku Gakkaishi*, **42**, 1179 (1978).
- M. Itoh, K. Sakai, and T. Nakamura, *Inorg. Chem.*, **21**, 3552 (1982).
- I. Barin and O. Knacke, *Thermochemical Properties of Inorganic Substances*, pp. 522, 887, Springer-Verlag, Heidelberg (1973).
- H. Hayashi, N. Hayashi, Z. Takehara, and A. Katagiri, *This Journal*, **136**, 2606 (1989).