

Human Anatomy & Physiology

Physiology fields

Okayama University

Year 2008

Barley plasma membrane intrinsic
proteins (PIP aquaporins) as water and
CO₂ transporters

Maki Katsuhara*

Yuko T. Hanba†

*Research Institute for Bioresources, Okayama University, kmaki@rib.okayama-u.ac.jp

†Center for Bioresource Field Science, Kyoto Institute of Technology

This paper is posted at eScholarship@OUDIR : Okayama University Digital Information Repository.

<http://escholarship.lib.okayama-u.ac.jp/physiology/1>

(Short review for PA)

Original Ms No. EJP-00262-2007

Revised (R-1) at Nov., 2007

Title:

Barley Plasma membrane Intrinsic Proteins (PIP Aquaporins) as water and CO₂ transporters.

Authors: **Maki Katsuhara¹ and Yuko T. Hanba²**

Addresses:

1 Research Institute for Bioresources, Okayama University, Kurashiki 710-0046, Japan

2 Center for Bioresource Field Science, Kyoto Institute of Technology, Kyoto 616-8354, Japan

Correspondings: **Maki Katsuhara**

kmaki@rib.okayama-u.ac.jp

TEL +81-86-434-1221 FAX +81-434-1249

Abstract

We identified barley aquaporins, and demonstrated that one, HvPIP2;1, transports water and CO₂. Regarding water homeostasis in plants, regulations of aquaporin expression were observed in many plants under several environmental stresses. Under salt stress, a number of plasma-membrane type aquaporins were down-regulated, which can prevent continuous dehydration resulting in cell death. The leaves of transgenic rice plants that expressed the largest amount of HvPIP2;1 showed a 40% increase in internal CO₂ conductance compared with leaves of wild-type rice plants. The rate of CO₂ assimilation also increased in the transgenic plants. The goal of our plant aquaporin research is to determine the key aquaporin species responsible for water and CO₂ transport, and to improve plant water relations, stress tolerance, CO₂ uptake/assimilation, and plant productivity via molecular breeding of aquaporins.

Keywords

Barley, CO₂, Plant aquaporins, Salt stress, Water transport

Introduction

Water uptake is an indispensable function of plant roots for the survival of terrestrial plants. If water uptake through the roots is reduced or blocked by water-related stress such as drought, salt stress (salt accumulation in soil inducing osmotic stress), or low temperatures, plant growth is seriously or lethally inhibited. Physiological and agronomical attention has been focused on the mechanism of root water uptake and water transport to the shoot, as this determines cell growth and plant yield. Since water flow is determined as the product of the motive force (V) and water permeability (L_p), V and/or L_p should be adjectively regulated to maintain the growth of plants under water-related stress. Water motive force is described as the water potential difference between the inside and outside of a root cell. For decades, researchers have investigated the biochemical and molecular biological regulatory mechanisms of the internal water potential in several ways: via accumulations of ions and compatible solutes, or the opening/closing of plant stomata. At the molecular level, however, almost nothing concerning water permeability was considered until Dr. P. Agre discovered the first aquaporin in 1992. Soon after this first aquaporin (AQP1) from erythrocytes was reported, plant aquaporins were identified. Not long following this, it was established that L_p of the plasma membrane and tonoplasts (vacuolar membrane) in plant cells mainly depends on two types of plant aquaporins: plasma membrane intrinsic proteins (PIPs) and tonoplast intrinsic proteins (TIPs), respectively, although the membrane lipid compositions and other membrane proteins can also affect water permeability. In the plant genome, more than 30 aquaporin genes have been detected [11,21]. Some of them specifically transport water molecules, but others can mediate other low molecular weight compounds such as CO_2 [7, 8, 28], silicon [19], boron [24], ammonia [9], and H_2O_2 [2].

To date, it has been established that aquaporins play a critical role in the transport of many essential molecules in plants. Except for light reception, aquaporins partially or mainly contribute to the transport of water, minerals, and CO_2 that are essential for plant life. In this article, two functions of plant aquaporins are described: i) the regulation of water transport in various environments, and ii) CO_2 transport. The authors have investigated barley aquaporins, because barley is a crop showing fairly good tolerance to drought, salt stress, and low temperatures compared to many other crops, including rice. Analyses of barley aquaporins are providing a good insight into the molecular mechanisms involved in the transport of water and other essential molecules in *Gramineous* crops.

Water transport of barley PIPs

The peripheral plasma membrane is composed of two major membranes of plant cells together with the membrane of the vacuole (tonoplast). These two major serial conducting parts are involved in cellular hydraulic conductivity. Because L_p of the tonoplast is basically higher than that of the plasma membrane according to the abundance of TIPs [27], and because L_p of the whole system principally obeys the L_p of the lower conducting part (that is, the plasma membrane), PIPs are the most important factors regarding the characteristics of cellular water uptake/water loss. This is the reason why PIPs have been the most intensively investigated in terms of the water-relations of plant cells under various environmental conditions.

Three PIPs: *HvPIP2;1*, *HvPIP1;3*, and *HvPIP1;5*, were first identified and barley PIPs were analyzed by the authors and collaborators [12]. Absolute amounts of these 3 transcripts were determined in the roots of barley seedlings. It was revealed that transcripts of *HvPIP2;1* (22 million copies/ μg total RNA) were ten-fold more abundant than those of *HvPIP1;3* (2 million copies/ μg total RNA) and *HvPIP1;5* (1 million copies/ μg total RNA). Water transport activity of *HvPIP2;1* and *HvPIP1;3* was assayed in a *Xenopus laevis* oocyte heterologous expression system. The former (*HvPIP2;1*) markedly increased the water permeability (P_f) of oocytes, but *HvPIP1;3* did not [15]. This result is consistent with the general feature of plant PIP aquaporins [3], whereby aquaporins of the PIP2-subfamily show marked water transport activity in the *Xenopus* oocyte system but PIP1 aquaporins do not.

Under water-related stress, aquaporins play an important role in plant osmotic and ion homeostasis [16,18,27]. Phosphorylation of certain serine residues activates some PIPs, and dephosphorylation rapidly closes them [26]. This aquaporin inactivation can prevent water loss from cells under drought or strong salt stress [16]. This activation/inactivation can be effective for short-term (within a few hours) adjustment of the cellular water balance.

During several hours to days of salt or osmotic stress, the down-regulation of several aquaporins including *HvPIP2;1* was observed in many plant species [10,12,29,32]. A decrease in the level of aquaporins and reduction of L_p can prevent cell death due to continuous dehydration and gain time for intracellular osmotic adjustment. Consistently, continuous over-expression of *HvPIP2;1* increased the water permeability of roots and raised the salt sensitivity of transgenic rice plants [14]. In such transgenic plants, a lower down-regulation of cellular water permeability might induce a

relatively higher rate of water loss from roots/shoots, and result in death under salt stress with induced osmotic stress and dehydration.

Recently, the authors and collaborators detected novel barley aquaporin genes (Table 1) in addition to those previously reported, and a total of 10 barley PIP genes (*HvPIP*s) were preliminarily analyzed (Table 2). Among them, 5 genes were classified into the sub-class *PIP1* and the others into the sub-class *PIP2* according to the sequence homology. Transcript amounts of *HvPIP1*s, except for *HvPIP1;2*, were lower than *HvPIP2*s. However, the *HvPIP1;2* transcript was markedly more abundant (> 10-fold) than *HvPIP2;1*. Down-regulation of *HvPIP1;2* and many *HvPIP2*s due to salt stress was observed. Water transport activities of some *HvPIP2*s were detected in the *Xenopus laevis* oocyte heterologous expression system. Co-injection of *HvPIP1;2* and *HvPIP2;1* increased the Pf of oocytes, but such an enhancement of Pf was not observed when *HvPIP1;2* cRNA was injected alone. This activation mechanism of “heteromerization” was first proposed in maize PIPs [6]. Interactions between *ZmPIP1*s and *ZmPIP2*s were clearly demonstrated in living maize cells using a FRET imaging system [31]. At present, however, there is no information suggesting whether heteromerization regulation is or is not involved in the regulation of aquaporin activity and cellular water transport in plant cells under salt or osmotic stress.

In addition to salt and osmotic stress, many environmental factors markedly change the expression of plant aquaporins. Nutritional deficiency in general decreases the expression of *Lp* and aquaporin genes in roots [4], however, specific genes were induced under special conditions, such as *Arabidopsis* *NIP5;1* by boron-deficiency [24] and *Arabidopsis* *TIP2;1* by nitrogen-deficiency [17]. Low temperature [13], phytohormones [23], and far-red light [22] also regulated the expression of plant aquaporins. These observations indicate that adequate regulation of water homeostasis via aquaporins is common in plant cells adjusting to various environments.

Aquaporins as CO₂ transporters

In mammalian cells, there were some indications that CO₂ could permeate AQP1 [5,20], but Yang et al. raised a serious question over this interpretation[30]. Mostly negative data regarding animal aquaporins mediating CO₂ transport were discussed in a recent conference (The 5th International Aquaporin Conference, Nara), suggesting that sufficient CO₂ transport can be achieved by simple diffusion across the lipid bilayer, because a high CO₂ pressure gradient between the inside and outside of

cells is generated by respiration in animal cells. However, the simple diffusion of CO₂ without an efficient transport system seems to be inadequate to maintain photosynthesis (CO₂ assimilation) in chloroplasts within mesophyll cells because the atmospheric CO₂ concentration is low (0.03 %) and CO₂ is rapidly consumed internally if CO₂-fixing enzymes (RubisCO) function properly. Aquaporin is one candidate facilitating CO₂ transport.

Photosynthesis is the most basic and important function of plants. Except for chemoautotrophic bacteria, all plants and animals, including humans, depend on photosynthetic products. On the one hand, stomata are one of the limiting factors for CO₂ uptake, and the regulation of stomatal conductance has been investigated for many years. On the other, internal conductance (g_i), conductance regarding CO₂ diffusion from stomata to chloroplasts, is known to be another limiting step. The limitation of photosynthesis by g_i is often greater than that by stomata. Although many factors are involved in g_i , recent research has indicated that aquaporins are probably the most important factor determining g_i . In the pioneering work of Terashima and Ono [25], a significant decline of g_i in the presence of HgCl₂, an inhibitor of most of the aquaporins, was demonstrated.

The authors demonstrated that one of the barley aquaporins, HvPIP2;1, transports CO₂ in addition to water. Because the generation of transgenic barley was not established, *HvPIP2;1* was introduced into rice plants to be analyzed *in planta*. The leaves of transgenic rice plants that expressed the greatest amount of HvPIP2;1 showed a 40% increase in g_i compared with the leaves of wild-type rice plants [8]. This was the first evidence of a direct relation between aquaporins and g_i (Fig.1). Although Uehlein et al. showed that membrane permeability to CO₂ increased in *Xenopus* oocytes expressing the tobacco aquaporin NtAQP1 [28], they did not perform the measurement of g_i . Recently, Flexas et al. confirmed an increase in g_i of leaves of transgenic plants over-expressing NtAQP1[7].

In transgenic rice leaves showing a high g_i , CO₂ assimilation also increased [8]. A rise in g_i is supposed to effectively promote the CO₂ supply to the photosynthetic center in chloroplasts and result in a high CO₂ assimilation rate. The same phenomenon was observed in tobacco plants, that is, the CO₂ assimilation rate increased in leaves over-expressing AtPIP1;2 (PIP1b) [1], or NtAQP1[28]. Recently, we also observed increases in CO₂ assimilation in tobacco leaves over-expressing ice plant (*Mesembryanthemum crystallinum*) aquaporins (unpublished data). These results suggest that CO₂ assimilation is commonly limited by g_i , which depends on aquaporins, and the enhancement of aquaporin activity is a potentially promising way to promote

plant CO₂ assimilation via improving g_i .

Future direction

All aquaporin genes in *Arabidopsis thaliana* and rice have been identified [11,21]. However, profiles (substrate specificity, spatial and developmental expression pattern, and stress responses) of most plant aquaporins are waiting to be clarified. Although some aquaporins (AtPIP1;2, NtAQP1, or HvPIP2;1) showed CO₂ transport activities, as described previously, many aquaporin species have not yet been analyzed and characterized from the view point of CO₂ transport at present. It is possible that one of such aquaporins can show a higher CO₂ permeability and be the most important for CO₂ assimilation. It is also necessary to establish which aquaporin species is the most crucial for water uptake *in planta*. Therefore, first, all aquaporin species should be investigated, analyzed, and profiled. Second, special aquaporin species should be selected as targets for molecular engineering to improve plant water relations, stress tolerance, CO₂ uptake/assimilation, and plant productivity.

Acknowledgements

The authors are grateful to Dr. Shibasaka and graduate students as collaborators. Transgenic rice studies involved collaboration with Dr. Hayashi and Dr. Hayakawa (Plantech Research Institute). This manuscript summarized work supported by the Bio Design Program, CREST of the Japanese Science and Technology Corporation, and the Program for the Promotion of Basic Research Activities for Innovative Biosciences (PROBRAIN) to MK.

References

1. Aharon R, Shahak Y, Wininger S, Bendov R, Kapulnik Y, Galili G (2003) Overexpression of a plasma membrane aquaporin in transgenic tobacco improves plant vigor under favorable growth conditions but not under drought or salt stress. *Plant Cell* 15: 439–447 doi: 10.1105/tpc.009225
2. Bienert GP, Møller ALB, Kristiansen KA, Schulz A, Møller IM, Schjoerring JK, Jahn TP (2007) Specific aquaporins facilitate the diffusion of hydrogen peroxide

- across membranes. *J Biol Chem* 282:1183–1192 doi: 10.1074/jbc.M603761200
3. Chaumont F, Barrieu F, Jung R, Chrispeels MJ (2000) Plasma membrane intrinsic proteins from maize cluster in two sequence subgroups with differential aquaporin activity. *Plant Physiology* **122**, 1025-1034.
 4. Clarkson DT, Carvajal M, Henzler T, Waterhouse RN, Smyth AJ, Cooke DT, Steudle E (2000) Root hydraulic conductance: diurnal aquaporin expression and the effects of nutrient stress. *J Exp Bot* 51: 61-70
 5. Cooper GJ, Boron WF (1998) Effect of PCMBMS on CO₂ permeability of *Xenopus* oocytes expressing aquaporin 1 or its C189S mutant. *American J Physiol Cell Physiol* 275: 1481–1486
 6. Fetter K, Wider VV, Moshelion M, Chaumont F (2004) Interactions between plasma membrane aquaporins modulate their water channel activity. *Plant Cell* 16: 215–228 doi: 10.1105/tpc.017194.
 7. Flexas J, Ribas-Carbó M, Hanson DT, Bota J, Otto B, Cifre J, McDowell N, Medrano H, Kaldenhoff R (2006) Tobacco aquaporin NtAQP1 is involved in mesophyll conductance to CO₂ in vivo. *Plant J* 48:427–439 doi: 10.1111/j.1365-3113X.2006.02879.x
 8. Hanba YT, Shibasaka M, Hayashi Y, Hayakawa T, Kasamo K, Terashima I, Katsuhara M (2004) Overexpression of the barley aquaporin HvPIP2;1 increases internal CO₂ conductance and CO₂ assimilation in the leaves of transgenic rice plants. *Plant Cell Physiol* 45: 521–529
 9. Jahn TP, Møller ALB, Zeuthen T, Holm LM, Klaerke DA, Mohsin B, Kuhlbratndt WK, Schjoerring JK (2004) Aquaporin homologues in plants and mammals transport ammonia. *FEBS Letters* **574**, 31–36. doi: 10.1016/j.febslet.2004.08.004
 10. Jang JY, Kim DG, Kim YO, Kim JS, Kang H (2004) An expression analysis of a gene family encoding plasma membrane aquaporins in response to abiotic stresses in *Arabidopsis thaliana*. *Plant Mol Biol* 54: 713–725
 11. Johanson U, Karlsson M, Johansson I, Gustavsson S, Sjövall S, Fraysse L, Weig AR, Kjellbom P (2001) The complete set of genes encoding major intrinsic proteins in *Arabidopsis* provides a framework for new nomenclature for major intrinsic proteins in plants. *Plant Physiol.* 126: 1358–1369
 12. Katsuhara M, Akiyama Y, Koshio K, Shibasaka M, Kasamo K (2002) Functional analysis of water channels in barley roots. *Plant Cell Physiol* 43: 885–893
 13. Katsuhara M, Chung GC, Sakurai J, Murai M, Izumi Y, Tsumuki H (2007) Low temperature and aquaporins, a molecular mechanism of water transport. *Cryobiol Cryotech* 53:21-32 (2007)

14. Katsuhara M, Koshio K, Shibasaka M, Hayashi Y, Hayakawa T, Kasamo K (2003) Over-expression of a barely aquaporin increased the shoot/root ratio and raised salt sensitivity in transgenic rice plants. *Plant Cell Physiol* 44: 1378–1383
15. Katsuhara, M., Shibasaka, M. Barley root hydraulic conductivity and aquaporins expression in relation to salt tolerance. *Soil Sci Plant Nutr* 53: 466–470 (2007)
16. Kjellbom P, Larsson C, Johansson I, Karlsson M, Johanson U (1999) Aquaporins and water homeostasis in plants. *Trend Plant Sci* 4: 308–314
17. Liu LH, Ludewig U, Gassert B, Frommer WB, von Wiren N (2003) Urea transport by nitrogen-regulated tonoplast intrinsic proteins in Arabidopsis. *Plant Physiol* 133: 1220–1228
18. Luu DT, Maurel C (2005) Aquaporins in a challenging environment: molecular gears for adjusting plant water status. *Plant Cell Environ* 28: 85–96 doi: 10.1111/j.1365-3040.2004.01295.x
19. Ma JF, Tamai K, Yamaji N, Mitani N, Konishi S, Ishiguro M, Katsuhara M, Murata Y, Yano M (2006) Silicon transporter in rice. *Nature* **440**, 688–691. doi: 10.1038/nature04590
20. Nakhoul NL, Bruce AD, Romero MF, Boron WF (1998) Effect of expressing the aquaporin aquaporin-1 on the CO₂ permeability of *Xenopus* oocytes. *Amer J Physiol Cell Physiol* 274: 543–548
21. Sakurai J, Ishikawa F, Yamaguchi T, Uemura M, Maeshima M (2005) Identification of 33 rice aquaporin genes and analysis of their expression and function. *Plant Cell Physiol* 46: 1568–1577 doi: 10.1093/pcp/pci172
22. Sato-Nara K, Nagasaka A, Yamashita H, Ishida J, Enju A, Seki M, Shinozaki K, Suzuki H (2004) Identification of genes regulated by dark adaptation and far-red light illumination in roots of *Arabidopsis thaliana*. *Plant Cell Environ* 27:1387–1394
23. Suga S, Komatsu S, Maeshima M (2002) Aquaporin isoforms responsive to salt and water stresses and phytohormones in radish seedlings. *Plant Cell Physiol* 43: 1229–1237
24. Takano J, Wada M, Ludewig U, Schaaf G, Wirén N, Fujiwara T (2006) The Arabidopsis major intrinsic protein NIP5;1 is essential for efficient boron uptake and plant development under boron limitation. *Plant Cell* 18: 1498–1509 doi: 10.1105/tpc.106.041640
25. Terashima I, Ono K (2002) Effects of HgCl₂ on CO₂ dependence of leaf photosynthesis: evidence indicating involvement of aquaporins in CO₂ diffusion across the plasma membrane. *Plant Cell Physiol* 43: 70–78

26. Törnroth-Horsefield S, Wang Y, Hedfalk K, Johanson U, Karlsson M, Tajkhorshid E, Neutze R, Kjellbom P (2006) Structural mechanism of plant aquaporin gating. *Nature* 439: 688–659 doi: 10.1038/nature04316
27. Tyerman SD, Bohnert H, Maurel C, Steudle E, Smith JAC (1999) Plant aquaporins: their molecular biology, biophysics and significance for plant water relations. *J Exp Bot* 50: 1055-1071.
28. Uehlein N, Lovisolo C, Siefritz F, Kalenhoff R (2003) The tobacco aquaporin NtAQP1 is a membrane CO₂ pore with physiological functions. *Nature* 425: 734–737
29. Vandeleur R, Niemietz C, Tilbrook J, Tyerman SD (2005) Roles of aquaporins in root responses to irrigation. *Plant Soil* 274:141–161 doi: 10.007/s11104-004-8070-z
30. Yang X, Fukuda N, van Hoek A, Matthay MA, Ma TH, Verkman AS (2000) Carbon dioxide permeability of aquaporin-1 measured in erythrocytes and lung of aquaporin-1 null mice and in reconstituted proteoliposomes. *J Bio Chem* 25: 2686–2692.
31. Zelazny E, Borst JW, Muylaert M, Batoko H, Hemminga MA, Chaumont F (2007) FRET imaging in living maize cells reveals that plasma membrane aquaporins interact to regulate their subcellular localization. *Pro Nat Acad Sci* 104: 12359–12364 doi: 10.1073/pnas.0701180104
32. Zhu C, Schraut D, Hartung W, Schäffner AR (2005) Differential responses of maize MIP genes to salt stress and ABA. *J Exp Bot* 56: 2971–2981 doi: 10.1093/jxb/eri294

Table 1
 Number of barley aquaporin genes estimated and analyzed

	Estimated gene number	Identified with RT-PCR	Full length cDNA isolated and sequenced	Quantitative analysis of transcript	Swelling assay performed
PIPs	10	10	10	10	5
TIPs	8	5	2	5	2
NIPs	3	1	1	1	1
SIPs	2	0	0	0	0

Putative aquaporin genes were estimated as contigs identified from the HarvEST barley database (<http://harvest.ucr.edu/>). Data from November 2007.

Table 2

Expression profile of barley PIP transcripts in roots

	<i>PIP1</i> subfamily					<i>PIP2</i> subfamily				
	<i>1;1</i>	<i>1;2</i>	<i>1;3</i>	<i>1;4</i>	<i>1;5</i>	<i>2;1</i>	<i>2;2</i>	<i>2;3</i>	<i>2;4</i>	<i>2;5</i>
Amount of transcript in control roots	+	+++++	++++	++	++	+++	++++	++	+++	+++
Amount of transcript under salt stress	→	↓	↓	↓	→	↓	↓	↓	→	→

Transcripts were absolutely quantified using real-time PCR, and 200 mM NaCl was added to the hydroponic solution as salt stress up to 24 h. +, >10⁵ copies/μg total RNA: ++, >10⁶ copies/μg total RNA: +++, >10⁷ copies/μg total RNA: +++++, >10⁸ copies/μg total RNA: ++++++, >10⁹ copies/μg total RNA: →, no change (comparison with no-stress control): ↓, down-regulated.

Figure legend

Fig.1

Transport of ambient CO₂ to the chloroplast through several barriers. 1, leaf surface with stomata; 2, intercellular space; 3, cell wall; 4, plasma membrane of mesophyll cell; 5, cytoplasm; 6, chloroplast. An aquaporin can mediate CO₂ transport in the plasma membrane (right path) via simple diffusion (left path). Internal conductance of CO₂ involves steps 2 to 6 mentioned above. Tissue and cellular structures are not to scale.




Fig.1