

Mathematical Journal of Okayama University

Volume 36, Issue 1

1994

Article 9

JANUARY 1994

Structure of Rings Satisfying Certain Polynomial Identities and Commutativity Theorems

Isao Mogami*

*Tsuyama College of Technology

Copyright ©1994 by the authors. *Mathematical Journal of Okayama University* is produced by
The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

**STRUCTURE OF RINGS SATISFYING CERTAIN
POLYNOMIAL IDENTITIES AND
COMMUTATIVITY THEOREMS**

ISAO MOGAMI

0. Introduction. Throughout, all rings will mean associative rings which are not necessarily commutative. Moreover, \mathbf{Z} will represent the ring of rational integers, and by $\mathbf{Z}\langle X, Y \rangle$ will be meant the free algebra over \mathbf{Z} in two indeterminates. For positive integers n_1, \dots, n_r their greatest common divisor is denoted by (n_1, \dots, n_r) .

In [5], Y. Kobayashi defined an additive map Φ of $\mathbf{Z}\langle X, Y \rangle$ to \mathbf{Z} , and indicated that for $f(X, Y) \in \mathbf{Z}\langle X, Y \rangle$, $\Phi(f(X, Y))$ is closely related with the commutativity of rings with 1 and satisfying the polynomial identity $f(X, Y) = 0$, where Φ will be defined later. In [6], he turned his attention to the fact that $\Phi((XY)^n - X^nY^n) = -n(n-1)/2$ for $n > 1$, and investigated the structure of $n(n-1)/2$ -torsion free rings with 1 and satisfying the polynomial identity $(XY)^n - X^nY^n = 0$. Coincidentally, he proved the following ([6, Theorem]): Let R be a ring with 1. If $E(R) = \{n \in \mathbf{Z} \mid n > 0 \text{ and } (xy)^n = x^n y^n \text{ for all } x, y \in R\}$ contains integers $n_1, \dots, n_r \geq 2$ such that $(n_1(n_1-1)/2, \dots, n_r(n_r-1)/2) = 1$ and some of n_i 's is even, then R is commutative. In connection with the above theorem, Y. Kobayashi and the present author raised respectively the following conjectures:

Conjecture 0.1 ([7, Conjecture 1]). Let R be a ring with 1. If $E(R)$ contains integers $n_1, \dots, n_r \geq 2$ such that R is $(n_1(n_1-1)/2, \dots, n_r(n_r-1)/2)$ -torsion free and some of n_i 's is even, then R is commutative.

Conjecture 0.2 ([16, Conjecture (I)]). Let R be a ring with 1. If for each $x, y \in R$, there exist integers $n_i \geq 2$ ($i = 1, \dots, r$) such that $(n_1(n_1-1)/2, \dots, n_r(n_r-1)/2) = 1$ and some of n_i 's is even and such that $(xy)^{n_i} = x^{n_i} y^{n_i}$ ($i = 1, \dots, r$), then R is commutative.

In [8] and [9], Y. Kobayashi gave partial affirmative answers to the above conjectures. In §2 and §4 of the present paper, those results will be improved more precisely and satisfactorily.

Meanwhile, J. Grosen [2] generalized some known commutativity theorems for a ring with 1 and satisfying certain polynomial identities

by assuming that the identities hold merely for the elements of a certain subset of the ring rather than for all elements of the ring. Almost all the results obtained in [2] have been improved and sharpened in [13]. In §3 of the present paper, we shall prove some commutativity theorems for a ring with 1 and satisfying polynomial identities of the form $(XY)^n - X^nY^n = 0$ merely for the elements of a certain subset of the ring.

Recently, W. Streb [17] gave a classification of non-commutative rings. H. Komatsu and H. Tominaga applied the classification to the proof of some commutativity theorems, in [11], [12], [13] and [14]. In our subsequent study, we shall use frequently several results obtained in [12] and [14], which will be summarized in §1 together with notations employed in the present paper.

1. Preliminaries. Throughout the present paper, R will represent a ring with 1. We use the following notations. Let M be a non-empty subset of R , and k a positive integer.

$C = C(R)$ = the center of R .

$D = D(R)$ = the commutator ideal of R .

$N = N(R)$ = the set of all nilpotent elements in R .

$N^* = N^*(R) = \{x \in R \mid x^2 = 0\}$.

$J = J(R)$ = the Jacobson radical of R .

$U = U(R)$ = the set of units in R .

Q = the intersection of the set of non-units in R with the set of quasi-regular elements in $R = (1 + U) \setminus U (\supseteq N \cup J)$.

$C_R(M)$ = the centralizer of M in R .

$\text{Ann}_R(M) = \{x \in R \mid xM = Mx = 0\}$.

As usual, for $x, y \in R$, let $[x, y]_1 = [x, y] = xy - yx$, and define, recursively $[x, y]_k = [[x, y]_{k-1}, y]$ for all $k > 1$.

$Z\langle X, Y \rangle$ = the free algebra over Z in the indeterminates X and Y .

$K = Z\langle X, Y \rangle[X, Y]Z\langle X, Y \rangle$.

K_k = the set of all $f(X, Y) \in K$ each of whose monomial terms is of length $\geq k$ (together with 0).

W = the set of all words in X and Y , namely products of factors each of which is X or Y (together with 1).

As is well-known, $K = K_2$ coincides with the kernel of the natural homomorphism of $Z\langle X, Y \rangle$ onto $Z[X, Y]$. Let $f(X, Y) = \sum f_{ij}(X, Y)$ be a polynomial in $Z\langle X, Y \rangle$, where $f_{ij}(X, Y)$ is a homogeneous polynomial with degree i in X and degree j in Y . Then we can easily see that $f(X, Y)$

is in K if and only if for each i, j , the sum of the coefficients of $f_{ij}(X, Y)$ equals zero.

Following [5], we denote by Φ the additive map of $\mathcal{Z}\langle X, Y \rangle$ to \mathcal{Z} defined as follows: For each monic monomial $X_1 \cdots X_r$ (X_i is either X or Y), $\Phi(X_1 \cdots X_r)$ is the number of pairs (i, j) such that $1 \leq i < j \leq r$ and $X_i = X, X_j = Y$. We can easily see that, for any $f(X, Y) \in \mathcal{Z}\langle X, Y \rangle$, $\Phi(f(X, Y))$ equals the coefficient of XY occurring in $f(1 + X, 1 + Y)$. Now, let $f(X, Y) \in K$. Then $f(1 + X, 1 + Y) \in K$, and so there exists $g(X, Y) \in K_3$ such that $f(1 + X, 1 + Y) = \Phi(f(X, Y))[X, Y] + g(X, Y)$.

Further, we put

$$e(k) = \begin{cases} k & \text{if } k \text{ is even,} \\ k - 1 & \text{if } k \text{ is odd.} \end{cases}$$

We consider the following conditions:

(S) For each $x, y \in R$, there exists $f(X, Y) \in K_3$ such that $[x, y] = f(x, y)$.

Q(k) If $x, y \in R$ and $k[x, y] = 0$ then $[x, y] = 0$.

By [12, Theorem 1.2, Proposition 1.6 and Proposition 1.7], we obtain the next

Theorem 1.1. *Let R be a non-commutative ring with 1. Then there exists a factorsubring of R which is of type a)¹, b), c), d)¹ or e)¹:*

a)¹ $\begin{pmatrix} \text{GF}(p) & \text{GF}(p) \\ 0 & \text{GF}(p) \end{pmatrix}$, where p a prime number.

b) $M_\sigma(K) = \left\{ \begin{pmatrix} \alpha & \beta \\ 0 & \sigma(\alpha) \end{pmatrix} \mid \alpha, \beta \in K \right\}$, where K is a finite field with a non-trivial automorphism σ .

c) A non-commutative division ring.

d)¹ A domain which is generated by 1 and a simple radical subring.

e)¹ A ring $B = \langle 1, x, y \rangle$ with 1 such that $D(B)$ is the heart of B and $x, y \in \text{Ann}_B(D(B))$.

Now, let $\begin{pmatrix} \alpha & \beta \\ 0 & \sigma(\alpha) \end{pmatrix}$ be an element of $M_\sigma(K)$. Let $K^\sigma = \{\gamma \in K \mid \sigma(\gamma) = \gamma\}$. Then

$$(1.1) \quad \begin{pmatrix} \alpha & \beta \\ 0 & \sigma(\alpha) \end{pmatrix}^k = \begin{cases} \begin{pmatrix} \alpha^k & (\sigma(\alpha^k) - \alpha^k)(\sigma(\alpha) - \alpha)^{-1}\beta \\ 0 & \sigma(\alpha^k) \end{pmatrix} & \text{if } \alpha \notin K^\sigma, \\ \begin{pmatrix} \alpha^k & k\alpha^{k-1}\beta \\ 0 & \alpha^k \end{pmatrix} & \text{if } \alpha \in K^\sigma. \end{cases}$$

This formula will be used repeatedly in §2 and §4.

By [12, Proposition 1.3(2), Lemma 1.4(1) and (4), and Proposition 1.7], we obtain

Lemma 1.2. *Let R be a ring with 1. If $xy \neq 0 = yx$ for some $x, y \in R$, then there exists a factorsubring of R which is of type a)¹ or e)¹.*

Lemma 1.3 ([12, Lemma 2.1]). *Let R be a ring satisfying (S) such that $D \subseteq N$. Then there hold the following:*

- (1) N is a commutative ideal of R .
- (2) $C_R(N^*)$ is a maximal commutative subring of R .
- (3) $\text{Ann}_R([N^*, R])$ is the largest commutative ideal of R and is contained in $C_R(N^*)$.
- (4) For any non-empty subset M of N , $R/\text{Ann}_R([M, R])$ has no non-zero nil ideals.
- (5) Let $c \in N$, $x \in R$, k a positive integer, and p a prime number.
 - (i) If $x^k[c, x] = 0 = [c, x]x^k$ then $[c, x] = 0$.
 - (ii) If $[c, x]_k = 0$ then $[c, x] = 0$.
 - (iii) If $[c, px] = 0 = [c, x^p]$, then $[c, x] = 0$.
 - (iv) If the additive order of $[c, x]$ is finite, then it is square-free.

The next is included in [14, Proposition 2.9(2)].

Lemma 1.4. *Let R be a non-commutative subdirectly irreducible ring satisfying (S). Suppose that R satisfies the identity $[(XY)^n - X^nY^n, X] = 0$ with some $n > 1$. Then R is isomorphic to some $M_\sigma(K)$.*

The next is included in [14, Lemma 2.10(2)].

Lemma 1.5. *Let n be a positive integer. Let $R = M_\sigma(K)$, and put $t = (|K| - 1)/(|K^\sigma| - 1)$. If R satisfies the identity $[(XY)^{n+1} - X^{n+1}Y^{n+1}, X] = 0$, then t divides n or $n + 1$.*

Theorem 1.6 ([12, Theorem 3.6]). *Let R be a ring with 1, and n a positive integer. Then the following conditions are equivalent:*

- 1) R satisfies the identities $[X^n, Y^n] = 0$ and $[X - X^m, Y - Y^m] = 0$ for some $m > 1$.
- 2) R satisfies (S) and the identity $[X^n, Y^n] = 0$.
- 3) R is a subdirect sum of rings each of which has one of the following

types:

- i) *A commutative ring.*
- ii) $M_\sigma(K)$, where $(|K| - 1)/(|K^\sigma| - 1)$ divides n .

The next is included in [14, Theorem 2.12(II)].

Theorem 1.7. *Let R be a ring with 1, and n a positive integer. If $k = n(n + 1)/2$ is odd, then the following conditions are equivalent:*

- 1) *R satisfies $Q(k)$ and the identity $(XY)^n - Y^n X^n = 0$.*
- 2) *R satisfies $Q(k)$ and the identity $(XY)^{n+1} - X^{n+1} Y^{n+1} = 0$.*
- 3) *R is a subdirect sum of rings each of which has one of the following*

types:

- i) *A commutative ring.*
- ii) $M_\sigma(K)$, where $(|K| - 1)/(|K^\sigma| - 1)$ divides $e(n + 1)$ and $2K = 0$.

2. On Conjecture 0.1. Given $x, y \in R$, we denote by $E(x, y)$ the set of integers $n > 1$ such that $(xy)^n = x^n y^n$; and $\tilde{E}(x, y) = E(x, y) \cap E(y, x)$. For a positive integer n , an element x of a module G is said to be *n -torsion free* if the order of x is infinite or relatively prime to n . Obviously, every element of G is n -torsion free if and only if $nx = 0$ implies $x = 0$ for any $x \in G$.

The purpose of this section is to give a complete answer to Conjecture 0.1. In [8], Kobayashi proved the following theorem which is a partial answer to Conjecture 0.1.

Theorem A. *Let R be a ring with 1. If for any $x, y \in R$, $\tilde{E}(x, y)$ contains (at least one) even integers n_1, \dots, n_s and odd integers n_{s+1}, \dots, n_r ($r \geq s \geq 1$) such that $(n_1, \dots, n_s, n_{s+1} - 1, \dots, n_r - 1)$ is 2 (or a multiple of 4) and $[x, y]$ is $(n_1(n_1 - 1)/2, \dots, n_r(n_r - 1)/2)$ -torsion free, then R is commutative.*

In connection with the above theorem, in [9], he determined the structure of $n(n - 1)/2$ -torsion free rings with 1 satisfying the identity $(xy)^n = x^n y^n$, when n is a positive even integer. Recently, this result has been generalized by Komatsu and Tominaga (see [14, Theorem 2.12]). The main theorems of this section can be stated as follows:

Theorem 2.1. *Let R be a ring with 1. Suppose that, for each $x, y \in$*

R , $\tilde{E}(x, y)$ contains n_1, \dots, n_s such that $(e(n_1), \dots, e(n_s)) \equiv 0 \pmod{4}$ and $[x, y]$ is $(n_1(n_1 - 1)/2, \dots, n_s(n_s - 1)/2)$ -torsion free. Then R is commutative.

Theorem 2.2. Let R be a ring with 1, and n a positive integer such that $n \equiv 2 \pmod{4}$. Then the following conditions are equivalent:

- 1) R satisfies $Q(n(n - 1)/2)$ and the identity $(XY)^n - X^n Y^n = 0$.
- 2) R satisfies $Q(n(n + 1)/2)$ and the identity $(XY)^n - Y^n X^n = 0$.
- 3) R satisfies $Q(n(n + 1)/2)$ and the identity $(XY)^{n+1} - X^{n+1} Y^{n+1} = 0$.
- 4) For each $x, y \in R$, $\tilde{E}(x, y)$ contains n_1, \dots, n_s and m_1, \dots, m_r such that $(e(n_1), \dots, e(n_s)) = n$ and $[x, y]$ is $(m_1(m_1 - 1)/2, \dots, m_r(m_r - 1)/2)$ -torsion free.
- 5) R is a subdirect sum of rings each of which has one of the following types:
 - i) A commutative ring.
 - ii) $M_\sigma(K)$, where K is a finite field of characteristic 2 with a non-trivial automorphism σ such that $(|K| - 1)/(|K^\sigma| - 1)$ divides $n/2$.

In preparation for proving our theorems, we state the next lemma.

Lemma 2.3. Let R be a ring with 1. Suppose that, for each $x, y \in R$, $\tilde{E}(x, y)$ contains m_1, \dots, m_r such that $[x, y]$ is $(m_1(m_1 - 1)/2, \dots, m_r(m_r - 1)/2)$ -torsion free. Then there hold the following:

- (1) $D \subseteq N$.
- (2) $2[N, R] = 0$, namely $2N \subseteq C$.
- (3) R satisfies (S).
- (4) R is completely reflexive, namely $xy = 0$ implies $yx = 0$ for any $x, y \in R$.
- (5) Let $a \in N$, and $x \in R$. If $n \in \tilde{E}(1 + a, x)$ then $[a, x^{e(n)}] = 0$.

Proof. In preparation for proving (1), we state three claims.

Claim 1. If $x, y \in R$ and $n \in \tilde{E}(x, y)$, then $y[x^n, y^{n-1}]y = 0$.

Proof. Actually, $y[x^n, y^{n-1}]y = yx^n y^n - y^n x^n y = y(xy)^n - (yx)^n y = 0$.

Claim 2. For each $x, y \in R$, and for each positive integer k , there exists a positive integer m and $f(X, Y) \in \mathbf{K}_k$ such that $m[x, y] = f(x, y)$

and $[x, y]$ is m -torsion free.

Proof. There exist positive integers m_1, \dots, m_r in $\tilde{E}(1+x, 1+y)$ such that $[x, y] = [1+x, 1+y]$ is m -torsion free, where $m = (m_1(m_1 - 1)/2, \dots, m_r(m_r - 1)/2)$. Then we can easily see that there exist $f_i(X, Y) \in K_3$ such that

$$\begin{aligned} 0 &= (1+x)^{m_i}(1+y)^{m_i} - \{(1+x)(1+y)\}^{m_i} \\ &= \frac{m_i(m_i - 1)}{2}[x, y] + f_i(x, y). \end{aligned}$$

Hence we obtain $m[x, y] = f(x, y)$ with some $f(X, Y) \in K_3$. Claim 2 is an easy consequence of this fact.

Claim 3. N is a commutative ideal of R .

Proof. Let $a \in N$, and $x \in R$. Obviously, $a^{2^s} = 0$ for some positive integer s . Now, choose $n_1 \in \tilde{E}(a, x)$, and inductively $n_{i+1} \in \tilde{E}(a^{n_1 \cdots n_i}, x^{n_1 \cdots n_i})$ ($i = 1, \dots, s-1$). Then $n_1 \cdots n_s$ is in $\tilde{E}(a, x)$ and $a^{n_1 \cdots n_s} = 0$, and so both ax and xa are in N . Now let $c \in N$, and suppose that $ac = 0$. Then $cxa = [c, xa] = 0$. If not, Claim 2 shows that there exists a positive integer m such that $0 \neq m[c, xa] = f(c, xa)$ with some $f(X, Y) \in K_{2n}$, where n is a positive integer such that $c^n = 0 = (xa)^n$. But this forces a contradiction that $f(c, xa) = 0$. We have thus seen that if $c \in N$ and $ac = 0$ then $cRa = 0$. Now, let $b \in N$, and $a^\nu = 0 = b^\nu$. Then, by the above, we see that $a^i Ra^j = 0$, provided $i + j \geq \nu$. Further, we see that $a^i Ra^j Ra^k = 0$, provided $i + j + k \geq \nu$. Continuing this procedure, we obtain eventually $(a)^\nu = 0$; similarly, $(b)^\nu = 0$. If $[a, b] \neq 0$ then Claim 2 shows that there exists a positive integer μ such that $0 \neq \mu[a, b] = g(a, b)$ with some $g(X, Y) \in K_{2\nu}$. But, as is easily seen, $g(a, b) = 0$. This contradiction shows that N is commutative, and therefore N forms a commutative ideal.

(1) Note that N is an ideal by Claim 3. It suffices to show that R/N is commutative. Let $x, y \in R$. Then, by Claim 1, $y[x^n, y^{n-1}]y = 0$ for some $n > 1$. Since $y[x^n, y^{n-1}]$ and $[x^n, y^{n-1}]y$ belong to N , we see that

$$[x^n, y^{2(n-1)}] = y^{n-1}[x^n, y^{n-1}] + [x^n, y^{n-1}]y^{n-1} \in N.$$

Hence R/N is commutative, by [3, Theorem].

Claim 4. Let $a \in N$, $x \in R$, and n a positive integer. If $x^n[a, x] = 0 = [a, x]x^n$, then $[a, x] = 0$.

Proof. It is easy to see that $[a, \langle x \rangle^{2n}] = 0$. Suppose, to the contrary, that $[a, x] \neq 0$. Then Claim 2 shows that there exists a positive integer m and $f(X, Y) \in K_{2n}$ such that $0 \neq m[a, x] = f(a, x)$. Since $\langle x \rangle^{2n} \subseteq C_R(a)$, N is commutative and $N^2 \subseteq C$ (Claim 3), we can easily see that $f(a, x) = 0$. But this is a contradiction.

Claim 5. If $a \in N$, $x \in R$, and $n \in \tilde{E}(1+a, x)$, then $nx[a, x^{n-1}]x = 0$, $(n-1)[a, x^n] = 0$ and $n(n-1)[a, x] = 0$.

Proof. Noting that $N^2 \subseteq C$ by Claim 3, we see that

$$nx[a, x^{n-1}]x = x[(1+a)^n, x^{n-1}]x = 0$$

by Claim 1, and

$$\begin{aligned} (n-1)[a, x^n] &= -(1+a)^{-1}\{(1+a)[x^n, (1+a)^{n-1}](1+a)\}(1+a)^{-1} \\ &= 0. \end{aligned}$$

From those above, we obtain

$$n(n-1)[a, x]x^n = n(n-1)[a, x^n]x - n(n-1)x[a, x^{n-1}]x = 0;$$

and similarly, $n(n-1)x^n[a, x] = 0$. Hence $n(n-1)[a, x] = 0$, by Claim 4.

(2) Let $a \in N$, and $x \in R$. Then $\tilde{E}(1+a, x)$ contains m_1, \dots, m_r such that $[a, x] = [1+a, x]$ is m -torsion free, where $m = (m_1(m_1-1)/2, \dots, m_r(m_r-1)/2)$. By Claim 5, $m_i(m_i-1)[a, x] = 0$ ($i = 1, \dots, r$); and so $2m[a, x] = 0$. Hence $2[a, x] = 0$.

(3) Let $x, y \in R$, and choose n in $\tilde{E}(x, y)$. Since $2D \subseteq 2N \subseteq C$ by (1) and (2), we see that

$$0 = 2y[x^n, y^{n-1}]y = 2(n-1)[x^n, y]y^n = 2n(n-1)x^{n-1}[x, y]y^n$$

by Claim 1. Similarly, for $n' \in \tilde{E}(x, 1+y)$, we obtain

$$2n'(n'-1)x^{n'-1}[x, y](1+y)^{n'} = 0.$$

Then, as is easily seen, $lx^k[x, y] = 0$ for some positive integers k, l . Repeating the above process, we see that $l'[x, y] = 0$ for some positive integer l' . Now, by Claim 2, there exists a positive integer m and $f(X, Y) \in K_3$ such that $m[x, y] = f(x, y)$ and $[x, y]$ is m -torsion free. Since the additive order of $[x, y]$ is finite, there exists an integer m' such that $[x, y] = m'm[x, y] = m'f(x, y)$.

(4) Obviously,

$$\{\epsilon_{11}(\epsilon_{12} + \epsilon_{22})\}^n - \epsilon_{11}^n(\epsilon_{12} + \epsilon_{22})^n = -\epsilon_{12} \neq 0$$

for any integer $n > 1$. Hence R has no factorsubring of type a)¹. On the other hand, R satisfies (S) by (3), and so R has no factorsubring of type e)¹. Hence, by Lemma 1.2, R is completely reflexive.

(5) By Claim 5, we see that $nx[a, x^{n-1}]x = 0 = (n-1)[a, x^n]$. If n is even then $[a, x^n] = 0$ by (2). On the other hand, if n is odd then $x[a, x^{n-1}]x = 0$ again by (2). Further, $x^2[a, x^{n-1}] = 0 = [a, x^{n-1}]x^2$, by (4). Now, Claim 4 shows that $[a, x^{n-1}] = 0$.

Proof of Theorem 2.1. By Lemma 2.3(1) and (3), $D \subseteq N$ and R satisfies (S). Hence $C_R(N)$ is commutative by Lemma 1.3(2). Now, let $a \in N$, and $x \in R$. Then $\tilde{E}(1+a, x)$ contains n_1, \dots, n_s such that $(\epsilon(n_1), \dots, \epsilon(n_s)) \equiv 0 \pmod{4}$ and $[a, x] = [1+a, x]$ is $(n_1(n_1-1)/2, \dots, n_s(n_s-1)/2)$ -torsion free. As is easily seen, $(\epsilon(n_1), \dots, \epsilon(n_s)) \equiv 0 \pmod{4}$ if and only if $(n_1(n_1-1)/2, \dots, n_s(n_s-1)/2)$ is even. Hence, in view of Lemma 2.3(2), we obtain $[a, x] = 0$. We have thus seen that $R = C_R(N)$, which is commutative.

Proof of Theorem 2.2. Obviously, 1) implies 4), and Theorem 1.7 shows that 2), 3) and 5) are equivalent. It suffices therefore to show that 4) \Rightarrow 5) \Rightarrow 1).

4) \Rightarrow 5). Suppose that R satisfies 4). Then, by Lemma 2.3(1) and (3), $D \subseteq N$ and R satisfies (S). Let $a \in N$, and $x \in R$. Then $\tilde{E}(1+a, x)$ contains n_1, \dots, n_s such that $(\epsilon(n_1), \dots, \epsilon(n_s)) = n$. Without loss of generality, we may assume that $n = -m_1\epsilon(n_1) - \dots - m_t\epsilon(n_t) + m_{t+1}\epsilon(n_{t+1}) + \dots + m_s\epsilon(n_s)$ with some $m_i \geq 0$. Put $k = m_1\epsilon(n_1) + \dots + m_t\epsilon(n_t)$ and $l = m_{t+1}\epsilon(n_{t+1}) + \dots + m_s\epsilon(n_s)$. By Lemma 2.3(5), $[a, x^{\epsilon(n_i)}] = 0$ ($i = 1, \dots, s$). Then we see that

$$[a, x^n]x^k = [a, x^{n+k}] = [a, x^l] = 0,$$

and similarly $x^k[a, x^n] = 0$. Hence $[a, x^n] = 0$, by Lemma 1.3(5)(i). Further, $C_R(N)$ is commutative by Lemma 1.3(2). Hence R satisfies the identity $[X^n, Y^n] = 0$. Now, we can apply Theorem 1.6 to see that R is a subdirect sum of a commutative ring and rings each of which is isomorphic to some $M_\sigma(K)$, where $(|K| - 1)/(|K^\sigma| - 1)$ divides n . Furthermore, by Lemma 2.3(2), $2R \subseteq C_R(N)$, which is commutative. Hence $4[R, R] = 0$,

and so we can easily see that $2K = 0$. Then $(|K| - 1)/(|K^\sigma| - 1)$ is odd and divides $n/2$.

5) \implies 1). It is easy to see that R satisfies $Q(n(n-1)/2)$. Suppose that $M_\sigma(K)$ is of type ii). Then, $\alpha^n \in K^\sigma$ for all $\alpha \in K$. Since, n is even and K is of characteristic 2, we can easily see that

$$\begin{pmatrix} \alpha & \beta \\ 0 & \sigma(\alpha) \end{pmatrix}^n = \begin{pmatrix} \alpha^n & 0 \\ 0 & \alpha^n \end{pmatrix}$$

for all $\alpha, \beta \in K$, by (1.1) in §1. Hence, $M_\sigma(K)$ satisfies the identity $(XY)^n - X^nY^n = 0$.

3. Commutativity theorems for rings satisfying the polynomial identities of the form $(XY)^n - X^nY^n = 0$ on certain subsets. In this section, we shall generalize some known commutativity theorems for a ring R satisfying the polynomial identities of the form $(XY)^n - X^nY^n = 0$ by assuming that the identities hold merely for the elements of a certain subset of R rather than for all elements of R .

Let k be a positive integer, and A a subset of R . We consider the following conditions:

$$P_0(k, A) \quad (xy)^k = x^k y^k \quad \text{for all } x, y \in A.$$

$$P_0^*(k, A) \quad (xy)^k = y^k x^k \quad \text{for all } x, y \in A.$$

The statements in the following theorem are included in [15, Theorem 2] and [18, Theorem 4], respectively.

Theorem B. *Suppose that a ring R with 1 satisfies $P_0(k, R)$ ($k = n, n + 2, n + 4$).*

(1) *If n is even, then R is commutative.*

(2) *If $x^4 \in C$ for all x in R , then R is commutative.*

More recently, Komatsu and Tominaga proved [13, Theorems 2.4 and 2.7] which encompass several results of Grosen [2]. From [13, Theorems 2.4 and 2.7], we readily obtain

Theorem C. (1) *Suppose that a ring R with 1 satisfies $P_0(k, R \setminus Q)$ ($k = m, m + 1, n, n + 1$). If R satisfies $Q((m, n))$, then R is commutative.*

(2) *Suppose that a ring R with 1 satisfies $P_0(n + 1, R \setminus Q)$ (or $P_0^*(n, R \setminus Q)$). If R satisfies $Q(n(n + 1))$, then R is commutative.*

Obviously, Theorem C(2) includes [10, Theorem 1(b) and Theorem 2(b)]. The first main theorem of this section is stated as follows:

Theorem 3.1. *Let R be a ring with 1. Let n_1, \dots, n_r be positive integers such that $(n_1(n_1 - 1)/2, \dots, n_r(n_r - 1)/2) = 1$. If R satisfies $P_0(n_i, R \setminus J)$ ($i = 1, \dots, r$), then R is commutative.*

In preparation for proving Theorem 3.1, we state the following two lemmas.

Lemma 3.2. *Let R be a ring with 1. Let k, m, n be non-negative integers, and $f: R \rightarrow R$ a function such that $f(x) = f(x+1)$ for all $x \in R$. If $f(x)(x+k)^m x^n = 0$ (or $x^n(x+k)^m f(x) = 0$) for all $x \in R$, then $(k+1)^{mn} f(x) = 0$. In particular, if $f(x)x^n = 0$ (or $x^n f(x) = 0$) for all $x \in R$, then $f(x) = 0$.*

Proof. Obviously,

$$\begin{aligned} 0 &= f(x)(x+k+1)^m(x+1)^n(x+k)^m x^{n-1} \\ &= (k+1)^m f(x)(x+k)^m x^{n-1}. \end{aligned}$$

Continuing this process, we get $(k+1)^{mn} f(x)(x+k)^m = 0$. Next, we obtain

$$\begin{aligned} 0 &= (k+1)^{mn} f(x)((x+k)+1)^m(x+k)^{m-1} \\ &= (k+1)^{mn} f(x)(x+k)^{m-1}. \end{aligned}$$

Continuing this process, we conclude that $(k+1)^{mn} f(x) = 0$.

Lemma 3.3. *Let R be a ring with 1. Suppose that R satisfies $P_0(n, R \setminus Q)$ ($n > 1$). Then, for each $u \in U$, $u^{n(n-1)} \in C$, and $D \subseteq N$. In particular, if R satisfies $P_0(k, R \setminus Q)$ ($k = n (\geq 1), n+2, n+4$), then $u^2 \in C$ for each $u \in U$.*

Proof. Let $u, v \in U$, and $x \in R \setminus Q$. Then

$$u[x^n, u^{n-1}]u = ux^n u^n - u^n x^n u = u(xu)^n - (ux)^n u = 0,$$

and so $[x^n, u^{n-1}] = 0$. In particular, $[v^n, u^{n-1}] = 0 = [u^n, v^{n-1}]$. Accordingly, $[v^n, u^{n(n-1)}] = 0 = [u^{n(n-1)}, v^{n-1}]$, whence $[u^{n(n-1)}, v] = 0$ follows.

Now, let $u \in U$, and $x \in R$. In case $u^{n-1}x \in Q$, by the above,

$$0 = [u^{n(n-1)}, 1 - u^{n-1}x] = -u^{n-1}[u^{n(n-1)}, x],$$

and so $[u^{n(n-1)}, x] = 0$. Similarly, in case $x \in Q$, we obtain $[u^{n(n-1)}, x] = 0$. Finally, we consider the case that $u^{n-1}x \notin Q$ and $x \notin Q$. Obviously, $x^{n-1} \notin Q$. Recalling that $[x^n, u^{n-1}] = 0$, we see that

$$\begin{aligned} (x^{n-1}u^{n-1}x)^n &= x^{n(n-1)}(u^{n-1}x)^n = x^{n(n-1)}u^{n(n-1)}x^n \\ &= x^{n^2}u^{n(n-1)}. \end{aligned}$$

On the other hand,

$$\begin{aligned} (x^{n-1}u^{n-1}x)^n &= x^{n-1}(u^{n-1}x^n)^{n-1}u^{n-1}x \\ &= x^{n-1}x^{n(n-1)}u^{(n-1)^2}u^{n-1}x \\ &= x^{n^2-1}u^{n(n-1)}x. \end{aligned}$$

Hence $x^{n^2-1}[u^{n(n-1)}, x] = 0$, and so $[u^{n(n-1)}, x] = 0$ by Lemma 3.2.

Now, it is easy to see that R satisfies the polynomial identity

$$\{(XY)^n - X^nY^n\}Z[(1 - X)^{n(n-1)}, (1 - Y)^{n(n-1)}] = 0.$$

But, no $M_2(\text{GF}(p))$, p a prime, satisfies the above identity, as a consideration of the following elements shows: $X = e_{11}$, $Y = e_{12} + e_{22}$, $Z = e_{21}$. Hence, $D \subseteq N$ by [1, Theorem 1] (or [4, Proposition 2]).

Noting that $(n(n-1), (n+2)(n+1), (n+4)(n+3)) = 2$, we can easily see the latter assertion.

Proof of Theorem 3.1. First, we shall show that J is commutative. Let $x, y \in J$. Then

$$\begin{aligned} 0 &= (1+x)^{n_i}(1+y)^{n_i} - \{(1+x)(1+y)\}^{n_i} \\ &= \frac{n_i(n_i-1)}{2}[x, y] + f_i(x, y), \end{aligned}$$

where $f_i(X, Y) \in K_3$. Since $(n_1(n_1-1)/2, \dots, n_r(n_r-1)/2) = 1$, we see that J satisfies the condition (S) (and $D \subseteq N$ by Lemma 3.3). By Lemma 3.3, $u^{n_i(n_i-1)} \in C$ for each $u \in U$ ($i = 1, \dots, r$), so that $u^2 \in C$. Now, let $a \in J$, and $d \in N^*$. Then $2d = 1 - (1-d)^2 \in C$, and $[d, a^2] = [d, (1-a)^2] + [2d, a] = 0$. Now, by Lemma 1.3(5)(iii) shows that $[d, a] = 0$. Hence $[N^*, J] = 0$. By Lemma 1.3(2), this implies that J is commutative.

Noting that J is a commutative ideal, we readily see that $[J, R]J = [J, RJ] = 0$. This enables us to see that if a is in J and x in R then

$$(xa)^{n_i} - x^{n_i}a^{n_i} = x\{(ax)^{n_i-1} - x^{n_i-1}a^{n_i-1}\}a \in x([a, x])a = 0.$$

Similarly, $(ax)^{n_i} - a^{n_i}x^{n_i} = 0$. We have thus seen that R satisfies $P_0(n_i, R)$. Repeating the argument employed at the opening of this proof, we see that R satisfies (S) (and $D \subseteq N$). Now, by Lemma 1.4, R is a subdirect sum of commutative rings and some $M_\sigma(K)$'s. Suppose, to the contrary, that $M_\sigma(K)$ appears as a factor of the subdirect sum. Then, by Lemma 1.5, $(|K| - 1)/(|K^\sigma| - 1)$ divides $n_i(n_i - 1)$ ($i = 1, \dots, r$), and so does 2. But this is impossible.

Corollary 3.4. *Let R be a ring with 1, and n a positive integer. If R satisfies $P_0(k, R \setminus J)$ ($k = n, n + 2, n + 4$), then R is commutative.*

Proof. It suffices to note that $(n(n - 1)/2, (n + 2)(n + 1)/2, (n + 4)(n + 3)/2) = 1$.

Theorem 3.1'. *Let R be a ring with 1. Let n_1, \dots, n_r be positive integers such that $(n_1(n_1 - 1)/2, \dots, n_r(n_r - 1)/2) = 1$. If R satisfies $P_0(n_i, R \setminus N)$ ($i = 1, \dots, r$), then R is commutative.*

Proof. Since $D \subseteq N$ by Lemma 3.3, N forms an ideal of R . Then, careful scrutiny of the proof of Theorem 3.1 shows that R is commutative.

Corollary 3.4'. *Let R be a ring with 1, and n a positive integer. If R satisfies $P_0(k, R \setminus N)$ ($k = n, n + 2, n + 4$), then R is commutative.*

Corollary 3.5. *Let R be a ring with 1.*

(1) *If there exist positive integers n_1, \dots, n_r with $(n_1(n_1 - 1)/2, \dots, n_r(n_r - 1)/2) = 1$ such that R satisfies $P_0(n_i, R)$ ($i = 1, \dots, r$), then R is commutative.*

(2) *If there exist positive integers m, n with $(m, n) = 1$ or 2 such that R satisfies $P_0(k, R)$ ($k = m, m + 1, n, n + 1$), then R is commutative.*

(3) *If there exists a positive integer n such that R satisfies $P_0(k, R)$ ($k = n, n + 2, n + 4$), then R is commutative.*

Needless to say, Theorem B is included in Corollary 3.5(3).

Now, by making use of Corollary 3.5, we shall prove the following two theorems, which are related with Theorem C.

Theorem 3.6. *Let R be a ring with 1. Suppose that R satisfies $P_0(k, R \setminus Q)$ ($k = n, n + 2, n + 4$).*

(1) *If n is even, then R is commutative.*

(2) *If $2[x, a] = 0$ implies $[x, a] = 0$ for each $a \in Q$ and $x \in R$, then R is commutative.*

Theorem 3.7. *Let R be a ring with 1. Suppose that R satisfies $P_0^*(k, R \setminus Q)$ ($k = n, n + 2, n + 4$). Then R is commutative.*

Proof of Theorem 3.6. In view of Corollary 3.5, it suffices to show that R satisfies $P_0(k, R)$ ($k = n, n + 2, n + 4$).

Let $u, v \in U$, and $x \in R$. Then $(vu)^2 \in C$, by Lemma 3.2. Hence

$$\begin{aligned} 0 &= (vu)^2 v^n u^n - v^{n+2} u^{n+2} = v^n (vu)^2 u^n - v^{n+2} u^{n+2} \\ &= v^{n+1} [u, v] u^{n+1}, \end{aligned}$$

and so $[u, v] = 0$. In particular, $[Q, u] = [u, 1 - Q] = 0$. From this we see that

$$(3.1) \quad (xu)^k = x^k u^k \quad (k = n, n + 2, n + 4).$$

(1) Noting that n is even and $[x, u^2] = 0$ by Lemma 3.3, from (3.1) we get

$$u^n x^n (xu)^2 = x^n u^n (xu)^2 = x^{n+2} u^{n+2} = u^n x^n x^2 u^2,$$

whence $x^{n+1}[x, u] = 0$ follows. By Lemma 3.2, $[x, u] = 0$. This proves that $Q \subseteq C$. Hence R satisfies $P_0(k, R)$.

(2) In view of (1), we may assume that n is odd. Then, noting that $[x, u^2] = 0$ by Lemma 3.3, from (3.1) we get

$$(xu)^2 x^n u^n = (xu)^{n+2} = x^{n+2} u^2 u^n = x u^2 x^{n+1} u^n,$$

whence $xu[x, u]x^n = 0$ follows. In this equation, replacing x by $x + 1$ and multiplying x^n from right, we obtain $[x, u](x + 1)^n x^n = 0$. By Lemma 3.2, $2^{n^2}[x, u] = 0$. Thus $[x, u] = 0$, and so $[x, Q] = 0$ by the hypothesis. Hence R satisfies $P_0(k, R)$.

Proof of Theorem 3.7. Obviously, R satisfies $P_0(k + 1, R \setminus Q)$ ($k = n, n + 2, n + 4$). By Lemma 3.3, $u^2 \in C$ for each $u \in U$. In case n is odd,

Theorem 3.6(1) guarantees the commutativity of R . Thus, henceforth, we may restrict our attention to the case that n is even. Let $u \in U$, and $x \in R$. Then, as was shown in the proof of Theorem 3.6, $[Q, u] = 0$. From this and $P_0^*(k, R \setminus Q)$, we see that

$$(3.2) \quad (xu)^k = u^k x^k \quad (k = n, n + 2, n + 4).$$

Noting that n is even and $[x, u^2] = 0$, from (3.2) we get

$$u^n x^n (xu)^2 = u^{n+2} x^{n+2} = u^n x^{n+2} u^2,$$

whence $x^{n+1}[x, u] = 0$ follows. By Lemma 3.2, $[x, u] = 0$. This proves that R satisfies $P_0^*(k, R)$, and also $P_0(k + 1, R)$. Hence R is commutative, by Corollary 3.5.

4. On Conjecture 0.2. In [8, Theorem 2], Y. Kobayashi proved the following theorem which gives an affirmative answer to Conjecture 0.2 in a somewhat weak form.

Theorem D. *Let R be a ring with 1. If for each $x, y \in R$, $\tilde{E}(x, y)$ contains integers n_1, \dots, n_r such that $(n_1(n_1 - 1)/2, \dots, n_r(n_r - 1)/2) = 1$ and some of n_i 's is even, then R is commutative.*

In this section, we shall prove a generaliation of Theorem D. We consider the following conditions:

(*) For each $x, y \in R$, there exist integers $k \geq 0, n > 1$ and words $w(x, y), w'(x, y) \in \mathbf{W}$ such that

$$w(x, y)\{(xy)^n - x^n y^n\}w'(x, y) = 0 = y^k\{(yx)^n - y^n x^n\}x^k.$$

(#) For each $x, y \in R$, there exist non-negative integers $r_1 \leq r_2 \leq r_3 \leq r_4 \leq r_5 \leq r_6 \leq r_7 \leq r_8$ with $1 < r_8$, positive integers n_i ($1 \leq i \leq r_8$), m_i ($r_2 + 1 \leq i \leq r_8$), l_i ($r_4 + 1 \leq i \leq r_8$), and words $w_i(x, y), w'_i(x, y) \in \mathbf{W}$ ($1 \leq i \leq r_8$) such that

$$(\#)_0 \quad (n_1(n_1 + 1)/2, \dots, n_{r_2}(n_{r_2} + 1)/2, m_{r_2+1}n_{r_2+1}, \dots, m_{r_4}n_{r_4}, l_{r_4+1}m_{r_4+1}n_{r_4+1}, \dots, l_{r_8}m_{r_8}n_{r_8}) = 1,$$

$$(\#)_1 \quad w_i(x, y)\{(xy)^{n_i} - y^{n_i}x^{n_i}\}w'_i(x, y) = 0 \quad (1 \leq i \leq r_1),$$

$$(\#)_2 \quad w_i(x, y)\{(yx)^{n_i} - x^{n_i}y^{n_i}\}w'_i(x, y) = 0 \quad (r_1 + 1 \leq i \leq r_2),$$

$$(\#)_3 \quad w_i(x, y)\{(x^{m_i}y^{m_i})^{n_i} - (y^{m_i}x^{m_i})^{n_i}\}w'_i(x, y) = 0 \quad (r_2 + 1 \leq i \leq r_3),$$

$$\begin{aligned}
 (\#)_4 \quad w_i(x, y)[x^{m_i}, y^{n_i}]w'_i(x, y) &= 0 & (r_3 + 1 \leq i \leq r_4), \\
 (\#)_5 \quad w_i(x, y)[x^{l_i}, (x^{m_i}y^{m_i})^{n_i}]w'_i(x, y) &= 0 & (r_4 + 1 \leq i \leq r_5), \\
 (\#)_6 \quad w_i(x, y)[x^{l_i}, (y^{m_i}x^{m_i})^{n_i}]w'_i(x, y) &= 0 & (r_5 + 1 \leq i \leq r_6), \\
 (\#)_7 \quad w_i(x, y)[y^{l_i}, (x^{m_i}y^{m_i})^{n_i}]w'_i(x, y) &= 0 & (r_6 + 1 \leq i \leq r_7), \\
 (\#)_8 \quad w_i(x, y)[y^{l_i}, (y^{m_i}x^{m_i})^{n_i}]w'_i(x, y) &= 0 & (r_7 + 1 \leq i \leq r_8).
 \end{aligned}$$

Now, the main theorem of this section is stated as follows:

Theorem 4.1. *Let R be a ring with 1. If R satisfies $(*)$ and $(\#)$, then R is commutative.*

According to Theorem 1.1, in order to complete the proof of Theorem 4.1, it suffices to prove the following two lemmas.

Lemma 4.2. *If R is of type a)¹, c) or d)¹, then R does not satisfy $(*)$.*

Lemma 4.3. *If R is of type b) or e)¹, then R does not satisfy $(\#)$.*

Proof of Lemma 4.2. First, assume that R is of type a)¹. Then, for any integers $k \geq 0$ and $n > 1$, we see that

$$e_{11}^k \{ (e_{11}(e_{12} + e_{22}))^n - e_{11}^n (e_{12} + e_{22})^n \} (e_{12} + e_{22})^k = -e_{12} \neq 0.$$

Hence R does not satisfy $(*)$.

Next, assume that R is of type c) or d)¹. Suppose, to the contrary, that R satisfies $(*)$. Now, let $x, y \in R$, and choose $k \geq 0$, $n > 1$ and $w(x, y), w'(x, y) \in \mathbf{W}$ such that

$$w(x, y)\{(xy)^n - x^n y^n\}w'(x, y) = 0 = y^k\{(yx)^n - y^n x^n\}x^k.$$

Since R is a domain, we see that $(xy)^n = x^n y^n$ and $(yx)^n = y^n x^n$. Therefore,

$$y[x^n, y^{n-1}]y = yx^n y^n - y^n x^n y = y(xy)^n - (yx)^n y = 0,$$

and hence $[x^n, y^{n-1}] = 0$. Now, [3, Theorem] forces a contradiction that R is commutative.

Proof of Lemma 4.3. First, assume that $R = M_\sigma(K)$. Let γ be a generating element of the multiplicative group of K , and put

$$x = \begin{pmatrix} \gamma & 0 \\ 0 & \sigma(\gamma) \end{pmatrix} \quad \text{and} \quad y = \begin{pmatrix} \gamma & 1 \\ 0 & \sigma(\gamma) \end{pmatrix}.$$

Suppose, to the contrary, that R satisfies $(\#)$. Then there exist non-negative integers $r_1 \leq r_2 \leq r_3 \leq r_4 \leq r_5 \leq r_6 \leq r_7 \leq r_8$ with $1 < r_8$, positive integers n_i ($1 \leq i \leq r_8$), m_i ($r_2 + 1 \leq i \leq r_8$), l_i ($r_4 + 1 \leq i \leq r_8$), and words $w_i(x, y)$, $w'_i(x, y) \in \mathbf{W}$ ($1 \leq i \leq r_8$) such that $(\#)_{0-(\#)_8}$ hold good. Since x and y are units in R , $(\#)_{1-(\#)_8}$ become

$$\begin{aligned} (xy)^{n_i} - y^{n_i}x^{n_i} &= 0 & (1 \leq i \leq r_1), \\ (yx)^{n_i} - x^{n_i}y^{n_i} &= 0 & (r_1 + 1 \leq i \leq r_2), \\ (x^{m_i}y^{m_i})^{n_i} - (y^{m_i}x^{m_i})^{n_i} &= 0 & (r_2 + 1 \leq i \leq r_3), \\ [x^{m_i}, y^{n_i}] &= 0 & (r_3 + 1 \leq i \leq r_4), \\ [x^{l_i}, (x^{m_i}y^{m_i})^{n_i}] &= 0 & (r_4 + 1 \leq i \leq r_5), \\ [x^{l_i}, (y^{m_i}x^{m_i})^{n_i}] &= 0 & (r_5 + 1 \leq i \leq r_6), \\ [y^{l_i}, (x^{m_i}y^{m_i})^{n_i}] &= 0 & (r_6 + 1 \leq i \leq r_7), \\ [y^{l_i}, (y^{m_i}x^{m_i})^{n_i}] &= 0 & (r_7 + 1 \leq i \leq r_8). \end{aligned}$$

Further, by making use of (1.1) in §1, we can easily see that for arbitrary positive integers l, m, n ,

$$\begin{aligned} (xy)^n - y^n x^n &= -\{(yx)^n - x^n y^n\} \\ &= (\sigma(\gamma^n) - \gamma^n)(\sigma(\gamma^{n+1}) - \gamma^{n+1})(\gamma^2 - \sigma(\gamma^2))^{-1} e_{12}, \\ (x^m y^m)^n - (y^m x^m)^n &= (\sigma(\gamma^{2mn}) - \gamma^{2mn})(\sigma(\gamma^m) - \gamma^m)(\sigma(\gamma^m) + \gamma^m)^{-1}(\gamma - \sigma(\gamma))^{-1} e_{12}, \\ &\text{provided } \gamma^{2m} \notin K^\sigma, \\ [x^m, y^n] &= (\sigma(\gamma^m) - \gamma^m)(\sigma(\gamma^n) - \gamma^n)(\gamma - \sigma(\gamma))^{-1} e_{12}, \\ [x^l, (x^m y^m)^n] &= -[y^l, (y^m x^m)^n] \\ &= (\sigma(\gamma^{2mn}) - \gamma^{2mn})(\sigma(\gamma^l) - \gamma^l)(\sigma(\gamma^m) + \gamma^m)^{-1}(\gamma - \sigma(\gamma))^{-1} \gamma^m e_{12}, \\ &\text{provided } \gamma^{2m} \notin K^\sigma, \\ [x^l, (y^m x^m)^n] &= -[y^l, (x^m y^m)^n] \\ &= (\sigma(\gamma^{2mn}) - \gamma^{2mn})(\sigma(\gamma^l) - \gamma^l)(\sigma(\gamma^m) + \gamma^m)^{-1}(\gamma - \sigma(\gamma))^{-1} \sigma(\gamma^m) e_{12}, \\ &\text{provided } \gamma^{2m} \notin K^\sigma. \end{aligned}$$

Therefore we obtain

$$\begin{aligned} \gamma^{n_i(n_i+1)} &\in K^\sigma & (1 \leq i \leq r_2), \\ \gamma^{2m_i n_i} &\in K^\sigma & (r_2 + 1 \leq i \leq r_4), \\ \gamma^{2l_i m_i n_i} &\in K^\sigma & (r_4 + 1 \leq i \leq r_8). \end{aligned}$$

Hence, by $(\#)_0$, we get $\gamma^2 \in K^\sigma$. But this is impossible.

Next, assume that R is of type e)¹. Then $\text{Ann}(D)$ contains a, b with $[a, b] \neq 0$. Put $x = 1 + a$ and $y = 1 + b$. Now, suppose, to the contrary, that R satisfies $(\#)$. Then there exist non-negative integers $r_1 \leq r_2 \leq r_3 \leq r_4 \leq r_5 \leq r_6 \leq r_7 \leq r_8$ with $1 < r_8$, positive integers n_i ($1 \leq i \leq r_8$), m_i ($r_2 + 1 \leq i \leq r_8$), l_i ($r_4 + 1 \leq i \leq r_8$), and words $w_i(x, y), w'_i(x, y) \in \mathbf{W}$ ($1 \leq i \leq r_8$) such that $(\#)_0 - (\#)_8$ hold good. For each positive integers l, m, n , we can easily see that

$$\begin{aligned} (xy)^n - y^n x^n &= (yx)^n - x^n y^n = \frac{n(n+1)}{2}[a, b], \\ (x^m y^m)^n - (y^m x^m)^n &= m^2 n [a, b], \\ [x^m, y^n] &= mn [a, b], \\ [x^l, (x^m y^m)^n] &= [x^l, (y^m x^m)^n] \\ &= -[y^l, (x^m y^m)^n] = -[y^l, (y^m x^m)^n] = lmn [a, b]. \end{aligned}$$

Recalling here that $\langle a, b \rangle [a, b] = 0 = [a, b] \langle a, b \rangle$, we readily obtain

$$\begin{aligned} \frac{n_i(n_i+1)}{2}[a, b] &= 0 & (1 \leq i \leq r_2), \\ m_i^2 n_i [a, b] &= 0 & (r_2 + 1 \leq i \leq r_4), \\ l_i m_i n_i [a, b] &= 0 & (r_4 + 1 \leq i \leq r_8). \end{aligned}$$

Hence $[a, b] = 0$ by $(\#)_0$, which is a contradiction.

Noting that $(xy)^n - x^n y^n = x\{(yx)^{n-1} - x^{n-1}y^{n-1}\}y$ for any positive integer n , we obtain the next as an immediate consequence of Theorem 4.1.

Corollary 4.4. *Let R be a ring with 1. Suppose that for each $x, y \in R$, there exist positive integers $r \leq s$ and $n_i > 1$ ($i = 1, \dots, s$) such that*

- 1) $(n_1(n_1 - 1)/2, \dots, n_s(n_s - 1)/2) = 1$,
- 2) $(xy)^{n_i} = x^{n_i}y^{n_i}$ ($i = 1, \dots, r$),
- 3) $(yx)^{n_i} = y^{n_i}x^{n_i}$ ($i = r, \dots, s$).

Then R is commutative.

Needless to say, Corollary 3.5 is a direct consequence of Corollary 4.4.

REFERENCES

- [1] H. E. BELL: On some commutativity theorems of Herstein, Arch. Math. **24** (1973), 34–38.
- [2] J. GROSEN: Rings satisfying polynomial identities or constraints on certain subsets, Thesis (University of California, Santa Barbara, 1988).
- [3] I. N. HERSTEIN: A commutativity theorem, J. Algebra **38** (1976), 112–118.
- [4] Y. HIRANO, Y. KOBAYASHI and H. TOMINAGA: Some polynomial identities and commutativity of s -unital rings, Math. J. Okayama Univ. **24** (1982), 7–13.
- [5] Y. KOBAYASHI: A note on commutativity of rings, Math. J. Okayama Univ. **23** (1981), 141–145.
- [6] Y. KOBAYASHI: The identity $(xy)^n = x^n y^n$ and commutativity of rings, Math. J. Okayama Univ. **23** (1981), 147–151.
- [7] Y. KOBAYASHI: Some polynomial identities and commutativity of rings. I, Proc. 14th Symp. on Ring Theory, Shinshu Univ. (1981), 1–8.
- [8] Y. KOBAYASHI: On two conjectures about commutativity of rings, Math. Japonica **29** (1984), 219–228.
- [9] Y. KOBAYASHI: On rings satisfying the identity $(xy)^n = x^n y^n$, Math. Japonica **29** (1984), 645–654; *ibid.* **31** (1986), 165–166.
- [10] H. KOMATSU and H. TOMINAGA: On commutativity of s -unital rings, Math. J. Okayama Univ. **28** (1986), 93–96.
- [11] H. KOMATSU and H. TOMINAGA: Chacron's condition and commutativity theorems, Math. J. Okayama Univ. **31** (1989), 101–120.
- [12] H. KOMATSU and H. TOMINAGA: On non-commutative algebras and commutativity conditions, Resultate Math. **18** (1990), 74–92.
- [13] H. KOMATSU and H. TOMINAGA: Commutativity theorems for rings with polynomial constraints on certain subsets, Bull. Austral. Math. Soc. **43** (1991), 451–462.
- [14] H. KOMATSU and H. TOMINAGA: Commutativity theorems for algebras and rings, Math. J. Okayama Univ. **33** (1991), 71–95.
- [15] I. MOGAMI: Note on commutativity of rings. III, Math. J. Okayama Univ. **23** (1981), 163–169.
- [16] I. MOGAMI: Note on commutativity of rings, Proc. 14th Symp. on Ring Theory, Shinshu Univ. (1981), 109–115.
- [17] W. STREB: Zur Struktur nichtkommutativer Ringe. Math. J. Okayama Univ. **31** (1989), 135–140.
- [18] C.-T. YEN: Some commutativity theorems for rings, Chinese J. Math. **9** (1981), 99–105.

TSUYAMA COLLEGE OF TECHNOLOGY
TSUYAMA 708, JAPAN

(Received December 17, 1992)