

Mathematical Journal of Okayama University

Volume 45, Issue 1

2003

Article 2

JANUARY 2003

Symbols of Pseudodifferential Operators Associated to Gevrey Kernel's Type

Mohammed Hazi*

*Ecole Normale Supérieure, France

Copyright ©2003 by the authors. *Mathematical Journal of Okayama University* is produced by
The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

Symbols of Pseudodifferential Operators Associated to Gevrey Kernel's Type

Mohammed Hazi

Abstract

In this article, we aim at proving the truthfulness of the inverse Theorem (1) of [5]. More precisely, we associated symbols of Gevrey type to pseudodifferential operators when the latter are given by their kernels.

KEYWORDS: Symbols of Gevrey, Kernels, Pseudo-differential operators.

Math. J. Okayama Univ. **45** (2003), 117–122

SYMBOLS OF PSEUDODIFFERENTIAL OPERATORS ASSOCIATED TO GEVREY KERNEL'S TYPE

MOHAMMED HAZI

ABSTRACT. In this article, we aim at proving the truthfulness of the inverse Theorem (1) of [5]. More precisely, we associated symbols of Gevrey type to pseudodifferential operators when the latter are given by their kernels.

1. INTRODUCTION

In [4], we gave a description of pseudodifferential operators when defined by their Kohn-Nirenberg's symbols or their kernels (cf. [2], [3]); and this in the C^∞ -case. It appeared that the two approaches are equivalent. In [1], it is shown that this equivalence remains also true in the analytic case. This paper is a continuation of [5], where we obtained that the definition by symbols implies the one by kernels. Here, we deal with the converse. And this, in the more fine context of Gevrey classes.

2. DEFINITIONS AND NOTATIONS

We recall three definitions as they were given in [5].

Definition 1. Let n be a non zero positive integer, Ω an open subset of \mathbb{R}^n and s any real number larger or equal to 1. A real function φ in $C^\infty(\Omega)$ is said of Gevrey class with order s if, for any compact subset $K \subset \Omega$, there exists constant $C_K > 0$ such that

$$(1) \quad \forall \alpha \in \mathbb{N}^n \quad \|D^\alpha f\| \leq C_K^{|\alpha|+1} (|\alpha|!)^s .$$

Definition 2. Given $n \in \mathbb{N}^*$, $m \in \mathbb{R}$, $s \geq 1$ and Ω an open subset of \mathbb{R}^n . We say that a real function $a = a(z, x)$ in $C^\infty(\Omega \times \mathbb{R}^n)$, is a symbol (or amplitude) of Gevrey type with class s on Ω if, and only if, it satisfies

For any compact subset $K \subset \Omega$, there exist three positive constants C_0 , C_1 , R such that

$$(2) \quad \left| D_\xi^\alpha D_z^\beta a(z, \xi) \right| \leq C_0 C_1^{|\alpha+\beta|} (|\alpha|!)^s (|\beta|!)^s \left(1 + |\xi|^2 \right)^{m-|\alpha|} ,$$

for any $z \in \mathbb{R}^n$, $\xi \in \mathbb{R}^n$ with, $|\xi| \geq R$, α and $\beta \in \mathbb{N}^n$.

We denote by ${}_{1,0}\mathcal{S}_{G^s}^m(\Omega \times \mathbb{R}^n)$ the set of such symbols.

Mathematics Subject Classification. 47G30.

Key words and phrases. Symbols of Gevrey, Kernels, Pseudo-differential operators.

Notice that, in general, a is supposed analytic in z . (This is done by taking $s = 1$ in the factor $(|\beta|!)^s$ corresponding to the variable z .)

Definition 3. We keep the notations of Definition 2. Let U be an open neighborhood of 0 in \mathbb{R}^n and m a positive real number. We say that a distribution $T = T(z, x)$, on $\Omega \times U$, is a Gevrey kernel of order m if, and only if, the following assertions are satisfied:

- a) The restriction f of T to $(U \setminus \{0\})$ is Gevrey of order s such that: For every compact K of Ω , there is an open neighborhood V of $U \setminus \{0\}$ and a scalar $C > 0$ such that

$$(3) \quad \left| D_x^\alpha D_z^\beta f(z, x) \right| \leq C^{|\alpha+\beta|+1} (|\alpha|!)^s (|\beta|!)^s \left(1 + |x|^2\right)^{-m-n-|\alpha|},$$

$$\forall (z, x) \in K \times V.$$

- b) The distribution T is of the form

$$(4) \quad T(z, \cdot) = P_{f_\theta}(z, \cdot) + \sum_{|\alpha| \leq m} C_\alpha(z) \delta^{(\alpha)}.$$

where $(C_\alpha)_\alpha$ is a family of Gevrey functions of order s in Ω and θ is a map of $C_0^\infty(U)$, verifying $\theta \equiv 1$ in a neighborhood of zero, while P_{f_θ} is a distribution given for any ψ in $C_0^\infty(U)$ by

$$(5) \quad \langle P_{f_\theta} f, \psi \rangle = \int_\Omega f(x) \left(\psi(x) - \sum_{|\alpha| \leq m} \frac{D^\alpha \psi(0)}{\alpha!} x^\alpha \theta(x) \right) dx$$

- c) If m is a positive integer, then, for any compact K in Ω , there is a scalar $C_K > 0$ such that:

For every α in \mathbb{N}^n with $|\alpha| = m$ and any $\varepsilon > 0$

$$(6) \quad \left| \int_{|x| \geq \varepsilon, x \in U} x^\alpha f(z, x) dx \right| \leq C_K, \forall z \in K.$$

The set of Gevrey kernels, so defined, is designated by ${}_{1,0}\mathcal{K}_{G^s}^m(\Omega \times \mathbb{R}^n)$.

3. THE MAIN RESULT

We prove the following result.

Theorem 4. Let m be a positive real number. If T is in ${}_{1,0}\mathcal{K}_{G^s}^m(\Omega \times \mathbb{R}^n)$, then there is a symbol f in ${}_{1,0}\mathcal{S}_{G^s}^m(\Omega \times \mathbb{R}^n)$ such that

$$T = \mathcal{F}^{-1} f$$

is of Gevrey type of order s on $\Omega \times U'$, where U' is an open 0 neighborhood in \mathbb{R}^n .

WE recall here that $\mathcal{F}^{-1}f$ denotes the image of f by the inverse of Fourier transform.

To prove this theorem we need the following lemma which is an analogous of the one in [1].

Lemma 5. *Let m be a positive real number, U and U' open neighborhoods of zero, in \mathbb{R}^n , such that $U' \subset \subset U$.*

- i) *Let T be a distribution on U , the restriction g of which, to $U \setminus \{0\}$, is C^∞ and satisfies*

$$(7) \quad \left| D_x^\beta g(x) \right| \leq C |x|^{m-n-|\beta|}, \quad |\beta| \leq m.$$

- ii) *We suppose $x^\alpha T$ that is an integrable on U , for $|\alpha| > m$.*
- iii) *If m is a positive integer, we suppose the existence of a map φ in $C_0^\infty(U)$, identically equal to 1 on U' and such that, for $|\alpha| = m$,*

$$(8) \quad \sup_{0 < \varepsilon < 1} |\langle x^\alpha T_\varepsilon(x), \varphi_\varepsilon(x) \rangle| \leq C,$$

with

$$\varphi_\varepsilon(x) = \varphi\left(\frac{x}{\varepsilon}\right).$$

Then, for every ψ in $C_0^\infty(U)$, there exists a scalar $M_\psi > 0$ such that

$$\left| D_\xi^\alpha (\widehat{\psi T}) (\xi) \right| \leq C M_\psi (1 + |\xi|)^{m-|\alpha|},$$

and this for every α in \mathbb{N}^n , $m \leq |\alpha| < m + 1$, and every ξ in \mathbb{R}^n .

Proof. Since α is in \mathbb{N}^n , we have, for $|\xi| \geq 1$,

$$\begin{aligned} D_\xi^\alpha (\widehat{\psi T}) (\xi) &= (-i)^{|\alpha|} (\widehat{x^\alpha \psi T}) (\xi) = (-i)^{|\alpha|} \langle x^\alpha \psi T, e^{ix\xi} \rangle \\ &= (-i)^{|\alpha|} (I_1 + I_2 + I_3) \end{aligned}$$

where

$$\begin{aligned} I_1 &= \langle x^\alpha T, \varphi_\varepsilon \psi \rangle, \\ I_2 &= \langle x^\alpha T, \varphi_\varepsilon \psi (e^{-ix\xi} - 1) \rangle, \\ I_3 &= \langle x^\alpha T, (1 - \varphi_\varepsilon) \psi e^{-ix\xi} \rangle. \end{aligned}$$

Let us examine these expressions one by one. First, fix

$$(9) \quad \varepsilon = \frac{1}{|\xi|}.$$

Using (7), (8) and (9), we get

$$|I_1| \leq C J |\xi|^{m-|\alpha|}$$

with a constant $J > 0$.

Concerning the estimation of I_2 , the definition of φ assures that

$$\left| e^{-ix\xi} - 1 \right| \leq |\xi| |x|.$$

This inequality and (7) yield to

$$|I_2| \leq C S |\xi|^{m-|\alpha|},$$

with a constant $S > 0$.

Concerning I_3 , we can write

$$I_3 = x^\alpha T \widehat{(1 - \varphi_\varepsilon)} \psi.$$

Then

$$\xi_j I_3 = -i \langle D_{x_j} (x^\alpha (1 - \varphi_\varepsilon) \psi T, e^{-ix\xi}) \rangle, \quad \forall j = 1, \dots, n.$$

Since the support of $1 - \varphi_\varepsilon$ is contained in $U \setminus U'$, we can find a scalar μ such that

$$0 < \mu \leq |\xi| |x|,$$

which permits to have

$$|\xi_j I_3| \leq C K |\xi|^{m-|\alpha|+1},$$

where K is a strictly positive scalar.

With these three estimations we obtain (12) in the case $|\xi| \geq 1$ considered.

As (12) remains obviously true when $|\xi| < 1$ the proof of Lemma 5 is finished. \square

The proof of Theorem 4 need also the following proposition.

Proposition 6. *Let U be an open 0-neighborhood in \mathbb{R}^n , a given α in \mathbb{N}^n and T an element of ${}_{1,0}\mathcal{K}_{G^s}^m(\Omega \times \mathbb{R}^n)$ such that its restriction f to $U \setminus \{0\}$, satisfies*

$$(10) \quad |D_x^\alpha f(x)| \leq C^{|\alpha|+1} (|\alpha|!) |x|^{-m-n-|\alpha|}.$$

For any map φ in $C_0^\infty(U)$ which is identically equal to 1 on an open 0-neighborhood $U' \subset\subset U$, we get

$$(11) \quad \left| D_\xi^\alpha (\widehat{\varphi T})(\xi) \right| \leq C^{|\alpha|+1} (|\alpha|!)^s (1 + |\xi|)^{m-|\alpha|}.$$

Proof. We define a function ψ on $C_0^\infty(U)$ as follows

$$(12) \quad \psi(x) = \begin{cases} 1 & \text{for } |x| \leq 1, \\ 0 & \text{for } |x| \geq 3. \end{cases}$$

For any β in \mathbb{N}^n such that $|\beta| = |\alpha|$ we get

$$\left| \xi^\alpha D^\alpha (\widehat{\varphi T})(\xi) \right| \leq |I_1| + |I_2|,$$

where

$$|I_1| = \left| \int_{\Omega} D_x^\alpha (x^\alpha (\varphi T)(x)) [1 - \psi(|\xi| x)] e^{-ix\xi} dx \right|,$$

$$|I_2| = \left| \int_{\Omega} D_x^\alpha (x^\alpha (\varphi T)(x)) \psi(|\xi| x) e^{-ix\xi} dx \right|.$$

Let us estimate these expressions. We have

$$|I_2| \leq \int_{|x| < \frac{3}{|\xi|}} |D_x^\alpha (x^\alpha (\varphi T)(x))| dx.$$

The Leibniz formula and (10) permits to get

$$|I_2| \leq \int_{|x| < \frac{3}{|\xi|}} \left| \sum_{v \leq \beta} \binom{\beta}{v} \partial^{\beta-v} T \partial^v (x^\alpha \varphi(x)) \right| dx$$

$$\leq \int_{|x| < \frac{3}{|\xi|}} \sum_{v \leq \beta} \binom{\beta}{v} |\partial^v (x^\alpha \varphi(x))| C^{|\beta-v|+1} (|\beta-v|)^s |x|^{-n-m-|\beta-v|} dx.$$

Whence

$$|I_2| \leq \int_{|x| < \frac{3}{|\xi|}} A_2 C^{|\beta|+1} (|\beta|)^s |x|^{-n-m} dx$$

$$(13) \quad \leq B_2 C^{|\beta|+1} (|\beta|)^s (1 + |\xi|)^m.$$

We proceed in the same way for $|I_1|$. We obtain

$$(14) \quad |I_1| \leq \int_{\frac{1}{|\xi|} < |x| < \frac{3}{|\xi|}} A_1 C^{|\beta|+1} (|\beta|)^s |x|^{-n-m} dx$$

$$\leq B_1 C^{|\beta|+1} (|\beta|)^s (1 + |\xi|)^m.$$

Notice that the scalars A_1 and A_2 depend on β while B_1 and B_2 depend on β , m and n . The constant C does not depend on β , nor n , nor m .

On the other hand, using (13) and (14), we easily obtain

$$\left| D_\xi^\alpha \widehat{(\varphi T)}(\xi) \right| \leq AC^{|\alpha|+1} (|\alpha|)^s (1 + |\xi|)^{m-|\alpha|}.$$

This inequality completes the proof of the Relation (11). □

The proof of Theorem 4 follows directly from the Lemma 5 and the Proposition 6.

REFERENCES

- [1] M. S. BAOUENDI, C. GOULAOUIC AND G. MÉTIVIER, *Kernels and symbols of analytic pseudodifferential operators*, J. Differ. Equations **48**(1983), 227–240.
- [2] L. BOUTET MONVEL AND P. KRÉE, *Pseudodifferential operators and Gevrey classes*, Annales Inst. F., Grenoble, 17.1, (1967), 295–323.
- [3] L. BOUTET DE MONVEL, *Opérateurs pseudodifférentiels analytiques et opérateurs d'ordre infini*, Annales Inst. Fourier, Grenoble, 22.3, (1972), 229–268.
- [4] M. HAZI, *Noyaux et symboles des opérateurs pseudodifférentiels en C^∞* , Mémoire de D.E.A, Ecole Polytechnique, Paris, (1982).
- [5] M. HAZI, *Kernels of pseudo-differential operators associated to Hörmander symbols of Gevrey type*, Arab Gulf Journal of Scientific Research, Manama, Volume 19. 1, (2001), 52–58.

MOHAMMED HAZI

DEPARTMENT OF MATHEMATICS

ÉCOLE NORMALE SUPÉRIEURE

16050 - KOUBA, ALGIERS, ALGERIA

e-mail address: mohamedhazi@hotmail.com, hazi@ens-kouba.dz

Fax: (213) 21 28 20 67

(Received November 14, 2002)