

Mathematical Journal of Okayama University

Volume 14, Issue 1

1969

Article 4

NOVEMBER 1969

A note on Schauder basis

Osamu Katsumata*

*Okayama University

Copyright ©1969 by the authors. *Mathematical Journal of Okayama University* is produced by
The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

A NOTE ON SCHAUDER BASIS

OSAMU KATSUMATA

Preliminary and notations. Let E denote a topological linear space over the real or complex scalar field. A sequence $(b_n)_{n=1,2,\dots}$ in E is called a basis of E if for any element x of E there exists a unique sequence $(t_n)_{n=1,2,\dots}$ of scalars such that $x = \sum_{n=1}^{\infty} t_n b_n$. The convergence of the series is in respect to the vector topology of E . If (b_n) is a basis of E and $x = \sum_{n=1}^{\infty} t_n b_n$, then the uniqueness of the expansion ensures that the correspondence $x \rightarrow t_n$ is a linear functional for each n . When all of these linear functionals are continuous, the basis $(b_n)_{n=1,2,\dots}$ is named Schauder basis. E with a basis being complete and metrizable is a sufficient condition for the basis to be Schauder (Arsove [1], p. 368). Also in $\mathcal{L}\mathcal{T}$ -space (strict inductive limit of Fréchet spaces) a basis is a Schauder basis (Arsove and Edwards [2], p. 113). In this note, we shall show that the some theorems known in Banach spaces also hold in certain Fréchet spaces.

For a topological linear space E , let us denote by E' its topological dual. The topologies $\sigma(E, E')$, $\mathcal{T}_b(E, E')$ (abbreviated $\mathcal{T}_b(E)$) are weak-, strong-topologies on E' , that is, the topologies deduced from the polars of all finite sets, of all bounded sets of E , respectively. The facts used in this note about topological linear spaces will be found for example in Bourbaki [4], Kelley [7], Köthe [8] or Robertsons [10]. We mean the biorthogonal pair $(x_n, f_n)_{n=1,2,\dots}$ in (E, E') as the sequences (x_n) in E and (f_n) in E' which satisfy $\langle x_m, f_n \rangle = \delta_{mn}$ (Kronecker's delta). Let us put

$$(1) \quad U_m(x) = \sum_{n=1}^m \langle x, f_n \rangle b_n, \quad V_m(x) = x - U_m(x)$$

for a biorthogonal pair (x_n, f_n) in (E, E') . Then, U_m and V_m are continuous operators in E . Moreover, U_m^* will mean the transpose of U_m defined by the relation $\langle x, U_m^*(f) \rangle = \langle U_m(x), f \rangle$ for all $x \in E$, $f \in E'$, and similarly for V_m^* . A sequence (x_n) in E is basic if and only if it is a Schauder basis for the closed linear hull (denoted by $[x_n]$) of the set $\{x_n; n=1, 2, \dots\}$.

Characterization of a basic sequence and its corollaries. In what follows, "completeness" will play the essential role. First, we shall prove a lemma.

Lemma. *Let E be a complete topological linear space. If $(T_n)_{n=1,2,\dots}$ is an equicontinuous sequence of linear operators which map E into E , then $E_0 = \{x; \lim_{n \rightarrow \infty} T_n(x) \text{ exists}\}$ is a closed linear subspace of E .*

Proof. It is clear that E_0 is a linear subspace of E from the linearity of each T_n . Let x_0 be any point of the closure of E_0 , then a net $(x_\alpha)_{\alpha \in A}$ in E_0 converges to x_0 , where for every $\alpha \in A$ there exists $y_\alpha \in E$ such that $\lim_{n \rightarrow \infty} T_n x_\alpha = y_\alpha$. For every V , a closed neighbourhood of 0, there exists a neighbourhood U of 0 which satisfies $U + U + \bar{U} \subset V$, and $T_n(U) \subset V$ for each n by equicontinuity. The net $(x_\alpha)_{\alpha \in A}$ converges to x_0 , so there exists $\delta \in A$ such that if $\alpha, \beta \geq \delta$ then $x_\alpha - x_\beta \in U$ and so $T_n(x_\alpha - x_\beta) \in T_n(U) \subset V$. This and the closedness of V imply $y_\alpha - y_\beta \in V$, $\alpha, \beta \geq \delta$, i. e. $(y_\alpha)_{\alpha \in A}$ to be a Cauchy net in E . Completeness assumption assures then the existence of an element $y_0 \in E$ to which (y_α) converges. The two nets on the same directed system A converge, therefore we can find α_1 in A such that both $y_{\alpha_1} - y_0$ and $x_0 - x_{\alpha_1}$ are in U . For this α_1 there exists an integer n_1 such that $T_n(x_{\alpha_1}) - y_{\alpha_1} \in \bar{U}$ for all n larger than n_1 . Then, $T_n(x_0) - y_0 = T_n(x_0) - T_n(x_{\alpha_1}) + T_n(x_{\alpha_1}) - y_{\alpha_1} + y_{\alpha_1} - y_0 \in U + U + \bar{U} \subset V$, $\geq n_1$. Hence, $T_n(x_0)$ converges to y_0 and we finish the proof.

Theorem. *Let E be a Fréchet space, $(b_n, f_n)_{n=1,2,\dots}$ a biorthogonal pair in (E, E') , and define $U_m(x) = \sum_{n=1}^m \langle x, f_n \rangle b_n$ for every $x \in E$ and every natural number m . Then, (b_n) is basic if and only if the sequence $(U_m)_{m=1,2,\dots}$ is pointwise bounded on $E_1 = [b_n]$ (A. Wilansky [13], p. 210).*

Proof. When (b_n) is basic, the sequence $U_m(x)$ converges to x for every $x \in E_1$ and $\{U_m(x); m=1, 2, \dots\}$ is a bounded set in E . Conversely, E_1 being a closed linear subspace of a Fréchet space, E_1 is itself a barrelled space. When we regard U_m as a pointwise bounded sequence of continuous linear operators from E_1 to E_1 , $\{U_m; m=1, 2, \dots\}$ is equicontinuous. Here we can apply the preceding lemma to obtain a closed linear subspace $\{x \in E_1; \lim_{m \rightarrow \infty} U_m(x) \text{ exists}\}$. As it contains all b_n 's, it coincides with E_1 , that is, $x = \sum_{n=1}^{\infty} \langle x, f_n \rangle b_n$ for all $x \in E_1$. This implies that (b_n) is a Schauder basis in E_1 .

In the following, metrizable of $(E', \mathcal{T}_b(E))$ is needed. This is equivalent to the condition that all bounded sets in E has countable co-base, i. e. there is a countable family of bounded sets such that every bounded set is contained in some member of the family.

Corollary. *Let E be a Fréchet space which has a countable co-base for all bounded sets. If E has a basis $(b_n)_{n=1,2,\dots}$, then there exists a sequence $(f_n)_{n=1,2,\dots}$ in E' biorthogonal to (b_n) , which is basic in $(E', \mathcal{T}_b(E))$.*

Proof. Because every basis in a Fréchet space is a Schauder basis, there exists a biorthogonal sequence (f_n) in E' to (b_n) . If we choose U_m as in (1) then the sequence is equicontinuous by the same reason as in the last part of the proof of the theorem, and the sequence of their transposes U_m^* is equicontinuous in respect to $\mathcal{T}_b(E)$. Next, for every $g \in E'$, $x \in E$ we have

$$\langle x, U_m^*(g) \rangle = \langle U_m(x), g \rangle \longrightarrow \langle x, g \rangle, \text{ as } m \longrightarrow \infty,$$

which means $\{U_m^*(g); m=1, 2, \dots\}$ is a $\sigma(E, E')$ bounded set. From this and the fact that E is a barrelled space, $\{U_m^*(g); m=1, 2, \dots\}$ becomes a $\mathcal{T}_b(E)$ bounded set for each $g \in E'$. Now, $(E', \mathcal{T}_b(E))$ is complete as the strong dual of a bornologic space E . We conclude therefore by the preceding theorem, that (f_n) is basic in $(E', \mathcal{T}_b(E))$.

As corollaries to the theorem, we shall mention two propositions.

Proposition 1 (M. M. Day [5], p. 70). *Let E be a Fréchet space which has a countable co-base for all bounded sets, and let $(b_n)_{n=1,2,\dots}$ be a basis and $(f_n)_{n=1,2,\dots}$ be a sequence in E' biorthogonal to (b_n) . The following conditions are equivalent:*

- (i) (f_n) is a $\mathcal{T}_b(E)$ Schauder basis.
- (ii) $V_m^* \longrightarrow 0$, as $m \longrightarrow \infty$, in respect to the strong operator topology.
- (iii) For every $f \in E'$ and for every bounded set B of E ,
 $\sup \{|f(x)|; x \in B \cap [b_n, b_{n+1}, \dots]\} \longrightarrow 0$, as $n \longrightarrow \infty$.
- (iv) $[f_n] = E'$ i. e. $\{f_n; n=1, 2, \dots\}$ is fundamental in $(E', \mathcal{T}_b(E))$.

Remark. (a) V_m^* means the transpose of V_m where V_m is defined as in (1). When for every $f \in E'$ $V_m^*(f) \longrightarrow 0$ in respect to $\mathcal{T}_b(E)$, we say that $V_m^* \longrightarrow 0$ in respect to the strong operator topology.

(b) A Schauder basis in a Banach space is called shrinking when it satisfies the condition that for every $f \in E'$,

$$\sup \{|f(x)|; x \in [b_n, b_{n+1}, \dots], \|x\| \leq 1\} \longrightarrow 0, \text{ as } n \longrightarrow \infty.$$

The property "shrinking" is related to the basis of a reflexive Banach space (cf. for instance James [6], Retherford [9] and Singer [12]).

(c) For a barrelled space X (in particular, for a Fréchet space), it

can be proved that if X' provides the topology of uniform convergence of all precompact sets then every sequence biorthogonal to a Schauder basis spans X' . Comparing this with the condition (iv), one will see the difference between a shrinking basis and a Schauder basis.

Proof. (i) \Rightarrow (ii): For every $x \in E, f \in E'$,

$$\begin{aligned} \langle x, V_m^*(f) \rangle &= \langle V_m(x), f \rangle = \langle x - \sum_{n=1}^{m-1} \langle x, f_n \rangle b_n, f \rangle \\ &= \langle x, f - \sum_{n=1}^{m-1} \langle b_n, f \rangle f_n \rangle. \end{aligned}$$

Accordingly, if (f_n) is a $\mathcal{T}_b(E)$ -basis then for every bounded set B of E we have $\sup_{x \in B} |\langle x, V_m^*(f) \rangle| \rightarrow 0 (m \rightarrow \infty)$, that is, $V_m^* \rightarrow 0 (m \rightarrow \infty)$ in respect to the strong operator topology.

(ii) \Rightarrow (iii): Since (b_n) is a Schauder basis, it is obvious that x is in $B \cap [b_n, b_{n+1}, \dots]$ when and only when $x = V_n(x)$ and $x \in B$. On the other hand, for every $f \in E'$

$$\begin{aligned} \sup \{ |\langle x, f \rangle|; x = V_n(x), x \in B \} &\leq \sup \{ |\langle V_n(x), f \rangle|; x \in B \} \\ &= \sup \{ |\langle x, V_n^*(f) \rangle|; x \in B \}. \end{aligned}$$

We see therefore (ii) implies (iii).

(iii) \Rightarrow (iv): For every $f \in E'$ and every bounded set B of E , if we set $B_1 = \bigcup_{m=1}^{\infty} V_m(B)$ then there holds

$$\begin{aligned} \sup_{x \in B} |\langle x, \sum_{n=1}^m \langle b_n, f \rangle f_n - f \rangle| &= \sup_{x \in B} |\langle V_m(x), f \rangle| \\ &\leq \sup_{x \in B_1} \{ |\langle x, f \rangle|; x = V_m(x) \}. \end{aligned}$$

(Note that $V_m^2 = V_m$.) As the union B_1 of the images of bounded set B by equicontinuous operators is a bounded set of E , this yields the implication (iii) \Rightarrow (iv).

(iv) \Rightarrow (i): This is a consequence of the preceding corollary.

Proposition 2. (I. Singer [11], p. 77). *Let E be a Fréchet space which has a countable co-base for all bounded sets. If E' contains a $\sigma(E, E')$ Schauder basis $(f_n)_{n=1,2,\dots}$ then E contains a Schauder basis $(b_n)_{n=1,2,\dots}$, and conversely.*

Proof. For the $\sigma(E, E')$ -Schauder basis (f_n) there exists a biorthogonal sequence of $\sigma(E, E')$ -continuous linear functionals (b_n) , which we may regard as a subset of E . Since $U_m^*(f)$ $\sigma(E, E')$ -converges to f , it follows that $U_m(x)$ $\sigma(E', E)$ -converges to x . Accordingly, $\{U_m(x); m=1, 2, \dots\}$ is $\sigma(E', E)$ -bounded, i. e. it is a bounded set of E . Now, noting that the linear hull of b_n 's is dense in E , the same argument used in the

proof of the theorem shows that the sequence (b_n) is a Schauder basis. Conversely, assume that (b_n) is a Schauder basis in E . Then, by definition, there exists a biorthogonal sequence (f_n) and for each $x \in E$ there holds $x = \sum_{n=1}^{\infty} \langle x, f_n \rangle b_n$. As $\langle x, f - \sum_{n=1}^m \langle b_n, f \rangle f_n \rangle = \langle x - \sum_{n=1}^m \langle x, f_n \rangle b_n, f \rangle$ for every $f \in E'$, we readily see that $\sum_{n=1}^m \langle b_n, f \rangle f_n \rightarrow \sigma(E, E')$ converges to f . Moreover, b_n 's being continuous linear functionals on $(E', \sigma(E, E'))$ and biorthogonal to (f_n) , the expansion is unique.

REFERENCES

- [1] M.G. ARSOVE: The Paley-Wiener theorem in metric linear spaces, Pacific Journal of Mathematics, 10 (1960), 365—379.
- [2] M.G. ARSOVE and R.E. EDWARDS: Generalized bases in topological linear spaces, Studia Mathematica, 19 (1960), 95—113.
- [3] S. BANACH: Théorie des opérations linéaires, Warszawa, 1933.
- [4] N. BOURBAKI: Elements de Mathématique, Espaces vectoriels topologiques, Hermann, Paris, 1955.
- [5] M.M. DAY: Normed linear spaces, Springer-Verlag, 1962.
- [6] R.C. JAMES: Bases and reflexivity of Banach spaces, Annals of Mathematics, 52 (1950), 518—527.
- [7] J.L. KELLEY and COAUTHORS: Linear topological spaces, D. Van Nostrand, New York, 1963.
- [8] G. KÖTHER: Topologische lineare Räume I, Springer-Verlag, 1960.
- [9] J.R. RETHERFORD: Bases, basic sequences and reflexivity of linear topological spaces, Mathematische Annalen, 164 (1966), 280—285.
- [10] A. ROBERTSON and W. ROBERTSON: Topological vector spaces, Cambridge University Press, 1964.
- [11] I. SINGER: Weak* bases in conjugate Banach spaces, Studia Mathematica, 21 (1961), 75—81.
- [12] I. SINGER: Basic sequences and reflexivity of Banach spaces, Studia Mathematica, 21 (1962), 351—369.
- [13] A. WILANSKY: Functional analysis, Blaisdell Publishing Company, 1964.

DEPARTMENT OF MATHEMATICS,
OKAYAMA UNIVERSITY

(Received April 17, 1969)