

Mathematical Journal of Okayama University

Volume 50, Issue 1

2008

Article 9

JANUARY 2008

On Fox Spaces and Jacobi Identities

Marek Golasinski*

Daciberg Lima Gonçalves†

Peter Wong‡

*Nicolaus Copernicus University

†Departamento De Matemática

‡Bates College

Copyright ©2008 by the authors. *Mathematical Journal of Okayama University* is produced by
The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

On Fox Spaces and Jacobi Identities

Marek Golasinski, Daciberg Lima Gonçalves, and Peter Wong

Abstract

In 1945, R. Fox introduced the so-called Fox torus homotopy groups in which the usual homotopy groups are embedded and their Whitehead products are expressed as commutators. A modern treatment of Fox torus homotopy groups and their generalization has been given and studied. In this note, we further explore these groups and their properties. We discuss co-multiplications on Fox spaces and Jacobi identities for the generalized Whitehead products and the T- Whitehead products.

KEYWORDS: Fox torus homotopy groups, generalized Whitehead products, Jacobi identity

Math. J. Okayama Univ. **50** (2008), 161–176**ON FOX SPACES AND JACOBI IDENTITIES**

MAREK GOLASIŃSKI, DACIBERG LIMA GONÇALVES AND PETER WONG

ABSTRACT. In 1945, R. Fox introduced the so-called Fox torus homotopy groups in which the usual homotopy groups are embedded and their Whitehead products are expressed as commutators. A modern treatment of Fox torus homotopy groups and their generalization has been given and studied. In this note, we further explore these groups and their properties. We discuss co-multiplications on Fox spaces and Jacobi identities for the generalized Whitehead products and the Γ -Whitehead products.

INTRODUCTION

In an attempt to give a geometric interpretation of the Whitehead product of [27], R. Fox [9, 10] introduced the so-called torus homotopy groups $\tau_n(X, x_0)$ of a space X . For each $n \geq 1$, the group $\tau_n(X, x_0)$ is completely determined by the classical homotopy groups $\pi_k(X, x_0)$ for $k \leq n$ and the Whitehead products of the elements of $\pi_k(X, x_0)$. In fact, for any given $\alpha \in \pi_m(X, x_0)$ and $\beta \in \pi_n(X, x_0)$, the Whitehead product $[\alpha, \beta]$ when considered as an element in $\tau_k(X, x_0)$, $k \geq n + m - 1$, is a commutator.

The Fox torus homotopy groups were given a modern interpretation and were generalized in [13]. Evaluation subgroups of τ_n were defined and their relationships with the classical Gottlieb groups were also studied. Moreover, it was shown in [13] that the generalized Whitehead product when embedded in the same spirit as [2] in a larger (and different) group is a commutator as well.

In 1954, M. Nakaoka and H. Toda [20] proved a Jacobi identity for the classical Whitehead products involving elements in higher homotopy groups. In fact, they gave two such proofs one of which made use of the Fox torus homotopy groups. We should point out that H. Suzuki (1954)[25], G. Whitehead (1954)[29], S.C. Chang (1954)[6], P. Hilton (1955)[16], H. Uehara and

Mathematics Subject Classification. Primary: 55Q05, 55Q15, 55Q91; secondary: 55M20.

Key words and phrases. Fox torus homotopy groups, generalized Whitehead products, Jacobi identity.

This work was conducted during the second and third authors' visits to the Faculty of Mathematics and Computer Science, Nicolaus Copernicus University, August 4 - 13, 2005 and July 29 - August 8, 2006. The second and third authors would like to thank the Faculty of Mathematics and Computer Science for its hospitality and support. The third author was supported in part by a grant from Bates College and the National Science Foundation.

W. Massey (1957)[26], and W. Barcus and M. Barratt (1958)[3] obtained the same or similar Jacobi identities, independently. Furthermore, Barcus and Barratt also derived a Jacobi identity in which one of the elements is in the fundamental group. In 1961, P. Hilton [17] published a short note extending the Jacobi identity by allowing elements from the fundamental group.

For the generalized Whitehead product¹, the Jacobi identity was first proved by D. Cohen (1957)[7, Section 5.8], in the language of carrier theory. It was pointed out by M. Arkowitz that such identity was also obtained in his doctoral dissertation (1960). When A, B and C are suspensions, the Jacobi identity was obtained independently by J. Boardman and B. Steer (1967)[5, Equation (4.4)], and by H. Ando (1968)[1, Theorem 5.4]. The most general form of the Jacobi identity was due to J.W. Rutter (1968)[23, 24] who presented two proofs of the Jacobi identity for the generalized Whitehead product, under the assumption that $\Sigma A, \Sigma B$, and ΣC are homotopy co-commutative. We should point out that J. Neisendorfer (1980) proved a Jacobi identity [21, Proposition 9.10], for the *external Samelson product* $\langle \cdot, \cdot \rangle$ for $[Z_1, G], [Z_2, G]$, and $[Z_3, G]$ where each Z_i is *co-abelian* (i.e., Z_i has weak category ≤ 1) and G is group-like. In fact, $\langle \cdot, \cdot \rangle$ is the same as the generalized Whitehead product [28, p. 467]. When $G = \Omega X$, Corollary 9.7² of [21] implies that $[Z, \Omega X] = [\Sigma Z, X]$ is an abelian group when Z is co-abelian. This means that ΣZ is homotopy co-commutative. Thus, the Jacobi identity of Neisendorfer's follows from that of Rutter's. Furthermore, there are examples due to W. Gilbert [11] (see also [8, Example 2.60, p. 70]) where a space X is not co-abelian while ΣX is homotopy co-commutative.

The objective of this paper is to further explore the Fox torus homotopy groups and their generalizations. In particular, Section 1 takes up the study of Fox torus homotopy groups of spheres, Eilenberg-MacLane spaces, and Fox spaces. Section 2 explores Fox spaces F_n and their co-multiplication sets $\mathcal{C}(F_n)$ for $n \geq 1$. Section 3 is concerned with Jacobi identities for the generalized Whitehead product studied in [7, 2, 18, 5, 1, 21, 23, 24, 22]. By using the generalized Fox torus groups studied in [13], we give a unified approach to the Jacobi identities of Rutter's [23] and of Hilton's [17]. In Section 4, we generalize the Fox torus groups to Γ -Fox torus groups for any co-grouplike space Γ . We improve the Jacobi identity in [22] for the Γ -Whitehead product.

¹The generalized Whitehead products of [1, 2, 7] are slightly different from the others although they are all defined based upon commutators.

²When Z is a connected *CW*-complex, this result was already obtained by I. Berstein and T. Ganea [4, Corollary 3.3].

1. FOX SPACES

First, we recall from [9, 10] the definition of the n -th Fox torus homotopy group of a pointed space X , for $n \geq 1$. Let x_0 be a basepoint of X , then

$$\tau_n(X, x_0) = \pi_1(X^{T^{n-1}}, \overline{x_0})$$

where $X^{T^{n-1}}$ denotes the space of unbased maps from the $(n-1)$ -torus T^{n-1} to X and $\overline{x_0}$ is the constant map at x_0 . When $n = 1$, $\tau_1(X, x_0) = \pi_1(X, x_0)$.

To re-interpret Fox's result, we showed in [13] that

$$\tau_n(X, x_0) \cong [\Sigma(T^{n-1} \sqcup *), X]$$

the group of homotopy classes of basepoint preserving maps from the reduced suspension of T^{n-1} adjoined with a distinguished point to X . As a result, we call $F_n = \Sigma(T^{n-1} \sqcup *)$ the n -th Fox space with $F_1 = \mathbb{S}^1$ the circle.

One of the main results of [10] is the following split exact sequence:

$$(1.1) \quad 0 \rightarrow \prod_{i=2}^n \pi_i(X, x_0)^{\alpha_i(n)} \rightarrow \tau_n(X, x_0) \overset{\leftarrow}{\rightleftarrows} \tau_{n-1}(X, x_0) \rightarrow 1$$

where $\alpha_i(n)$ is the binomial coefficient $\binom{n-2}{i-2}$.

With the isomorphism $\tau_{n-1}(\Omega X) \cong \prod_{i=2}^n \pi_i(X, x_0)^{\alpha_i(n)}$ shown in [13, Theorem 1.1], (1.1) becomes

$$(1.2) \quad 0 \rightarrow \tau_{n-1}(\Omega X) \rightarrow \tau_n(X) \overset{\leftarrow}{\rightleftarrows} \tau_{n-1}(X) \rightarrow 1$$

or equivalently

$$0 \rightarrow [\Sigma F_{n-1}, X] \rightarrow [F_n, X] \overset{\leftarrow}{\rightleftarrows} [F_{n-1}, X] \rightarrow 1.$$

Using (1.1), the following is easy to verify.

Proposition 1.1. *For any pointed space X with basepoint x_0 , $\tau_k(X, x_0) = 1$, for all $k \leq n$ if and only if $\pi_k(X, x_0) = 1$, for all $k \leq n$. Moreover, $\tau_n(X, x_0) \cong \pi_n(X, x_0)$ if and only if X is $(n-1)$ -connected.*

For any $n \geq 1$, the n -th Fox space F_n has the homotopy type of $\Sigma T^{n-1} \vee \mathbb{S}^1$. In fact, the natural isomorphism

$$[F_n, X] \cong [\Sigma F_{n-1}, X] \times [F_{n-1}, X]$$

from the split exact sequence (1.2) yields:

Proposition 1.2. *For any integer $n > 1$, we have a homotopy equivalence*

$$F_n \simeq \Sigma F_{n-1} \vee F_{n-1}.$$

As already pointed out in [10], one can define a Hurewicz homomorphism $\rho_n : \tau_n(X, x_0) \rightarrow H_n(X; \mathbb{Z})$, where \mathbb{Z} is the group of integers. It follows from Proposition 1.1 that the (absolute) Hurewicz Isomorphism Theorem holds for the torus homotopy groups as in the classical case.

Example 1. Let $X = \mathbb{S}^n$, the n -sphere. For all $k \leq n$, $\tau_k(\mathbb{S}^n) = \pi_k(\mathbb{S}^n)$. Using (1.1), we have $\tau_{n+1}(\mathbb{S}^n) \cong \pi_{n+1}(\mathbb{S}^n) \rtimes \pi_n(\mathbb{S}^n)$. More generally, we use (1.2) to obtain

$$\tau_m(\mathbb{S}^n) \cong \tau_{m-1}(\Omega\mathbb{S}^n) \rtimes \tau_{m-1}(\mathbb{S}^n).$$

Example 2. Let $X = K(\pi, n)$ be an Eilenberg-MacLane space of type (π, n) for some abelian group π and $n \geq 2$. It follows from (1.1) that

$$(1.3) \quad \tau_m(X) \cong \prod_{i=2}^m \pi_i(X, x_0)^{\alpha_i(m)} \rtimes \tau_{m-1}(X).$$

Since $X = K(\pi, n)$, we get $\tau_m(X) \cong \pi_m(X) = 1$ for $m < n$ and $\tau_n(X) = \pi_n(X) = \pi$. For $k \geq 1$, (1.3) yields

$$\tau_{n+k}(X) \cong \pi^{\alpha_n(n+k)} \rtimes \tau_{n+k-1}(X).$$

The semi-direct product structure comes from the Whitehead products which are all trivial since $X = K(\pi, n)$. Hence, we conclude that $\tau_{n+k}(X)$ is a direct sum, i.e.,

$$\tau_{n+k}(X) \cong \bigoplus_{j=0}^k \pi_n(X)^{\alpha_n(n+j)} \cong \pi^{\sum_{j=0}^k \binom{n-2+j}{n-2}}$$

for any $k \geq 0$. Using a straightforward application of the Pascal's triangles one obtains that

$$\sum_{j=0}^k \binom{n-2+j}{n-2} = \sum_{j=0}^k \binom{n-2+j}{j} = \binom{n+k-1}{k}.$$

Therefore, for any space X of type $K(\pi, n)$, we have

$$\tau_m(X) = \begin{cases} 1, & \text{if } m < n; \\ \pi^{\binom{m-1}{m-n}}, & \text{if } m \geq n. \end{cases}$$

If $n = 1$, that is, $X = K(\pi, 1)$, then it is easy to show that $\tau_k(X) \cong \pi$, for all $k \geq 1$.

Next, we show that the wreath product \wr appears in the second torus homotopy group of the Fox space F_n .

Example 3. For any positive integer n , consider the n -th Fox space F_n . It was shown in [13] that F_n has the homotopy type of a bouquet of spheres. More precisely,

$$F_n \simeq \bigvee_{i=1}^n (\mathbb{S}^i)^{\gamma_i(n)},$$

where $\gamma_i(n)$ is the binomial coefficient $\binom{n-1}{i-1}$. It follows that $\pi_1(F_n) \cong \mathbb{Z}$ and

$$\pi_2(F_n) \cong \pi_2(\tilde{F}_n) \cong \bigoplus_{j=-\infty}^{\infty} (\mathbb{Z}^{n-1})_j,$$

where \tilde{F}_n denotes the universal cover of F_n . Now, (1.1) leads to the wreath product

$$\tau_2(F_n) \cong \pi_2(F_n) \rtimes \tau_1(F_n) \cong \mathbb{Z}^{n-1} \wr \mathbb{Z}$$

since the action of π_1 on the universal cover is the translation along the real line which covers the only copy of \mathbb{S}^1 in F_n .

Moreover, $F_m \simeq \bigvee_{i=1}^m (\mathbb{S}^i)^{\gamma_i(m)}$ is a homotopy retract of $F_n \simeq \bigvee_{i=1}^n (\mathbb{S}^i)^{\gamma_i(n)}$ provided that $m \leq n$. Therefore, we derive:

Proposition 1.3. *Let $m \leq n$. Then, there is an inclusion of groups*

$$\tau_k(F_m) \subseteq \tau_k(F_n)$$

for $k \geq 1$.

Remark 1. The calculation of the torus homotopy groups of an Eilenberg-MacLane space X of type $K(\pi, n)$ in Example 2 can be done directly as follows. Note that $F_m \simeq \bigvee_{i=1}^m (\mathbb{S}^i)^{\binom{m-1}{i-1}}$. Therefore for $m \geq n$, the number of n -spheres in F_m is precisely $\binom{m-1}{n-1} = \binom{m-1}{m-n}$. The fact that the Whitehead products all vanish implies that $\tau_m(X)$ is isomorphic to the direct sum of $\binom{m-1}{m-n}$ copies of π . We should point out that the co-multiplication of F_m does not yield a different group structure on $\tau_m(X)$ because X has no non-trivial homotopy groups in dimensions other than n .

2. GENERALIZED FOX TORUS GROUPS AND CO-MULTIPLICATION

In this section, we further explore the Fox spaces and examine co-multiplications on them. The functor τ_n was generalized in [13] as follows.

Definition 1. Let X be a space and $x_0 \in X$. For any space W , the W -Fox group of X is defined to be

$$\tau_W(X, x_0) = [\Sigma(W \sqcup *), X].$$

It is clear that τ_W reduces to τ_n when $W = T^{n-1}$.

We can further generalize Proposition 1.2 by the following construction. For any pointed W , we let

$$\mathfrak{F}(W) = \Sigma W \vee W.$$

Then, for any $n \geq 2$,

$$\mathfrak{F}_n(W) = \Sigma \mathfrak{F}_{n-1}(W) \vee \mathfrak{F}_{n-1}(W)$$

or

$$\mathfrak{F}_n(W) = (\mathbb{S}^1 \wedge \mathfrak{F}_{n-1}(W)) \vee \mathfrak{F}_{n-1}(W)$$

with $\mathfrak{F}_1(W) = W$. Certainly, $\mathfrak{F}_n(\mathbb{S}^1) \simeq F_n$. Furthermore, we can easily check that

$$[\mathfrak{F}(W), X] = [W, A_{\mathfrak{F}}(X)]$$

for any pointed space W where $A_{\mathfrak{F}}(X) = X \times \Omega X$.

The definition of τ_W prompts another generalization of the Fox space as follows.

Definition 2. For any $n \geq 2$, we let

$$\hat{\mathfrak{F}}_n(W) = \Sigma(W^{n-1} \sqcup *).$$

It follows that when $W = \mathbb{S}^1$, we have $F_n = \hat{\mathfrak{F}}_n(\mathbb{S}^1) \simeq \mathfrak{F}_n(\mathbb{S}^1)$.

Consider the suspension co-multiplication on $\hat{\mathfrak{F}}_n(\mathbb{S}^1)$. While $\hat{\mathfrak{F}}_n(\mathbb{S}^1)$ and $\mathfrak{F}_n(\mathbb{S}^1)$ are homotopy equivalent, they are not as co- H -spaces. To see that compare $[\hat{\mathfrak{F}}_2(\mathbb{S}^1), F_2]$ with $[\mathfrak{F}_2(\mathbb{S}^1), F_2]$.

Then, $[\hat{\mathfrak{F}}_2(\mathbb{S}^1), F_2] = \tau_2(F_2) \cong (\bigoplus_{j=-\infty}^{\infty} (\mathbb{Z})_j) \rtimes \mathbb{Z} = \mathbb{Z} \wr \mathbb{Z}$, by the calculation in Example 3. On the other hand, $\mathfrak{F}_2(\mathbb{S}^1) = \Sigma \mathbb{S}^1 \vee \mathbb{S}^1 = \mathbb{S}^2 \vee \mathbb{S}^1$. It follows that

$$[\mathfrak{F}_2(\mathbb{S}^1), F_2] \cong [\mathbb{S}^2 \vee \mathbb{S}^1, \mathbb{S}^2 \vee \mathbb{S}^1] \cong [\mathbb{S}^2, \mathbb{S}^2 \vee \mathbb{S}^1] \oplus [\mathbb{S}^1, \mathbb{S}^2 \vee \mathbb{S}^1] \cong (\bigoplus_{j=-\infty}^{\infty} (\mathbb{Z})_j) \oplus \mathbb{Z}.$$

Following e.g., [12, Proposition 1.1], the kernel of the obvious epimorphism

$$[\mathfrak{F}_n(\mathbb{S}^1), \mathfrak{F}_n(\mathbb{S}^1) \vee \mathfrak{F}_n(\mathbb{S}^1)] \rightarrow [\mathfrak{F}_n(\mathbb{S}^1), \mathfrak{F}_n(\mathbb{S}^1) \times \mathfrak{F}_n(\mathbb{S}^1)]$$

is in 1-1 correspondence with the set $\mathcal{C}(\mathfrak{F}_n(\mathbb{S}^1))$ of co-multiplications on the co- H -space $\mathfrak{F}_n(\mathbb{S}^1)$ for $n \geq 1$. Because $\mathfrak{F}_n(\mathbb{S}^1) \simeq F_n \simeq \bigvee_{i=1}^n (\mathbb{S}^i)^{\gamma_i(n)}$, we get the epimorphism

$$\begin{aligned} & \left[\bigvee_{i=1}^n (\mathbb{S}^i)^{\gamma_i(n)}, \bigvee_{i=1}^n (\mathbb{S}^i)^{\gamma_i(n)} \vee \bigvee_{i=1}^n (\mathbb{S}^i)^{\gamma_i(n)} \right] \rightarrow \left[\bigvee_{i=1}^n (\mathbb{S}^i)^{\gamma_i(n)}, \right. \\ & \left. \bigvee_{i=1}^n (\mathbb{S}^i)^{\gamma_i(n)} \times \bigvee_{i=1}^n (\mathbb{S}^i)^{\gamma_i(n)} \right]. \end{aligned}$$

Next, it holds $\pi_1(\mathfrak{F}_n(\mathbb{S}^1)) \cong \mathbb{Z}$, $\pi_k(\mathfrak{F}_n(\mathbb{S}^1)) \cong \pi_k(\bigvee_{j=-\infty}^{\infty} \bigvee_{i=2}^n (\mathbb{S}^i)^{\gamma_i(n)})$ and $\pi_1(\mathfrak{F}_n(\mathbb{S}^1) \vee \mathfrak{F}_n(\mathbb{S}^1)) \cong \mathbb{Z} * \mathbb{Z}$, $\pi_k(\mathfrak{F}_n(\mathbb{S}^1) \vee \mathfrak{F}_n(\mathbb{S}^1)) \cong \pi_k(\bigvee_{j \in \mathbb{Z} * \mathbb{Z}} \bigvee_{i=2}^n ((\mathbb{S}^i)^{\gamma_i(n)} \vee (\mathbb{S}^i)^{\gamma_i(n)}))$ for $k \geq 2$, where $\mathbb{Z} * \mathbb{Z}$ is the free group on two generators.

In particular, for $\mathfrak{F}_1(\mathbb{S}^1) \simeq \mathbb{S}^1$ and $\mathfrak{F}_2(\mathbb{S}^1) \simeq \mathbb{S}^2 \vee \mathbb{S}^1$, we get the epimorphisms

$$\mathbb{Z} * \mathbb{Z} \longrightarrow \mathbb{Z} \oplus \mathbb{Z}$$

and

$$\bigoplus_{j \in \mathbb{Z} * \mathbb{Z}} (\mathbb{Z} \oplus \mathbb{Z})_j \oplus (\mathbb{Z} * \mathbb{Z}) \longrightarrow \bigoplus_{j \in \mathbb{Z} \oplus \mathbb{Z}} (\mathbb{Z} \oplus \mathbb{Z})_j \oplus (\mathbb{Z} \oplus \mathbb{Z}).$$

Hence, we can derive:

Corollary 2.1. *There are bijections $\mathcal{C}(\mathfrak{F}_1(\mathbb{S}^1)) \cong (\mathbb{Z} * \mathbb{Z})'$ and $\mathcal{C}(\mathfrak{F}_2(\mathbb{S}^1)) \cong (\mathbb{Z} * \mathbb{Z})'$, where $(\mathbb{Z} * \mathbb{Z})'$ denotes the commutator subgroup of $\mathbb{Z} * \mathbb{Z}$ which is an infinitely generated group.*

Remark 2. Observe that $\pi_k(\bigvee_{j \in J} \bigvee_{i=2}^n (\mathbb{S}^i)^{\gamma_i(n)}) \cong \text{colim}_{J'} \pi_k(\bigvee_{j' \in J'} \bigvee_{i=2}^n (\mathbb{S}^i)^{\gamma_i(n)})$ and

$$\pi_k(\bigvee_{j \in J} \bigvee_{i=2}^n ((\mathbb{S}^i)^{\gamma_i(n)} \vee (\mathbb{S}^i)^{\gamma_i(n)})) \cong \text{colim}_{J'} \pi_k(\bigvee_{j' \in J'} \bigvee_{i=2}^n ((\mathbb{S}^i)^{\gamma_i(n)} \vee (\mathbb{S}^i)^{\gamma_i(n)})),$$

where $J' \subseteq J$ runs over all finite subsets of J . Hence, we can make use of Hilton's result [16] so that the set of multiplications $\mathcal{C}(\mathfrak{F}_n(\mathbb{S}^1))$ might be expressed by means of homotopy groups of appropriate spheres for $n \geq 1$.

Similar to the Fox space, the co-multiplication of $\hat{\mathfrak{F}}_n(W)$ that yields $\tau_{W^{n-1}}(X) = [\Sigma(W^{n-1} \sqcup *), X] = [\hat{\mathfrak{F}}_n(W), X]$ is given by the following split exact sequence which can be deduced from [13, Theorem 3.1]:

$$(2.1) \quad 1 \rightarrow [\Sigma P_{n-1}(W), X] \rightarrow \tau_{W^{n-1}}(X) \overset{\leftarrow}{\rightarrow} \tau_{W^{n-2}}(X) \rightarrow 1$$

which is the same as

$$1 \rightarrow [\Sigma P_{n-1}(W), X] \rightarrow [\hat{\mathfrak{F}}_n(W), X] \overset{\leftarrow}{\rightarrow} [\hat{\mathfrak{F}}_{n-1}(W), X] \rightarrow 1.$$

Here, $P_n(W) = W^n/W^{n-1}$ is the n -fold pinched space of W .

Hence, $[\hat{\mathfrak{F}}_n(W), X] \cong [\Sigma P_{n-1}(W), X] \rtimes [\hat{\mathfrak{F}}_{n-1}(W), X]$ is the semi-direct product with respect to a natural action

$$[\hat{\mathfrak{F}}_{n-1}(W), X] \times [\Sigma P_{n-1}(W), X] \rightarrow [\Sigma P_{n-1}(W), X].$$

In particular, for $X = \hat{\mathfrak{F}}_{n-1}(W) \vee \Sigma P_{n-1}(W)$, by the natural bijection $[\hat{\mathfrak{F}}_{n-1}(W), X] \times [\Sigma P_{n-1}(W), X] \cong [\hat{\mathfrak{F}}_{n-1}(W) \vee \Sigma P_{n-1}(W), X]$, the identity map id_X is sent to the corresponding co-action

$$\alpha : \Sigma P_{n-1}(W) \rightarrow \hat{\mathfrak{F}}_{n-1}(W) \vee \Sigma P_{n-1}(W).$$

Furthermore, the natural bijection $[\Sigma P_{n-1}(W), X] \times [\hat{\mathfrak{F}}_{n-1}(W), X] \cong [\hat{\mathfrak{F}}_n(W), X]$ for any space X yields a homotopy equivalence

$$\hat{\mathfrak{F}}_n(W) \simeq \Sigma P_{n-1}(W) \vee \hat{\mathfrak{F}}_{n-1}(W).$$

Now, write $\nu : \Sigma P_{n-1}(W) \rightarrow \Sigma P_{n-1}(W) \vee \Sigma P_{n-1}(W)$ for the suspension co-multiplication on $\Sigma P_{n-1}(W)$ and $\nu_{n-1} : \hat{\mathfrak{F}}_{n-1}(W) \rightarrow \hat{\mathfrak{F}}_{n-1}(W) \vee \hat{\mathfrak{F}}_{n-1}(W)$ for the co-multiplication on $\hat{\mathfrak{F}}_{n-1}(W)$.

Then, the maps

$$\nu_n^1 : \hat{\mathfrak{F}}_{n-1}(W) \xrightarrow{\nu_{n-1}} \hat{\mathfrak{F}}_{n-1}(W) \vee \hat{\mathfrak{F}}_{n-1}(W) \hookrightarrow \hat{\mathfrak{F}}_n(W) \vee \hat{\mathfrak{F}}_n(W)$$

and

$$\begin{aligned} \nu_n^2 : \Sigma P_{n-1}(W) &\xrightarrow{\nu} \Sigma P_{n-1}(W) \vee \Sigma P_{n-1}(W) \xrightarrow{\alpha \vee 1} \hat{\mathfrak{F}}_{n-1}(W) \vee \Sigma P_{n-1}(W) \vee \Sigma P_{n-1}(W) \\ &\hookrightarrow \hat{\mathfrak{F}}_n(W) \vee \hat{\mathfrak{F}}_n(W) \end{aligned}$$

lead to the suspension co-multiplication

$$\nu_n : \hat{\mathfrak{F}}_n(W) \longrightarrow \hat{\mathfrak{F}}_n(W) \vee \hat{\mathfrak{F}}_n(W)$$

on the space $\hat{\mathfrak{F}}_n(W)$. Thus, we have shown:

Theorem 2.2. *For any $n > 1$, the co-multiplication on the homotopy type $\hat{\mathfrak{F}}_n(W) \simeq \Sigma P_{n-1}(W) \vee \hat{\mathfrak{F}}_{n-1}(W)$ of the space $\hat{\mathfrak{F}}_n(W)$ that corresponds to the group structure on $\tau_{W^{n-1}}$ is given by the map constructed above*

$$\nu_n : \hat{\mathfrak{F}}_n(W) \longrightarrow \hat{\mathfrak{F}}_n(W) \vee \hat{\mathfrak{F}}_n(W).$$

Recall from [13] that the n -th Fox space $F_n = \hat{\mathfrak{F}}^n(\mathbb{S}^1)$ has the same homotopy type, as pointed spaces, of the pinched torus T^n/T^{n-1} and by Proposition 2.2, $F_n \simeq \Sigma F_{n-1} \vee F_{n-1}$. Hence, in light of Theorem 2.2, we derive:

Corollary 2.3. *The co-multiplication on the homotopy type $F_n \simeq \Sigma F_{n-1} \vee F_{n-1}$ of the Fox space F_n that corresponds to the group structure on τ_n is given by Theorem 2.2 for $W = \mathbb{S}^1$.*

We end this section by noting that $P_n(W)$ is not homotopy equivalent to $\hat{\mathfrak{F}}^n(W)$ in general. For example, $P_2(T^2) = T^4/T^2$ whereas $\hat{\mathfrak{F}}^2(T^2) = F_3$.

3. JACOBI IDENTITIES FOR GENERALIZED WHITEHEAD PRODUCTS

In [17], P. Hilton gave a Jacobi identity for the Whitehead product allowing elements of the fundamental group. While the classical Jacobi identity (for elements in higher homotopy groups) of [20] has been generalized for the generalized Whitehead product, Hilton's result [17] has not. In this section, we give a unified approach to the Jacobi identities of Rutter's and Hilton's using the generalized Fox torus groups studied in [13]. We extend

Rutter's result by allowing $\Sigma C = \mathbb{S}^1$ or $\Sigma B = \Sigma C = \mathbb{S}^1$, thereby generalizing Hilton's result for the generalized Whitehead product. When one of the spaces is co- H , we can define the generalized Whitehead product via a group action similar to the treatment of Rutter's in [23] except that Rutter assumed one of the spaces to be a suspension.

Now let A and B be pointed spaces. Via the projections $p_A : ((A \times B) \sqcup *) \rightarrow A$ and $p_B : ((A \times B) \sqcup *) \rightarrow B$, we can regard $[\Sigma A, X]$ and $[\Sigma B, X]$ as subgroups of $\tau_{A \times B}(X)$. Following [18], recall that for any $\alpha \in [\Sigma A, X]$ and $\beta \in [\Sigma B, X]$, the generalized Whitehead product of α and β is the element $[\alpha, \beta] = [K']$, where $K' : \Sigma(A \wedge B) \rightarrow X$ is induced by the map

$$K = f' \cdot g' \cdot f'^{-1} \cdot g'^{-1} : \Sigma(A \times B) \rightarrow X$$

which, when restricted to $\Sigma A \vee \Sigma B$, is homotopic to a constant. Here, $f : \Sigma A \rightarrow X$, $g : \Sigma B \rightarrow X$ are maps representing α and β , respectively and $f' = f \Sigma p_A, g' = g \Sigma p_B$. Using the co-multiplication of $\Sigma(A \times B)$, K is a well-defined map. It is easy to observe that $[\alpha, \beta] = -(\Sigma t)^*[\beta, \alpha]$, where $t : A \wedge B \rightarrow B \wedge A$ is the switching equivalence of smash products.

In [13], we gave another interpretation of the generalized Whitehead product as follows³.

Theorem 3.1. *Let $\alpha \in [\Sigma A, X]$ and $\beta \in [\Sigma B, X]$. Then, the image of $[\alpha, \beta]$ in $\tau_{A \times B}(X)$ given by the homotopy class of the composite*

$$\Sigma((A \times B) \sqcup *) \rightarrow \Sigma(A \times B) \rightarrow \Sigma(A \wedge B) \xrightarrow{K'} X$$

is the commutator of the images of α and β in $\tau_{A \times B}(X)$.

For any pointed spaces A and B with disjoint unions $A \sqcup *_1$ and $B \sqcup *_2$, we have the following identification

$$(A \sqcup *_1) \wedge (B \sqcup *_2) \approx (A \times B) \sqcup *_3,$$

where $*_3$ corresponds to $(A \sqcup *_1) \vee (B \sqcup *_2)$ in the quotient $(A \sqcup *_1) \wedge (B \sqcup *_2)$. Since, by definition, $\tau_W(X) = [\Sigma(W \sqcup *), X]$, it follows that the generalized Whitehead product takes the form of

$$\tau_A(X) \times \tau_B(X) \rightarrow \tau_{A \times B}(X).$$

Hence, for $\alpha \in [\Sigma(A \sqcup *_1), X]$ and $\beta \in [\Sigma(B \sqcup *_2), X]$, the generalized Whitehead product $[\alpha, \beta] \in [\Sigma((A \sqcup *_1) \wedge (B \sqcup *_2)), X] = \tau_{A \times B}(X)$ is a commutator.

³This differs from Theorem 4.1 of [13] in which the generalized Whitehead product of [2] was used.

Thus, the following diagram

$$(3.1) \quad \begin{array}{ccc} \tau_A(X) \times \tau_B(X) & \xrightarrow{(-,-)} & \tau_{A \times B}(X) \\ \uparrow & & \uparrow \\ [\Sigma A, X] \times [\Sigma B, X] & \xrightarrow{GWP} & [\Sigma(A \wedge B), X] \end{array}$$

is commutative.

By using the canonical maps $(A \sqcup *_1) \rightarrow A$ and $(B \sqcup *_2) \rightarrow B$, the image of $[\alpha, \beta]$ given by the composite map in Theorem 3.1 factors through $\tau_A(X) \times \tau_B(X)$. Furthermore, when $A = T^{m-1}$ and $B = T^{n-1}$, the generalized Whitehead product takes the form of

$$GWP : \tau_m(X) \times \tau_n(X) \rightarrow \tau_{m+n-1}(X).$$

This means that the generalized Whitehead product gives rise to a *Whitehead type product* on the Fox torus homotopy groups $\{\tau_k(X)\}$. Moreover, if we denote by WP the classical Whitehead product $\pi_m(X) \times \pi_n(X) \rightarrow \pi_{m+n-1}(X)$ then we have the following commutative diagram

$$(3.2) \quad \begin{array}{ccc} \tau_m(X) \times \tau_n(X) & \xrightarrow{GWP} & \tau_{m+n-1}(X) \\ j_m \times j_n \uparrow & & \uparrow j_{m+n-1} \\ \pi_m(X) \times \pi_n(X) & \xrightarrow{WP} & \pi_{m+n-1}(X), \end{array}$$

where j_k is the inclusion $\prod_{i=2}^k \pi_i(X, x_0)^{\alpha_i} \hookrightarrow \tau_k(X)$ in the split exact sequence (1.1) restricted to the only copy $\pi_k(X)$.

For any group G , Hall [15, p. 150, 10.2.1.4] established the following *Jacobi identity* for elements $a, b, c \in G$:

$$(\star) \quad (b^{-1}, a, c)^b (c^{-1}, b, a)^c (a^{-1}, c, b)^a = 1,$$

where $(x, y, z) = ((x, y), z)$, (x, y) denotes the commutator $xyx^{-1}y^{-1}$ and $x^y = yxy^{-1}$ for $x, y, z \in G$. Furthermore, it can be easily shown

$$(\star\star) \quad ((y^{-1}, x), z)^y = ((x, y), z)((x, y), (z^{-1}, y))^z.$$

To state our Jacobi identity for the generalized Whitehead product, we first recall from [14, Theorem 8.22] the following:

Lemma 3.2. *The natural transformations $A \wedge B \wedge C \rightarrow (A \wedge B) \wedge C$ and $A \wedge B \wedge C \rightarrow A \wedge (B \wedge C)$ are homotopy equivalences provided that A, B and C have the homotopy type of pointed compactly generated Hausdorff spaces.*

Now, we prove the main result of this section. We shall write the operation in $\tau_{A \times B \times C}(X)$ additively.

Theorem 3.3. *Let A, B and C be pointed spaces with the homotopy type of compactly generated Hausdorff spaces. Suppose that $\alpha \in [\Sigma A, X]$, $\beta \in [\Sigma B, X]$ and $\gamma \in [\Sigma C, X]$ and denote by $\bar{\alpha}, \bar{\beta}$, and $\bar{\gamma}$ the respective images in $\tau_{A \times B \times C}(X)$. Then*

(1)

$$(\Sigma t_{213})^* [[\bar{\beta}^{-1}, \bar{\alpha}], \bar{\gamma}]^{\bar{\beta}} + (\Sigma t_{321})^* [[\bar{\gamma}^{-1}, \bar{\beta}], \bar{\alpha}]^{\bar{\gamma}} + (\Sigma t_{132})^* [[\bar{\alpha}^{-1}, \bar{\gamma}], \bar{\beta}]^{\bar{\alpha}} = 1,$$

where t_{ijk} is the appropriate twisting function for $i, j, k = 1, 2, 3$.

(2) *If $\Sigma A, \Sigma B$ and ΣC are homotopy co-commutative co- H -spaces, then*

$$[[\alpha, \beta], \gamma] + (\Sigma t_{312})^* [[\gamma, \alpha], \beta] + (\Sigma t_{231})^* [[\beta, \gamma], \alpha] = 0.$$

(3) *If each of $\Sigma A, \Sigma B$, and ΣC is either homotopy co-commutative or equal to S^1 then*

$$(\Sigma t_{213})^* \beta \cdot [[\beta^{-1}, \alpha], \gamma] + (\Sigma t_{321})^* \gamma \cdot [[\gamma^{-1}, \beta], \alpha] + (\Sigma t_{132})^* \alpha \cdot [[\alpha^{-1}, \gamma], \beta] = 1$$

where

$$\xi \cdot \eta = \begin{cases} \eta, & \text{if } \xi \in [\Sigma W, X] \text{ and } W \neq S^0; \\ \eta^\xi, & \text{if } \xi \in \pi_1(X). \end{cases}$$

Proof. The first assertion (1) follows from the fact that the generalized Whitehead product $[-, -]$ in $\tau_{A \times B \times C}(X)$ coincides with the group theoretic commutator and the Jacobi identity (\star) together with the appropriate twisting functions.

For (2), we may assume, based upon the discussion above, that $\alpha, \beta, \gamma \in \tau_{A \times B \times C}(X)$. Thence, by (\star), we get

$$((\beta^{-1}, \alpha), \gamma)^\beta ((\gamma^{-1}, \beta), \alpha)^\gamma ((\alpha^{-1}, \gamma), \beta)^\alpha = 1$$

in the group $\tau_{A \times B \times C}(X)$. But, by means of ($\star\star$), it holds $((\beta^{-1}, \alpha), \gamma)^\beta = ((\alpha, \beta), \gamma)((\alpha, \beta), (\gamma^{-1}, \beta))^\gamma$. Because of a homomorphism $\Sigma((A \times C) \wedge B) \approx (A \times C) \wedge \Sigma B$, both elements (α, β) and (γ^{-1}, β) lie in the image of the abelian group $[\Sigma((A \times C) \wedge B), X]$ in the group $\tau_{A \times B \times C}(X)$ and we deduce that $((\alpha, \beta), (\gamma^{-1}, \beta))^\gamma = 1$. Similarly, $((\beta, \gamma), (\alpha^{-1}, \gamma))^\alpha = ((\gamma, \alpha), (\beta^{-1}, \alpha))^\beta = 1$. Finally, in light of Theorem 3.1, the following equation

$$[[\alpha, \beta], \gamma] + (\Sigma t_{231})^* [[\beta, \gamma], \alpha] + (\Sigma t_{312})^* [[\gamma, \alpha], \beta] = 0$$

holds in the image of the abelian group $[\Sigma(A \wedge B \wedge C), X]$ in the group $\tau_{A \times B \times C}(X)$ and (2) is proven. This is the same as [23, Theorem 1] but our spaces need not be CW complexes.

To prove (3), we note that the convention $\xi \cdot \eta$ is valid for the following reasons. When ξ is not in $\pi_1(X)$, the image of η^ξ is conjugation in the group τ and this will yield ξ under the assumption that $\eta \in [\Sigma B, X]$ and ΣB is co-commutative. On the other hand, if $\xi \in \pi_1(X)$ then η^ξ is given by the classical action of $\pi_1(X)$. Now, (3) is simply the same as (1) when pulled

back to $[\Sigma A \wedge B \wedge C, X]$. This is a generalization of formula (12) of Hilton's [17].

□

Note that the obvious pointed projection $A \sqcup * \rightarrow S^0$ leads to the inclusion $\pi_1(X) \subseteq \tau_A(X)$ for any nonempty topological space A and a pointed space X . In fact, it has been shown in [13] that $\tau_A(X) \cong [\Sigma A, X] \rtimes \pi_1(X)$ for any pointed spaces A and X . Furthermore, this can be deduced from the following split exact sequence of [13, Theorem 3.1]

$$(3.3) \quad 1 \rightarrow [(V \times W)/V, \Omega X] \rightarrow \tau_{V \times W}(X) \xrightarrow{\leftarrow} \tau_V(X) \rightarrow 1$$

which generalizes that of Fox.

When V is a point and $W = A$, we have $\tau_A(X) \cong [\Sigma A, X] \rtimes \pi_1(X)$. The splitting (3.3) gives the classical action of $\pi_1(X)$ on $[\Sigma A, X]$. Furthermore, when $V = A$ and $W = B$, this splitting gives rise to an action of $[\Sigma A, X] (\subseteq \tau_A(X))$ on $[(A \times B)/A, \Omega X]$. Since $[(A \times B)/A, \Omega X] = [\Sigma((A \times B)/A), X] \cong [\Sigma(A \wedge B), X] \rtimes [\Sigma B, X]$, we have an action of $[\Sigma A, X]$ on $[\Sigma(A \wedge B), X] \rtimes [\Sigma B, X]$.

Next, we give an alternative definition of the generalized Whitehead product when one of the spaces is co- H , thereby generalizing that of Rutter's [23].

When B is a co- H space, it follows from the proof of [13, Theorem 3.1] that $\Sigma((A \times B)/A) \simeq \Sigma(B \vee (A \wedge B))$ is co-commutative so that $[\Sigma((A \times B)/A), X]$ is abelian and $[\Sigma((A \times B)/A), X] \cong [\Sigma(A \wedge B), X] \times [\Sigma B, X]$. Given $\alpha \in [\Sigma A, X]$ and $\beta \in [\Sigma B, X]$, we define

$$\alpha * \beta := \alpha \cdot \tilde{\beta}$$

as the action of α on the image $\tilde{\beta}$ of β by the inclusion $[\Sigma B, X] \hookrightarrow [\Sigma(A \wedge B), X] \rtimes [\Sigma B, X]$.

Then, the image of the generalized Whitehead product $[\alpha, \beta]$ is given by the following relation in the group $\tau_{A \times B}(X)$:

$$(3.4) \quad \overline{\alpha * \beta} = \overline{[\alpha, \beta]} + \tilde{\beta}.$$

Evidently, if A is co- H , then one can define a similar action $\beta * \alpha$ so that (3.4) becomes $\overline{\beta * \alpha} = \overline{[\beta, \alpha]} + \tilde{\alpha}$. These actions are analogous to the actions given in [23, Theorem 2].

4. Γ -FOX GROUPS AND JACOBI IDENTITIES

In [22], N. Oda defined a generalization of the generalized Whitehead product (Γ -Whitehead product) by replacing the suspension Σ with the smash product with a co-grouplike space Γ . A Jacobi identity was given

in [22, Theorem 2.11] for $[\Gamma \wedge A, X]$, $[\Gamma \wedge B, X]$ and $[\Gamma \wedge C, X]$. In the case when $\Gamma = \mathbb{S}^1$, Oda established the same Jacobi identity as Rutter under the assumption that A, B and C be co- H spaces. Thus, Rutter's result does not follow from Oda's since Oda's hypotheses imply that $\Sigma A, \Sigma B$ and ΣC are necessarily homotopy co-commutative but a space Z need not be co- H while ΣZ is homotopy co-commutative.

We end the paper by showing that the Fox group $\tau_W(X)$ can easily be generalized by replacing Σ with Γ and thus we can obtain the same kind of Jacobi identities as in the last section. These identities generalize those obtained in [22].

Definition 3. Let X be a space and $x_0 \in X$ and Γ a co-grouplike space. For any space W , the W - Γ -Fox group of X is defined to be

$$\tau_M^\Gamma(X) = \tau_W^\Gamma(X, x_0) = [\Gamma \wedge (W \sqcup *), X].$$

Note that

$$\tau_W^\Gamma(X) = [\Gamma(W \sqcup *), X] = [(W \sqcup *) \wedge \Gamma, X] = [(W \sqcup *), X_*^\Gamma],$$

where X_*^Γ denotes the space of basepoint preserving maps from Γ to X . Following the notation of [22], we write $\Gamma W = \Gamma \wedge W$.

Theorem 4.1. Let Γ be a co-grouplike space and B a well-pointed space. Then for any pointed space X and any space A , the following sequence

$$(4.1) \quad 1 \rightarrow [\Gamma((A \times B)/A), X] \rightarrow \tau_{A \times B}^\Gamma(X) \overset{\leftarrow}{\rightleftarrows} \tau_A^\Gamma(X) \rightarrow 1$$

is split exact.

Proof. The space B is well-pointed, so we can consider the split cofibration

$$A \sqcup * \overset{\leftarrow}{\rightleftarrows} (A \times B) \sqcup * \rightarrow (A \times B)/A.$$

The corresponding Barrett-Puppe sequence yields the following short split-exact sequence

$$(4.2) \quad 1 \rightarrow [(A \times B)/A, Z] \rightarrow [(A \times B) \sqcup *, Z] \overset{\leftarrow}{\rightleftarrows} [A \sqcup *, Z] \rightarrow 1$$

of pointed sets for any pointed space Z . For $Z = X_*^\Gamma$ using the adjoint isomorphism, (4.2) yields the desired short split-exact sequence. \square

In particular, we have the following:

Corollary 4.2. Let W be a well-pointed space. Then the split cofibration $* \sqcup * \rightarrow W \sqcup * \rightarrow W$ leads to the isomorphism

$$\tau_W^\Gamma(X) \cong [\Gamma W, X] \times [\Gamma, X].$$

Given $\alpha \in [\Gamma A, X], \beta \in [\Gamma B, X]$, the Γ -Whitehead product $[\alpha, \beta]_{\Gamma} \in [\Gamma(A \wedge B), X]$ is defined (see [22]) in the same way as the generalized Whitehead product GWP by simply replacing Σ by Γ . The co-multiplication of Γ allows the proof of Theorem 4.1 of [13] to remain valid when GWP is replaced by $[-, -]_{\Gamma}$. Thus, the proof of Theorem 4.1 of [13] yields the following Γ -analog of Theorem 3.1.

Theorem 4.3. *Let $\alpha \in [\Gamma A, X]$ and $\beta \in [\Gamma B, X]$. Then, the image of $[\alpha, \beta]_{\Gamma}$ in $\tau_{A \times B}^{\Gamma}(X)$ given by the homotopy class of the composite*

$$\Gamma((A \times B) \sqcup *) \rightarrow \Gamma(A \times B) \rightarrow \Gamma(A \wedge B) \xrightarrow{K'_{\Gamma}} X$$

is the commutator of the images of α and β in $\tau_{A \times B}^{\Gamma}(X)$.

Finally, we observe that the proof of Theorem 3.3 remains valid when we replace Σ with Γ and the generalized Whitehead product with the Γ -Whitehead product. Thus, we have:

Theorem 4.4. *Let A, B and C be pointed spaces with the homotopy type of compactly generated Hausdorff spaces and let Γ be a co-grouplike space. Suppose $\alpha \in [\Gamma A, X], \beta \in [\Gamma B, X]$ and $\gamma \in [\Gamma C, X]$ and denote by $\bar{\alpha}, \bar{\beta}$, and $\bar{\gamma}$ the respective images in $\tau_{A \times B \times C}^{\Gamma}(X)$. Here $[-, -]_{\Gamma}$ denotes the corresponding Γ -Whitehead product. Then*

(1)

$$(\Gamma t_{213})^* [[\bar{\beta}^{-1}, \bar{\alpha}]_{\Gamma}, \bar{\gamma}]_{\Gamma}^{\bar{\beta}} + (\Gamma t_{321})^* [[\bar{\gamma}^{-1}, \bar{\beta}]_{\Gamma}, \bar{\alpha}]_{\Gamma}^{\bar{\gamma}} + (\Gamma t_{132})^* [[\bar{\alpha}^{-1}, \bar{\gamma}]_{\Gamma}, \bar{\beta}]_{\Gamma}^{\bar{\alpha}} = 1.$$

(2) *If $\Gamma A, \Gamma B$ and ΓC are homotopy co-commutative co- H -spaces, then*

$$[[\alpha, \beta]_{\Gamma}, \gamma]_{\Gamma} + (\Gamma t_{312})^* [[\gamma, \alpha]_{\Gamma}, \beta]_{\Gamma} + (\Gamma t_{231})^* [[\beta, \gamma]_{\Gamma}, \alpha]_{\Gamma} = 0.$$

(3) *If each of $\Gamma A, \Gamma B$, and ΓC is either homotopy co-commutative or equal to Γ then*

$$(\Gamma t_{213})^* \beta \cdot [[\beta^{-1}, \alpha]_{\Gamma}, \gamma]_{\Gamma} + (\Gamma t_{321})^* \gamma \cdot [[\gamma^{-1}, \beta]_{\Gamma}, \alpha]_{\Gamma} + (\Gamma t_{132})^* \alpha \cdot [[\alpha^{-1}, \gamma]_{\Gamma}, \beta]_{\Gamma} = 1$$

where

$$\xi \cdot \eta = \begin{cases} \eta, & \text{if } \xi \in [\Gamma W, X] \text{ and } \Gamma W \neq \Gamma; \\ \eta^{\xi}, & \text{if } \xi \in [\Gamma, X]. \end{cases}$$

Remark 3. For the Γ -Whitehead product, one can give an alternative definition in terms of a group action similar to (3.4) under the hypothesis that one of the spaces is co- H .

ACKNOWLEDGEMENT

The authors are very grateful to Professor N. Oda for his very careful reading of the previous version of this paper and all valuable comments and suggestions. We also thank Professor M. Arkowitz for bringing [8] to our attention regarding spaces X of weak category > 1 while ΣX is homotopy co-commutative.

REFERENCES

- [1] Ando, H., On the generalized Whitehead products and the generalized Hopf invariant of a composition element, *Tôhoku Math. J. (2)* **20** (1968), 516–553.
- [2] Arkowitz, M., The generalized Whitehead product, *Pacific J. Math.* **12** (1962), 7–23.
- [3] Barcus, W. and Barratt, M., On the homotopy classification of the extensions of a fixed map, *Trans. Amer. Math. Soc.* **88** (1958), 57–74.
- [4] Berstein, I. and Ganea, T., On the homotopy-commutativity of suspensions, *Illinois J. Math.* **6** (1962), 336–340.
- [5] Boardman, J. and Steer, B., On Hopf invariants, *Comment. Math. Helv.* **42** (1967), 180–221.
- [6] Chang, S., On Jacobi identity, (Chinese) *Acta Math. Sinica* **4** (1954), 365–379.
- [7] Cohen, D.E., Products and carrier theory, *Proc. London Math. Soc.* **7** (1957), 219–248.
- [8] Cornea, O., Lupton, G., Oprea, J., and Tanré, D., *Lusternik-Schnirelmann category*, Mathematical Surveys and Monographs, **103**, American Mathematical Society, Providence, RI, 2003.
- [9] Fox, R., Torus homotopy groups, *Proc. Nat. Acad. Sci. (U.S.A.)* **31** (1945), 71–74.
- [10] Fox, R., Homotopy groups and torus homotopy groups, *Ann. Math.* **49** (1948), 471–510.
- [11] Gilbert, W., Some examples for weak category and conilpotency, *Illinois J. Math.* **12** (1968), 421–432.
- [12] Golasinski, M. and Gonçalves D.L., Comultiplications of the wedge of two Moore spaces, *Colloq. Math.* **76** (1998), no. 2, 229–242.
- [13] Golasinski, M., Gonçalves D.L., and Wong, P., Generalizations of Fox homotopy groups, Whitehead products, and Gottlieb groups, *Ukrain. Math. Zh.* **57** (2005), no. 3, 320–328 (translated in *Ukrainian Math J.* **57** (2005) no. 3, 382–393).
- [14] Gray, B., *Homotopy Theory. An Introduction to Algebraic Topology*, Academic Press, New York, San Francisco, London (1975).
- [15] Hall, M., Jr. *Theory of groups*, New York (1959).
- [16] Hilton, P., On homotopy groups of union of spheres, *J. London Math. Soc.* **30** (1955), 154–172.
- [17] Hilton, P., Note on the Jacobi identity for Whitehead products, *Proc. Cambridge Philos. Soc.* **57** (1961), 180–182.
- [18] Hilton, P., “Homotopy Theory and Duality,” Gordon and Breach Science Publishers, New York-London-Paris 1965 x+224 pp.
- [19] Milnor, J.W., On spaces having the homotopy type of a CW -complex, *Trans. Amer. Math. Soc.*, **90** (1959), 272–280.
- [20] Nakaoka, M. and Toda, H., On Jacobi identity for Whitehead products, *J. Inst. Polytech. Osaka City Univ. Ser. A* **5** (1954), 1–13.

- [21] Neisendorfer, J., Primary homotopy theory, *Mem. Amer. Math. Soc.* **25** (1980), no. 232, iv+67 pp.
- [22] Oda, N., A generalization of the Whitehead product, *Math. J. Okayama Univ.* **39** (1997), 113–133 (1999).
- [23] Rutter, J.W., Two theorems on Whitehead products, *J. London Math. Soc.*, **43** (1968), 509–512.
- [24] Rutter, J.W., Correction to “Two theorems on Whitehead products”, *J. London Math. Soc.*, (2), **1** (1969), 20.
- [25] Suzuki, H., A product in homotopy theory, *Tôhoku Math. J. (2)* **6** (1954), 78–88.
- [26] Uehara, H. and Massey, W., The Jacobi identity for Whitehead products, *Algebraic geometry and topology* (A symposium in honor of S. Lefschetz), pp. 361–377, Princeton University Press, Princeton, N. J., 1957.
- [27] Whitehead, J.H.C., On adding relations to homotopy groups, *Ann. Math.* **42** (1941), 400–428.
- [28] Whitehead, G., “Elements of Homotopy Theory,” Springer-Verlag, New York, 1978.
- [29] Whitehead, G., On mappings into group-like spaces, *Comment. Math. Helv.* **28** (1954), 320–328.

MAREK GOLASIŃSKI
FACULTY OF MATHEMATICS AND COMPUTER SCIENCE
NICOLAUS COPERNICUS UNIVERSITY
CHOPINA 12/18, 87-100 TORUŃ
POLAND

e-mail address: marek@mat.uni.torun.pl

DACIBERG LIMA GONÇALVES
DEPARTAMENTO DE MATEMÁTICA - IME - USP
CAIXA POSTAL 66.281 - CEP 05311-970
SÃO PAULO - SP
BRASIL

e-mail address: dlgoncal@ime.usp.br

PETER WONG
DEPARTMENT OF MATHEMATICS
BATES COLLEGE
LEWISTON, ME 04240
U.S.A.

e-mail address: pwong@bates.edu

(Received February 6, 2007)

(Revised May 4, 2007)