

# *Acta Medica Okayama*

---

*Volume 54, Issue 3*

2000

*Article 3*

JUNE 2000

---

## The perineuronal proteoglycan surface coat in the adult rat brain, with special reference to its reactions to Gomori's ammoniacal silver.

Ryutaro Endo\*

Shinichiro Murakami†

Yu Masuda‡

Takehito Taguchi\*\*

Aiji Ohtsuka††

Kazunori Nishizaki‡‡

Takuro Murakami§

\*Okayama University,

†Okayama University,

‡Okayama University,

\*\*Okayama University,

††Okayama University,

‡‡Okayama University,

§Okayama University,

# The perineuronal proteoglycan surface coat in the adult rat brain, with special reference to its reactions to Gomori's ammoniacal silver.\*

Ryutaro Endo, Shinichiro Murakami, Yu Masuda, Takehito Taguchi, Aiji Ohtsuka, Kazunori Nishizaki, and Takuro Murakami

## Abstract

The present study showed that many neurons in the adult rat brain possessed a perineuronal sulfated proteoglycan surface coat which reacted to cationic iron colloid and aldehyde fuchsin. This surface coat was stained supravivally with Ehrlich's methylene blue and doubly stained with Ehrlich's methylene blue and aldehyde fuchsin. The surface coat was also stained with Gomori's ammoniacal silver and doubly stained with Gomori's ammoniacal silver and cationic iron colloid. The surface coat was usually expressed together with a nerve cell surface glycoprotein net detectable with lectin *Wisteria floribunda* agglutinin. These findings indicate that the perineuronal proteoglycan surface coat is identical to Cajal's superficial reticulum and contains some collagenous elements. It was further demonstrated that collagenase digestion erased Gomori's ammoniacal silver impregnation within the perineuronal proteoglycan surface coat.

**KEYWORDS:** brain, extracellular matrix, perineuronal proteoglycans, cell surface glycoproteins

---

\*PMID: 10925735 [PubMed - indexed for MEDLINE]

Copyright (C) OKAYAMA UNIVERSITY MEDICAL SCHOOL

## The Perineuronal Proteoglycan Surface Coat in the Adult Rat Brain, with Special Reference to its Reactions to Gömöri's Ammoniacal Silver

Ryutaro ENDO, Shinichiro MURAKAMI<sup>a</sup>, Yu MASUDA, Takehito TAGUCHI<sup>a</sup>, Aiji OHTSUKA<sup>a</sup>, Kazunori NISHIZAKI and Takuro MURAKAMI<sup>a\*</sup>

Departments of Otolaryngology and <sup>a</sup>Anatomy, Okayama University Medical School, Okayama 700-8558, Japan

The present study showed that many neurons in the adult rat brain possessed a perineuronal sulfated proteoglycan surface coat which reacted to cationic iron colloid and aldehyde fuchsin. This surface coat was stained supravivally with Ehrlich's methylene blue and doubly stained with Ehrlich's methylene blue and aldehyde fuchsin. The surface coat was also stained with Gömöri's ammoniacal silver and doubly stained with Gömöri's ammoniacal silver and cationic iron colloid. The surface coat was usually expressed together with a nerve cell surface glycoprotein net detectable with lectin *Wisteria floribunda* agglutinin. These findings indicate that the perineuronal proteoglycan surface coat is identical to Cajal's superficial reticulum and contains some collagenous elements. It was further demonstrated that collagenase digestion erased Gömöri's ammoniacal silver impregnation within the perineuronal proteoglycan surface coat.

**Key words:** brain, extracellular matrix, perineuronal proteoglycans, cell surface glycoproteins

**O**ur previous histochemical and electron microscopic studies showed that many neurons in the central nervous systems of humans, dogs, cats, rats, mice and other animals possess a marked perineuronal sulfated proteoglycan surface coat which is stained with cationic iron colloid and aldehyde fuchsin (1-16). Additional experiments (8-10, 17) also showed that the surface coat reacts to Golgi's or Kopsch's silver nitrate and is identical to Golgi's reticular coating (18, 19). Recently, we also demonstrated that the perineuronal proteoglycan surface coat in the mouse brain is stained supravivally with Ehrlich's methylene blue, impregnated with Gömöri's

ammoniacal silver and digested by collagenase (20, 21).

The present study reinvestigated the adult rat brain and confirmed our recent findings concerning the mouse brain. The retrosplenial cortex formed the material of the present study, as it contains many neurons with well developed perineuronal proteoglycan surface coats (7, 8, 12, 13, 20, 21).

### Materials and Methods

Adult male Wistar rats weighing about 250 g were anesthetized with ethyl ether. After ligation of their thoracic aorta, they were treated as follows.

**Staining with cationic iron colloid, ammoniacal silver, aldehyde fuchsin or lectin *Wisteria floribunda* agglutinin.** The animals were perfused through the ascending aorta with 15 ml of Ringer's solution and with 15 ml of 4% paraformaldehyde in a 0.1 M cacodylate buffer (pH 7.2). Each brain was then isolated, and 1-2 mm-thick blocks traversing the retrosplenial cortex were prepared with a vibratome. These blocks were refixed in the 4% paraformaldehyde fixative, embedded in paraffin, and cut into 10-15  $\mu$ m-thick sections.

The sections were deparaffinized with xylene and stained with our fine cationic iron colloid with pH values of 1.0-1.5 (22), Gömöri's ammoniacal silver (23, 24), Fujita's aldehyde fuchsin (25), or lectin *Wisteria floribunda* agglutinin (26). They were then embedded in balsam.

Cationic iron colloid stainability was demonstrated by a Prussian blue reaction. In the lectin labeling, peroxidase activity was demonstrated with diaminobenzidine. Counter-staining with nuclear fast red or Mayer's hematoxylin was sometimes done prior to the balsam-

---

\* To whom correspondence should be addressed.

embedding. Controls for lectin-labeled sections consisted of adjacent sections treated with a phosphate buffer containing no agglutinin.

**Double staining with ammoniacal silver and cationic iron colloid.** Some sections stained with Gömöri's ammoniacal silver were successively stained with fine cationic iron colloid (pH 1.0-1.5) (20). They were then counter-stained with nuclear fast red and embedded in balsam.

**Supravital staining with methylene blue.** The animals were perfused through the ascending aorta with 5 ml of Ringer's solution, 15 ml of 0.25% Ehrlich's methylene blue in 0.9% NaCl, and 10 ml of saturated ammonium picrate (27). The brains were then isolated, and 1-2 mm-thick blocks traversing the retrosplenial cortex were prepared with a vibratome. These blocks were frozen and cut into 15-20  $\mu$ m-thick sections and embedded in glycerol.


**Double staining with methylene blue and aldehyde fuchsin, or methylene blue and lectin *Wisteria floribunda* agglutinin.** Some sections from the supravital methylene blue-stained (methylene blue/ammonium picrate-treated) samples were refixed with 4% paraformaldehyde in a 0.1 M cacodylate buffer (pH 7.2) for 10 min, stained with Fujita's aldehyde fuchsin or lectin *Wisteria floribunda* agglutinin, and embedded in balsam (20). Counter-staining with nuclear fast red was sometimes done prior to balsam-embedding.

**Collagenase digestion.** Some sections were rinsed with distilled water and incubated with 40 U/ml collagenase (*Cl. histolyticum*) (Elastin Products Co., MO, USA) in distilled water for 60 min at 37 °C (28). These sections were impregnated with Gömöri's ammoniacal silver (21). Control sections were incubated in distilled water containing no enzyme and stained with ammoniacal silver.

**Light microscopy.** All sections prepared as described above were observed with a transmission light microscope (Olympus BX 50 or BH2).


## Results

**Staining with cationic iron colloid, ammoniacal silver, aldehyde fuchsin or lectin *Wisteria floribunda* agglutinin.** Many neurons (15-20% of the neurons) in the retrosplenial cortex possessed a surface coat which reacted to the cationic iron colloid (Fig. 1), Gömöri's ammoniacal silver (Fig. 2),



**Fig. 1** Retrosplenial cortex of an adult rat brain stained with cationic iron colloid (pH 1.5), treated for a Prussian blue reaction, and counter-stained with nuclear fast red. A perineuronal proteoglycan surface coat is demonstrated by a Prussian blue reaction (single arrowhead). **Upper Inset** shows a retrosplenial section stained with aldehyde fuchsin and nuclear fast red. The core protein of the perineuronal proteoglycan surface coat reacts to aldehyde fuchsin (single arrowhead). **Lower Inset** shows a retrosplenial section of an adult rat brain treated with lectin *Wisteria floribunda* agglutinin. Some neurons express cell surface glycoproteins with terminal *N*-acetylgalactosamine reactive to lectin *Wisteria floribunda* agglutinin (single arrowhead).  $\times 1,200$ ; Insets,  $\times 1,200$ .

Fujita's aldehyde fuchsin (Fig. 1 Upper Inset), or lectin *Wisteria floribunda* agglutinin (Fig. 1 Lower Inset). In each case, the surface coat appeared as a fine meshwork surrounding the nerve cell bodies and their main processes (Figs. 1, 2). In the control sections for lectin labeling, no neurons were labeled with lectin.



**Fig. 2** Retrosplenial cortex of an adult rat brain stained with Gömöri's ammoniacal silver. The perineuronal proteoglycan surface coat contains some collagenous elements reactive to ammoniacal silver (single arrowhead). **Inset** shows a retrosplenial section treated with Gömöri's ammoniacal silver and cationic iron colloid. The perineuronal proteoglycan surface coat is doubly stained with ammoniacal silver and cationic iron colloid (double arrowheads).  $\times 1,200$ ; Inset,  $\times 900$ .

**Double staining with ammoniacal silver and cationic iron colloid.** In the retrosplenial cortex, neurons stained with ammoniacal silver always reacted to the cationic iron colloid. More precisely, the surface coat was constantly stained with ammoniacal silver and cationic iron colloid (Fig. 2 Inset).

**Supravital staining with methylene blue.** Supravital methylene blue staining by vascular perfusion preferentially stained the nerve cell surfaces (Fig. 3).



**Fig. 3** Retrosplenial cortex supravitaly stained with Ehrlich's methylene blue. The surface coat is stained supravitaly with methylene blue (single arrowhead). **Inset** shows the retrosplenial cortex successively stained with Ehrlich's methylene blue and aldehyde fuchsin. The surface coat is stained doubly with methylene blue and aldehyde fuchsin (double arrowheads).  $\times 500$ ; Inset,  $\times 500$ .

Thus, 10–15% of the neurons in the retrosplenial cortex possessed a surface coat which reacted to methylene blue. The surface coat thus stained with methylene blue showed a meshwork structure surrounding the cell bodies and main processes of the nerve cells (Fig. 3).

Neither perivascular reticular fibers nor glial cell nuclei reacted to methylene blue. Elastic fibers around the arterial vessels also showed no reaction to methylene blue.

**Double staining with methylene blue and**

**aldehyde fuchsin.** In the retrosplenial cortex, the neurons supravitaly stained with methylene blue always reacted to aldehyde fuchsin (Fig. 3 Inset). More precisely, the surface coat was constantly doubly stained with methylene blue and aldehyde fuchsin.

**Double staining with methylene blue and lectin *Wisteria floribunda* agglutinin.** The neurons supravitaly stained with methylene blue were not always labeled with lectin *Wisteria floribunda* agglutinin (Fig. 4); 5% of the neurons stained with methylene blue were not labeled with lectin *Wisteria floribunda* agglutinin, and 5% of the neurons labeled with this lectin were not stained with methylene blue.


**Collagenase digestion.** In the collagenase-treated sections, the perineuronal surface coat was not stained with Gömöri's ammoniacal silver (Fig. 5A). In the control sections, the perineuronal surface was well stained with ammoniacal silver (Fig. 5B).

## Discussion

Recent histochemical, immunohistochemical and biochemical studies of humans and various animals, including mice, have shown that mature as well as immature central nervous tissues express many proteoglycans and that some of these proteoglycans form an extracellular matrix to cover certain neuron subsets (26, 29-43).

The present study, together with our previous studies of humans, dogs, cats, mice and other animals, including lower vertebrates such as reptiles and fish (5-7, 9, 12, 14-16, 20), confirmed that such perineuronal proteoglycan surface coats are stained with cationic iron colloid and aldehyde fuchsin and that cationic iron colloid and aldehyde fuchsin stain the sulfate groups and core proteins of the proteoglycans, respectively. Distribution of the neurons with this proteoglycan surface coat in the mouse brain and spinal cord was compared with that of the neurons whose cell surfaces were labeled with lectin *Wisteria floribunda* agglutinin (5-8, 12, 14-16).

Our previous studies of human, rat and mouse brain samples (8-10, 17) showed that the perineuronal proteoglycan surface coat is usually stained with Golgi's or Kopsch's silver nitrate and is identical to Golgi's reticular coating (18, 19). Our previous studies of human, rat and mouse brain samples showed furthermore that the silver staining of the surface coat is erased by hyaluronidase digestion and not by chondroitinase ABC/keratanase/heparitinase digestions, indicating that Golgi's or Kopsch's




**Fig. 4** Retrosplenial cortex successively stained with methylene blue, lectin *Wisteria floribunda* agglutinin and nuclear fast red. The surface coat is usually stained doubly with methylene blue and lectin agglutinin (double arrowheads). In some nerve cells, the surface coat is stained only with methylene blue (single arrowhead) or lectin agglutinin.  $\times 1,000$ .

silver nitrate stains the core proteins of proteoglycans (8, 9, 17).

The present study together with our previous studies of the mouse brain (20, 21) showed that the perineuronal proteoglycan is stained with Gömöri's ammoniacal silver and that it can be stained with this ammoniacal silver and cationic iron colloid. The present study also showed that Gömöri's ammoniacal silver staining is erased by collagenase digestion. These findings indicate that the proteoglycan surface coat contains some additional collagenous elements (reactive to Gömöri's ammoniacal





**Fig. 5** A retrosplenial section incubated with collagenase and stained with Gömöri's ammoniacal silver. **B** shows a control section stained with silver without preliminary collagenase incubation. In **A**, no neurons react to Gömöri's ammoniacal silver. In contrast, **B** shows that many neurons react to the ammoniacal silver (single arrowhead). The single arrow indicates the reticular network surrounding the vascular walls. **A** and **B**,  $\times 400$ .

silver) linked to the cell surfaces.

The present study showed further that the perineuronal proteoglycan surface coat is stained supravivally with Ehrlich's methylene blue and that it is stained doubly with this methylene blue and aldehyde fuchsin. These findings indicate that the perineuronal proteoglycan surface coat is identical to Cajal's superficial reticulum or red superficial (44), or réseaux péricellulaires (45). Golgi's reticular coating and Cajal's pericellular reticulum are defined as perineuronal meshwork structures which are demonstrated by means of Golgi's or Kopsch's silver nitrate or

Ehrlich's methylene blue (8, 19, 44, 45). Many researchers have stained adult rat or mouse brain sections with lectin *Vicia villosa*, soybean or *Wisteria floribunda* agglutinin and have discovered another perineuronal meshwork structure reactive to these lectins which detected the cell surface glycoproteins (26, 46-59). Distribution of the neurons with such a lectin-labeled meshwork in the rat brain and spinal cord has been compared with that of the neurons with surface coats reactive to Mowry's colloidal iron hydroxide (50, 51, 60). Brückner *et al.* (50, 51) and Blümcke *et al.* (54) showed by means of electron

microscopes that the lectin-binding sites were scattered throughout the perineuronal tissue spaces.

The present study showed that the neurons reactive to methylene blue were not always labeled with lectin *Wisteria floribunda* agglutinin and that the neurons reactive to this lectin were not always stained with methylene blue. Similar findings were obtained in our previous double staining of human, rat and mouse brain samples with lectin (lectin *Wisteria floribunda*, *Vicia villosa* or soybean agglutinin) and cationic iron colloid (5-9, 12, 14-16). We believe that the nerve cell surfaces are stained only with methylene blue or iron colloid when the cell surface glycoproteins are occupied thoroughly by the perineuronal proteoglycans; that the cell surfaces are stained solely with lectin agglutinin when the cell surface glycoproteins are thoroughly free from the perineuronal proteoglycans; and that the cell surface are doubly stained with methylene blue and lectin agglutinin or with iron colloid and lectin agglutinin when the cell surface glycoproteins are reacting in part to the perineuronal proteoglycans (20).

Recently, Müller (61) supravitaly stained an adult mouse brain with methylene blue and reported that this dye is attracted to anionic sites or sulfate groups of the perineuronal proteoglycans. However, our experiments indicated that methylene blue and Gömöri's ammoniacal silver react to the collagenous elements of the perineuronal proteoglycan surface coat (20, 21).

**Acknowledgements.** We are grateful to Mr. H Kusano for his help in tissue preparations and stainings.

## References

- Murakami T, Taguchi T, Ohtsuka A and Kikuta A: Neurons with strongly negative-charged surface-coats in adult rat brain as detected by staining with cationic iron colloid. *Arch Histol Cytol* (1993) **56**, 13-21.
- Murakami T, Taguchi T and Ohtsuka A: The occurrence in the human brain of neurons with strongly negative-charged proteoglycans. *Arch Histol Cytol* (1993) **56**, 23-26.
- Murakami T, Tsubouchi Y, Tsubouchi M, Ohtsuka A, Taguchi T and Kikuta A: The occurrence of rat spinal cord neurons with strongly negative-charged surface coats. *Arch Histol Cytol* (1993) **56**, 501-504.
- Murakami T, Tsubouchi M, Tsubouchi Y, Taguchi T and Ohtsuka A: The occurrence of neurons with strongly negatively charged surface coats in mammalian, avian, reptilian, amphibian and piscine brains. *Acta Med Okayama* (1994) **48**, 195-197.
- Murakami T, Ohtsuka A and Taguchi T: Neurons with intensely negatively charged extracellular matrix in the human visual cortex. *Arch Histol Cytol* (1994) **57**, 509-522.
- Murakami T, Hitomi S, Ohtsuka A and Taguchi T: Neurons with perineuronal sulfated proteoglycans in the human visual cortex, with special reference to their reactions to lectins. *Arch Histol Cytol* (1995) **58**, 357-364.
- Murakami T, Ohtsuka A, Taguchi T and Piao DX: Perineuronal sulfated proteoglycans and dark neurons in the brain and spinal cord: A histochemical and electron microscopic study of newborn and adult mice. *Arch Histol Cytol* (1995) **58**, 557-565.
- Murakami T, Ohtsuka A and Ono K: Neurons with perineuronal sulfated proteoglycans in the mouse brain and spinal cord: Their distribution and reactions to lectin *Vicia villosa* agglutinin and Golgi's silver nitrate. *Arch Histol Cytol* (1996) **59**, 219-231.
- Murakami T, Ohtsuka A and Piao DX: Perineuronal sulfated proteoglycans in the human brain are identical to Golgi's reticular coating. *Arch Histol Cytol* (1996) **59**, 233-237.
- Murakami T, Murakami T, Mahmut N, Hitomi S and Ohtsuka A: Dark and light neurons in the human brain, with special reference to their reaction to Golgi's silver nitrate, luxol fast blue MBS and azocarmine G. *Arch Histol Cytol* (1997) **60**, 265-274.
- Murakami T, Su WD, Hong LJ, Piao DX, Ohtsuka A and Sano K: Demonstration of tissue-reacted cationic iron colloid with protein silver and gold chloride (Japanese text with English abstract). *Okayama Igakkai Zasshi* (1997) **109**, 151-156 (in Japanese).
- Murakami T, Murakami T, Hong LJ, Su WD, Piao DX, Mahmut N and Ohtsuka A: Perineuronal sulfated proteoglycans and cell surface glycoproteins in adult and newborn mouse brains, with special reference to their postnatal developments. *Arch Histol Cytol* (1997) **60**, 347-354.
- Ohtsuka A and Murakami T: Dark neurons in the mouse brain: An investigation into the possible significance of variable appearance within a day and their relation to negatively charged cell coats. *Arch Histol Cytol* (1996) **59**, 79-85.
- Tsubouchi Y, Tsubouchi M, Hitomi S, Ohtsuka A and Murakami T: Perineuronal sulfated proteoglycans in the adult rat brain: Histochemical and electron microscopic studies. *Acta Med Okayama* (1996) **50**, 237-241.
- Tsubouchi M, Tsubouchi Y, Hitomi S, Ohtsuka A and Murakami T: Perineuronal sulfated proteoglycans, cell surface glycoproteins and dark neurons in the cingulate cortex of newborn and adult rats. *Acta Med Okayama* (1997) **50**, 313-317.
- Hitomi S, Su WD, Hong LJ, Ohtsuka A and Murakami T: Perineuronal sulfated proteoglycans and cell surface glycoproteins in the visual cortex of adult and newborn cats. *Acta Med Okayama* (1997) **51**, 295-299.
- Murakami T and Ohtsuka A: Perineuronal sulfated proteoglycans in the human brain as doubly stained with Golgi's silver nitrate and cationic iron colloid; in *Recent Advances in Microscopy of Cells, Tissues and Organs*, Motta PM ed, Antonio Delfino Editore, Rome (1997) pp211-214.
- Golgi C: Intorno all struttura delle cellule nervose. *Arch Ital Biol* (1898) **30**, 60-71.
- Lipsky NG: On the structure of nerve cells, by C. Golgi. Historical note. *J Microsc* (1989) **155**, 3-7.
- Murakami T, Murakami T, Sato H, Mubarak WA, Ohtsuka A and Abe K: Perineuronal nets of proteoglycans in the adult mouse brain, with special reference to their reactions to Gömöri's ammoniacal silver and Ehrlich's methylene blue. *Arch Histol Cytol* (1999) **62**, 71-81.
- Murakami T, Murakami T, Su WD, Ohtsuka A, Abe K, Ninomiya Y: Perineuronal nets of proteoglycans in the adult mouse brain are digested by collagenase. *Arch Histol Cytol* (1999) **62**, 199-204.
- Murakami T, Taguchi T, Ohtsuka A, Sano K, Kaneshige T, Owen RL and Jones AL: A modified method of fine-granular cationic iron colloid



- preparation: Its use in the light and electron microscopic detection of anionic sites in the rat kidney glomerulus and certain other tissues. *Arch Histol Cytol* (1986) **49**, 12-23.
23. Gömöri G: Silver impregnation of reticulum in paraffin sections. *Am J Pathol* (1937) **13**, 993-1002.
  24. Romeis B: Impragnierung des argyrophilen und kollagenen Bindegewebes mit Metallsalzen; in *Mikroskopische Technik*, Romeis B ed, 16 Auflage, R Oldenbourg, Munchen-Wien (1968) pp376-387.
  25. Fujita T: Histological studies on the neuro-insular complexes in the pancreas of some mammals. *Z Zellforsch* (1959) **50**, 94-109.
  26. Hartig W, Brauer K and Bruckner G: *Wisteria floribunda* agglutinin-labelled nets surround parvalbumin-containing neurons. *Neuroreport* (1992) **3**, 869-872.
  27. Romeis B: Darstellung mittels supravitaler Methylenblaufarbung; in *Mikroskopische Technik*, Romeis B ed, 16 Auflage, R Oldenbourg, Munchen-Wien (1968) pp486-495.
  28. Yoshikawa T, Nishida K, Doi T, Inoue H, Ohtsuka A, Taguchi T and Murakami T: Negative charges bound to collagen fibrils in the rabbit articular cartilage: A light and electron microscopic study using cationic colloidal iron. *Arch Histol Cytol* (1997) **60**, 435-443.
  29. Rambourg A, Neutra M and Leblond CP: Presence of a "cell coat" rich in carbohydrate at the surface of cells in the rat. *Anat Rec* (1966) **154**, 41-71.
  30. Brauer K, Werner L and Leibnitz L: Perineuronal nets of glia. *J Hirnforsch* (1982) **23**, 701-708.
  31. Nakagawa F, Schulte BA and Spicer SS: Selective cytochemical demonstration of glycoconjugate-containing terminal N-acetylgalactosamine on some brain neurons. *J Comp Neurol* (1986) **243**, 280-290.
  32. Oohira A, Matsui F, Matsuda M and Shoji R: Developmental change in the glycosaminoglycan composition of the rat brain. *J Neurochem* (1986) **47**, 588-593.
  33. Hendry SH, Jones EG, Hockfield S and McKay RD: Neuronal populations stained with the monoclonal antibody Cat-301 in the mammalian cerebral cortex and thalamus. *J Neurosci* (1988) **8**, 518-542.
  34. Bertolotto A, Rocca G and Schiffer D: Chondroitin 4-sulfate proteoglycans forms an extracellular network in human and rat central nervous system. *J Neurol Sci* (1990) **100**, 113-123.
  35. Herndon ME and Lander AD: A diverse set of developmentally regulated proteoglycans is expressed in the rat central nervous system. *Neuron* (1990) **4**, 949-961.
  36. Bignami A, Asher R and Perides G: Co-localization of hyaluronic acid and chondroitin sulfate proteoglycan in rat cerebral cortex. *Brain Res* (1992) **579**, 173-177.
  37. Fujita SC and Kudo J: A novel member of the family of perineuronal antigens associated with subpopulations of central neurons in the rat. *Exp Brain Res* (1992) **88**, 345-354.
  38. Celio MR: Perineuronal nets of extracellular matrix around parvalbumin-containing neurons of the hippocampus. *Hippocampus* (1993) **3**, 55-60.
  39. Margolis RK, Rauch U, Maurel P and Margolis RU: Neurocan and phosphacan: Two major nervous tissue-specific chondroitin sulfate proteoglycans. *Perspect Devel Neurobiol* (1996) **3**, 273-290.
  40. Ruoslahti E: Brain extracellular matrix. *Glycobiology* (1996) **6**, 489-492.
  41. Nishizuka M, Ikeda S, Arai Y, Maeda N and Noda M: Cell surface-associated extracellular distribution of a neural proteoglycan, 6B4 proteoglycan/phosphacan, in the olfactory epithelium, olfactory nerve, and cells migrating along the olfactory nerve in chick embryos. *Neurosci Res* (1996) **24**, 345-355.
  42. Wintergerst ES, Faissner A and Celio MR: The proteoglycan DSD-1-PG occurs in perineuronal nets around parvalbumin-immunoreactive interneurons of the rat cerebral cortex. *Int J Devel Neurosci* (1996) **14**, 249-255.
  43. Yamada H, Fredette B, Shitara K, Hagihara K, Miura R, Ranscht B, Stallcup WB and Yamaguchi Y: The brain chondroitin sulfate proteoglycan brevican associates with astrocytes ensheathing cerebellar glomeruli and inhibits neurite outgrowth from granule neurons. *J Neurosci* (1997) **17**, 7784-7795.
  44. Ramon Y Cajal S: La red superficial de las cellulas nerovas centrales. *Rev Trimest Micrograf (Madrid)* (1897) **3**, 163-178.
  45. Ramon Y Cajal S: Reseaux pericellulaires; in *Histologie du System nerveux de l'Homme et des Vertebres* (translated by Azoulay L), Vol. I, Maloine, Paris (1909-1911) pp155-158.
  46. Kosaka T and Heizmann CW: Selective staining of a population of parvalbumin-containing GABAergic neurons in the rat cerebral cortex by lectins with specific affinity for terminal N-acetylgalactosamine. *Brain Res* (1989) **183**, 158-163.
  47. Drake CT, Mulligan KA, Wimpey TL, Hendrickson A and Chavkin C: Characterization of *Vicia villosa* agglutinin-labeled GABAergic interneurons in the hippocampal formation and in acutely dissociated hippocampus. *Brain Res* (1991) **554**, 176-185.
  48. Hartig W, Brauer K, Bigl V and Bruckner G: Chondroitin sulfate proteoglycan-immunoreactivity of lectin-labeled perineuronal nets around parvalbumin containing neurons. *Brain Res* (1994) **635**, 307-311.
  49. Luth HJ, Fischer J and Celio MR: Soybean lectin binding neurons in the visual cortex of the rat contain parvalbumin and are covered by glial nets. *J Neurocytol* (1992) **21**, 211-221.
  50. Bruckner G, Brauer K, Hartig W, Wolff JR, Rickmann MJ, Derouiche A, Delpech B, Girard N, Oertel WH and Reichenbach A: Perineuronal nets provide a polyanionic, glia-associated form of microenvironment around certain neurons in many parts of the rat brain. *Glia* (1993) **8**, 183-200.
  51. Bruckner G, Hartig W, Kacza J, Seeger J, Welt K and Brauer K: Extracellular matrix organization in various regions of rat brain grey matter. *J Neurocytol* (1996) **25**, 333-346.
  52. Celio MR and Chiquet-Ehrismann R: 'Perineuronal nets' around cortical interneurons expressing parvalbumin are rich in tenascin. *Neurosci Lett* (1993) **162**, 137-140.
  53. Seeger G, Brauer K, Hartig W and Bruckner G: Mapping of perineuronal nets in the rat brain stained by colloidal iron hydroxide histochemistry and lectin cytochemistry. *Neuroscience* (1994) **58**, 371-388.
  54. Blumcke I, Eggli P and Celio MR: Relationship between astrocytic processes and "perineuronal nets" in rat neocortex. *Glia* (1995) **15**, 131-140.
  55. Koppe G, Bruckner G, Hartig W, Delpech B and Bigl V: Characterization of proteoglycan-containing perineuronal nets by enzymatic treatments of rat brain sections. *Histochem J* (1997) **29**, 11-20.
  56. Koppe G, Bruckner G, Brauer K, Hartig W and Bigl V: Developmental patterns of proteoglycan-containing extracellular matrix in perineuronal nets and neuropil of the postnatal rat brain. *Cell Tissue Res* (1997) **288**, 33-41.
  57. Naegele JR and Katz LC: Cell surface molecules containing N-acetylgalactosamine are associated with basket cells and neuroglia-form cells in cat visual cortex. *J Neurosci* (1990) **10**, 540-557.
  58. Derouiche A, Hartig W, Brauer K and Bruckner G: Spatial relationship of lectin-labeled extracellular matrix and glutamine synthetase-immunoreactive astrocytes in rat cortical forebrain regions. *J Anat* (1996) **189**, 363-372.
  59. Seeger T, Luth HJ, Winkelmann E and Brauer K: Distribution patterns of *Wisteria floribunda* agglutinin binding sites and parvalbumin-

- immunoreactive neurons in the human visual cortex: A double-labeling study. *J Hirnforsch Hirnforsch* (1996) **37**, 351-366.
60. Brauer K, Brückner G, Leibnitz L and Werner L: Structural and cytochemical features of perineuronal glial nets in the rat brain. *Acta Histochem* (1984) **74**, 53-60.
61. Müller T: Methylene blue supravital staining: An evaluation of its applicability to the mammalian brain and pineal gland. *Histol Histopathol* (1998) **13**, 1019-1026.
- 

Received November 5, 1999; accepted January 17, 2000.