

Acta Medica Okayama

Volume 54, Issue 6

2000

Article 6

DECEMBER 2000

Immunophilin ligands prevent H₂O₂-induced apoptotic cell death by increasing glutathione levels in neuro 2A neuroblastoma cells.

Ken-ichi Tanaka*

Naoko Fujita†

Masato Asanuma‡

Norio Ogawa**

*Okayama University,

†Okayama University,

‡Okayama University,

**Okayama University,

Immunophilin ligands prevent H₂O₂-induced apoptotic cell death by increasing glutathione levels in neuro 2A neuroblastoma cells.*

Ken-ichi Tanaka, Naoko Fujita, Masato Asanuma, and Norio Ogawa

Abstract

We examined the effects of FK506 and its non-immunosuppressive derivative, GPI1046, on H₂O₂-induced reduction of cell viability and apoptotic cell death in Neuro 2A cells. Our results suggest that the protective properties of GPI1046 against H₂O₂-induced reduction of cell viability are equipotent with those of FK506 and may be mediated by increased intracellular concentrations of glutathione (GSH). In addition, both FK506 and GPI1046 prevented apoptotic cell death in Neuro 2A cells, although the antiapoptotic effect of FK506 was somewhat stronger than that of GPI1046. These findings suggest that non-immunosuppressive immunophilin ligands such as GPI1046 might be potentially useful in treatment of neurodegenerative diseases without serious side effects such as immune deficiency.

KEYWORDS: hydrogen peroxide, immunophilin ligands, apoptosis, glutathione, FK506, GPI1046

*PMID: 11132921 [PubMed - indexed for MEDLINE]

Copyright (C) OKAYAMA UNIVERSITY MEDICAL SCHOOL

Immunophilin Ligands Prevent H₂O₂-Induced Apoptotic Cell Death by Increasing Glutathione Levels in Neuro 2A Neuroblastoma Cells

Ken-ichi TANAKA*, Naoko FUJITA, Masato ASANUMA and Norio OGAWA

Department of Neuroscience, Institute of Molecular and Cellular Medicine, Okayama University Medical School, Okayama 700-8558, Japan

We examined the effects of FK506 and its non-immunosuppressive derivative, GPI1046, on H₂O₂-induced reduction of cell viability and apoptotic cell death in Neuro 2A cells. Our results suggest that the protective properties of GPI1046 against H₂O₂-induced reduction of cell viability are equipotent with those of FK506 and may be mediated by increased intracellular concentrations of glutathione (GSH). In addition, both FK506 and GPI1046 prevented apoptotic cell death in Neuro 2A cells, although the anti-apoptotic effect of FK506 was somewhat stronger than that of GPI1046. These findings suggest that non-immunosuppressive immunophilin ligands such as GPI1046 might be potentially useful in treatment of neurodegenerative diseases without serious side effects such as immune deficiency.

Key words: hydrogen peroxide, immunophilin ligands, apoptosis, glutathione, FK506, GPI1046

It is believed that the immunosuppressive properties of immunophilin ligands do not need their neurotrophic effects so that their non-immunosuppressive derivatives retain potent neurotrophic effects (1-9) comparable to those of their mother compounds such as tacrolimus (FK506) or cyclosporin A (CsA). For instance, it has been shown that non-calcineurin (CaN) binding analogues of FK506, such as GPI1046 and V10367, which are not immunosuppressive, still promote neurite outgrowth (2, 3) and prevent cell death (10, 11). Furthermore, the neurotrophic actions of immunophilin ligands are restricted to damaged neurons, in contrast to neurotrophic factors, which also elicit neurite outgrowth in naive neurons (6). Several properties of immunophilin ligands, including their neurotrophic activities, are

mediated by inhibition of rotamase activity rather than by inhibition of CaN activity (2, 3). Therefore, CaN inhibition by FK506 is probably not essential for protection against cell damage, although inhibition of both CaN and rotamase could contribute to the neuroprotective effects of FK506 (1, 3, 6, 12). We previously reported that, by increasing intracellular glutathione (GSH) levels in undifferentiated NG108-15 hybridoma cells, GPI1046 had neuroprotective effects equal to those of FK506 against H₂O₂-induced cell damage (11). Since GSH is the most abundant low-molecular-weight thiol within the cell, the GSH system has a rapid and great capacity for the prevention of reactive oxygen species (ROS)-induced cell damage and thus plays an important role in the detoxification of ROS in the brain (13, 14). However, several researchers have postulated that non-immunosuppressive immunophilin ligands do not have any beneficial effects (15-17). Therefore, it is not enough to understand that non-immunosuppressive immunophilin ligands are useful drugs for treating neurodegenerative diseases.

In the present study, we examined the neuroprotective effects of FK506 and its non-immunosuppressive derivative, GPI1046 [3-(3-pyridyl)-1-propyl (2S)-1-(3, 3-dimethyl-1, 2-dioxopentyl)-2-prolidine carboxylate], on H₂O₂-induced apoptotic cell death (18) in Neuro 2A cells which are a mouse neuroblastoma cell line. Undifferentiated NG108-15 cells mainly displays characteristics of glial lineages (19, 20), and we have already confirmed the neuroprotective effects of immunophilin ligands in undifferentiated NG108-15 cells (11). Therefore, we used neuronal cell lines to clarify the effects of immunophilin ligands on neuronal lineages. In addition, we also examined the effects of these immunophilin ligands on GSH contents in naive Neuro 2A cells.

* To whom correspondence should be addressed.

Materials and Methods

Neuro 2A cells were obtained from the Institute for Fermentation (IFO, Osaka, Japan), and stocks were grown using a method described previously with a minor modification (21, 22). Briefly, the cells were continuously cultured in Dulbecco's Modified Eagle Medium containing, in addition to 4 mM L-glutamine, 10% fetal bovine serum and penicillin (100 U/ml)-streptomycin (100 μ g/ml) (all purchased from GIBCO BRL, Rockville, MD, USA). For experiments, cells were plated onto 96-well plates at a density of 1.6×10^4 cells/ml to assess cell viability. In addition, cells were plated onto 8-chamber slides at 1.6×10^4 cells/ml (NNI, Rochester, NY, USA) to investigate apoptotic cell death, and they were plated separately onto 6-well plates at 1.3×10^5 cells/ml to determine GSH contents. All cultures were maintained at 37 °C in a gas mixture of 5%–95% CO₂-air. Neuro 2A cells were grown for 4–5 days and then used in the present studies.

In the assessment of cell viability, cells were treated with immunophilin ligands, FK506 (Fujisawa, Osaka, Japan) or GPI1046, at a final concentration of 1, 10 or 100 nM, to evaluate their protective effects against H₂O₂-induced cell damage. Following a 24 h-incubation after the treatment with immunophilin ligands, the cell culture medium was replaced with a fresh medium, and then cells were treated with H₂O₂ at a final concentration of 300 μ M. Following a 24 h-incubation after H₂O₂ challenge, cell viability was determined by a 2-(4-lodophenyl)-3-(4-nitrophenyl)-5-(2, 4-disulfophenyl)-2H-tetrazolium (WST-1) cell counting kit (Dojin Lab, Kumamoto, Japan). FK506 or GPI1046 was dissolved in dimethyl sulfoxide (DMSO). The final concentration of DMSO was maintained at $\leq 0.1\%$. The cell viability in each well of the 96-well plate was determined by quantitative colorimetric assay with WST-1 (11, 23), a modification of the standard MTT assay. Viability was expressed as a percentage of the cell viability from each 0.1% DMSO-treated control culture in the non-H₂O₂ exposed group.

In addition, the morphological changes in the cells were observed by fluorescence microscopy after staining with Hoechst 33342 (Molecular Probes, Eugene, OR, USA) or propidium iodide (PI; Molecular Probes, Eugene, OR, USA). Cells were fixed with 4% paraformaldehyde (in 0.1 M phosphate buffer) for 10 min. After washing with 10 mM phosphate-buffered saline

(PBS), Hoechst 33342 (20 μ g/ml) or PI (4 μ g/ml) was added to the cells for 3 min. Slides were rinsed briefly with PBS, air-dried, and then mounted in anti-fluorescein fading medium (PermaFluor; Lipshaw Co., Pittsburgh, PA, USA). After staining with Hoechst 33342 (emission, above 420 nm; excitation, 330–385 nm) or PI (emission, above 580 nm; excitation, 520–550 nm), cells were assessed as having undergone apoptosis or necrotic cell death by fluorescence microscopy.

GSH levels were determined using the enzymatic recycling method of Tietze (24) with some modifications. For preparation of the GSH sample, Neuro 2A cells were grown for 24 h in the presence or absence of immunophilin ligands. Cells were homogenized in 0.1 M phosphate buffer (pH 7.4) and then were added at equivalent volumes to 10% trichloroacetic acid and cellular homogenates. Acid extracts were mixed with 0.01 M phosphate buffer (pH 7.4, 580 μ l), NADPH (4 mM, 50 μ l) and glutathione reductase (6 U/ml, 100 μ l), and incubated for 5 min at 37 °C. 3-Carboxy-4-nitrophenyl disulfide (10 mM, 50 μ l) was added just before reading absorbance. The formation of 2-nitro-5-thiobenzoic acid was measured at 450 nm for 6 min. Total GSH was determined from a standard curve constructed using known amounts of GSH.

Data are presented as mean \pm SEM. Differences between groups were tested for statistical significance using one-way analysis of variance (ANOVA) followed by a *post hoc* Dunnett's test ($P < 0.05$).

Results

We examined the protective effects of immunophilin ligands against H₂O₂-induced cell damage. Exposure to H₂O₂ (300 μ M) for 24 h reduced cell viability to $51.4 \pm 2.3\%$ in 0.1% DMSO-treated Neuro 2A cells (Fig. 1). However, pretreatment with FK506 and GPI1046 increased cell viability in a dose-dependent manner (Fig. 1). In particular, a high dose of both FK506 and GPI1046 (100 nM) significantly improved H₂O₂-induced reduction of cell viability in contrast to that of the 0.1% DMSO-treated group (Fig. 1). In addition, the occurrence of apoptosis or necrotic cell death were differentiated by morphological observation after staining with Hoechst 33342 or PI. By means of fluorescence microscopy, Neuro 2A cells positively stained with Hoechst 33342 showed condensed and fragmented nuclei typical of H₂O₂-induced apoptotic cell death in the 0.1% DMSO-




Fig. 1 Viability of Neuro 2A cells was measured by WST-1 colorimetric assay. Neuro 2A cells were maintained in the presence or absence of FK506 or GPI1046 (final concentrations of 1, 10, 100 nM) for 24 h and then of 300 μ M of H₂O₂. After an additional 24 h, cell damage was assessed by WST-1 assay. Each value is the mean \pm SEM (n = 12) expressed as a percentage of control cultures treated with autoclaved water instead of H₂O₂. *P < 0.05 compared with 0.1% DMSO-treated control cultures in non-H₂O₂ exposed group.

treated group (Fig. 2). However, both FK506 and GPI1046 (100 nM) prevented H₂O₂-induced apoptotic cell death, although the anti-apoptotic effect of FK506 was somewhat stronger than that of GPI1046 (Fig. 2). On the other hand, H₂O₂ did not induce necrotic cell death since PI staining was not investigated in any experimental groups (data not shown).

To clarify the mechanism of the protective effects of immunophilin ligands, we determined the effects of FK506 and GPI1046 on GSH contents in naive Neuro 2A cells. Treatment with FK506 (100 nM) or GPI1046 (100 nM) significantly increased GSH contents to 180.7% or 169.6%, respectively, compared with 0.1% DMSO treatment (Fig. 3).

Discussion

Oxidative stress plays an important role in the pathogenesis of neurodegenerative diseases, such as Alzheimer's disease, Parkinson's disease, and stroke, due to the ROS (25-27). Therefore, antioxidant properties may contribute to the neuroprotective effects of drugs on

neurodegenerative diseases. To control the harmful effects of oxidative stress, efficient detoxification pathways have been identified in various cells. Of the various antioxidant systems in the brain, the GSH system is particularly important in controlling cellular redox states and is the primary defense mechanism for removal of peroxide from the brain (13, 14, 28). Furthermore, cellular GSH has been found to be important in modulating oxidative stress-induced apoptosis in several cell lines (18, 29-31). Moreover, apoptosis has been prevented by increasing intracellular GSH levels in several models of apoptosis (29-31). In the case of neuronal lineages such as neuroblastoma or neurons, GSH has also protected against oxidative stress-induced apoptotic cell death (32-35). We have reported previously that the increasing effects of GPI1046 on the level of intracellular GSH are equipotent to those of FK506 in undifferentiated NG108-15 cells (11). In the present study, we also demonstrated that both GPI1046 and FK506 enhanced intracellular GSH contents in Neuro 2A cells. Furthermore, the efficacy of FK506 and GPI1046 for Neuro 2A cells (neuronal lineages) was almost equipotent to that for undifferentiated NG108-15 cells (glial lineages). In addition, both FK506 and GPI1046 had neuroprotective properties against H₂O₂-induced reduction of cell viability and apoptotic cell death. However, the anti-apoptotic effect of FK506 was somewhat stronger than that of GPI1046, although the neuroprotective effect of GPI1046 on H₂O₂-induced reduction of cell viability was similar to that of FK506. Nevertheless, our present and previous findings (11) are almost in agreement with those of Steiner's previous studies (3, 4, 6, 8).

On the other hand, CaN has been implicated in the signal transduction pathway leading to Ca²⁺-dependent apoptosis (36). Recent studies have shown that anti-CD3-induced apoptosis is mediated by Fas-ligand production, which is regulated by the CaN-activated nuclear factor of activated T-cells (NF-AT) (37) and the interaction between Fas and Fas-ligand (38-40). CaN has also been found to be activated by the dephosphorylation of Bad and nitric oxide synthase (41, 42). Furthermore, CaN-induced apoptosis is due to activation of the cytochrome c/caspase-3 pathway in neurons (43). Therefore, CaN appears to participate in apoptosis in various ways, and inhibition of CaN activity may be important for neuroprotective properties of immunosuppressive immunophilin ligands, such as FK506, against CaN-related apoptosis (36, 44, 45). Therefore, our results indicate




Fig. 2 H₂O₂ (300 μ M)-induced apoptotic cell death in Neuro 2A cells with Hoechst 33342 staining. Neuro 2A cells were maintained in the presence or absence of FK506 or GPI1046 (final concentrations of 100 nM) for 24 h and then of 300 μ M of H₂O₂ for 24 h. Panel A, 0.1% DMSO-treated + non-H₂O₂ exposed group; Panel B, 0.1% DMSO-treated + H₂O₂ exposed group; Panel C, FK506 (100 nM)-treated + H₂O₂ exposed group; Panel D, GPI1046 (100 nM)-treated + H₂O₂ exposed group.

that H₂O₂-induced apoptotic cell death in Neuro 2A cells may be dependent not only on the GSH depletion pathway but also the CaN pathway.

In conclusion, our present results suggest that the cellular protective properties of GPI1046, a non-immunosuppressive FK506 derivative, is equipotent to that of FK506 and may be mediated by increasing intracellular GSH levels. Moreover, both FK506 and GPI1046 prevented apoptotic cell death in Neuro 2A

cells, although the anti-apoptotic effect of FK506 was somewhat stronger than that of GPI1046. Thus, non-immunosuppressive immunophilin ligands, such as GPI1046, might have a potentially beneficial effect on neurodegenerative diseases, particularly since they do not cause serious side effects such as immune deficiency. In addition, the GSH-activated effect could be regarded as a beneficial property of neuroprotective drugs.




Fig. 3 Effects of FK506 and GPI1046 on GSH contents in Neuro 2A cells. Neuro 2A cells were grown for 24 h in the presence or absence of FK506 or GPI1046 (1, 10, 100 nM). Each bar represents the mean \pm SEM (n = 6). * P < 0.05 compared with 0.1% DMSO-treated control cultures.

Acknowledgments. This work was supported in part by Grants-in-Aid for Scientific Research on Priority Area and Scientific Research (C) from the Japanese Ministry of Education, Science, Sports and Culture, and Grants-in-Aid for Comprehensive Research on Aging and Health, for Research on Brain Science, and for Research on Specific Diseases from the Japanese Ministry of Health and Welfare.

References

- Gold BG: FK506 and the role of immunophilins in nerve regeneration. *Mol Neurobiol* (1997) **15**, 285-306.
- Gold BG, Zeleny-Pooley M, Wang M-S, Chaturvedi P and Armistead DM: A nonimmunosuppressant FKBP-12 ligand increases nerve regeneration. *Exp Neurol* (1997) **147**, 269-278.
- Steiner JP, Connolly MA, Valentine HL, Hamilton GS, Dawson TM, Hester L and Snyder SH: Neurotrophic actions of nonimmunosuppressive analogues of immunosuppressive drugs FK506, rapamycin and cyclosporin A. *Nat Med* (1997) **3**, 421-428.
- Steiner JP, Hamilton GS, Rose DT, Valentine HL, Guo H, Connolly MA, Liang S, Ramsey C, Li JH, Huang W, Howorth P, Soni R, Fuller M, Sauer H, Nowotnik AC and Suzdak PD: Neurotrophic immunophilin ligands stimulate structural and functional recovery in neurodegenerative animal models. *Proc Natl Acad Sci USA* (1997) **94**, 2019-2024.
- Gold BG, Zeleny-Pooley M, Chaturvedi P and Wang M-S: Oral administration of a nonimmunosuppressant FKBP-12 ligand speeds nerve regeneration. *Neuroreport* (1998) **9**, 553-558.
- Snyder SH, Sabatini DM, Lai MM, Steiner JP, Hamilton GS and Suzdak PD: Neural actions of immunophilin ligands. *Trends Pharmacol Sci* (1998) **19**, 21-26.
- Costantini LC, Chaturvedi P, Armistead DM, McCaffrey PG, Deacon TW and Isacson O: A novel immunophilin ligands: Distinct branching effects on dopaminergic neurons in culture and neurotrophic actions after oral administration in an animal model of Parkinson's disease. *Neurobiol Dis* (1998) **5**, 97-106.
- Sauer H, Francis JM, Jiang H, Hamilton GS and Steiner JP: Systemic treatment with GPI 1046 improves spatial memory and reverses cholinergic neuron atrophy in the medial septal nucleus of aged mice. *Brain Res* (1999) **842**, 109-118.
- Herdegen T, Fischer G and Bold BG: Immunophilin ligands as a novel treatment of neurological disorders. *Trends Pharmacol Sci* (2000) **21**, 3-5.
- Costantini LC and Isacson O: Immunophilin ligands and GDNF enhance neurite branching or elongation from developing dopamine neurons in culture. *Exp Neurol* (2000) **164**, 60-70.
- Tanaka K, Fujita N, Yoshioka M and Ogawa N: Immunosuppressive and non-immunosuppressive immunophilin ligands improve H₂O₂-induced cell damage by increasing glutathione levels in NG108-15 cells. *Brain Res* (in press).
- Ogawa N: Immunophilin ligands: Possible neuroprotective agents. *NeuroSci News* (1999) **2**, 28-34.
- Cooper AJ and Kristal BS: Multiple roles of glutathione in the central nervous system. *Biol Chem* (1997) **378**, 793-802.
- Hall AG: Review: The role of glutathione in the regulation of apoptosis. *Eur J Clin Invest* (1999) **29**, 238-245.
- Bavetta S, Hamlyn PJ, Burnstock G, Lieberman AR and Anderson PN: The effects of FK506 on dorsal column axons following spinal cord injury in adult rats: Neuroprotection and local regeneration. *Exp Neurol* (1999) **158**, 382-393.
- Harper S, Bilsland J, Young L, Bristow L, Boyce S, Mason G, Rigby M, Hewson L, Smith D, O'Donnell R, O'Connor D, Hill RG, Evans D, Swain C, Williams B and Hefti F: Analysis of the neurotrophic effects of GPI-1046 on neuron survival and regeneration in culture and *in vivo*. *Neuroscience* (1999) **88**, 257-267.
- Winter C, Schenkel J, Bürger E, Eickmeier C, Zimmermann M and Herdegen T: The immunophilin ligand FK506, but not GPI-1046, protects against neuronal death and inhibits c-Jun expression in the substantia nigra pars compacta following transection of the rat medial forebrain bundle. *Neuroscience* (2000) **95**, 753-762.
- Mielke K, Damm A, Yang DD and Herdegen T: Selective expression of JNK isoforms and stress-specific JNK activity in different neural cell lines. *Mol Brain Res* (2000) **75**, 128-137.
- Higashida H, Kato T, Kano-Tanaka K, Okuya M, Miyake A and Tanaka T: Proliferation and synapse formation of neuroblastoma glioma hybrid cells: Effect of glial maturation factor. *Brain Res* (1981) **214**, 287-299.
- Ma W, Pancrazio JJ, Coulombe M, Dumm J, Sathanoori R, Barker JL, Kowtha VC, Stenger DA and Hickman JJ: Neuronal and glial epitopes and transmitter-synthesizing enzymes appear in parallel with membrane excitability during neuroblastoma \times glioma hybrid differentiation. *Dev Brain Res* (1998) **106**, 155-163.
- Pangalos MN, Shioi J and Robakis NK: Expression of the chondroitin sulfate proteoglycans of amyloid precursor (appican) and amyloid precursor-like protein 2. *J Neurochem* (1995) **65**, 762-769.
- Adan R, van der Kraan M, Doornbos RP, Bar PR, Burbach JP and Gispen WH: Melanocortin receptors mediate alpha-MSH-induced stimulation of neurite outgrowth in neuro 2A cells. *Mol Brain Res* (1996) **36**, 37-44.
- Iwata-Ichikawa E, Kondo Y, Miyazaki I, Asanuma M and Ogawa N: Glial cells protect neurons against oxidative stress via transcriptional up-regulation of the glutathione synthesis. *J Neurochem* (1999) **72**, 2334-2344.
- Tietze F: Enzymic method for quantitative determination of nanogram

- amounts of total and oxidized glutathione: Applications to mammalian blood and other tissues. *Anal Biochem* (1969) **27**, 502-522.
25. Jenner P and Olanow CW: Understanding cell death in Parkinson's disease. *Ann Neurol* (1998) **44** (Suppl 1), S72-S84.
 26. Behl C: Alzheimer's disease and oxidative stress: Implications for novel therapeutic approaches. *Prog Neurobiol* (1999) **57**, 301-323.
 27. Love S: Oxidative stress in brain ischemia. *Brain Pathol* (1999) **9**, 119-131.
 28. Dringen R, Kussmaul L, Gutterer JM, Hirrlinger J and Hamprecht B: The glutathione system of peroxide detoxification is less efficient in neurons than in astroglial cells. *J Neurochem* (1999) **72**, 2523-2530.
 29. Beaver JP and Waring P: A decrease in intracellular glutathione concentration precedes the onset of apoptosis in murine thymocytes. *Eur J Cell Biol* (1995) **68**, 47-54.
 30. Sandstrom PA, Mannie MD and Buttke TM: Inhibition of activation-induced death in T cell hybridomas by thiol antioxidants: Oxidative stress as a mediator of apoptosis. *J Leukoc Biol* (1994) **55**, 221-226.
 31. Watson RW, Rotstein OD, Nathens AB, Dackiw AP and Marshall JC: Thiol-mediated redox regulation of neutrophil apoptosis. *Surgery* (1996) **120**, 150-158.
 32. De Laurenzi V, Melino G, Savini I, Annicchiarico-Petruzzelli M, Finazzi-Agro A and Avigliano L: Cell death by oxidative stress and ascorbic acid regeneration in human neuroectodermal cell lines. *Eur J Cancer* (1995) **31**, 463-466.
 33. Gabby M, Tauber M, Porat S and Simantov R: Selective role of glutathione in protecting human neuronal cells from dopamine-induced apoptosis. *Neuropharmacology* (1996) **35**, 571-578.
 34. Anderson CP, Tsai JM, Meek WE, Liu R-M, Tang Y, Forman HJ and Reynolds CP: Depletion of glutathione by buthionine sulfoxine is cytotoxic for human neuroblastoma cell lines via apoptosis. *Exp Cell Res* (1999) **246**, 183-192.
 35. Stokes AH, Lewis DY, Lash LH, Jerome WG IIIrd, Grant KW, Aschner M and Vrana KE: Dopamine toxicity in neuroblastoma cells: Role of glutathione depletion by L-BSO and apoptosis. *Brain Res* (2000) **858**, 1-8.
 36. Mukerjee N, McGinnis KM, Park YH, Gnegy ME and Wang KK: Caspase-mediated proteolytic activation of calcineurin in thapsigargin-mediated apoptosis in SH-SY5Y neuroblastoma cells. *Arch Biochem Biophys* (2000) **379**, 337-343.
 37. Latinis KM, Norian LA, Eliason SL and Koretzky GA: Two NFAT transcription factor binding sites participate in the regulation of CD95 (Fas) ligand expression in activated human T cells. *J Biol Chem* (1997) **272**, 31427-31434.
 38. Shi YF, Sahai BM and Green DR: Cyclosporin A inhibits activation-induced cell death in T-cell hybridomas and thymocytes. *Nature* (1989) **339**, 625-626.
 39. Brunner T, Mogil RJ, LaFace D, Yoo NJ, Mahboubi A, Echeverri F, Martin SJ, Force WR, Lynch DH, Ware CF and Green DR: Cell-autonomous Fas (CD95)/Fas-ligand interaction mediates activation-induced apoptosis in T-cell hybridomas. *Nature* (1995) **373**, 441-444.
 40. Dhein J, Walczak H, Baumler C, Debatin KM and Krammer PH: Autocrine T-cell suicide mediated by APO-1/. *Nature* (1995) **373**, 438-441.
 41. Dawson TM, Steiner JP, Dawson VL, Dinerman JL, Uhl GR and Snyder SH: Immunosuppressant FK506 enhances phosphorylation of nitric oxide synthase and protects against glutamate neurotoxicity. *Proc Natl Acad Sci USA* (1993) **90**, 9808-9812.
 42. Wang HG, Pathan N, Ethell IM, Krajewski S, Yamaguchi Y, Shibasaki F, McKeon F, Bobo T, Franke TF and Reed JC: Ca²⁺-induced apoptosis through calcineurin dephosphorylation of BAD. *Science* (1999) **284**, 339-343.
 43. Nakatsuka H, Ohta S, Tanaka J, Toku K, Kumon Y, Maeda N, Sakanaka M and Sakaki S: Release of cytochrome c from mitochondria to cytosol in gerbil hippocampal CA1 neurons after transient forebrain ischemia. *Brain Res* (1999) **849**, 216-219.
 44. Liu J, Farmer JD Jr, Lane WS, Friedman J, Weissman I and Schreiber SL: Calcineurin is a common target of cyclophilin-cyclosporin A and FKBP-FK506 complexes. *Cell* (1991) **66**, 807-815.
 45. Fruman DA, Mather PE, Burakoff SJ and Bierer BE: Correlation of calcineurin phosphatase activity and programmed cell death in murine T cell hybridomas. *Eur J Immunol* (1992) **22**, 2513-2517.

Received September 13, 2000; accepted September 28, 2000.