

Acta Medica Okayama

Volume 21, Issue 4

1967

Article 3

AUGUST 1967

Localization of TCA cycle dehydrogenases in the mitochondria

Kozo Inaba*

Akira Wakabayashi[†]

Takuzo Oda[‡]

*Okayama University,

[†]Okayama University,

[‡]Okayama University,

Localization of TCA cycle dehydrogenases in the mitochondria*

Kozo Inaba, Akira Wakabayashi, and Takuzo Oda

Abstract

The site of localization of TCA cycle dehydrogenases in mitochondria has been investigated by observing the dehydrogenase activities and fine structure of the fractionated samples after freezing and thawing or sonication of beef heart and rat liver mitochondria. 1. In the sonicated mitochondria, activities of malic and isocitric dehydrogenases were highest in the supernatant fraction centrifuged at 198,000 x g for 60 minutes, while the specific activity of a-ketoglutaric dehydrogenase was higher in the fluffy or residue fraction. The distribution of the activity of pyruvic dehydrogenase was similar to that of a-ketoglutaric dehydrogenase. 2. In a sucrose density gradient fractionation of the fluffy fraction obtained by centrifugation of sonicated mitochondria at 198,000 x g for 60 minutes, the activities of malic and pyruvic dehydrogenase were observed in the top (or low density) layer in the form of fine particles, while that of a-ketoglutaric dehydrogenase was observed in the middle (or medium density) layers in the form of aggregates of fine particles and membranous fragments. 3. In the samples fractionated after freezing and thawing of mitochondria, which were considered to be a relatively mild disruption, the specific activity of a-ketoglutaric dehydrogenase was higher in the residue (submitochondria) fraction than that in the supernatant fraction (centrifuged at 144,000 x g, 30 minutes), and the activity of malic dehydrogenase still remained significantly high in the residue fraction. 4. It was deduced that the TCA cycle dehydrogenases could be localized in the matrix of the mitochondria by a loose binding to the inner membrane.

*PMID: 4230848 [PubMed - indexed for MEDLINE] Copyright ©OKAYAMA UNIVERSITY MEDICAL SCHOOL

Acta Med. Okayama 21, 167—176 (1967)

LOCALIZATION OF TCA CYCLE DEHYDROGENASES IN THE MITOCHONDRIA

Kozo INABA, Akira WAKABAYASHI and Takuzo ODA

*Department of Biochemistry, Cancer Institute, Okayama University
Medical School (Director: Prof. T. Oda)*

Received for publication, July 29, 1967

It has already been clarified by many investigators^{1,2} that TCA cycle dehydrogenases are mainly located in mitochondria. However, it remains still obscure as to the exact sites of localization of these enzymes in the mitochondria. It has been generally considered that the TCA cycle dehydrogenases, except succinic dehydrogenase (EC 1. 3. 99. 1) may be localized in the matrix of mitochondria as they are easily solubilized by mechanical disruption of mitochondria^{2,3}. GREEN and his collaborators reported, however, that these enzymes may be localized on the outer membrane of mitochondria⁴.

In the present paper, the sites of localization of TCA cycle dehydrogenases in mitochondria were investigated by observing the dehydrogenase activities and structures of the samples fractionated by differential centrifugation and sucrose density gradient after severe or mild disruption of mitochondria.

MATERIALS AND METHODS

Preparation of mitochondria : Mitochondria were isolated from rat liver or beef heart by the modification of the method of HOGEBOM⁵ or CRANE *et al.*⁶, respectively.

Sonication of mitochondria : Mitochondria (70 mg protein) were suspended in 10ml of 0.25 M sucrose solution containing 0.01 M Tris-HCl buffer (pH 7.4) and 0.7mg of vitamin E, and frozen at -25°C for 24 hours before sonication. The mitochondria were thawed under tap water and sonicated immediately for 5 minutes at maximum intensity with a sonicator (Kaijo Electric Co., 20 KC, 7 mm tip at 150 W in air).

Freezing-thawing of mitochondria : Mitochondria (100 mg protein) were suspended in 10 ml of 0.05 M sucrose solution containing 0.01 M Tris-

* This work was supported by a PHS research grant (GM 10538) from the General Medical Sciences, National Institutes of Health, U.S.A. This article was presented at the 38th General Meeting of the Japanese Biochemical Society, Fukuoka, Japan, 1965.

HCl buffer, pH 7.4, and frozen in dry ice-acetone and thawed under tap water. The freezing and thawing were repeated three times.

Centrifugal fractionation of disrupted mitochondria : Mitochondria disrupted with sonication (S_0) or the supernatant of frozen-thawed mitochondria centrifuged at 11,000 x g for 15 minutes (S_0) were centrifuged at 33,000 x g for 15 minutes, and separated into supernatant (S_1) and residue (R_1). The S_1 was centrifuged at 144,000 x g for 30 minutes, and separated into supernatant (S_2) and residue (R_2). The supernatant (S_2) was finally centrifuged at 198,000 x g for 60 minutes and separated into supernatant (S_3), fluffy layer (F_3) and residue (R_3). The fluffy layer (F_3) was further fractionated on a 0.1 to 0.6 M sucrose density gradient at 160,000 x g for 60 min.

Assay of dehydrogenase activities in the fraction : The activity of ferricyanide linked α -ketoglutaric dehydrogenase (EC 1.2.4.2) in the fractions was determined by the method of MASSAY⁷.

Ferricyanide linked pyruvic dehydrogenase (EC 1.2.4.1) activity in the fractions was measured by a modification of the method of MASSAY⁸ in which pyruvate was replaced with α -ketoglutarate as substrate and thiamine pyrophosphate (2.5×10^{-4} M) was used as supplement.

The activity of NADP-linked isocitric dehydrogenase (EC 1.1.1.42) in the fraction was estimated by the method of PLAUT and SUNG⁸ in the presence of potassium cyanide (10^{-3} M).

The activity of malic dehydrogenase (EC 1.1.1.37) in the fraction was measured by the method of OCHOA⁹ in the presence of 10^{-3} M potassium cyanide.

Determination of protein : Protein was estimated by the method of LOWRY *et al.*¹⁰ or the method of GORNALL *et al.*¹¹.

Electron microscopy of the fractionated samples : Electron microscope observation was made on the samples negatively stained with 1 % potassium phosphotungstate, pH 7.0¹².

RESULTS

Distribution of TCA cycle dehydrogenase activities in sonicated mitochondrial fractions : Distributions of TCA cycle dehydrogenase activities in sonicated beef heart mitochondria and rat liver mitochondria are shown in Table 1 and Table 2, respectively. The protein recovery was 62.5 or 74.5 % in the supernatant fraction (S_1) of the sonicated beef heart or rat liver mitochondria centrifuged at 33,000 x g for 15 minutes, respectively. The majority of activities of TCA cycle dehydrogenases was recovered in the S_1 fraction except pyruvic dehydrogenase which was considerably inactivated by the centrifugal

Localization of TCA Cycle Dehydrogenases in Mitochondria

169

Table 1 TCA cycle dehydrogenase activities in the fractions from sonicated beef heart mitochondria

Fraction	Protein recovery %	Fe (CN) ⁼⁶ -Pyruvic DH.		Fe (CN) ⁼⁶ - α -KG DH.		NADP-Isocit. DH.		Malic DH.	
		S. A. *	T. A. **	S. A. *	T. A. **	S. A. *	T. A. **	S. A. *	T. A. **
Sonicated Mt S ₀	100.0	40	9,280	106	24,600	420	97,400	31	7,200
33,000xg, 15' S ₁	62.5	11	1,600	150	21,800	320	46,500	48	6,950
R ₁	41.0	10	950	80	7,600	96	9,100	0	
144,000xg, 30' S ₂	25.3	25	1,460	160	9,450	705	41,600	115	6,800
R ₂	26.7	3	186	50	3,100	41	2,500	0	
198,000xg, 60' S ₃	15.6	21	762	96	3,500	1000	36,300	110	4,000
F ₃	3.9	16	118	240	1,800	568	4,200	371	2,745
R ₃	4.9	37	418	360	4,100	158	1,900	0	

* : S. A., specific activity, μ mole/min/mg protein. ** : T. A., total activity, μ mole/min.

Table 2 TCA cycle dehydrogenase activities in the fractions from sonicated rat liver mitochondria

Fraction	Protein recovery %	Fe (CN) ⁼⁶ -Pyruvic DH.		Fe (CN) ⁼⁶ - α -KG DH.		NADP-Isocit. DH.		Malic DH.	
		S. A. *	T. A. **	S. A. *	T. A. **	S. A. *	T. A. **	S. A. *	T. A. **
Sonicated Mt S ₀	100.0	43	12,900	37	11,100	45	13,500	4.0	1,200
33,000xg, 15' S ₁	74.5	20	4,480	31	6,950	50	11,200	3.2	720
R ₁	14.2	0		29	1,230	22	936	0	
144,000xg, 30' S ₂	52.2	32	5,030	41	6,450	60	9,430	4.2	660
R ₂	12.4	0		11	410	10	372	0	
198,000xg, 60' S ₃	38.5	15	1,740	36	4,180	114	13,200	4.7	545
F ₃	7.7	6	121	64	1,480	60	1,390	4.6	106
R ₃	7.2	21	454	22	475	20	432	4.6	99

* : S. A., specific activity, μ mole/min/mg protein. ** : T. A., total activity, μ mole/min.

separation of sonicated mitochondria. The majority of activities of TCA cycle dehydrogenases in the S₁ fraction was also recovered in the supernatant fraction (S₂) on the centrifugation at 144,000 x g for 30 minutes. The S₂ fraction was further centrifuged at 198,000 x g for 60 minutes, and separated into residue (R₃), fluffy layer (F₃) and supernatant (S₃). The recovery of total activity of all TCA cycle dehydrogenases in both heart and liver mitochondria, except α -ketoglutaric dehydrogenase in beef heart mitochondria, was highest in the supernatant fraction (S₃). The specific activities of pyruvic and α -ketoglutaric dehydrogenases were highest in R₃ or F₃ fraction.

As shown in Fig. 1, F_3 and R_3 fractions were composed of particles and membranous structures. The F_3 fraction was further fractionated on a 0.1 M to 0.6 M sucrose density gradient into five layer fractions which were designated as F_3 -d₁ (top or 0.1 M sucrose layer) to F_3 -d₅ (bottom or 0.6 M sucrose layer). Activities of pyruvic, α -ketoglutaric, and malic dehydrogenases were determined on each of these fractions (Table 3) and electron microscope observation was made (Figs. 2 to 6).

Table 3 Malic, pyruvic and α -ketoglutaric dehydrogenase activities in the density gradient fractions of the fluffy layer* from sonicated beef heart mitochondria

Fraction	Protein recovery %	Malic DH.		Fe(CN) ₅ -Pyruvic DH.		Fe(CN) ₅ - α -KG DH.	
		S. A. **	T. A. ***	S. A. **	T. A. ***	S. A. **	T. A. ***
F_3	100	190	3,990	60	1,260	230	4,830
F_3 -d ₁	47	139	1,390	44	440	65	650
F_3 -d ₂	33	63	442	0		73	520
F_3 -d ₃	18	34	129	0		227	865
F_3 -d ₄	7	0		0		210	328
F_3 -d ₅	2	0		0		0	

*: Fluffy layer of 198,000 xg, for 60 min. **: S. A., specific activity, m μ mole/min/mg protein. ***: T. A., total activity, m μ mole/min.

The activity of malic dehydrogenase was mainly recovered in the F_3 -d₁ fraction, while that of α -ketoglutaric dehydrogenase was mainly recovered in the F_3 -d₃ fraction. Although the activity of pyruvic dehydrogenase was considerably inactivated with this procedure, the activity observed was mainly recovered in the F_3 -d₁ fraction.

The F_3 -d₁ fraction contained mainly of small particles measuring approximately 40 to 100 Å in diameter and the F_3 -d₃ fraction contained fairly large particles or aggregates of particles (200 to 500 Å) and small membranous fragments (300 to 600 Å in diameter).

Distribution of α -ketoglutaric and malic dehydrogenase activities in the frozen-thawed mitochondrial fraction: A further attempt has been made to obtain some clues about the binding of TCA cycle dehydrogenases to the mitochondrial inner membrane. After freezing-thawing of mitochondria, which was considered as a relatively mild disruption, the mitochondria were centrifuged at 11,000 x g for 15 minutes, and the supernatant (S_0) was further centrifuged at 33,000 x g for 15 minutes, and separated into supernatant (S_1) and residue (R_1). The supernatant (S_1) was then centrifuged at 144,000 x g for 30 minutes, and separated into the supernatant (S_2) and residue (R_2).

Table 4 shows α -ketoglutaric and malic dehydrogenase activities in these

Table 4 Malic and α -ketoglutaric dehydrogenase activities in the fraction from frozen-thawed rat liver mitochondria

Fraction	Protein recovery %	Malic dehydrogenase		Fe (CN) $^{\ominus 6}$ - α -KG dehydrogenase	
		S. A. *	T. A. **	S. A. *	T. A. **
11,000xg, 15' S ₀	100	20	3,940	122	24,020
33,000xg, 15' S ₁	61	20	2,400	187	22,430
R ₁	20	14	545	121	4,720
144,000xg, 30' S ₂	56	21	2,310	166	18,250
R ₂	8	7	105	335	5,330

* : S. A', specific activity, μ mole/min/mg protein ** : T. A., total activity, μ mole/min

fractions. In this relatively mild disruption of mitochondria, the specific activity of α -ketoglutaric dehydrogenase was somewhat higher in the R₂ fraction, which was regarded as inner membrane fraction, than in the S₂ fraction. The α -ketoglutaric dehydrogenase in R₂ fraction was solubilized in supernatant fraction by repeated washing of the R₂ fraction. On the other hand, the majority of malic dehydrogenase activity was recovered in S₂ fraction.


DISCUSSION


The analytical data summarized in Table 5 and Fig. 7 indicate that the degree of the release of TCA cycle dehydrogenases is consistent with that of the rupture of the membranous envelopes; however, there seems to exist a difference in the mode of binding of these dehydrogenases to the mitochondrial membrane.

Table 5 The dissociability of TCA cycle dehydrogenase in mitochondria

Dissociability	Dehydrogenase
Sonic non-dissociable	Succinic DH., (NADH ₂ DH.)
dissociable	Pyruvic, Isocitric, α -KG, Malic Dehydrogenases
Freezing-thawing	
dissociable (incompletely)	α -KG DH., Pyruvic DH.
(almost completely)	Malic DH., Isocitric DH.
Hypotonic dissociable	(Cytochrome c in part), (Adenylate Kinase)

This assumption may be supported by the finding that the specific activity of α -ketoglutaric dehydrogenase was highest in the membrane fraction (R₂) obtained from frozen-thawed mitochondria. Electron microscope observation revealed that the submitochondrial fraction (R₂) was composed mainly of membrane fragments derived from the inner membrane containing the elementary particles¹³.





- Fig. 1 Electron micrograph of fluffy (F_3) fraction obtained from sonicated beef heart mitochondria (PTA negative staining). ($\times 55,000$)
- Fig. 2 Electron micrograph of F_3 -d₁ fraction obtained from F_3 fraction by sucrose density gradient. ($\times 55,000$)
- Fig. 3 Electron micrograph of F_3 -d₂ fraction obtained from F_3 fraction by sucrose density gradient. ($\times 55,000$)
- Fig. 4 Electron micrograph of F_3 -d₃ fraction obtained from F_3 fraction by sucrose density gradient. ($\times 55,000$)
- Fig. 5 Electron micrograph of F_3 -d₃ fraction. (same specimen as in Fig. 4). ($\times 55,000$)
- Fig. 6 Electron micrograph of F_3 -d₄ fraction obtained from F_3 fraction by sucrose density gradient. ($\times 55,000$)

The α -ketoglutaric dehydrogenase in the R_2 fraction can be solubilized by repeated washing of the R_2 fraction. Malic dehydrogenase, on the other hand, is easily solubilized by freezing-thawing of mitochondria. However, hypotonic treatment does not release TCA cycle dehydrogenases in any significant amount, although it does adenylate kinase and secondary phosphate transferases, which are supposed to be localized between the outer and the inner membranes of mitochondria¹⁴.




Fig. 7 Localization of TCA cycle dehydrogenases and electron transfer chain in the mitochondria

Recently, some informations have been obtained about the isolation of outer membrane and the localization of enzymes on the outer membrane of mitochondria. GREEN and his collaborators^{4,15-17} reported on the isolation and properties of the mitochondrial outer membrane, in which the activities of pyruvic dehydrogenase complex and citric cycle enzymes were observed. Further, it has been reported that rotenone insensitive $NADH_2$ -cytochrome c reductase (EC 1. 6. 2. 1) and cytochrome b^5 , of which α -band differs somewhat from that in endoplasmic reticulum, are contained in the mitochondrial outer membrane^{18,19}. SCHNEITMAN *et al*²⁰ demonstrated monoamine oxidase (EC 1. 4. 3. 4) to be a specific enzyme marker for the mitochondrial outer membrane.

The data presented in the present paper suggest that TCA cycle dehydrogenases seem to be localized in the matrix, in which α -ketoglutaric dehydrogenase may be loosely bound to the inner membrane while malic dehydrogenase may be of a soluble form. This assumption is compatible with the fact that externally added $NADH_2$ is scarcely oxidized by intact mitochondria as it cannot enter into the intact mitochondria, while it is most rapidly oxidized by inner membrane fragments, whose matrix side of the membrane is exposed to the reaction medium.

It is suggested that the localization of TCA cycle dehydrogenases in a close relation to the electron transfer chain is rational for the smooth operation of oxi-

dative phosphorylation in the mitochondria.

SUMMARY

The site of localization of TCA cycle dehydrogenases in mitochondria has been investigated by observing the dehydrogenase activities and fine structure of the fractionated samples after freezing and thawing or sonication of beef heart and rat liver mitochondria.

1. In the sonicated mitochondria, activities of malic and isocitric dehydrogenases were highest in the supernatant fraction centrifuged at 198,000 x g for 60 minutes, while the specific activity of α -ketoglutaric dehydrogenase was higher in the fluffy or residue fraction. The distribution of the activity of pyruvic dehydrogenase was similar to that of α -ketoglutaric dehydrogenase.

2. In a sucrose density gradient fractionation of the fluffy fraction obtained by centrifugation of sonicated mitochondria at 198,000 x g for 60 minutes, the activities of malic and pyruvic dehydrogenase were observed in the top (or low density) layer in the form of fine particles, while that of α -ketoglutaric dehydrogenase was observed in the middle (or medium density) layers in the form of aggregates of fine particles and membranous fragments.

3. In the samples fractionated after freezing and thawing of mitochondria, which were considered to be a relatively mild disruption, the specific activity of α -ketoglutaric dehydrogenase was higher in the residue (submitochondria) fraction than that in the supernatant fraction (centrifuged at 144,000 x g, 30 minutes), and the activity of malic dehydrogenase still remained significantly high in the residue fraction.

4. It was deduced that the TCA cycle dehydrogenases could be localized in the matrix of the mitochondria by a loose binding to the inner membrane.

REFERENCES

1. SCHNEIDER, W. C. and HOGEBOM, G. H. : Biochemistry of cellular particles. *Ann. Rev. Biochem.* 25, 201, 1956
2. HOGEBOM, G. H. and SCHNEIDER, W. C. : Sonic disintegration of isolated liver mitochondria. *Nature* 166, 302, 1950
3. GREEN, D. E. and FLEISCHER, S. : The mitochondrial system of enzymes. Metabolic pathway, Vol. I, p. 41, Academic Press, New York and London, 1960
4. ALLMANN, D. W. and BACHMAN, E. : The outer membrane of the mitochondria. *Fed. Proc.* 24, 425, 1965
5. HOGEBOM, G. H. : Fractionation of cell components of animal tissues. Methods in Enzymology, Vol. I, p. 16, Academic Press, New York, 1955
6. CRANE, F. L., GLENN, J. L. and GREEN, D. E. : Studies on the electron transfer system IV. The electron transfer particle. *Biochim. Biophys. Acta* 22, 475, 1956
7. MASSAY, V. : The composition of the α -ketoglutarate dehydrogenase complex. *Biochim. Biophys. Acta* 38, 447, 1960

8. PLAUT, G. W. E. and SUNG, S. C. : Diphosphopyridine nucleotide isocitric dehydrogenase from animal tissues. *Methods in Enzymology*. Vol. I, p.710, Academic Press, New York, 1955
9. OCHOA, S. : Malic dehydrogenase from pig heart. *Methods in Enzymology*. Vol. I, p. 735, Academic Press, New York, 1955
10. LOWRY, O. H., ROSEBROUGH, N. J., FARR, A. L. and RANDALL, R. J. : Protein measurement with the folin phenol reagent. *J. Biol. Chem.* 193, 265, 1951
11. GORNALL, A. G., BARDAWILL, C. J. and DAVID, M. M. : Determination of serum proteins by means of the biuret reaction. *J. Biol. Chem.* 177, 751, 1949
12. ODA, T. and NISHI, Y. : Fundamental structure and function of mitochondrial membrane. *J. Electron Microscopy* 12, 290, 1963
13. FERNÁNDEZ-MORÁN, H., ODA, T., BLAIR, P. V. and GREEN, D. E. : A macromolecular repeating unit of mitochondrial structure and function. *J. Cell Biol.* 22, 63, 1964
14. KLINGENBERG, M. and PFAFF, E. : Structure and functional compartmentation in mitochondria. I. E. G., No. 1, Sci. Memo #418, 1965
15. GREEN, D. E., BACHMANN, E. and ALLMANN, D. : Isolation and properties of the outer membrane of beef heart mitochondrion. *Arch. Biochem. Biophys.* 115, 172, 1966
16. BACHMANN, E., ALLMANN, D. and Green, D. E. : The membrane systems of the mitochondrion I. The S fraction of the outer membrane of beef heart mitochondria. *Arch. Biochem. Biophys.* 115, 153, 1966
17. ALLMANN, D., BACHMANN, E. and GREEN, D. E. : The membrane system of the mitochondrion II. The K fraction of the outer membrane of beef heart mitochondria. *Arch. Biochem. Biophys.* 115, 165, 1966
18. PARSONS, D. F., WILLIAMS, G. R., THOMPSON, W., WILLSON, D. and CHANCE, B. : Improvements in the procedure for purification of mitochondrial outer and inner membrane. Comparison of the outer membrane with smooth endoplasmic reticulum. I. E. G., No. 1, Sci. Memo #649, 1966; *Biochem. Biophys. Acta Library* by American Elsevier, to be published.
19. SOTTOCASS, G., KUYLENSTIERNA, B. and ERNSTER, L. : An electron transport system associated with the outer membrane of liver mitochondria. I. E. G., No. 1, Sci. Memo #652, 1966
20. SCHEITMAN, C., ERWIN, V. and GREENAWALT, J. : The submitochondrial localization of monoamine oxidase - an enzyme marker for the outer membrane of rat liver mitochondria. I. E. G., No. 1, Sci. Meamo #686, 1966; *J. Cell Biol.*, in press.