

Acta Medica Okayama

Volume 59, Issue 6

2005

Article 4

DECEMBER 2005

A comparison of the signal pathways between the TNF alpha- and oridonin-induced murine L929 fibrosarcoma cell death.

Jian Huang* Lijun Wu[†] Shin-ichi Tashiro[‡]
Satoshi Onodera** Takashi Ikejima^{††}

*﻿Shenyang Pharmaceutical University,

[†]﻿Shenyang Pharmaceutical University,

[‡]﻿Showa Pharmaceutical University,

**﻿Showa Pharmaceutical University,

^{††}﻿Shenyang Pharmaceutical University,

A comparison of the signal pathways between the TNF alpha- and oridonin-induced murine L929 fibrosarcoma cell death.*

Jian Huang, Lijun Wu, Shin-ichi Tashiro, Satoshi Onodera, and Takashi Ikejima

Abstract

Oridonin, an active component isolated from *Rabdosia rubescences*, has been reported to have antitumor effects. In this study, we compared the signal transduction pathways between TNFalpha-and oridonin-induced L929 cell death. Oridonin and TNFalpha initiated apoptotic morphologic changes, but DNA fragmentation was found in TNFalpha-treated L929 cells but not in oridonin-treated ones. The pan-caspase inhibitor (z-VAD-fmk), caspase-8 inhibitor (z-IETD-fmk) and caspase-3 inhibitor (z-DEVD-fmk) augmented oridonin-and TNFalpha-induced cell death. However, the caspase-9 inhibitor (z-LEHD-fmk) only increased oridonin-induced L929 cell death. Moreover, poly (ADPribose) polymerase (PARP) was cleaved in oridonin-treated L929 cells but not in the TNFalpha-treated groups, and the caspase-3 inhibitor (z-DEVD-fmk) failed to inhibit PARP cleavage. These results showed that only oridonin-induced L929 cell death required PARP degradation in a caspase-3 independent manner. In addition, oridonin increased the ratio of Bax/Bcl-2 protein expression, but TNFalpha did not. TNFalpha induced p38 and ERK activation, whereas oridonin triggered only ERK activation. We also investigated the effect of oridonin on intracellular TNFalpha expression, and found that oridonin augmented endogenous pro-TNFalpha expression and its upstream protein I κ B phosphorylation. These results indicated that although oridonin promoted endogenous pro-TNFalpha expression, a great difference existed between the signal pathways through which TNFalpha-and oridonin-induced cell death.

KEYWORDS: oridonin, caspase, Bax/Bcl-2, MAPK, I κ B

*PMID: 16418769 [PubMed - indexed for MEDLINE]

Copyright (C) OKAYAMA UNIVERSITY MEDICAL SCHOOL

Original Article

A Comparison of the Signal Pathways between the TNF α - and Oridonin-induced Murine L929 Fibrosarcoma Cell DeathJian Huang^{a,b}, Lijun Wu^b, Shin-ichi Tashiro^c,
Satoshi Onodera^c, and Takashi Ikejima^{a,*}^aChina-Japan Research Institute of Medical and Pharmaceutical Sciences,^bDepartment of Phytochemistry, Shenyang Pharmaceutical University, Shenyang, 110016, P. R. China, and^cDepartment of Clinical and Biomedical Science, Showa Pharmaceutical University, Machida, Tokyo 194–8543, Japan

Oridonin, an active component isolated from *Rabdosia rubescens*, has been reported to have antitumor effects. In this study, we compared the signal transduction pathways between TNF α - and oridonin-induced L929 cell death. Oridonin and TNF α initiated apoptotic morphologic changes, but DNA fragmentation was found in TNF α -treated L929 cells but not in oridonin-treated ones. The pan-caspase inhibitor (z-VAD-fmk), caspase-8 inhibitor (z-IETD-fmk) and caspase-3 inhibitor (z-DEVD-fmk) augmented oridonin- and TNF α -induced cell death. However, the caspase-9 inhibitor (z-LEHD-fmk) only increased oridonin-induced L929 cell death. Moreover, poly (ADP-ribose) polymerase (PARP) was cleaved in oridonin-treated L929 cells but not in the TNF α -treated groups, and the caspase-3 inhibitor (z-DEVD-fmk) failed to inhibit PARP cleavage. These results showed that only oridonin-induced L929 cell death required PARP degradation in a caspase-3 independent manner. In addition, oridonin increased the ratio of Bax/Bcl-2 protein expression, but TNF α did not. TNF α induced p38 and ERK activation, whereas oridonin triggered only ERK activation. We also investigated the effect of oridonin on intracellular TNF α expression, and found that oridonin augmented endogenous pro-TNF α expression and its upstream protein I κ B phosphorylation. These results indicated that although oridonin promoted endogenous pro-TNF α expression, a great difference existed between the signal pathways through which TNF α - and oridonin-induced cell death.

Key words: oridonin, caspase, Bax/Bcl-2, MAPK, I κ B

Tumor necrosis factor- α (TNF α), a pleiotropic cytokine, is an important mediator in many biologic responses in different cell types through binding to either 55- or 75-KD membrane receptors [1, 2]. These biological responses include the inhibition of cell growth and induction of differentiation and apoptosis [3–

5]. *In vitro*, TNF α induces cell death in a variety of malignant cells. Among these cell lines, it is widely believed that murine L929 fibrosarcoma cells are significantly sensitive to TNF α . Some researchers have reported that caspases do not induce L929 cell death, but protect L929 against cell death in the TNF α -treated L929 cells [6].

Oridonin is a diterpenoid isolated from *Rabdosia rubescens*, which has been reported to have various pharmacological and physiological effects such as anti-

Received February 4, 2005; accepted June 30, 2005.

*Corresponding author. Phone: +86-24-2384463; Fax: +86-24-2384463

E-mail: ikejimat@vip.sina.com (T. Ikejima)

inflammatory, anti-bacterial and anti-tumor effects [7-8]. Our previous study showed that oridonin had cytotoxic effects on L929 cells, and that caspase inhibitors also did not protect against oridonin-induced L929 cell death, but instead made the L929 cells more sensitive to oridonin. Therefore, we compared the signal pathways between the TNF α - and oridonin-induced murine L929 fibrosarcoma cell death.

It is well known that the caspase family, mitochondria, and MAPK family signal transduction pathways play an important role in regulating apoptosis. In this study, we further investigated the signal transduction pathways mentioned above to analyze the relationship between TNF α - and oridonin-induced L929 cell death. The results showed that oridonin was different from TNF α in many aspects of induction of L929 cell death though oridonin-promoting endogenous pro-TNF α expression.

Materials and Methods

Regents. Oridonin was obtained from the Beijing Institute of Biological Products (Beijing, China). The structure of oridonin was assigned by comparing the chemical and spectral data ($^1\text{H-NMR}$, $^{13}\text{C-NMR}$) with those reported in the literature. The purity of the oridonin was measured by HPLC and determined to be 99.2%, and oridonin was resolved in the medium. Human TNF α was prepared from PMAL-C₂-TNF α /JM109 (*E. coli*) in our laboratory. Pan-caspase inhibitor (z-VAD-fmk), caspase-3 inhibitor (z-DEVD-fmk), caspase-9 inhibitor (z-LEHD-fmk) and caspase-8 inhibitor (z-IETD-fmk) were purchased from Enzyme Systems (Livermore, CA, USA). ERK inhibitor (PD98059), p38 inhibitor (SB203580), and JNK inhibitor (SP600125) were obtained from Calbiochem (La Jolla, CA, USA). Polyclonal antibodies against I κ B phospho-I κ B, β -actin, and horseradish peroxidase-conjugated secondary antibodies (goat-anti-rabbit and goat-anti-mouse) were purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Anti-Bcl-2 and Bax were from Oncogene Research Products (Boston, MA, USA).

Cell Culture. The murine fibrosarcoma cells (L929#CRL-2148) were purchased from American Type Culture Collection (ATCC, Manassas, VA, USA). The cells were cultured in RPMI-1640 medium (GIBCO, NYC, NY, USA.) supplemented with 10% FBS (Shengma Yuanheng, Beijing, China), 100 mg/l streptomycin, 100 IU/ml penicillin, and 0.03% L-glutamine and

maintained at 37 °C with 5% CO₂ in a humidified atmosphere.

Cytotoxicity Assay. The L929 cells were incubated at 1×10^5 cells/well in 96-well plates (NUNC, Roskilde, Denmark). The cells were incubated with z-VAD-fmk, z-IETD-fmk, z-DEVD-fmk, z-LEHD-fmk, cyclosporine A, PD98059, SB203580, or SP600125 at given concentrations for 1 h, and then treated with TNF α for different time periods. Cell growth was measured with a 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium (MTT) assay. The percentage of cell growth inhibition was calculated as follows:

$$\text{Cell death (\%)} = \frac{[A_{570}(\text{control}) - A_{570}(\text{oridonin})]}{A_{570}(\text{control})} \times 100$$

Observation of Morphological Changes.

The L929 cells were divided into 2 groups and placed on culture plates and cultured for 18 h. One group was treated with the control medium and the other group was treated with TNF α (20 ng/ml) and the cellular morphology was observed using phase contrast microscopy (Leica, Wetzlar, Germany).

Nuclear Damage Observed by Hoechst 33258 Staining. Apoptotic nuclear morphology was assessed using Hoechst 33258. The cells were fixed with 3.7% paraformaldehyde for 30 min at room temperature and then washed and stained with 167 $\mu\text{mol/l}$ Hoechst 33258 at 37 °C for 30 min. The cells were washed and resuspended in PBS for morphologic observation using fluorescence microscopy (Leica, Wetzlar, Germany).

DNA fragmentation assay. The L929 cells (1×10^6) were collected by centrifugation at $150 \times g$ for 5 min, and washed with PBS. The cells were pelleted and suspended in 10 mM Tris (pH 7.4), 10 mM EDTA (pH 8.0), and 0.5% Triton X-100 and kept at 4 °C for 10 min. The supernatant was incubated with 20 mg/ml RNase A (2 μl) and 20 mg/ml proteinase K (2 μl) at 37 °C for 1 h, then kept in 0.5 M NaCl (20 μl) and isopropanol (120 μl) at -20 °C overnight, and then centrifuged at $15,000 \times g$ for 15 min. DNA were dissolved in TE buffer [10 mM Tris (pH 7.4), 10 mM EDTA (pH 8.0)] and subjected to 2% agarose gel electrophoresis at 50 V for 40 min and stained with ethidium bromide.

LDH Activity-Based Cytotoxicity Assays. The LDH (lactate dehydrogenase) activity was assessed using a standardized kinetic determination (Zhongsheng LDH kit, Beijing, China). The LDH activity was

measured in both floating dead cells and viable adherent cells. Floating cells were collected from the culture medium by centrifugation ($240 \times g$) at 4°C for 5 min, and the LDH obtained from the pellets was used as an index of apoptotic cell death (LDHp). The LDH released in the culture medium (extracellular LDH, or LDHe) was used as an index of necrotic death, and the LDH present in the adherent viable cells as intracellular LDH (LDHi). The percentages of apoptotic and necrotic cell death were calculated as follows:

$$\text{Apoptosis (\%)} = \frac{\text{LDHp}}{\text{LDHp} + \text{LDHi} + \text{LDHe}} \times 100$$

$$\text{Necrosis (\%)} = \frac{\text{LDHe}}{\text{LDHp} + \text{LDHi} + \text{LDHe}} \times 100$$

Western Blot Analysis. The L929 cells were treated with TNF α (20 ng/ml) for different time periods. Both adherent and floating cells were collected, then Western-blot analysis was performed as previously described, with some modification. Briefly, the cell pellets were resuspended in lysis buffer consisting of Hepes 50 mmol/l pH 7.4, Triton-X 100 1%, sodium orthovanada 2 mmol/l, sodium fluoride 100 mmol/l, edetic acid 1 mmol/l, PMSF 1 mmol/l, aprotinin (Sigma, MO, USA) 10 mg/l and leupeptin (Sigma) 10 mg/l and lysed at 4°C for 60 min. After $13,000 \times g$ centrifugation for 15 min, the protein content of the supernatant was determined by a protein assay reagent (Bio-Rad, Hercules, CA, USA). The protein lysates were separated by electrophoresis in 12% SDS polyacrylamide gel and blotted onto a nitrocellulose membrane. Proteins were detected using polyclonal antibody and visualized using anti-rabbit IgG conjugated with peroxidase (HRP) and 3,3-diaminobenzidine tetrahydrochloride (DAB) as the HRP substrate.

Statistical Analysis of the Data. The data are expressed as means \pm S.D. Statistical comparisons were made by student's *t*-test. $p < 0.05$ was considered significant.

Results

Cytotoxic Effects of TNF α and oridonin on L929 cells. To investigate the cytotoxic effects of oridonin and TNF α against L929 cells, the L929 cells were cultured with oridonin or TNF α at various doses for 24 h. We found that both oridonin and TNF α induced L929 cell death in a dose-dependent manner.

Oridonin 25 μM or TNF α 20 ng/ml caused almost 50% of the L929 cell deaths (Figs. 1A, B).

Characterization of TNF α - and oridonin-induced L929 cell death. To characterize the oridonin- and TNF α -induced L929 cell death, we examined the morphologic changes and DNA fragmentation. When L929 cells were cultured with oridonin 25 μM or 20 ng/ml TNF α for 18 h, marked morphologic changes were observed as compared with the medium control group. The oridonin- or TNF α -treated L929 cells underwent contraction and became round in shape, with shrunken nuclei (Figs. 2 C, J, E, H), but the untreated cells did not show these apoptotic changes (Fig. 2A). Morphologic changes were further confirmed by Hoechst 33258 staining of the cell nuclei. In the control group, the nuclei of the L929 cells were round and homogeneously stained (Fig. 2B), but the 25 μM oridonin- or 20 ng/ml TNF α -treated cells showed marked nuclear fragmentation (Figs. 2D, F).

DNA regular fragmentation is another hallmark of typical apoptosis [9]. In our study, DNA ladders were observed in TNF α -treated but not in oridonin-treated L929 cells by agarose electrophoresis (Fig. 3).

In order to further investigate whether TNF α and oridonin also initiated necrosis in L929 cells, the rates of lactate dehydrogenase (LDH) released from viable cells, floating dead cells, and the culture medium were measured. In the presence of oridonin 5–50 μM , the number




Fig. 1 Cytotoxic effects of oridonin or TNF α on L929 cells. The cells were treated with various doses of oridonin (A) or TNF α (B) for 24 h. The cell death rate was measured by MTT assay. $n = 3$, Mean \pm S.D.




Fig. 2 Oridonin- and $\text{TNF}\alpha$ -induced the morphological changes in L929 cells. The cellular morphologic changes were observed after the cells were cultured with $25\ \mu\text{M}$ oridonin ($\times 200$ magnification C, $\times 400$ magnification J) or $20\ \text{ng/ml}$ $\text{TNF}\alpha$ ($\times 200$ magnification E, $\times 400$ magnification H), or medium ($\times 200$ magnification A) for 18 h. Morphologic changes of the nuclei were also observed with Hoechst 33258 staining after the cells were cultured with $25\ \mu\text{M}$ oridonin ($\times 200$ magnification D), $20\ \text{ng/ml}$ $\text{TNF}\alpha$ ($\times 200$ magnification F) or medium ($\times 200$ magnification B) for 18 h.

of necrotic cells increased from 5.5% to 32.3% (Fig. 4A). Moreover, in the presence of $\text{TNF}\alpha$ 10–400 ng/ml, the number of necrotic cells increased from 8.8% to 37.4% (Fig. 4B). Therefore, we concluded that both $\text{TNF}\alpha$ and oridonin induced cell death through the balance between apoptosis and necrosis.

The caspases protected against $\text{TNF}\alpha$ - and oridonin-induced cell death. It is well known that caspases are required in apoptosis. Some researchers have also reported that caspase inhibitors render L929 cells more sensitive to $\text{TNF}\alpha$ [10, 11]. We further compared the function of caspase inhibitors in $\text{TNF}\alpha$ -




Fig. 3 Oligonucleosomal DNA fragmentation analysis with Oridonin- and TNF α -treated L929 cells on agarose gel electrophoresis. After the cells were treated with 20 ng/ml TNF α (A) or 25 μ M oridonin (B) for 0, 12, 18 and 24 h. The DNA was separated on 2% agarose gel. M: Marker.





Fig. 4 Characterization of cell death induced by oridonin or TNF α in L929 cells. The cells were treated with oridonin (A) or TNF α (B) with various doses for 24 h. The cell death rate was measured by an LDH activity-based assay. $n = 3$, mean \pm S.D.

and oridonin-induced L929 cell death. In the TNF α - and oridonin-treated L929 cells, the inhibitor of caspase-3, z-DEVD-fmk, the inhibitor of caspase-8, z-IETD-fmk, and the pan-caspase inhibitor, z-VAD-fmk, augmented the cell death to a different degree. The caspase-9 inhibitor, z-LEHD-fmk, had almost no effect on the TNF α -induced L929 cell death, but it made the cells even more sensitive to oridonin. In the 25 μ M oridonin-treated L929 cells, the caspase-9 inhibitor, z-LEHD-fmk, augmented the cell death ratio from 42% (cultured with oridonin alone) to 69% at 24 h (Figs. 5A, B). In the LDH-based assay, we found that the caspase inhibitors increased both apoptosis and necrosis in the oridonin-treated L929 cells (Fig. 5 C). The results suggest that the caspases play protective roles against TNF α - or


oridonin-induced L929 cell death, but caspase-9 was only effective on oridonin-induced L929 cell death.

Bcl-2/Bax involvement in oridonin-induced but not in TNF α -induced L929 cell death.

It is reported that the Bcl-2 family of proteins plays a role in either the inhibition or promotion of apoptotic cell death [12]. We also examined the expression of the Bcl-2 family protein, Bcl-2 and Bax, in oridonin- and TNF α -treated L929 cells by Western-blot analysis. The results showed that the treatment of L929 cells with 25 μ M oridonin decreased the Bcl-2 expression and increased the Bax expression, but in the TNF α -treated L929 cells, the Bax and Bcl-2 expressions did not change with the time period (Fig. 6), indicating that only the Bcl-2/Bax ratio was involved in the oridonin-induced



A



C

L929 cell death.

MAPK modulation in TNF α - and oridonin-induced L929 cell death. MAPK signaling pathways have been shown to be implicated in cell growth and cell death [13]. To compare the regulation of the MAPK cascades in TNF α - and oridonin-induced cell death, specific inhibitors for p38 (SB203580), JNK (SP600125) and ERK (PD98059), were applied to evaluate the function of MAPK in both the TNF α - and oridonin-induced L929 cell death. After incubation with 20 ng/ml TNF α or 25 μ M oridonin for various periods of time, 5 μ M SB 203580 and PD98059 markedly

	Cell death (%)
TNF α	51.2 \pm 0.9
z-IETD-fmk	13.2 \pm 2.1
TNF α + z-IETD-fmk	76.2 \pm 2.2**
z-DEVD-fmk	10.3 \pm 1.8
TNF α + z-DEVD-fmk	80.7 \pm 0.9**
z-LEHD-fmk	6.7 \pm 0.9
TNF α + z-LEHD-fmk	50.8 \pm 1.6
z-VAD-fmk	9.7 \pm 1.3
TNF α + z-VAD-fmk	87.9 \pm 1.7**

B

Fig. 5 Effects of caspase inhibitors on oridonin- and TNF α -induced L929 cell death. The cells were treated with 25 μ M oridonin (A) or 20 ng/ml TNF α (B) for 24 h in the presence of z-IETD-fmk 5 μ M (caspase-8), z-DEVD-fmk 5 μ M (caspase-3), z-LEHD-fmk 5 μ M (caspase-9) or z-VAD-fmk 5 μ M (pan-caspase), and the cell death rate was measured by MTT assay. The cells were treated as in (A), and the cell death rate was measured by LDH activity-based assay (C). n = 3, Mean \pm S.D. **, P < 0.01

reduced TNF α -induced cell death, but the PD98059 inhibitory function declined with the time period. However, in the oridonin-treated L929 cells, only PD98059 effectively reversed the cell death (Fig. 7). The results showed that in the TNF α - and oridonin-treated L929 cells, ERK had the same functions of mediating the cell death, but p38 merely regulated the TNF α -induced L929 cell death.

PARP regulation in both the TNF α - and oridonin-induced L929 cell death. To investigate the role of PARP, a downstream regulator, the PARP inhibitor (DPQ) was applied to access the PARP





Fig. 6 Western-blot analysis of Bax and Bcl-2 protein expression in oridonin- and TNF α -treated L929 cells. The cells were treated with 25 μ M oridonin (A) or 20 ng/ml TNF α (B) for various time periods. The cell lysates were separated by 12% SDS-PAGE electrophoresis and the protein expression was detected by Western-blot analysis.



	Cell death (%)		
	12	18	24 (h)
TNF α	40.2 \pm 2.1	50.3 \pm 3.2	69.4 \pm 1.2
TNF α + PD98059	23.0 \pm 2.6**	46.2 \pm 1.5	69.0 \pm 1.4
TNF α + SB203580	21.3 \pm 2.8**	35.3 \pm 2.1**	36.3 \pm 0.8**
TNF α + SP600125	45.8 \pm 0.7	60.6 \pm 2.1	74.9 \pm 2.3

B

Fig. 7 Effects of MAPK inhibitors on oridonin- and TNF α -treated L929 cell death. The cells were treated with 25 μ M oridonin (A) or 20 ng/ml TNF α (B) and MAPK inhibitors including 5 μ M SP600125 (JNK), 5 μ M SB203580 (p38) or 5 μ M PD98059 (ERK) for various time periods. Cell death was measured by MTT assay. n = 3, Mean \pm S.D., **p < 0.01 vs TNF α or oridonin group.




Fig. 8 Effects of PARP inhibitor DPQ on oridonin- and TNF α -induced L929 cell death. The cells were treated with 25 μ M oridonin or 20 ng/ml TNF α in the presence of the PARP inhibitor DPQ with various doses for 18 h. Cell death was measured by the MTT assay n = 3, Mean \pm S.D., **p < 0.01 vs oridonin (alone) group.

function in both the oridonin- and TNF α -induced L929 cell death. It was found that 20 μ M DPQ reduced the oridonin-induced cell death from 60.1 to 41.2% at 18 h; however, DPQ almost had no inhibitory effect on the TNF α -induced L929 cell death (Fig. 8). In order to further confirm the PARP participation in the oridonin-induced cell death, we examined the PARP protein expression by Western-blot analysis. The results showed that after 18 h treatment with 25 μ M oridonin, 116 KDa PARP protein was cleaved to the 85 KDa fragment, and that the caspase-3 inhibitor, z-DEVD-fmk, failed to inhibit the PARP cleavage (Fig. 9), but in the 20 ng/ml TNF α -treated group, we did not detect the PARP degradation, indicating that PARP was required for the oridonin-induced L929 cell death but not for the TNF α -induced cell death in a caspase-3 independent manner.

Oridonin up-regulated intracellular pro-TNF α expression in L929 cells. We also examined the effect of oridonin on the intracellular pro-TNF α expression by Western-blot analysis. Exposure of the L929 cells to 25 μ M oridonin for 14 h initiated the pro-TNF α expression. In order to further confirm this, we also analyzed the I κ B expression and I κ B phosphorylation in oridonin-treated L929 cells. The NF- κ B transcription factor plays an essential role in the expression of proinflammatory cytokines such as TNF α and IL-1. NF- κ B activity is controlled by the inhibitory I κ B protein, which masks the nuclear localization sequence in the Rel homology domain of NF- κ B, thereby sequestering NF- κ B in a latent state in the cytoplasm [14-16]. The results showed that treatment with 25 μ M oridonin for 12 h reduced the I κ B expression and increased I κ B




Fig. 9 Western-blot analysis of PARP protein expression in oridonin- and TNF α -treated L929 cells. The cells were treated with 25 μ M oridonin (A) or 20 ng/ml TNF α (B) for various time periods. The cell lysates were separated by 12% SDS-PAGE electrophoresis, and the protein expression was detected by Western-blot analysis.




Fig. 10 Western-blot analysis of endogenous pro-TNF α protein expression in oridonin- treated L929 cells. The cells were treated with 25 μ M oridonin for various time periods. The cell lysates were separated by 12% SDS-PAGE electrophoresis, and the protein expression was detected by Western-blot analysis.

phosphorylation (Fig. 10). Thus, these results suggested that oridonin up-regulated the TNF α expression through activation of the transcription factor NF- κ B.

Discussion

In this study we compared the signal pathways between the TNF α - and oridonin-induced murine L929 fibrosarcoma cell death, and the results showed that differences exist between the oridonin- and TNF α -induced cell death signal pathways. Although we observed the apoptotic changes of cellular morphology in both the oridonin- and TNF α -treated groups, DNA fragmentation did not occur in the oridonin-treated group, but did in the TNF α -treated group. DNA fragmentation, a hallmark of apoptosis, is the internucleosomal DNA degradation result, which a Ca²⁺, Mg²⁺-dependent deoxyribonuclease I (DNase I) accounts for. In addition, several isoforms of DNase I exist, and some of them are unable to degrade the DNA into the apoptotic ladder, but cause a fuzzy pattern of digested nuclear DNA when apoptosis occurs [17]. Based on the results of our experiment, we speculated that unlike TNF α , oridonin might initiate the activation of the isoform(s) of DNase I, and this kind of DNase I digested the DNA into smaller-sized fragments.

Caspases are a family of cysteases, which cleave protein substrates after their Asp residues. Caspases appear to be involved in regulating the activation of apoptotic signal transmission, and are divided into 2 classes based on the lengths of their N-terminal prodomains, including upstream caspases such as caspase-2, -8, and -10, and downstream caspases such as caspase-3, -6, and -7 [18]. In addition, previous reports have demonstrated that the caspase inhibitors augment TNF α -induced cell death in L929 cells and in RAW246.7 macrophages [19]. In L929 cells, it has been inferred that caspases act as a negative regulator of reactive oxygen intermediate (ROI) [6]. In this study, the inhibitor of caspase-3, z-DEVD-fmk, the inhibitor of caspase-8, z-IETD-fmk, and the pan-caspase inhibitor, z-VAD-fmk augmented the cell death in both the TNF α - and oridonin-treated L929 cells; however, the caspase-9 inhibitor, z-LEHD-fmk, only made the L929 cells more sensitive to oridonin, indicating that caspase-9 was only activated in oridonin-induced L929 cell death. Bcl-2 family members are characterized by the fact that they contain at least one of four Bcl-2 homology domains

(BH1-BH4). Some of these proteins, such as Bax, function to promote apoptosis, whereas others like Bcl-2 inhibit apoptosis [12]. In this study, only oridonin up-regulated the Bax expression and down-regulated the Bcl-2 expression in L929 cells. In mitochondria, when apoptosis occurred, cytochrome *c* is released into the cytoplasm, and then cytochrome *c*, Apaf-1 and procaspase-9 form a complex. Activated caspase-9 cleaved the caspase-3 proenzyme [20]. PARP is a nuclear enzyme that responds to DNA damage, facilitates DNA repair, and promotes cell survival [21]. When apoptosis occurs, it is cleaved to an 85-KDa fragment by caspase-3 [22]. In this study, the PARP inhibitor DPQ reduced oridonin-induced L929 cell death, but had no inhibitory effect on TNF α -induced L929 cell death. In addition, an 85-KDa fragment was also observed after 12 h treatment with oridonin, and the caspase-3 inhibitor failed to suppress PARP degradation, indicating that PARP was involved in oridonin-induced L929 cell death but not in TNF α -induced L929 cell death. Together, all these results suggested that unlike in TNF α -induced cell death, oridonin-induced L929 cell death required the up-regulation of Bax, caspase-9 activation and PARP cleavage, and that some protease(s) different from caspase-3 might be responsible for the degradation of PARP.

The MAPK family members, including extracellular signal-regulated protein kinase (ERK), c-Jun N-terminal kinase (JNK) and p38, play important roles in apoptosis and necrosis. The ERK pathway is predominantly activated by mitogens through a Ras-dependent mechanism, which is required for cell proliferation and differentiation; however, JNK and p38 are activated by pro-inflammatory cytokines and various environmental stresses [23, 24]. In this study, the results showed that in both the TNF α - and oridonin-treated L929 groups, ERK had the same functions of mediating cell death, but p38 only mediated TNF α -induced L929 cell death. Activation of the distinct MAPK subtype cascade depends on the types of cells and the stimuli, and the functional role of each MAPK subtype may be different according to the cell types [13]. Therefore, ERK was activated as the same functional pathway in both the TNF α - and oridonin-induced L929 cell death; however, the p38 signal cascade might be activated as different signal pathways involved in the TNF α -treated L929 cells.

TNF α is a pleiotropic cytokine which is involved in inflammatory and immune responses. The nuclear factor

NF- κ B plays a crucial role in pro-TNF α gene expression mediated by agents such as TNF α and PMA. I κ B is phosphorylated and targeted to the 26S proteasome complex, resulting in the release and nuclear translocation of NF- κ B [14–16]. We investigated the effect of oridonin on TNF α expression, and the results showed that oridonin increased pro-TNF α expression and I κ B phosphorylation, and reduced I κ B expression, indicating that oridonin up-regulated TNF α expression through activating the transcription factor NF- κ B. Endogenous pro-TNF α might be cleaved and secreted into medium, then TNF α might be bound to its receptors and promote oridonin-induced cell death. However, in our study, oridonin induced L929 cell death in very different ways from TNF α , so we speculated that the pro-TNF α was not cleaved or secreted into the medium.

In summary, we compared the signal pathways between the TNF α - and oridonin-induced murine L929 fibrosarcoma cell deaths and found that oridonin increased the endogenous pro-TNF α expression. However, there was a considerable difference between the TNF α - and oridonin-induced L929 cell death signal pathways. Moreover, oridonin is a diterpenoid compound and easily enters the cytoplasm, so target proteins may exist for oridonin to bind with in the upstream of the cell death signaling pathway.

References

- Smith CA and Goodwin RG: The TNF receptor superfamily of cellular and viral proteins: activation, costimulation, and death. *Cell* (1994) 76: 959–962.
- Beutler B and van Huffel C: Unraveling function in the TNF ligand and receptor families. *Science* (1994) 264: 667–668.
- Vassalli P: The pathophysiology of tumor necrosis factors. *Annu Rev Immunol* (1992) 10: 411–452.
- Fiers W: Biologic therapy with TNF; in *Biologic therapy of cancer*, De Vita VT, Jr, Hellman S and Rosenberg SA eds, 2nd Ed, JP Lippincott Co., New York (1995) pp 295–327.
- Wallach D, Boldin M, Varfolomeev E, Beyaert R, Vandenebeele P and Fiers W: Cell death induction by receptors of the TNF family: Towards a molecular understanding. *FEBS Lett* (1997) 410: 96–106.
- Goossens V, Grooten JK, De Vos K and Fiers W: Direct evidence for tumor necrosis factor-induced mitochondrial reactive oxygen intermediates and their involvement in cytotoxicity. *Proc Natl Acad Sci USA* (1995) 92: 8115–8119.
- Osawa K, Yasuda H, Maruyama T, Morita H, Takeya K and Itokawa H: An investigation of diterpenes from the leaves of *Rabdosia trichocarpa* and their antibacterial activity against oral microorganisms. *Chem Pharm Bull* (1994) 42: 922–925.
- Fuji K, Node M, Sai M, Fujita E, Takeda S and Unemi N: Terpenoids. LIII. Antitumor activity of trichorabdals and related compounds. *Chem Pharm Bull* (1989) 37: 1472–1476.
- Mizukami S, Kikuchi K, Higuchi T, Urano Y, Mashima T, Tsuruo T and Nagano T: Imaging of caspase-3 activation in HeLa cells stimulated with etoposide using a novel fluorescent probe. *FEBS Lett* (1999) 453: 356–360.
- Vercammen D, Beyaert R, Denecker G, Goossens V, Van Loo G, Declercq W, Grooten J, Fiers W and Vandenebeele P: Inhibition of caspases increases the sensitivity of L929 cells to necrosis mediated by tumor necrosis factor. *J Exp Med* (1998) 187: 1477–1485.
- Los M, Mozoluk M, Ferrari D, Stepczynska A, Stroh C, Renz A, Herceg Z, Wang ZQ and Schulze-Osthoff K: Activation and Caspase-mediated Inhibition of PARP: A Molecular Switch between Fibroblast Necrosis and Apoptosis in Death Receptor Signaling. *Mol Biol Cell* (2002) 13: 978–988.
- Bernardi P, Broekemeier KM and Pfeiffer DR: Recent progress on regulation of the mitochondrial permeability transition pore; a cyclosporin-sensitive pore in the inner mitochondrial membrane. *J Bioenerg Biomembr* (1994) 26: 509–517.
- Schaeffer HJ and Weber MJ: Mitogen-activated protein kinases: specific messages from ubiquitous messengers. *Mol Cell Bio* (1999) 19: 2435–2444.
- Baeuerle PA: I κ B-NF- κ B structures: at the interface of inflammation control. *Cell* (1998) 95: 729–731.
- Huxford T, Huang DB, Malek S and Ghosh G: The crystal of the I κ B α /NF- κ B complex reveals mechanism of NF- κ B inactivation. *Cell* (1998) 95: 759–770.
- Jacobs MD and Harrison SC: Structure of an I κ B α /NF- κ B complex. *Cell* (1998) 95: 749–758.
- Peitsch MC, Polzar B, Stephan H, Crompton T, MacDonald HR, Mannherz HG and Tschopp J: Characterization of the endogenous deoxyribonuclease involved in nuclear DNA degradation during apoptosis (programmed cell death). *EMBO J* (1993) 12: 371–377.
- Nicholson DW, Ali A, Thornberry NA, Vaillancourt JP, Ding CD, Gallant M, Gareau Y, Griffin PR, Labelle M and Lazebnik YA: Identification and Inhibition of the ICE/CED-3 Protease Necessary for Mammalian Apoptosis. *Nature* (1995) 376: 37–43.
- Kim SO and Han J: Pan-caspase inhibitor zVAD enhances cell death in RAW246.7 macrophages. *J Endotoxin Res* (2001) 7: 292–296.
- Srinivasan A, Li F, Wong A, Kodandapani L, Smidt R Jr, Krebs JF, Fritz LC, Wu JC and Tomaselli KJ: Bcl-xL Functions Downstream of Caspase-8 to Inhibit Fas- and Tumor Necrosis Factor Receptor 1-induced Apoptosis of MCF7 Breast Carcinoma Cells. *J Biol Chem* (1998) 273: 4523–4529.
- Yu SW, Wang H, Poitras MF, Coombs C, Bowers WJ, Federoff HJ, Poirier GG, Dawson TM and Dawson VL: Mediation of poly (ADP-Ribose) polymerase-1-dependent cell death, by apoptosis-inducing factor. *Science* (2002) 297: 259–263.
- Mandal M, Maggirwar SB, Sharma N, Kaufmann SH, Sun SC and Kumar R: Bcl-2 Prevents CD95 (Fas/APO-1)-induced Degradation of Lamin B and Poly (ADP-ribose) Polymerase and Restores the NF- κ B Signaling Pathway. *J Biol Chem* (1996) 271: 30354–30359.
- Zoratti M and Szabo I: The Mitochondrial Permeability Transition. *Biochim Biophys* (1995) 1241: 139–176.
- Pastorino JG, Chen ST, Tafani M, Snyder JW and Farber JL: The overexpression of bax produces cell death upon induction of the mitochondrial permeability transition. *J Biol Chem* (1998) 273: 7770–7775.