

Acta Medica Okayama

Volume 35, Issue 4

1981

Article 7

OCTOBER 1981

14q12 translocation in a non-Burkitt lymphoma.

Kanji Miyamoto*

Kyoichi Hayashi†

Teruhiko Tsubota‡

Toshio Tanaka**

*Okayama University,

†Okayama University,

‡Okayama University,

**Okayama University,

14q12 translocation in a non-Burkitt lymphoma.*

Kanji Miyamoto, Kyoichi Hayashi, Teruhiko Tsubota, and Toshio Tanaka

Abstract

Chromosome analysis was performed on cells from a patient of null cell lymphoma, well-differentiated type. A 14q12 translocation was observed in all the banded cells. In addition, there were multiple chromosome abnormalities. This case will be useful in considering the significance of the 14q1(1-3) translocation in malignant lymphoma disease.

KEYWORDS: malignant lymphoma, chromosome analysis, 14q12 translocation.

— BRIEF NOTE —

14q12 TRANSLOCATION IN A NON-BURKITT LYMPHOMA

Kanji MIYAMOTO, Kyoichi HAYASHI*
Teruhiko Tsubota* and Toshio TANAKA**

*Division of Pathology, Cancer Institute (Director: Prof. J. Sato); * Department of Medicine
(2nd Clinic); and ** Pathology Section, Central Laboratories,
Okayama University Medical School, Okayama.*

Received March 9, 1981



Abstract. Chromosome analysis was performed on cells from a patient of null cell lymphoma, well-differentiated type. A 14q12 translocation was observed in all the banded cells. In addition, there were multiple chromosome abnormalities. This case will be useful in considering the significance of the 14q1(1-3) translocation in malignant lymphoma disease.

Key words : malignant lymphoma, chromosome analysis, 14q12 translocation.

Abnormalities of chromosome 14 have been found frequently in various lymphoid malignancies including conventional malignant lymphomas, African and non-African Burkitt lymphomas (1-3), and immunodeficiency diseases (4-6). The break point was band 14q32 in most cases of lymphoid malignancies (1-3). On the other hand, a band which has another break point, *i.e.*, 14q1(1-3), was reported in some cases such as ataxia teleangiectasia (4-6). Recently, Fukuhara *et al.* described 14q1(2-3) translocation in patients with poorly differentiated lymphoma, histiocytic lymphoma, Hodgkin's disease and T-cell lymphoma (7). The present study describes a 14q12 translocation in a patient with non-Burkitt lymphoma of null cell type.

A 65-year-old man was admitted to the Department of Medicine, Okayama University Hospital, in March 1980 with a diagnosis of malignant lymphoma of diffuse, well-differentiated lymphocytic type (Fig. 1). The patient was treated with adriamycin, Vincristine, endoxan and prednisolone, and complete remission was achieved. Seven months after the first admission, however, the patient relapsed with generalized lymphadenopathy. Treatment with combination chemotherapy was given, but the patient died of leukemic conversion three months after the second admission.

Chromosome analysis was performed on peripheral blood leukocytes after relapse. At that time, the white blood cell count was 42,900/mm³ with 89% of lymphocytes (Fig. 2); the surface marker was of null cell type. Peripheral blood was cultured for 24 h without phytohemagglutinin. Chromosomes were prepared



using the air-dry method, banded with quinacrine mustard, and stained with conventional Giemsa.

The number of chromosomes ranged from 46 to 49 with a bimodal number of 47 and 48. All of 10 banded cells had a 14q12 translocation. The exact origin of extra chromosome material which attached to band 14q12 could not be determined. The main karyotype was 48,XY,1q+,3q+,+i(3p),-6,-10,t(14;?)(q12;?),+16,17p-,+18,+mar (Fig. 3). The marker chromosome appeared to include 14pter→14q12.

In the present study, we demonstrated a 14q12 translocation in a null cell lymphoma, although the origin of the translocated segment could not be identified. In addition, there were multiple chromosome abnormalities as shown in Fig. 3. This case will be useful in considering the significance of 14q translocation in malignant lymphoma disease.

REFERENCES

1. McCaw, B. K., Epstein, A. L., Kaplan, H. S. and Hecht, F.: Chromosome 14 translocation in African and North American Burkitt's lymphoma. *Int. J. Cancer* **19**, 482-486, 1977.
2. Fukuhara, S. and Rowley, J. D.: Chromosome 14 translocations in non-Burkitt lymphomas. *Int. J. Cancer* **22**, 14-21, 1978.
3. Douglass, E. C., Magrath, I. M., Lee, E. C. and Whang-Peng, J.: Cytogenetic studies in non-African Burkitt lymphoma. *Blood* **55**, 148-155, 1980.
4. McCaw, B. K., Hecht, F., Harnden, D. G. and Teplitz, R. L.: Somatic rearrangement of chromosome 14 in human lymphocytes. *Proc. Natl. Acad. Sci. USA* **72**, 2071-2075, 1975.
5. Levitt, R., Pierre, R. V., White, W. L. and Siekert, R. G.: Atypical lymphoid leukemia in ataxia telangiectasia. *Blood* **52**, 1003-1011, 1978.
6. Saxon, A., Stevens, R. H. and Golde, D. W.: Helper and suppressor T-lymphocyte leukemia in ataxia telangiectasia. *N. Engl. J. Med.* **300**, 700-704, 1979.
7. Fukuhara, S., Ueshima, Y., Shirakawa, S. and Uchino, H.: Cytogenetic approaches to the classification of lymphoid malignancy: A proposal of t(14q) type lymphoid malignancies and the phenotype. *Acta Haematol. Jpn.* **44**, 267 (Abst.), 1981 (in Japanese).

Fig. 1. A lymph node showing completely effaced nodal structure. H.E., x40.

Fig. 2. Peripheral blood showing a cluster of relatively well-differentiated lymphocytes. May-Grünwald-Giemsa, x100.

Fig. 3. (Below the Figs. 1,2) Q-banding karyotype of a cell: 48,XY,1q+,3q+,+i(3p),-6,-10,t(14;?)(q12;?),+16,17p-,+18,+mar. The arrow indicates a 14q12 translocation. A pair No. 14 from another cell is shown for comparison in the insert.