

# *Acta Medica Okayama*

---

*Volume 63, Issue 1*

2009

*Article 2*

FEBRUARY 2009

---

## Nutrition and Depressive Symptoms in Community-dwelling Elderly Persons in Japan

Junko Oishi\*

Hiroyuki Doi<sup>†</sup>

Norito Kawakami<sup>‡</sup>

\*Department of Epidemiology, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, [junko@md.okayama-u.ac.jp](mailto:junko@md.okayama-u.ac.jp)

<sup>†</sup>Department of Epidemiology, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences,

<sup>‡</sup>Department of Mental Health, Graduate School of Medicine and Faculty of Medicine, the University of Tokyo,

Copyright ©1999 OKAYAMA UNIVERSITY MEDICAL SCHOOL. All rights reserved.

# Nutrition and Depressive Symptoms in Community-dwelling Elderly Persons in Japan

Junko Oishi, Hiroyuki Doi, and Norito Kawakami

## Abstract

A cross-sectional study was conducted to evaluate the association between dietary intake and depressive symptoms in community-dwelling elderly persons. Five-hundred elderly persons aged 65 to 75 years were randomly selected from the population of O city in Okayama Prefecture, Japan. Among 401 respondents (response rate, 80.0%), data from 279 (133 males, 146 females) who completed the questionnaire concerning dietary intake and depressive state were analyzed using logistic regression. Each of 17 dietary intakes was classified into tertiles: high intake, moderate intake, and low intake. The Center for Epidemiology Studies Depression scale (CES-D) was used. Among males, the observed odds ratios (the 95% confidence intervals) for the depressive state were 0.36 (95% CI:0.13-0.98) in the highest tertile of carotene intake, 0.33 (95% CI:0.12-0.93) in the highest tertile of vitamin C intake, 0.29 (95% CI:0.10-0.85) in the highest tertile of carbohydrate intake, and 0.33 (95% CI:0.12-0.92) in the medium tertile of vitamin E intake. Among females, similar results were observed, but these results were not statistically significant. The results suggested that carbohydrate, carotene, and vitamin C intakes are associated with lowering depressive symptoms among elderly persons dwelling in communities in Japan.

**KEYWORDS:** dietary intake, depressive symptoms, elderly persons, cross-sectional study, gender

## Original Article

**Nutrition and Depressive Symptoms in Community-dwelling Elderly Persons in Japan**Junko Oishi<sup>a\*</sup>, Hiroyuki Doi<sup>a</sup>, and Norito Kawakami<sup>b</sup><sup>a</sup>*Department of Epidemiology, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama 700-8558, Japan, and*<sup>b</sup>*Department of Mental Health, Graduate School of Medicine and Faculty of Medicine, the University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan*

A cross-sectional study was conducted to evaluate the association between dietary intake and depressive symptoms in community-dwelling elderly persons. Five-hundred elderly persons aged 65 to 75 years were randomly selected from the population of O city in Okayama Prefecture, Japan. Among 401 respondents (response rate, 80.0%), data from 279 (133 males, 146 females) who completed the questionnaire concerning dietary intake and depressive state were analyzed using logistic regression. Each of 17 dietary intakes was classified into tertiles: high intake, moderate intake, and low intake. The Center for Epidemiology Studies Depression scale (CES-D) was used. Among males, the observed odds ratios (the 95% confidence intervals) for the depressive state were 0.36 (95% CI: 0.13–0.98) in the highest tertile of carotene intake, 0.33 (95% CI: 0.12–0.93) in the highest tertile of vitamin C intake, 0.29 (95% CI: 0.10–0.85) in the highest tertile of carbohydrate intake, and 0.33 (95% CI: 0.12–0.92) in the medium tertile of vitamin E intake. Among females, similar results were observed, but these results were not statistically significant. The results suggested that carbohydrate, carotene, and vitamin C intakes are associated with lowering depressive symptoms among elderly persons dwelling in communities in Japan.

**Key words:** dietary intake, depressive symptoms, elderly persons, cross-sectional study, gender

**N**on-clinical depression and depressive disorders are common mental health problems among the elderly [1]. Depression is a strong contributor to poor physical health, lower levels of activities of daily living (ADL), poor quality of life (QOL), and suicide among older populations [2, 3]. On the other hand, dietary habits, nutrient intakes and aging processes are interrelated and are of particular importance among the elderly [4, 5]. Elderly persons are likely

to have unbalanced diets, so dietary intake and nutrition greatly influence their health [6].

Recent studies have shown the association between the intake of various nutrients and depression, including fatty acids [7–9], cholesterol [10–12], tryptophan [13], vitamin D [14, 15], vitamin C [16, 17], vitamin E [18, 19], homocysteine [20–22], folate [20–22] and vitamin B12 [2, 22]. These findings have been also replicated in the elderly to some extent. Lipids are associated with depression: Low plasma cholesterol was associated with depressive symptoms in elderly men [10], while another study reported that hyperlipidemia was associated with depression in

Received July 11, 2008; accepted September 10, 2008.

\*Corresponding author. Phone:+81-86-223-7151; Fax:+81-86-235-7178  
E-mail:junko@md.okayama-u.ac.jp (J. Oishi)

older women in a rural community [23]. The ratio of omega-3 to omega-6 fatty acids decreased among depressed elderly subjects [8], while another study failed to find the association between the dietary intake of omega-3 fatty acid and depression [9]. Vitamins are next associated with depression: Increased plasma ascorbic acid levels were associated with low levels of depression in the elderly aged 60–80 years old [16, 17]. Plasma alpha-tocopherol levels were also reported to be lower among the elderly with major depression compared with healthy controls [18], although this finding was not replicated in female elderly subjects [19]. Vitamin D deficiency [15] and low levels of folate [22] were associated with deteriorated mood in older adults. However, these findings are still inconsistent, depending on gender. In addition, there are few studies from Japan to test or replicate the association of dietary intake with depression in the elderly [23]. Nutritional intake varies among countries; particularly in Japan, people consume low fat and high salt. Such country-specific patterns of nutritional intake may affect the association between diet and depression in the elderly.

In this study, we conducted a cross-sectional study to investigate the association between dietary intake and depressive symptoms (*i. e.*, non-clinical depression) in a random sample of the elderly aged 65 or older living in a community of Japan, by estimating specific nutritional intake using a validated semi-quantitative Food Frequency Questionnaire (FFQ).

## Subjects and Methods

**Subjects.** A self-administered questionnaire which included items and questions related to dietary habits and depression was mailed to a random sample of 500 people aged from 65 to 75 years who were living in O City in Okayama Prefecture on September 1st, 2001. A small pre-test was conducted before the survey, which indicated that people aged over 75 years had a great difficulty in completing the FFQ. Thus, we restricted the sample to those aged 65–75 years. Responses were obtained from 401 people (response rate, 80.0%). After excluding respondents with missing responses to sex, age, depressive score or FFQ, data from 279 respondents (133 males and 146 females) were analyzed.

The design and procedure of the study were

reviewed and approved by the Ethics Committee on Epidemiologic Research of the Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences on February 19th 2002.

### Measures.

#### 1. Dietary intake

Dietary intake was determined using a semi-quantitative FFQ [24, 25]. The respondents were asked to report the frequency of intake of 31 food items during the year prior to the study (based on 8 categories) and the usual portion size of each item (based on 3 categories). The amount of intakes per day for 17 nutrients (total energy, protein, fat, carbohydrates, fiber, calcium, retinol, carotene, vitamin A, vitamin C, vitamin D, vitamin E, salt, cholesterol, saturated fatty acid, mono-unsaturated fatty acid, and poly-unsaturated fatty acid) were calculated based on the reported frequency of intake and portion size [24, 25]. When a respondent did not report the frequency of intake or portion size in the FFQ, the frequency of a food item was classified as “scarcely or never”, since we assumed that he/she did eat the food item and thus skipped the item. In addition, any missing data on portion size of a food item was calculated by the mean item score of that food item among other respondents. Intake of each nutrient was categorized into tertiles (high, medium, and low intake) among the entire study population.

#### 2. Depressive symptoms

Depressive symptoms were assessed using the Japanese version of the Center for Epidemiology Studies Depression scale (CES-D) [26, 27]. The CES-D consists of 20 self-reported measures of symptoms experienced in the last week and was scored on a scale of 0 to 3 points. A CES-D score of 16 was used as the cut-off value [26, 27].

#### 3. Other covariates

Other covariates examined included sex, age, chronic diseases, body mass index (BMI), and social support. Average (standard deviation, SD) age was 68.8 (2.93) for men and 68.8 (2.64) for women. Average (SD) BMI was 22.6 (2.75) for men and 23.2 (3.19) for women. Chronic disease was defined as having a present medical history of any of the following 8 chronic diseases based on self-report: hypertension, cerebrovascular disease, heart disease, kidney disease, diabetic mellitus, gout, liver disease, and cancer (prevalence of chronic conditions, 50% for men

and 40% for women). Social support was measured by using a 4-item scale asking whether the respondent “has people around who understand you”, “has people around who enjoy your happiness with you”, “has people who advise you when needed”, and “has people who listen to you when you are distressed about something”. We asked the degree of satisfaction for each item (1–4 points), then summed these item scores into a total score, and categorized the participants into 3 groups (high, medium, and low) (36%, 21%, and 43%, respectively, for men; 38%, 23%, and 39%, respectively, for women). Information about activities of daily living (ADL) was also collected by questionnaire for the daily activities of taking a bath, dressing, using the toilet, going to the toilet, going to and getting out of bed, eating, household matters, and going out. All subjects under analysis had no difficulty in pursuing any activity.

**Statistical analysis.** We compared the prevalence of depressive symptoms (16 or greater of CES-D score) among the tertiles for each nutrition intake, separately for men and women. The lowest tertile was the reference category. Moreover, we estimated the odds ratios and their 95% confidence intervals (95%

CI) of depressive symptoms for high and medium categories of each nutrition intake, with the lowest tertile as the reference category, adjusting for age, chronic diseases, BMI, and social support, using multiple logistic regression analysis. All statistical analyses were performed using the SPSS 11.0 software program.

## Results

Nutritional intake and depressive scores are presented by sex in Table 1. The intake of calcium, carotene and vitamin C was significantly higher in women than in men ( $p < 0.05$ ).

In men, there was a negative association between the intakes of carbohydrate, carotene, and vitamin C with depressive symptoms (Table 2,  $p$  for trend  $< 0.05$ ). The observed odds ratios of depressive symptoms were significantly lower in the highest tertile of carbohydrate intake, carotene intake, and vitamin C intake, compared with the lowest tertile. The odds ratio was also significantly lower in the medium tertile of vitamin E intake, while the trend was only marginally significant ( $p < 0.10$ ).

**Table 1** Nutritional intake and depressive symptoms (CES-D scores) in men and women aged 65–75 years old living in a community in Japan: Means with standard deviations (SDs)

	Men (n = 133)		Women (n = 146)		<i>p</i> -value
	Mean	SD	Mean	SD	
CES-D score	12.8	6.4	13.0	6.6	
Total energy (kcal)	2,544.0	730.9	2,530.9	561.5	0.766
Protein (g)	114.4	33.1	113.8	25.9	0.857
Fat (g)	84.2	28.9	82.1	22.5	0.501
Carbohydrates (g)	333.3	89.9	342.2	69.9	0.354
Fiber (g)	6.1	1.6	6.4	1.4	0.127
Calcium (mg)	824.8	221.9	876.0	187.9	0.038*
Retinol (mg)	1,154.0	1,114.3	1,001.6	975.5	0.224
Carotene (ug)	5,225.3	1,466.2	5,607.1	1,417.4	0.028*
Vitamin A (IU)	5,820.4	3,233.3	5,630.4	2,883.4	0.604
Vitamin C (mg)	207.9	71.4	225.1	65.6	0.037*
Vitamin D (IU)	157.8	50.0	151.9	41.4	0.286
Vitamin E (mg)	11.8	3.3	12.0	2.7	0.476
Salt (g)	17.7	4.8	17.7	3.8	0.983
Cholesterol (mg)	447.7	142.0	438.3	114.3	0.544
Saturated fatty acid (g)	17.1	5.7	17.0	4.6	0.955
Mono-unsaturated fatty acid (g)	24.8	8.4	24.3	6.7	0.579
Poly-unsaturated fatty acid (g)	18.0	5.2	17.9	4.0	0.778

\* $p < 0.05$  (Student *t* test).

**Table 2** Association between 17 nutrient intake and depressive symptoms among community-dwelling elderly men in Japan (n = 133): Prevalence, odds ratios and 95% confidence intervals for tertiles of nutrition intake #.

Nutrient intake	Depressive symptoms			Odds ratio	(95% Confidence Interval)	p value for trend
	No.	(CES-D >= 16)				
		No	%			
<b>Total Energy</b>						
Low	44	15	34	1.00		0.154
Medium	45	16	36	1.25	(0.47-3.28)	
High	44	8	18	0.44	(0.15-1.30)	
<b>Protein</b>						
Low	44	15	34	1.00		0.279
Medium	45	14	31	1.04	(0.39-2.78)	
High	44	10	23	0.56	(0.20-1.58)	
<b>Fat</b>						
Low	44	17	39	1.00		0.110
Medium	45	13	29	0.84	(0.31-2.25)	
High	44	9	21	0.42	(0.15-1.20)	
<b>Carbohydrate</b>						
Low	44	19	43	1.00		0.024*
Medium	45	12	27	0.59	(0.22-1.56)	
High	44	8	18	0.29	(0.10-0.85)*	
<b>Fiber</b>						
Low	44	18	41	1.00		0.075
Medium	45	12	27	0.69	(0.26-1.83)	
High	44	9	21	0.39	(0.14-1.10)	
<b>Calcium</b>						
Low	44	15	34	1.00		0.462
Medium	45	13	29	1.24	(0.45-3.42)	
High	44	11	25	0.67	(0.24-1.85)	
<b>Retinol</b>						
Low	44	15	34	1.00		0.782
Medium	45	10	22	0.45	(0.16-1.29)	
High	44	14	32	1.19	(0.43-3.29)	
<b>Carotene</b>						
Low	44	15	43	1.00		0.043*
Medium	45	10	22	0.52	(0.19-1.45)	
High	44	10	23	0.36	(0.13-0.98)*	
<b>Vitamin A</b>						
Low	44	15	34	1.00		0.600
Medium	45	13	29	0.67	(0.23-1.70)	
High	44	11	25	0.78	(0.28-2.17)	
<b>Vitamin C</b>						
Low	44	19	43	1.00		0.034*
Medium	45	11	24	0.55	(0.21-1.47)	
High	44	9	21	0.33	(0.12-0.93)*	
<b>Vitamin D</b>						
Low	44	13	30	1.00		0.829
Medium	45	14	31	1.27	(0.46-3.45)	
High	44	12	27	0.89	(0.32-2.46)	
<b>Vitamin E</b>						
Low	44	19	43	1.00		0.058
Medium	45	10	22	0.33	(0.12-0.92)*	
High	44	10	23	0.39	(0.14-1.08)	

Table 2 Continued from opposite page

Nutrient intake	Depressive symptoms			Odds ratio	(95% Confidence Interval)	<i>p</i> value for trend
	No.	(CES-D >= 16)				
		No	%			
<b>Salt</b>						
Low	44	16	36	1.00		0.180
Medium	45	14	31	0.81	(0.31–2.12)	
High	44	9	21	0.49	(0.17–1.39)	
<b>Cholesterol</b>						
Low	44	13	30	1.00		0.954
Medium	45	15	33	1.70	(0.60–4.85)	
High	44	11	25	0.98	(0.34–2.82)	
<b>Saturated fatty acid</b>						
Low	44	14	32	1.00		0.288
Medium	45	16	36	1.66	(0.62–4.50)	
High	44	9	21	0.54	(0.19–1.59)	
<b>Mono-unsaturated fatty acid</b>						
Low	44	16	36	1.00		0.129
Medium	45	14	31	0.93	(0.35–2.54)	
High	44	9	21	0.44	(0.15–1.25)	
<b>Poly-unsaturated fatty acid</b>						
Low	44	16	36	1.00		0.111
Medium	45	14	31	0.86	(0.32–2.31)	
High	44	9	21	0.42	(0.15–1.21)	

# Multiple logistic regression adjusting for age, chronic diseases, BMI and social support, with the “low” category as a reference.

\*  $p < 0.05$ .

In women, any nutrient intake was not significantly associated with depressive symptoms, compared with the lowest tertile. (Table 3).

## Discussion

In this study, we found that carbohydrate, carotene, and vitamin C intake were significantly associated with a reduced prevalence of depressive symptoms among men. However, among women, the intake of any nutrient under study was not significantly associated with depressive symptoms.

The observed associations of carotenoid and vitamin C, as well as the marginally significant association of vitamin E, with a reduced prevalence of depressive symptoms among elderly men are similar to those from previous studies [16–19, 29–31]. A higher consumption level of fruits and vegetable has been associated with both better mental health and better physical health [28]. A history of attempted suicide was reported to be associated with a low level of antioxidant vitamins and carotenoids [29]. A sufficient level

of vitamin A is required for adult central nervous system function [30]. Vitamin A derivatives, retinoids, are involved in a complex signaling pathway. The major functional implication of retinoic signaling that may alter the expression of components of the serotonergic neurotransmitter system has been suggested in depression [30, 31]. In addition, a previous study found a significant decrease in plasma ascorbic acid levels among patients with major depression as compared to control subjects [32]. A positive association was reported between changes in the plasma ascorbic acid level and changes in mood [16, 17]. Depressive patients have a lower serum Vitamin E concentration than controls [18]. The association between vitamin E and lower depressive symptoms was observed in elderly men, but not in elderly women [19]. Oxidative stress, an imbalance between antioxidants and prooxidants has been suggested to be involved in the pathophysiology of major depression [34]. The brain, including the cell membrane, contains a lot of lipid. Antioxidative defense by carotenoids, vitamin C and vitamin E may protect the brain

**Table 3** Association between 17 nutrient intake and depressive symptoms among community-dwelling elderly women in Japan (n = 146): Prevalence, odds ratios and 95% confidence intervals for tertiles of nutrition intake #.

Nutrient intake	Depressive symptoms			Odds ratio	(95% Confidence Interval)	p value for trend
	No.	(CES-D >= 16)				
		No	%			
<b>Total Energy</b>						
Low	48	18	38	1.00		0.439
Medium	49	14	29	0.69	(0.26-1.80)	
High	49	13	27	0.68	(0.25-1.82)	
<b>Protein</b>						
Low	48	16	33	1.00		0.831
Medium	49	15	31	1.26	(0.48-3.31)	
High	49	14	29	1.11	(0.42-2.94)	
<b>Fat</b>						
Low	48	19	40	1.00		0.476
Medium	49	13	27	0.65	(0.25-1.68)	
High	49	13	27	0.70	(0.26-1.89)	
<b>Carbohydrate</b>						
Low	48	19	40	1.00		0.114
Medium	49	15	31	0.70	(0.27-1.81)	
High	49	11	22	0.44	(0.16-1.22)	
<b>Fiber</b>						
Low	48	19	40	1.00		0.205
Medium	49	16	33	0.78	(0.31-1.98)	
High	49	10	20	0.53	(0.20-1.41)	
<b>Calcium</b>						
Low	48	16	33	1.00		0.459
Medium	49	19	39	1.42	(0.56-3.57)	
High	49	10	20	0.67	(0.24-1.83)	
<b>Retinol</b>						
Low	48	19	40	1.00		0.797
Medium	49	9	18	0.50	(0.18-1.34)	
High	49	17	35	1.14	(0.44-2.94)	
<b>Carotene</b>						
Low	48	21	44	1.00		0.172
Medium	49	13	27	0.55	(0.21-1.40)	
High	49	11	22	0.52	(0.20-1.35)	
<b>Vitamin A</b>						
Low	48	19	40	1.00		0.999
Medium	49	10	20	0.52	(0.20-1.37)	
High	49	16	33	1.00	(0.39-2.58)	
<b>Vitamin C</b>						
Low	48	20	42	1.00		0.117
Medium	49	14	29	0.55	(0.22-1.43)	
High	49	11	22	0.47	(0.18-1.22)	
<b>Vitamin D</b>						
Low	48	17	35	1.00		0.625
Medium	49	10	20	0.56	(0.21-1.53)	
High	49	18	37	1.27	(0.50-3.20)	
<b>Vitamin E</b>						
Low	48	18	38	1.00		0.568
Medium	49	14	29	0.70	(0.27-1.83)	
High	49	13	27	0.76	(0.29-1.98)	



Table 3 Continued from opposite page

Nutrient intake	Depressive symptoms			Odds ratio	(95% Confidence Interval)	<i>p</i> value for trend
	No.	(CES-D >= 16)				
		No	%			
<b>Salt</b>						
Low	48	17	35	1.00		0.704
Medium	49	15	31	0.85	(0.33–2.19)	
High	49	13	27	0.83	(0.32–2.19)	
<b>Cholesterol</b>						
Low	48	16	33	1.00		0.857
Medium	49	15	31	1.01	(0.39–2.62)	
High	49	14	29	0.92	(0.34–2.43)	
<b>Saturated fatty acid</b>						
Low	48	17	35	1.00		0.925
Medium	49	15	31	1.25	(0.49–3.20)	
High	49	13	27	0.94	(0.35–2.54)	
<b>Mono-unsaturated fatty acid</b>						
Low	48	18	38	1.00		0.591
Medium	49	14	25	0.86	(0.34–2.20)	
High	49	13	27	0.76	(0.29–2.05)	
<b>Poly-unsaturated fatty acid</b>						
Low	48	19	40	1.00		0.486
Medium	49	12	25	0.62	(0.24–1.58)	
High	49	14	29	0.71	(0.27–1.86)	

# Multiple logistic regression adjusting for age, chronic diseases, BMI and social support, with the "low" category as a reference.

\*  $p < 0.05$ .

from oxidative stress and thus may prevent depressive symptoms among the elderly. The elderly are particularly likely to suffer from undernutrition and to have a low dietary intake of antioxidants. The elderly may be more vulnerable than younger people to oxidative stress leading to neural damage including depression.

In men, carbohydrate intake was significantly associated with a reduced prevalence of depressive symptoms. Depressed subjects tend to consume more carbohydrates in their diets than non-depressed individuals [35], and they show heightened preference for sweet carbohydrate high in fat during depressed episodes [36]. A high carbohydrate intake increases the brain uptake of tryptophan, which in turn stimulates the synthesis of serotonin [37, 38]. Aged people are more sensitive to glycaemic changes [39]. Sufficient carbohydrates may be beneficial in decreasing depressive symptoms among the elderly.

In women, the intake of any nutrients under study was not significantly associated with a reduced prevalence of depression. The intake of nutrients was significantly different in men and women. However, the

patterns of the relationships were in general similar to those observed for men. The gender difference in diet intake may affect the findings. Women took in more carotenoid and vitamin C than men. The eating habits and food choices of women were healthier and more closely followed current health recommendations [40]. This may have attenuated the association of carotenoid and vitamin C with depressive symptoms in this sample of elderly women. In addition, women may take more vitamin supplements than men. In this study, we did not obtain information on the supplement intake. If women took more supplements of carotenoid or vitamin C, the association between the dietary intake of these nutrients and depression could be smaller.

In this study, the association between fatty acids and depressive symptoms was not significant. Taking DHA or EPA has been reported to decrease depressive symptoms in previous studies [7]. These fatty acids are important elements in the cell membrane. The FFQ uses in this study may be less sensitive to quantitative measurement of different fatty acid intakes. In the future, it will be necessary conduct a

study on the relationship between these fatty acids and depression among the elderly.

A major limitation of the present study is its cross-sectional nature. The observed associations between several nutrients and depressive symptoms may reflect the effects of depression on nutritional intake. For example, while they were not always significant, lower intakes of nutrients other than cholesterol were generally associated with the symptoms of depression, which may be attributable to poor appetite as a symptom of chronic depression. The findings of the present cross-sectional study should be interpreted with a caution, since they do not infer causal relationships. The possible effects of carbohydrate, carotene, and vitamin C intake on lowering depressive symptoms should be reexamined in a future prospective study and/or an intervention study.

In summary, our findings suggest that carotene and vitamin C intake may be associated with a reduction in depressive symptoms in elderly men. An interventional study is needed to confirm the effects of antioxidant nutrition and other dietary intake in the prevention of depression.

**Limitations.** The limitation due to the cross-sectional study design was mentioned earlier. Further methodological limitations of this study must also be considered. First, information on dietary intake and depressive symptoms was based on a self-administered questionnaire. People who are interested in health and food are able to recall their dietary intake more readily than others, and therefore there may have been an information bias in the questionnaire responses. Second, the study focused on elderly persons dwelling in the community. The response rate was high at 80%. Missing data, however, resulted in a decrease in the number of subjects for analysis. In a future study, a face-to-face interview, instead of a mailed questionnaire, should be used to collect detailed and correct information on dietary intake.

## References

- Shimizu H, Kawakami N, Haratani T, Iwata N and Kitamura T: The epidemiological study of the social age and the occurrence of depression of the elderly in the community: *Advances in Aging and Health Research* 1997, Foundation of Longevity Science. Higashiura-cho Aichi (1998) pp 99–108 (in Japanese).
- Rogers PJ: A healthy body a healthy mind: long-term impact of diet on mood and cognitive function. *Proc Nutr Soc* (2001) 60: 135-143.
- Conwell Y and Brent D: Suicide and aging 1: patterns of psychiatric diagnosis. *Int Psychogeriatr* (1995) 7: 149–164.
- McCormack P: Under nutrition in the elderly population living at home in the community: a review of the literature. *J Adv Nurs* (1997) 26: 856–863.
- Lesourd BM: Nutrition and immunity in the elderly: modification of immune responses with nutritional treatments. *Am J Clin Nutr* (1997) 66: 478S–484S.
- Marshall TA, Stumbo PJ, Warren JJ and Xie XJ: Inadequate nutrient intakes are common and are associated with low diet variety in rural community-dwelling elderly. *J Nutr* (2001) 131: 2192–2196.
- Stoll AL, Severus WE, Freeman MP, Rueter S, Zboyan HA, Diamon E, Cress KK and Marangell LB: Omega-3 fatty acids in bipolar disorder: a preliminary double-blind, placebo-controlled trial. *Arch Gen Psychiatry* (1999) 56: 407–412.
- Tiemeier H, van Tuijl HR, Hofman A, Kiliaan AJ and Breteler MM: Plasma fatty acid composition and depression are associated in the elderly: the Rotterdam Study. *Am J Clin Nutr* (2003) 78: 40–46.
- Hakkarainen R, Partonen T, Haukka J, Virtamo J, Albanes D and Lönnqvist J: Is low dietary intake of omega-3 fatty acids associated with depression? *Am J Psychiatry* (2004) 161: 567–569.
- Morgan RE, Palinkas LA, Barrett-Connor EL and Wingard DL: Plasma cholesterol and depressive symptoms in older men. *Lancet* (1993) 341: 75–79.
- Muldoon MF, Manuck SB and Matthews KA: Lowering cholesterol concentrations and mortality: a quantitative review of primary prevention trials. *BMJ* (1990) 310: 309–314.
- Kaplan JR, Shively CA, Fontenot MB, Morgan TM, Howell SM, Manuck SB, Muldoon MF and Mann JJ: Demonstration of an association among dietary cholesterol, central serotonergic activity, and social behaviour in monkeys. *Psychosom Med* (1994) 56: 479–484.
- Neumeister A, Praschak-Rieder N, Hasselmann B, Vitouch O, Rauh M, Barocka A and Kasper S: Effects of tryptophan depletion in fully remitted patients with seasonal affective disorder during summer. *Psychol Med* (1998) 28: 257–264.
- Lansdowne ATG and Provost SC: Vitamin D3 enhances mood in healthy subjects during winter. *Psychopharmacology (Berl)* (1998) 135: 319–323.
- Wilkins CH, Sheline YI, Roe CM, Birge SJ and Morris JC: Vitamin D deficiency is associated with low mood and worse cognitive performance in older adults. *Am J Geriatr Psychiatry* (2006) 14: 1032–1040.
- Smith A, Clark R, Nutt D, Haller J, Hayward S and Perry K: Antioxidant vitamins and mental performance of the elderly. *Human Psychopharmacology-Clinical and Experimental* (1999) 14: 459–471.
- Smith AP, Clark RE, Nutt DJ, Haller JG, Hayward S and Perry K: Vitamin C, mood and cognitive functioning in the elderly. *Nutrition Neuroscience* (2000) 2: 249–256.
- Maes M, De Vos N, Pioli R, Demedts P, Wauters A, Neels H and Christophe A: Lower serum vitamin E concentration in major depression: Another marker of lowered antioxidant defenses in that illness. *J Affect Disord* (2000) 58: 241–246.
- Tiemeier H, Hofman A, Kiliaan AJ, Meijer J and Breteler MMB: Vitamin E and depressive symptoms are not related. *The Rotterdam Study. J Affect Disord* (2002) 72: 79–83.
- Bottiglieri T, Laundry M, Crellin C, Toone BK, Carney MW and

- Reynolds EH: Homocysteine, folate, methylation and monoamine metabolism in depression. *J Neurol Neurosurg Psychiatry* (2000) 69: 228–232.
21. Huang RF, Hsu YC, Lin HL and Yang FL: Folate depletion and elevated plasma homocysteine promote oxidative stress in rat livers. *J Nutr* (2001) 131: 33–38.
  22. Folstein M, Liu T, Peter I, Buell J, Scott T and Qiu WW: The homocysteine hypothesis of depression. *Am J Psychiatry* (2007) 164: 861–867.
  23. Miyaji N, Higashi A, Ozasa K, Watanabe Y, Aoike A and Kawai K: Depression, health status and lifestyles of residents of a rural community. *Nippon Koushu Eisei Zasshi (Japanese Journal of Public Health)* (1994) 41: 452–460 (in Japanese).
  24. Takatsuka N, Kurisu Y, Nagata C, Owaki A, Kawakami N and Shimizu H: Validation of simplified diet history questionnaire. *J Epidemiol* (1997) 7: 33–41.
  25. Shimizu H, Ohwaki A, Kurisu Y, Takatsuka N, Ido M, Kawakami N, Nagata C and Inaba S: Validity and reproducibility of a quantitative food frequency questionnaire for a cohort study in Japan. *Jpn J Clin Oncol* (1999) 29: 38–44.
  26. Radloff LS: The CES-D scale. A self-report depression scale for research in the general population. *Applied Psychol Measure* (1977) 1: 385–401.
  27. Shima S, Kano T, Kitamura T and Asai M: New self-rating scales for depression. *Clin Psychiatry* (1985) 27: 717–723 (in Japanese).
  28. Myint PK, Welch AA, Bingham SA, Surtees PG, Wainwright NW, Luben RN, Wareham NJ, Smith RD, Harvey IM, Day NE and Khaw KT: Fruit and vegetable consumption and self-reported functional health in men and women in the European Prospective Investigation into Cancer-Norfolk (EPIC-Norfolk): a population-based cross-sectional study. *Public Health Nutr* (2007) 10: 34–41.
  29. Li Y and Zhang J: Serum concentrations of antioxidant vitamins and carotenoids are low in individuals with a history of attempted suicide. *Nutr Neurosci* (2007) 10: 51–58.
  30. O'Reilly K, Bailey SJ and Lane MA: Retinoid-mediated regulation of mood: possible cellular mechanisms. *Exp Bio Med (Maywood)* (2008) 233: 251–258.
  31. Tafti M and Ghyselinck NB: Functional implication of the vitamin A signaling pathway in the brain. *Arch Neurol* (2007) 64: 1706–1711.
  32. Khanzode SD, Dakhale GN, Khanzode SS, Saoji A and Palasodkar R: Oxidative damage and major depression: the potential antioxidant action of selective serotonin re-uptake inhibitors. *Redox Rep* (2003) 8: 365–370.
  33. Tabet N, Birks J and Grimley EJ: Vitamin E for Alzheimer's disease. *Cochrane Database Syst Rev* (2000) CD002854.
  34. Owen AJ, Batterham MJ, Probst YC, Grenyer BF and Tapsell LC: Low plasma vitamin E levels in major depression: diet or disease? *Eur J Clin Nutr* (2005) 59: 304–306.
  35. Christensen L and Somers S: Comparison of nutrient intake among depressed and nondepressed individuals. *Int J Eat Disord* (1996) 20: 105–109.
  36. Christensen L: The effect of food intake on mood. *Clin Nutr* (2001) 20: 161–166.
  37. Rogers PJ: Food, mood and appetite. *Nutr Res Rev* (1995) 8: 243–269.
  38. Wurtman RJ and Wurtman JJ: Brain serotonin, carbohydrate-craving, obesity and depression. *Obes Res* (1995) 3 Suppl 4: 477S–480S.
  39. Bourre JM: Effects of nutrients (in food) on the structure and function of the nervous system: update on dietary requirements for brain. Part 2: macronutrients. *J Nutr Health Aging* (2006): 386–399.
  40. Baker AH and Wardle J: Sex differences in fruit and vegetable intake in older adults. *Appetite* (2003) 40: 269–275.

