

Acta Medica Okayama

Volume 47, Issue 1

1993

Article 11

FEBRUARY 1993

Solitary pancreatic metastasis from renal cell carcinoma.

Hitoshi Takeuchi* Eiji Konaga[†] Masao harano[‡]
Kazuhiko Watanabe** Yoshiaki Takeuchi^{††}
Masahiro Hara^{‡‡} Shouhei Mano[§]

*Iwakuni National Hospital,

[†]Iwakuni National Hospital,

[‡]Iwakuni National Hospital,

**Iwakuni National Hospital,

^{††}Iwakuni National Hospital,

^{‡‡}Iwakuni National Hospital,

[§]Iwakuni National Hospital,

Solitary pancreatic metastasis from renal cell carcinoma.*

Hitoshi Takeuchi, Eiji Konaga, Masao harano, Kazuhiko Watanabe, Yoshiaki Takeuchi, Masahiro Hara, and Shouhei Mano

Abstract

A rare case of resectable solitary pancreatic metastasis from a renal cell carcinoma is reported. The patient was a 57-year-old man who presented with epigastralgia. He had undergone a radical nephrectomy of the right side 30 months previously. The diagnosis of pancreatic metastasis was based on the patient's past history and angiographic demonstration of typical hypervascular tumor staining. Histological examination was confirmatory. The patient was successfully treated by pancreaticoduodenectomy followed by alpha-interferon administration. As of 6 months after surgery, he remains well.

KEYWORDS: renal cell carcinoma, pancreatic metastasis

*PMID: 8460557 [PubMed - indexed for MEDLINE]

Copyright (C) OKAYAMA UNIVERSITY MEDICAL SCHOOL

Solitary Pancreatic Metastasis from Renal Cell Carcinoma

Hitoshi Takeuchi*, Eiji Konaga, Masao Harano, Kazuhiko Watanabe, Yoshiaki Takeuchi, Masahiro Hara^a and Shouhei Mano^b

Department of Surgery, ^aDepartment of Urology and ^bDepartment of Pathology, Iwakuni National Hospital, 2-5-1 Kuroiso-cho, Iwakuni, Yamaguchi 740, Japan

A rare case of resectable solitary pancreatic metastasis from a renal cell carcinoma is reported. The patient was a 57-year-old man who presented with epigastralgia. He had undergone a radical nephrectomy of the right side 30 months previously. The diagnosis of pancreatic metastasis was based on the patient's past history and angiographic demonstration of typical hypervascular tumor staining. Histological examination was confirmatory. The patient was successfully treated by pancreaticoduodenectomy followed by α -interferon administration. As of 6 months after surgery, he remains well.

Key words : renal cell carcinoma, pancreatic metastasis

Renal cell carcinoma (RCC) is a malignant tumor with a high metastatic potential. It frequently spreads to the lung, lymph node, liver, bone, adrenal gland, and contralateral kidney. The pancreas, however, is rarely involved, and metastatic foci in the pancreas are usually unresectable. We report a case of resectable solitary pancreatic metastasis from a right RCC, and a review of relevant literature.

The patient, aged 55 years presented with macrohematuria in November, 1989. Ultrasonography revealed a 10-cm tumor with irregular hypoechogenic internal echoes in the right kidney (Fig. 1a). Renal angiography of the right side showed a hypervascular tumor in the lower portion of the kidney. Vena cavernography demonstrated thrombosis in the vena cava arising from the right renal vein (Fig. 2a). The serum levels of the tumor markers, CEA and CA19-9, were within normal limits. Right radical nephrectomy and partial vena cavectomy were performed based on a diagnosis of renal cell carcinoma. The resected right kidney weighed 750 g and contained a yellow tumor 10-cm in diameter. Histological examination showed clear cell carcinoma (Fig. 3a).

In August, 1991, the patient was rehospitalized for epigastric pain. Ultrasonography showed a solid, 6-cm hypoechogenic mass with irregular internal echoes in the head of the pancreas. The main pancreatic duct was dilated (Fig. 1b). Celiac angiography demonstrated a hypervascular mass in the head of the pancreas (Fig. 2b). The tumor markers, CEA, CA19-9, SPAN-I, AFP and PIVKA-II, were within normal limits. There was no clinical syndrome due to hormonal hypersecretion. A metastatic pancreas tumor was suspected, rather than an islet cell tumor or acinar cell tumor on the basis of the past history and the findings of investigations. At surgery, a hard mass was found in the head of the pancreas, and pancreaticoduodenectomy was performed. The pathological specimen consisted of neoplastic cells with the same characteristic appearance as the clear cell carcinoma of the right kidney (Fig. 3b). The patient was treated with α -interferon and has been well for 6 months since the operation.

Approximately 3.7-4.5% of all malignant pancreatic tumors are metastatic and 0.9-1.7% originate from RCC (1, 2), which is an aggressive tumor that directly invades surrounding tissues and metastasizes to distant organs via the lymphatic system and blood vessels. The most com-

*To whom correspondence should be addressed.




Fig. 1 Ultrasonographs of primary (a) and metastatic (b) tumors. (a): A 10-cm tumor with irregular hypoechoic internal echoes is demonstrated in the right kidney; (b): The lesion in the head of the pancreas shows the same features as the primary lesion.




Fig. 2 Angiograms of primary (a) and metastatic (b) tumors. (a): Selective right renal angiography shows a lower pole tumor with extensive neovascularity. (b): Celiac angiography shows a hypervascular tumor in the head of the pancreas.




Fig. 3 Pathologic findings of primary (a) and metastatic (b) tumors. (a): Primary lesion in the right kidney shows a renal cell carcinoma of the clear cell type. The growth pattern is alveolar and tumor cells are uniform with clear cytoplasm and relatively uniform small nuclei. (HE \times 200) (b): Metastatic lesion in the pancreas shows the same features as the primary lesion. (HE \times 200)

Table 1 Cases of resected pancreatic metastasis from renal cell carcinoma

| No. | Age (Year) | Sex | Author | Primary site | Metastatic site | Size of metastases | Panc. Surg. (Int. after op.) | Outcome (after pancreatectomy) |
|-----|------------|-----|---------------------------------|--------------|-----------------|--------------------|------------------------------|--------------------------------|
| 1 | 50 | M | Marquand (1971) | Right | Head | 5 cm | PD (Simultaneous) | Alive (28 mos) |
| 2 | 66 | M | Guttman (1972) | Right | Head | 1.5 cm | TP (13 years) | Died (23 days) |
| 3 | 60 | M | Hermanutz (1977) | Left | Head | Fist-size | PD (14 years) | Died (19 days) |
| 4 | 43 | F | Saxon ⁽⁷⁾ (1980) | Bilateral | Head | 5 cm | PD (6 mos) | Alive (12 mos) |
| 5 | 76 | M | Yazaki (1981) | Left | Body | 1.5 cm | PD (Simultaneous) | Alive (9 mos) |
| 6 | 58 | F | Weerdenoburg (1984) | Right | Total | Multiple | TP (11 years) | Alive (10 mos) |
| 7 | 55 | M | Skaarup (1984) | Bilateral | Head and tail | Two lesions | TP (2 years) | Died (15 years) |
| 8 | 72 | M | Kishimoto (1985) | Left | Total | 15 cm | TP (Simultaneous) | Died (6 mos) |
| 9 | 66 | F | Audisio (1985) | Right | Head | 8 cm | PD (20 years) | Alive (12 mos) |
| 10 | 71 | F | Amemiya (1988) | Right | Head | 2 cm | PR (Simultaneous) | Alive (22 mos) |
| 11 | 66 | F | Hirano (1988) | Right | Body | 2.5 cm | DP (10 years) | Alive (2 mos) |
| 12 | 60 | F | Sharma (1988) | Left | Tail | 6 cm | DP (Simultaneous) | Alive (12 mos) |
| 13 | 67 | F | Carini (1988) | Bilateral | Head | 7 cm | PD (13 years) | Alive (13 mos) |
| 14 | 49 | F | Iwanami (1989) | Left | Body and tail | 4 cm 0.8 cm | DP (6 years) | Unknwn |
| 15 | 53 | F | Simpson (1989) | Left | Body and tail | 5 cm | DP (17 years) | Alive (24 mos) |
| 16 | 70 | F | Temellini ⁽⁹⁾ (1989) | Left | Head | 5 cm | PD (25 years) | Alive (12 mos) |
| 17 | 39 | M | Strijk (1989) | Right | Tail | 4 cm | DP (7 years) | Alive (12 mos) |
| 18 | 59 | M | Gohji (1990) | Bilateral | Tail | 9 cm | DP (2.5 years) | Alive (12 mos) |
| 19 | 66 | F | Yamamoto (1991) | Left | Body and tail | 3 cm 5 cm | DP (17 years) | Alive (2 mos) |
| 20 | 57 | H | Takeuchi (1992) | Right | Head | 6 cm | PD (2 years) | Alive (6 mos) |

Panc. Surg.: pancreatic surgery, Int. after op.: interval after operation for primary tumor, mos: months, PD: pancreatoduodenectomy, DP: distal pancreatectomy, PR: partial resection.

mon sites of metastasis are lung (35 %), lymph node (34 %), liver (33 %), bone (32 %), adrenal gland (19 %), contralateral kidney (11 %), brain (6 %), heart (5 %), spleen (5 %), bowel (4 %), and skin (3 %) (3). The pancreas is rarely affected: pancreatic metastases have been found in only 1.3-3 % at autopsy (4, 5). Klugo *et al.* (6) reporting on 101 patients with metastatic RCC, noted pancreatic metastases in only 2.8 %. Metastases

localized solely in the pancreas were even rarer, and metastatic foci were rarely resectable. A review of the literature through 1992 revealed only 20 cases of resectable RCC metastasis localized in the pancreas including our own case. The patients included nine men and 11 women ranging in age from 39-76 years (mean, 60.2 years). The primary site was the right kidney in nine patients, the left kidney in seven, and both kidneys in four. The metastatic

site was the head of the pancreas in nine patients, the head and tail in one, the body in two, the body and tail in three, the tail in three, and the entire pancreas in two. In 15 of the 20 cases, the pancreatic metastases were detected 6 months to 25 years after nephrectomy. The resected tumors were from 1.5–15-cm in diameter; histologically, all were of the clear cell subtype (Table 1).

This patient was fortunately treated by pancreaticoduodenectomy followed by α -interferon administration. Prognosis of metastatic renal cell carcinoma is poor. According to Saxon (7), the 5-year survival rate is 29–35 % even after successful surgical resection. Other treatments, including irradiation, chemotherapy, and hormonal therapy have been no more effective than surgical resection. The response rate to α -interferon therapy is 15–20 % (8), and tumor regression has been confirmed. Surgical resection alone or, whenever possible, in conjunction with interferon therapy confers the best chance for recovery at present.

The longest period to the detection of pancreatic metastasis is 25 years after nephrectomy (9), indicating the need for a careful long-term follow-up of patients who have undergone surgery for RCC. As shown in this case, standard serum tumor markers were not useful for the early detection of recurrence of RCC. However, several noninvasive imaging techniques, including ultrasonography, computed tomography, and magnetic resonance imaging may detect metastases in a reasonably early stage.

Ultrasonography, in particular, can be performed repeatedly at low cost and no risk.

References

1. Whittington R, Moylan DJ, Dobelbower RR and Kramer S: Pancreatic tumours in patients with previous malignancy. *Clin Radiol* (1982) **33**, 297–299.
2. Opocher E, Galeotti F, Spina GP, Battaglia G and Hernandez C: La diagnosi dei tumori secondari del pancreas: Analisi di 13 casi. *Minerva Med* (1982) **73**, 577–581.
3. Paulson DF, Perez CA and Anderson T: Cancer of the kidney and ureter; in *Cancer, Principles and Practice of Oncology*, Devita, Hellman and Resenberg eds, JB Lippincott, Philadelphia (1985) pp 895–913.
4. Willis RA: *Pathology of tumours*, Butterworths, London (1967) pp 163–190.
5. Tongio J, Peruta O, Wenger JJ and Warter P: Métastases duodénales et pancréatiques du néphroépithéliome: A propos de quatre observations. *Ann Radiol* (1977) **20**, 641–647.
6. Klugo RC, Detmers M, Stiles RE, Talley RW and Cerny JC: Aggressive versus conservative management of stage IV renal cell carcinoma. *J Urol* (1977) **118**, 244–246.
7. Saxon A, Gootesman J and Doolas A: Bilateral hypernephroma with solitary pancreatic metastasis. *J Surg Oncol* (1980) **13**, 317–322.
8. Muss HB: Interferon therapy of metastatic renal cell cancer. *Semi Surg Oncol* (1988) **4**, 199–203.
9. Temellini F, Bavosi M, Lamarra M, Quagliarini P and Giuliani F: Pancreatic metastasis 25 years after nephrectomy for renal cancer. *Tumori* (1989) **75**, 503–504.

Received April 8, 1992; accepted November 30, 1992.