

PUBLISHED VERSION

Kambor, Joachim; Maltman, Kim

[Strange quark mass from flavor breaking in hadronic t decays](#) Physical Review D, 2000; 62(9):093023

© 2000 American Physical Society

<http://link.aps.org/doi/10.1103/PhysRevD.62.093023>

PERMISSIONS

<http://publish.aps.org/authors/transfer-of-copyright-agreement>

“The author(s), and in the case of a Work Made For Hire, as defined in the U.S. Copyright Act, 17 U.S.C.

§101, the employer named [below], shall have the following rights (the “Author Rights”):

[...]

3. The right to use all or part of the Article, including the APS-prepared version without revision or modification, on the author(s)' web home page or employer's website and to make copies of all or part of the Article, including the APS-prepared version without revision or modification, for the author(s)' and/or the employer's use for educational or research purposes.”

27th March 2013

27th March 2013

<http://hdl.handle.net/2440/11178>

Strange quark mass from flavor breaking in hadronic τ decays

Joachim Kambor*

Institut für Theoretische Physik, Universität Zürich, CH-8057 Zürich, Switzerland

Kim Maltman†

*Department of Mathematics and Statistics, York University, 4700 Keele St., Toronto, Ontario, Canada M3J 1P3
and Special Research Centre for the Subatomic Structure of Matter, University of Adelaide, Adelaide, Australia 5005*

(Received 15 June 2000; published 11 October 2000)

The strange quark mass is extracted from a finite energy sum rule (FESR) analysis of the flavor-breaking difference of light-light and light-strange quark vector-plus-axial-vector correlators, using spectral functions determined from hadronic τ decay data. We point out problems for existing FESR treatments associated with potentially slow convergence of the perturbative series for the mass-dependent terms in the OPE over certain parts of the FESR contour, and show how to construct alternate weight choices which not only cure this problem, but also (1) considerably improve the convergence of the integrated perturbative series, (2) strongly suppress contributions from the region of s values where the errors on the strange current spectral function are still large and (3) essentially completely remove uncertainties associated with the subtraction of longitudinal contributions to the experimental decay distributions. The result is an extraction of m_s with statistical errors comparable to those associated with the current experimental uncertainties in the determination of the CKM angle, V_{us} . We find $m_s(1 \text{ GeV}) = 158.6 \pm 18.7 \pm 16.3 \pm 13.3 \text{ MeV}$ (where the first error is statistical, the second due to that on V_{us} , and the third theoretical).

PACS number(s): 12.15.Ff, 11.55.Hx, 12.38.-t, 13.35.Dx

I. INTRODUCTION

The light quark masses, m_s , m_u+m_d , are among the least well determined of the fundamental parameters of the standard model and, as such, have been the subject of much recent attention, in both the QCD sum rule [1–18] and lattice [19–22] communities.

Recent attempts to extract m_u+m_d and m_s via sum rule analyses of, in the former case, the light quark (ud) pseudo-scalar correlator [1] and, in the latter case, the light-strange (us) scalar [2,3,5,9] or pseudoscalar [8] correlators suffer from the problem that the relevant spectral functions are not fully determined experimentally in the region required for the analyses.

Analyses based on vector current correlators involving various pieces of the light quark electromagnetic (EM) current suffer from analogous problems. In the case of Narison's sum rule based on the difference of the flavor 33 (isovector) and 88 (hypercharge, or isoscalar) correlators [4], the G -parity-based identification of the 33 and 88 contributions to the EM hadroproduction cross section, which would allow the difference of 33 and 88 spectral functions to be determined from experimental data, is valid only in the absence of isospin breaking (IB). The high degree of cancellation (to the level of 10–15%) between the 33 and 88 spectral integrals makes the analysis rather sensitive to the neglect of IB [7]. This sensitivity is compounded by the fact that a sum rule determination of the corrections required to remove the 38 contributions from the experimental data shows that, for rea-

sons which are easily understood [7], the dominant corrections, associated with the ω contribution to the nominal 88 spectral function [7,23], are larger than one would naively expect.¹ The necessity of determining the IB corrections theoretically thus prevents one from working with a sum rule whose spectral side is determined solely by experimental data.

A similar problem exists for the sum rule based on the difference of 33 and ss vector current correlators [17], since the portion of the EM hadroproduction cross section associated with the ss part of the EM spectral function is not an experimental observable. In Ref. [17], it is assumed to be given by the cross section for the production of the various ϕ resonances. This approximation, while no doubt a reasonable one, is exactly valid only if both (1) the Zweig rule is 100% satisfied and (2) the ϕ resonances are all pure flavor $\bar{s}s$ states. The close cancellation (to the $\sim 15\%$ level) between the 33 and ss spectral integrals again makes the analysis sensitive to even small (few %) Zweig rule violations (ZRVs). To illustrate this sensitivity, let us take the deviation from ideal mixing in the vector meson sector as a measure of the natural scale of ZRVs,² and consider a scenario in which ZRVs occur dominantly in the mass matrix and not in the vacuum-to-vector-meson matrix elements of the vector cur-

¹The central value $m_s(1 \text{ GeV}) = 176 \text{ MeV}$ [17], obtained neglecting IB corrections, is reduced to 146 MeV when one applies the IB corrections obtained in the sum rule analysis of Ref. [23].

²From Ref. [27] one has that the vector meson mixing angle is either 36° or 39° , depending on whether one uses the linear or quadratic mass formula.

*Email address: kambor@physik.unizh.ch

†Email address: maltman@fewbody.phys.yorku.ca

rents. The strange (light) quark part of the EM current then couples only to the strange (light) part of any given resonance. If the flavor content of a given ϕ resonance is $\alpha\bar{s}s + \beta(\bar{u}u + \bar{d}d)/\sqrt{2}$ (with $\alpha \approx 1$ and β small), the ratio of the square of the full EM ϕ decay constant to that of the decay constant describing the coupling only to the ss part of the EM current is then $\approx 1 - \sqrt{2}\beta/\alpha$. For either the linear or quadratic versions of mixing this ratio is less than 1; including ZRV corrections will thus increase the ss spectral function and hence lower the extracted value of m_s . Taking, to be specific, the case that the radius of the circular part of the finite energy sum rule (FESR) contour is $(1.6 \text{ GeV})^2$, we find that, using an identical method of analysis and identical higher dimensional condensate values to those employed in Ref. [17] [and including, for completeness, the small IB isovector contribution to the $\phi(1020)$ EM decay constant determined in Ref. [23]], the central value of $m_s(1 \text{ GeV})$ obtained ignoring IB and ZRVs [17] (196 MeV) is lowered to 177 MeV (108 MeV) for the linear (quadratic) cases, respectively. We stress that the point of this exercise is not to attempt a realistic estimate of ZRV corrections but rather to point out that, given the scale at which such violations are already *known* to occur, the uncertainties in the extraction of m_s associated with the neglect of ZRVs are large and, moreover, cannot be significantly reduced without a major improvement in our theoretical understanding of the precise nature and magnitude of ZRVs.³

In light of the fact that, in each of the analyses above, it is not possible to work with sum rules for which the hadronic spectral function is determined entirely by experimental data, we will, in this paper, instead construct FESRs based on the flavor-breaking difference between the sum of the ud vector and axial vector correlators and the corresponding sum of us correlators, for which, up to $s = m_\tau^2$, the spectral function can be taken from experimental hadronic τ decay data [24,16]. The rest of the paper is organized as follows. In Sec. II we provide a brief review, and discuss the practical difficulties to be overcome in arriving at a reliable implementation of this approach. In Sec. III we describe a construction which leads to FESRs which successfully overcome these difficulties, and in Sec. IV we give numerical details and discuss our results.

II. FLAVOR-BREAKING SUM RULES INVOLVING HADRONIC τ DECAY DATA

For a general correlator, $\Pi(s)$, with a cut beginning at $s = s_{th}$ and running along the timelike real axis, one obtains

³In Ref. [17], the agreement of the 33–88 and 33– ss determinations of m_s obtained ignoring IB and ZRVs, respectively, was taken as evidence against the size of the IB corrections obtained in Ref. [23]. Note, however, that (1) within errors, the latter result is compatible with either the IB-corrected or -uncorrected 33–88 determination and (2) two inverse moment sum rule determinations of the 6th order chiral low-energy constant, \mathcal{Q} , one based on the 33–88 [25] and one on the $\bar{s}u$ -33 correlator difference [26], are brought into almost perfect agreement once the IB corrections of Ref. [23] are applied to the former analysis.

from Cauchy's theorem, defining the spectral function, as usual, by $\rho \equiv \text{Im } \Pi/\pi$, the general FESR relation

$$\int_{s_{th}}^{s_0} ds \rho(s) w(s) = \frac{-1}{2\pi i} \oint_{|s|=s_0} ds \Pi(s) w(s) \quad (1)$$

where $w(s)$ is any function analytic in the region of the contour, C , consisting of the union of the circle of radius s_0 in the complex s plane and the lines above and below the physical cut, running from s_{th} to s_0 .

As is well known, the ratios of ud and us inclusive hadronic τ decay widths to the τ electronic decay width,

$$R_\tau^{ij} \equiv \frac{\Gamma[\tau^- \rightarrow \nu_\tau \text{ hadrons}_{ij}(\gamma)]}{\Gamma[\tau^- \rightarrow \nu_\tau e^- \bar{\nu}_e(\gamma)]}, \quad (2)$$

where (γ) indicates additional photons or lepton pairs, and $ij = ud, us$ labels the flavors of the relevant portion of the hadronic weak current, can be expressed as weighted integrals over the relevant spectral functions. Equation (1) then allows these ratios to be recast into a form appropriate for the use of techniques based on the operator product expansion (OPE) and perturbative QCD [28–32]. Letting $J_{ij;V,A}^\mu$ be the usual vector and axial vector currents with flavor content ij , and defining the scalar $J=0,1$ parts of the corresponding correlators by

$$\begin{aligned} i \int d^4x e^{iq \cdot x} \langle 0 | T(J_{ij;V,A}^\mu(x) J_{ij;V,A}^\nu(0)^\dagger) | 0 \rangle \\ \equiv (-g^{\mu\nu} q^2 + q^\mu q^\nu) \Pi_{ij;V,A}^{(1)}(q^2) + q^\mu q^\nu \Pi_{ij;V,A}^{(0)}(q^2), \end{aligned} \quad (3)$$

one has

$$\begin{aligned} R_\tau^{ij} &= 12\pi^2 S_{EW} |V_{ij}|^2 \int_0^{m_\tau^2} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2}\right)^2 \\ &\times \left[\left(1 + 2\frac{s}{m_\tau^2}\right) \rho_{ij}^{(1)}(s) + \rho_{ij}^{(0)}(s) \right] \\ &= 6\pi S_{EW} |V_{ij}|^2 i \oint_{|s|=m_\tau^2} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2}\right)^2 \\ &\times \left[\left(1 + 2\frac{s}{m_\tau^2}\right) \Pi_{ij}^{(0+1)}(s) - 2\frac{s}{m_\tau^2} \Pi_{ij}^{(0)}(s) \right], \end{aligned} \quad (4)$$

where $\Pi_{ij}^{(J)} \equiv \Pi_{ij;V}^{(J)} + \Pi_{ij;A}^{(J)}$, $\rho_{ij}^{(J)}(s)$ are the corresponding spectral functions, $S_{EW} = 1.0194$ represents the leading electroweak corrections [33], and V_{ij} are the usual Cabibbo-Kobayashi-Maskawa (CKM) matrix elements. Since $m_\tau^2 \sim 3 \text{ GeV}^2$, the second expression in Eq. (4) is amenable to evaluation using the OPE. Dividing both the hadronic and OPE expressions by $|V_{ij}|^2$, and taking the difference of the $ij = ud$ and us cases, one arrives at a flavor-breaking FESR

$$\begin{aligned}
 & \int_0^1 dy [w_{L+T}(y)\Delta\rho^{(0+1)}(s) + w_L(y)\Delta\rho^{(0)}(s)] \\
 &= \frac{-1}{2\pi i} \oint_{|y|=1} dy [w_{L+T}(y)\Delta\Pi^{(0+1)}(s) \\
 & \quad + w_L(y)\Delta\Pi^{(0)}(s)] \quad (5)
 \end{aligned}$$

where $y \equiv s/m_\tau^2$, $\Delta\Pi^{(J)} \equiv \Pi_{ud}^{(J)} - \Pi_{us}^{(J)}$, $\Delta\rho^{(J)} \equiv \rho_{ud}^{(J)} - \rho_{us}^{(J)}$, and w_{L+T} , w_L refer to the longitudinal-plus-transverse [$(J=0)+(J=1)$, or ‘‘ $L+T$ ’']] and ‘‘longitudinal’’ [$(J=0)$] kinematic weights $w_{L+T}(y) \equiv (1-y)^2(1+2y)$ and $w_L(y) \equiv -2y(1-y)^2$, respectively. The mass-independent ($D=0$) piece of the correlator difference $\Delta\Pi^{(J)}$ on the OPE side of the sum rule, Eq. (5), of course vanishes by construction. In the limit that we neglect $m_{u,d}^2$ and $\alpha_s m_{u,d} m_s$ relative to m_s^2 , moreover, the $D=2$ terms in the OPE representation of $\Pi_{V+A;ij}^{(J)}$ become simply proportional to m_s^2 . Were the OPE representations of both the $L+T$ and longitudinal contributions above to be well converged at scale m_τ^2 , Eq. (5) would thus allow a determination of m_s in terms of the difference of experimental non-strange and strange decay number distributions.

The perturbative series for the integrated $D=2$ longitudinal contribution in Eq. (5), however, turns out not to be convergent at the scale $s_0 = m_\tau^2$ [11,12], creating a serious problem for the analysis in the absence of an experimental separation of transverse and longitudinal spectral contributions. This separation is straightforward at low s but experimentally problematic above 1 GeV^2 .⁴

Our inability to treat the OPE representation of the longitudinal contributions in a reliable manner thus creates difficult-to-quantify uncertainties for any FESR involving significant longitudinal spectral contributions. Existing analyses are included in this category since, for example, the central value for the difference of non-strange and strange spectral integrals from the analysis of Refs. [14,16],

$$\Delta^{00} \equiv \frac{R_\tau^{ud}}{|V_{ud}|^2} - \frac{R_\tau^{us}}{|V_{us}|^2} = 0.394 \pm 0.137, \quad (6)$$

corresponds to $L+T$, longitudinal and higher dimension condensate contributions which are 0.184, 0.155 and 0.055, respectively.

⁴In Ref. [11], an attempt was made to circumvent this problem by assuming the validity, even in the region of non-convergence, of a relation between the integrated longitudinal OPE vector and axial vector $D=2$ contributions valid in the region of convergence of the OPE representations of both. If true, this would allow the longitudinal strange axial integral to be obtained from the longitudinal strange vector integral. The latter can be obtained using the model strange scalar spectral function of Ref. [5]. Using appropriately-weighted FESRs for the strange pseudoscalar channel, we have now been able to test this assumption, and demonstrate that it is, in fact, incorrect.

Another practical problem is the close cancellation between the rescaled us and ud spectral integrals for the sum rules above, based on the kinematic weights, w_{L+T} and w_L . In the analysis of Refs. [14,16], for example, the cancellation is to the $\sim 10\%$ level, making the results very sensitive to both small variations in the input parameters and the sizable experimental errors ($\sim 20\text{--}30\%$) on the strange decay number distribution above the K^* region. Two features of the analysis of Refs. [14,16] illustrate the former sensitivity. First, Refs. [14,16] employ $|V_{us}| = 0.2218 \pm 0.0016$; cf. the 1998 Particle Data Group (PDG98) [27] value 0.2196 ± 0.0023 . Though compatible within errors, the squares of the two central values differ by $\sim 2\%$; use of the PDG98 value *decreases* the flavor-breaking difference, Δ^{00} , by 17%. Since one cannot reliably employ the OPE representation of the longitudinal contributions, moreover, the longitudinal spectral contribution (which is dominated, at the $\sim 80\%$ level, by the K pole term) must be subtracted; the shift in the inferred $L+T$ contribution (used to determine m_s) is thus even larger (36%). Similarly, use of the PDG98 value $f_K = 113.0 \pm 1.0 \text{ MeV}$ in place of the ALEPH determination, $f_K = 111.5 \pm 2.5 \text{ MeV}$ lowers the inferred $L+T$ contribution to Δ^{00} by a further 12%. The combined impact on the central value for m_s is thus extremely large, though the two central values are, of course, compatible within the (large) errors quoted in Refs. [14,16]. The *relative* size of the residual statistical errors as a fraction of the resulting Δ^{00} is, of course, also significantly increased by such a decrease in Δ^{00} . It is thus highly desirable to choose, in place of the kinematic weights, weights which produce a less close cancellation between the ud and us spectral integrals. The easiest way to accomplish this goal is to choose weight functions which fall off more rapidly through the region of the excited strange resonances. This has the happy consequence of also suppressing contributions from the region where both the errors on the strange spectral distribution are large and the transverse-longitudinal separation is experimentally difficult.

The final difficulty to be dealt with is theoretical. Suppose we are able to solve the longitudinal-transverse separation problem, and thus work with FESRs involving only the $L+T$ part of the flavor breaking difference:

$$\Pi(Q^2) \equiv \Pi_{ud,V+A}^{(1+0)} - \Pi_{us,V+A}^{(1+0)}. \quad (7)$$

The leading ($D=2$) m_s -dependent terms in the OPE representation of Π are [10]

$$\begin{aligned}
 [\Pi(Q^2)]_{D=2} &= -\frac{3}{2\pi} \frac{m_s^2(Q^2)}{Q^2} \\
 & \quad \times \left[1 + \frac{7}{3} a(Q^2) + (19.9332) a(Q^2)^2 + \dots \right] \\
 & \equiv -\frac{3}{2\pi} \frac{m_s^2(Q^2)}{Q^2} \sum_{k=0} g_k a(Q^2)^k, \quad (8)
 \end{aligned}$$

with $a(Q^2) = \alpha_s(Q^2)/\pi$ and $m_s(Q^2)$ the running coupling and running strange quark mass, both at scale $\mu^2 = Q^2$

TABLE I. OPE convergence of the ‘‘contour improved’’ $D=2$ contributions, $g_k A_k^{[w_{L+T}^N]}(m_\tau^2)$, as a function of the contour improved order, k , for the spectral weights, $w_{L+T}^N(y) = (1-y)^{N+2}(1+2y)$, assuming geometric growth of coefficients beyond $\mathcal{O}(\alpha_s^2)$. All entries have been rescaled by the corresponding entry for $k=0$.

Weight	$k=0$	$k=1$	$k=2$	$k=3$	$k=4$	$k=5$	$k=6$	$k=7$	$k=8$	$k=9$	$k=10$
w_{L+T}^0	1	0.143	-0.007	-0.145	-0.237	-0.286	-0.294	-0.272	-0.233	-0.187	-0.141
w_{L+T}^1	1	0.209	0.100	-0.027	-0.143	-0.232	-0.287	-0.308	-0.300	-0.272	-0.233
w_{L+T}^2	1	0.257	0.187	0.076	-0.048	-0.143	-0.260	-0.324	-0.357	-0.359	-0.339

$=-s$, in the modified minimal subtraction (\overline{MS}) scheme. The ratio of $\mathcal{O}(a)$ and $\mathcal{O}(a^2)$ coefficients in Eq. (8) is rather large (8.5), signaling potentially slow convergence [with $\alpha_s(m_\tau^2) = 0.334$ [24], the ratio of the $\mathcal{O}(a^2)$ and $\mathcal{O}(a)$ terms is 0.90 at $\mu^2 = m_\tau^2$, and >1 for μ^2 below ~ 2.2 GeV]. In recent analyses [14–16], this potential problem is brought under (apparent) control using the method of ‘‘contour improvement’’ [31]. In this method, the logarithms in Π are first summed [as has already been done in Eq. (8)] by choosing the renormalization scale equal to Q^2 at each point on the circle $|s| = s_0$. The integrals

$$A_k^{[w_{L+T}]}(s_0) = \frac{-1}{2\pi i} \oint_{|s|=s_0} ds \left[\frac{m(Q^2)^2}{Q^2} \right] a(Q^2)^k w_{L+T}(y),$$

$$y = s/s_0, \quad (9)$$

are then evaluated numerically, using the known 4-loop forms for the running mass and coupling. The OPE side of the $L+T$ part of the conventional τ decay sum rule then reduces to a linear combination of the $A_k^{[w_{L+T}]}(m_\tau^2)$, $k = 0, 1, 2$, with the index k giving the ‘‘contour-improved order.’’ Both the convergence and the residual scale dependence of the resulting truncated series are significantly improved by this procedure [12,16]. Since, relative to an expansion in terms of $a(\mu^2)$, for some fixed scale μ^2 , contour improvement represents a resummation of the perturbative series, it is possible that this improvement is physically meaningful.

Unfortunately, it turns out that the apparent improvement is not a general one, but rather the result of an accidental suppression of the $k=2$ integral. To see this, let us, for illustrative purposes, imagine that the unknown coefficients, g_k , for $k \geq 3$, in Eq. (8) grow geometrically, i.e., $g_k = (19.9332)[19.9332/(7/3)]^{k-2}$, $k \geq 3$.⁵ We then evaluate

⁵Note that Refs. [14–16] employ a form of the $L+T$ FESR in which the OPE integral has been partially integrated once in order to re-express it in terms of the difference of $L+T$ ud and us Adler functions. The contour-improved series for the Adler function version differs term by term from that based on the direct correlator difference. Though the agreement of the sums of the two versions to second order is excellent, the reader should bear in mind that the relative size of the terms of different order is not the same in the two cases.

$A_k^{[w_{L+T}]}(s_0)$ for $k=0, \dots, 10$ and $s_0 = m_\tau^2$, where $w_{L+T}^N(y) = w_{L+T}(y)[1-y]^N$, $N=0,1,2$, are the ‘‘spectral weights’’ employed in the analyses of Refs. [14–16]. The results of this exercise, rescaled in each case by the corresponding $k=0$ value, are displayed in Table I. In columns 2–4 we see the apparently favorable convergence of the $k=0,1,2$ terms already discussed. The results of the remaining columns, however, show that the smallness of the $k=2$ term is not the result of a favorable resummation (which would lead also to improved convergence for the remainder of the series) but rather a consequence of the fact that $A_k^{[w_{L+T}]}(m_\tau^2)$ has a zero as a function of k rather close to $k=2$. The magnitudes of the $k \geq 3$ terms are such that truncation of the series at $k=2$ would produce a significant theoretical error, one much larger in magnitude than the size of the $k=2$ term.⁶ The contour improved analysis employing FESRs based on the spectral weights thus has potentially significant theoretical uncertainties.

In light of the problems discussed above for those FESRs based on the spectral weights, w_{L+T}^N , our goal in the next section will be to construct alternate weights which lead to FESRs which bring these problems under control.

III. CONSTRUCTION OF ALTERNATE WEIGHT FUNCTIONS

We begin our search for an alternate choice of weight function by attempting to understand the source of the potential slow convergence of the contour-improved series noted above. The goal will be to find a weight such that, even were the unknown g_k , $k \geq 3$, to grow geometrically, as assumed above, the tail of the contour-improved series would be small relative to the known terms, in contrast to the behavior shown in Table I for the series corresponding to the spectral weights, w_{L+T}^N . If we succeed in doing so, the reliability of the standard approach, in which the truncation error is taken to be given by the size of the last known term (in

⁶One should bear in mind that, were one to work with the Adler function version of the $L+T$ FESR, the assumption of geometric growth of the coefficients of the Adler function difference is not the same as the assumption of geometric growth of the coefficients of the correlator difference itself. The potential convergence problem, however, may also be demonstrated to exist in the former case.

this case, $k=2$), will, of course, be improved regardless of the actual behavior of the unknown g_k . We will then attempt to simultaneously impose conditions which reduce the impact of the experimental errors.

To study the source of the slow convergence of the contour-improved series, it is useful to consider the behavior of the factor $f_k(Q^2) \equiv m(Q^2)^2 a(Q^2)^k g_k$, appearing in the integrand of $g_k A_k^{[w]}(s_0)$, on the contour $|s|=s_0$. Let $w(y)$, $y=s/s_0$, be any analytic function real on the real s axis, and $Q^2 = -s_0 \exp(i\varphi)$ ($\varphi=0, \pi$ thus correspond to timelike and spacelike points, respectively). One then has

$$g_k A_k^{[w]}(s_0) = \frac{1}{\pi} \int_0^\pi d\varphi \operatorname{Re}\{f_k(Q^2) w(\exp(i\varphi))\}. \quad (10)$$

The behavior of $\operatorname{Re}(f_k)$ and $\operatorname{Im}(f_k)$ as a function of φ , for $s_0 = m_\tau^2$ and $k=0, \dots, 10$, is shown in Fig. 1. We observe that both $\operatorname{Re}(f_k)$ and $\operatorname{Im}(f_k)$ have zeros on the circle $|s|=m_\tau^2$, and that these zeros move with the order k . Moreover, while $\operatorname{Re}(f_k)$ (slowly) decreases with increasing k for all angles φ , the magnitude of $\operatorname{Im}(f_k)$ is sizable in the region $\varphi \geq \pi/2$ even for $k \geq 5$. This slow convergence in the backwards (spacelike) direction is the origin of the slow convergence of the $k \geq 3$ tails of the integrated series shown in Table I, since the factor $(1-y)^{N+2}$ entering the weight w_{L+T}^N has maximum modulus at the spacelike point on the contour, and is more and more sharply peaked in the backward direction as N increases. In addition, the behavior of $\operatorname{Re}(f_2)$ and $\operatorname{Im}(f_2)$ happens to be just such that, combined with the changes of sign of the real and imaginary parts of w_{L+T}^N , there is a very strong cancellation in the integral over φ (particularly so for the case $N=0$). This strong cancellation is the origin of the ‘‘accidental’’ suppression of the magnitude of the $k=2$ term. As we have already seen in Table I, it is potentially dangerous to use weights for which the integrals $A_k^{[w]}(s_0)$ are small for a particular k (or for a small number of values of k) *only due to such cancellations*. Higher order contributions can then easily be large again, thereby spoiling the seemingly good convergence of the first few terms of the contour-improved series.

The behavior of the $\operatorname{Re}(f_k)$ and $\operatorname{Im}(f_k)$ displayed in Fig. 1 allows one not only to understand the origin of the potential convergence problem but also to construct alternate sum rules which avoid it. From Fig. 1 it is evident that convergence can be improved by avoiding weights which are large in the spacelike direction. The results of Ref. [34] also indicate that, for the FESR framework to be reliable at scales $\sim m_\tau^2$, it is necessary for the weight function to have a zero at $s=s_0$ ($y=1$).⁷

We have found two approaches useful for implementing these constraints. The first involves the use of polynomials with ‘‘shepherd’’ zeros, i.e., zeros either on, or near, the

⁷Such a zero suppresses contributions from the OPE representation in the region near the timelike real axis where, at scales $\sim m_\tau^2$ and below, data show that it breaks down [34].




FIG. 1. The real and imaginary parts of f_k , $k=0, \dots, 10$, at scale m_τ^2 , where k labels the power of α_s . Here f_k is defined explicitly in the text.

regions of the contour one wishes to suppress. The second involves the construction of weights, w_p , with $\operatorname{Im}(w_p)$ peaked on the contour at angles $\varphi \leq \pi/2$, thereby avoiding large contributions from $\operatorname{Im}(f_k)$, $k > 1$ (see Fig. 1). A convenient and effective choice is to take $\operatorname{Im}(w_p)$ to have a Gaussian form on the contour. Choosing the width of the Gaussian to be 10° and the center to be $\varphi = \varphi_p$, good convergence of the $k \geq 3$ tail of the integrated series can be obtained for any $20^\circ \leq \varphi_p \leq 90^\circ$. Technically, these profiles can be well represented using polynomials of degree $K \approx 20$:

$$w_p(y) = \sum_{i=0}^K a_i y^i. \quad (11)$$

The coefficients a_i are determined, upon normalizing $\operatorname{Im}(w_p)$ such that $w_p(0)=1$, by the Fourier integrals

TABLE II. OPE convergence of the ‘‘contour improved’’ $D=2$ contributions, $g_k A_k^{[w]}(m_\tau^2)$, as a function of the contour improved order, k , for the weights, w_{10} , \hat{w}_{10} , and w_{20} , assuming geometric growth of coefficients beyond $\mathcal{O}(\alpha_s^2)$. All entries have been rescaled by the corresponding entry for $k=0$.

Weight	$k=0$	$k=1$	$k=2$	$k=3$	$k=4$	$k=5$	$k=6$	$k=7$	$k=8$	$k=9$	$k=10$
w_{20}	1	0.262	0.213	0.143	0.073	0.018	-0.017	-0.033	-0.034	-0.027	-0.016
w_{10}	1	0.232	0.165	0.092	0.032	-0.008	-0.030	-0.038	-0.038	-0.035	-0.032
\hat{w}_{10}	1	0.248	0.193	0.125	0.064	0.019	-0.009	-0.023	-0.026	-0.024	-0.020

$$a_0 = 1, \quad a_k = \frac{2}{\pi} \int_0^\pi d\varphi \operatorname{Im}[w_p(\varphi)] \sin(k\varphi), \quad k = 1, \dots, K. \quad (12)$$

To summarize, given the problems discussed above with those FESRs involving the spectral weights, $w_{L+T}^N(y)$, we would like to find, if possible, an alternate weight choice, $w(y)$:

(1) such that $w(y)$ is strongly suppressed in the region above $s \sim 1 \text{ GeV}^2$, in order to (a) reduce the degree of cancellation between the ud and us spectral integrals, (b) reduce the impact of the large experimental errors in the us spectral distribution above the K^* region, and (c) minimize the role of the longitudinal subtraction which must, at present, be performed theoretically, and

(2) such that $w(y)$ emphasizes those regions of the contour $|s|=s_0$ for which the convergence of the $D=2$ series is favorable.

It is, of course, not *a priori* obvious that there exist $w(y)$ having the desired properties. We have, however, succeeded in constructing several polynomial weights which do.⁸ Since, as we will see below, the resulting weights do not contain $w_{L+T}(y)$ as a factor, the approach is less inclusive than the analysis employing $w_{L+T}(y)$ [12,16], but it has the advantage of being theoretically cleaner.

The strategy involving shepherd zeros can be implemented with the zeros either on or off the contour. The first weight we have constructed satisfying the criteria above has all zeros on the contour, and is given by

$$w_{10}(y) = [1-y]^4 [1+y]^2 [1+y^2] [1+y+y^2] \\ = 1 - y - y^2 + 2y^5 - y^8 - y^9 + y^{10}. \quad (13)$$

The absence of $\mathcal{O}(y^3, y^4)$ terms, which suppresses $D=8, 10$ contributions, is an additional positive feature of this weight. The fourth order zero at $y=1$ and second order zero at $y=-1$ provide the desired suppressions of the timelike and

⁸An important further restriction results from the observation that, in the FESR framework, higher dimension contributions are suppressed only by inverse powers of s_0 ; in order to avoid generating potentially large, and unknown, higher dimension contributions, therefore, the coefficients of the polynomials we construct should all be comparable in magnitude to the leading coefficient, $a_0=1$. We have chosen to implement this constraint by keeping all coefficients less than ~ 2 in magnitude.

spacelike regions. An alternate family of weights still having a fourth order zero at $y=1$, but with the remaining zeros moved off the contour and at a distance r from the origin, is

$$\hat{w}(r, \cos \theta_1, \cos \theta_2, y) \\ = [1-y]^4 \left[1 + \frac{y}{r} \right]^2 \left[1 + 2\frac{y}{r} \cos \theta_1 + \frac{y^2}{r^2} \right] \\ \times \left[1 + 2\frac{y}{r} \cos \theta_2 + \frac{y^2}{r^2} \right] \quad (14)$$

(θ_1 and θ_2 give the angular positions of the pairs of off-contour complex conjugate zeros corresponding to the last two factors, with respect to the spacelike direction). The choice $(r, \cos \theta_1, \cos \theta_2) = (1.2, 0.5, 0.1)$ produces a second solution to the constraints above, one whose biggest coefficient is $a_1 = -4/3$. We denote this solution by

$$\hat{w}_{10}(y) = \hat{w}(1.2, 0.5, 0.1, y). \quad (15)$$

In the approach based on weights which have imaginary parts with a Gaussian profile on the contour, we choose a basis of such weights having different centers, ϕ_p . As noted above, as long as all the ϕ_p lie in the interval $20^\circ \leq \phi_p \leq 90^\circ$, all of the corresponding integrated $D=2$ perturbative series will be under control. We then form linear combinations of these weights having different ϕ_p in such a way as to construct a new weight which not only retains this good convergence, but at the same time has a zero of sufficiently high order at $y=1$ to strongly suppress contributions to the spectral integral from the region $y > 0.5$. The weight of this type which most successfully satisfies the criteria discussed above has a rapid high- s falloff produced by a 6th order zero at $y=1$, a largest coefficient $a_4 = 2.087$, and is given by

$$w_{20}(y) = (1-y)^6 [1 + 4.2451y + 9.4682y^2 + 14.4155y^3 \\ + 16.4589y^4 + 14.6598y^5 + 10.2818y^6 + 5.5567y^7 \\ + 2.1157y^8 + 0.3520y^9 - 0.2065y^{10} - 0.2154y^{11} \\ - 0.1040y^{12} - 0.03040y^{13} - 0.0045y^{14}]. \quad (16)$$

The (vastly) improved convergence of the $k \geq 3$ tail of the integrated $D=2$ series for the weights w_{10} , \hat{w}_{10} and w_{20} is

displayed in Table II. The entries, as in Table I, have been rescaled by the corresponding $k=0$ value, and hence correspond to the ratios, $g_k A_k^{[w]}(m_\tau^2)/A_0^{[w]}$. The results also show that an estimate of the truncation error given by the magnitude of the $k=2$ term is, for the new weights, almost certainly a very conservative one. We will demonstrate, in the next section, that the suppression of the high- s region of the spectrum produced by the new weights is also sufficient to significantly reduce the impact of the experimental errors.

IV. NUMERICAL ANALYSIS AND RESULTS

In performing the numerical analysis of the FESRs constructed above, we employ the ALEPH data for the non-strange and strange number distributions⁹ and PDG98 values for f_K , f_π , $|V_{ud}|$ and $|V_{us}|$. As noted above, the weights have been chosen in such a way that, although theoretical input is required in order to subtract the longitudinal contributions to the experimental number distributions, and hence obtain the $L+T$ spectral functions, the effect of this subtraction on the final value of m_s is negligible. We will quantify this statement below. Once the $L+T$ spectral function has been determined, it is a straightforward matter to evaluate the weighted $L+T$ spectral integrals. The choice of steeply falling weights ensures that the strange spectral integrals are dominated by the K and K^* contributions, for which the experimental errors are much smaller than those of the rest of the strange number distribution. This plays a major role in reducing the impact of experimental errors on the final extracted value of m_s . To get a realistic determination of these errors it is important to separate correlated and uncorrelated errors and also to take into account the strong correlations between the spectral integrals involving different weights.

The nature of the longitudinal subtraction differs significantly in the low- s and high- s (~ 1 GeV) regions. For low s , the π and K pole subtractions are experimentally unambiguous. For high s (the resonance region), the longitudinal contributions are proportional to $(m_s \pm m_u)^2$, $(m_d \pm m_u)^2$, for us , ud , respectively, and hence dominated by the us contributions. The longitudinal us vector contribution is inferred from the strange scalar spectral function of Ref. [5]. This procedure is consistent provided the value of m_s resulting from the present analysis is compatible with that from the strange scalar channel [9], which it turns out to be. The longitudinal us axial vector contribution is similarly inferred from the spectral function of the strange pseudoscalar channel. The latter is obtained by fixing the excited resonance decay constants of a sum-of-resonances spectral ansatz through matching of the hadronic and OPE sides of a family of ‘pinch-weighted’ FESRs, in analogy to the analysis of

⁹The 1998 tabulation of the nonstrange data receives a small overall normalization correction as a result of the shift in R_τ^{us} between the preliminary 1998 and final 1999 analyses. We thank Shaomin Chen for bringing this point to our attention.

Ref. [35].¹⁰ The input value of m_s required for this analysis should, in principle, be determined iteratively. We have, however, employed as input the value of m_s obtained from the strange scalar analysis of Ref. [9], $m_s(1 \text{ GeV}) = 159 \pm 11$ MeV. This turns out to be consistent with our final result for m_s . Moreover, for the steeply falling weights employed in our analysis, the sum of the high- s V and A longitudinal subtractions is at the $<0.1\%$ level of the us spectral integral and, hence, at the $<1\%$ level in the $ud-us$ difference. As such, even were our evaluation to be in error by 100%, the effect on m_s would be completely negligible on the scale of the other errors present in the analysis.

On the OPE side, we retain contributions up to and including $D=8$. The leading $D=2$ term was given above.

The $D=4$ contribution is [30,10]

$$[\Pi(Q^2)]_{(D=4)} = \frac{2}{Q^4} \left[(m_l \langle \bar{l}l \rangle - I_s) \left(1 - a(Q^2) - \frac{13}{3} a(Q^2)^2 \right) + \frac{3}{7\pi^2} m_s^4(Q^2) \left(\frac{1}{a(Q^2)} - \frac{7}{12} \right) \right], \quad (17)$$

where I_s is the usual renormalization group (RG) invariant modification of the non-normal-order strange quark condensate [36], m_l is the average of the light u, d masses, and $\langle \bar{l}l \rangle$ is the light (u, d) condensate. We use the quark mass ratios determined from the chiral perturbation (ChPT) analyses of Ref. [37], the GMO relation $2m_l \langle \bar{l}l \rangle = -f_\pi^2 m_\pi^2$, and the range of values $0.7 < \langle \bar{s}s \rangle / \langle \bar{l}l \rangle < 1$ [2,3] for the ratio of condensates. The contour integration are performed as described below.

For the $D=6$ contribution we employ a rescaled version of the vacuum saturation approximation (VSA). From the results of Ref. [30], one finds

¹⁰The corresponding procedure works very well in the isovector vector channel, where the results can be checked against the well-known experimental spectral function [35]. A similar statement is true even in channels with strongly attractive interactions near threshold, for which the spectral function will be poorly represented *near threshold* by the tail of a Breit-Wigner resonance form with ‘conventional’ s -dependent width. For example, using the value of m_s obtained from the strange scalar channel analysis as input and redoing the strange scalar channel analysis, using now a sum-of-resonances spectral ansatz in place of the more realistic ansatz of Ref. [5], one finds that the ansatz of Ref. [5] is well reproduced in the region of the dominant $K_0^*(1430)$ peak. One can also use this approach to check the self-consistency between the assumed longitudinal contributions and the output m_s value in kinematic-weight-based analysis of Refs. [14,16]. It turns out that the high- s longitudinal contributions assumed are more than a factor of 2 smaller than would be expected based on the extracted value of m_s . If one employs the PDG98 values for $|V_{us}|$ and f_K , as discussed above, however, the assumed longitudinal contribution becomes compatible within the errors assigned to it in Ref. [14,16].

TABLE III. The extracted value of $m_s(1 \text{ GeV}^2)$ in MeV as a function of s_0 for the weight w_{10} having no $D=8,10$ contributions.

s_0 (GeV ²)	2.35	2.55	2.75	2.95	3.15
$m_s(1 \text{ GeV}^2)$ (MeV)	153.2	159.0	162.2	163.4	163.2

$$[\Pi(Q^2)]_{(D=6)} = \frac{64\pi\rho\alpha_s}{81Q^6} [\langle \bar{l}l \rangle^2 - \langle \bar{s}s \rangle^2], \quad (18)$$

where ρ represents a multiplicative rescaling of the VSA estimate. The analogous rescaling has been determined empirically for the isovector vector channel and the isospin-breaking vector 38 correlator, and found to be ~ 5 in both cases [38,23]. For the weights employed in our analysis, it turns out that the integrated $D=6$ contributions are very small. We are, therefore, able to employ the very conservative estimate $\rho=5 \pm 5$ for the degree of VSA violation without significantly affecting the overall theoretical error. The combination $\rho\alpha_s\langle \bar{q}q \rangle^2$ in Eq. (18) is to be understood as an effective RG-invariant combination for the evaluation of the OPE contour integrals.

Finally, for the $D=8$ contribution, we assume

$$[\Pi(Q^2)]_{(D=8)} = \frac{C_8}{Q^8}. \quad (19)$$

For w_{10} this term does not contribute to the integrated OPE; for w_{20} and \hat{w}_{10} , the value of the effective RG-invariant condensate combination, C_8 , is to be determined as part of the analysis.

As noted above, the OPE contour integration (for all D) are performed using the contour improvement prescription. Four-loop versions of the running mass and coupling are employed. To be specific, we have solved analytically for the running mass and coupling using the 4-loop truncated versions of the β [39] and γ [40] functions, with the value determined in nonstrange hadronic τ decays, $\alpha_s(m_\tau^2) = 0.334 \pm 0.022$ [24], as input. Following conventional practice, we take the error associated with the truncation of the perturbative series for the Wilson coefficient of the $D=2$ term at $\mathcal{O}(a^2)$ to be equal to the value of the last [$\mathcal{O}(a^2)$] contribution retained. In light of the discussion above we consider this to represent an extremely conservative estimate.

From the point of view of uncertainties on the OPE side, the w_{10} sum rule is favored over the \hat{w}_{10} and w_{20} sum rules for three reasons: (1) it has no $D=8,10$ contributions, (2) it has the smallest truncation error, and (3) it has the smallest errors associated with uncertainties in the input values of the $D=4$ and $D=6$ condensates.¹¹ In Table III we display, as a

¹¹Combining the errors associated with truncation, the condensate input values, and the uncertainty on $\alpha_s(m_\tau^2)$ in quadrature, the resulting errors in m_s are 7.7%, 8.2% and 8.4% for w_{10} , \hat{w}_{10} and w_{20} , respectively.

function of s_0 , the extracted values of $m_s(1 \text{ GeV}^2)$ obtained from the w_{10} sum rule, analyzed neglecting contributions of dimension 12 and higher. Central values have been used for all input on the OPE side and for the experimental spectral data. For the analysis to be self-consistent, the extracted value of m_s should be independent of s_0 . This will be true for s_0 sufficiently large that the $D \geq 12$ contributions are negligible. As s_0 is decreased, the extracted m_s values should eventually deviate from a constant, signaling the growth of the higher dimension terms. From the table we see that the range $2.75 \text{ GeV}^2 < s_0 < 3.15 \text{ GeV}^2$ provides an extremely good window of stability. In view of the falloff beginning around $s_0 \sim 2.55 \text{ GeV}^2$, we will work in the range $s_0 \geq 2.55 \text{ GeV}^2$ in the discussions which follow. It is worth stressing that the central values obtained from w_{20} and \hat{w}_{10} sum rules, though having slightly larger theoretical errors, are nonetheless completely consistent with those above: in the window $2.55 \text{ GeV}^2 \leq s_0 \leq 3.15 \text{ GeV}^2$, one finds that the range of solutions for $m_s(1 \text{ GeV}^2)$ lies between 156 and 161 MeV for w_{20} , 158 and 164 MeV for \hat{w}_{10} , and, as we saw already in Table III, 159 and 163 MeV for w_{10} . In contrast, the w_{L+T} sum rule, for which the longitudinal subtraction is important, and the $D=2$ convergence is not well under control, yields a range between 161 and 184 (with, moreover, inconsistent solutions for C_8).

From the point of view of the impact of the errors present in existing experimental data, the theoretically favored w_{10} weight is, unfortunately, no longer the favored one. The reason is that, although the impact of the errors in the high- s region of the us spectrum has been strongly suppressed by the rapid falloff of the weights employed, the $ud-us$ cancellation is still rather close (e.g., at $s_0 = m_\tau^2$, to the level of 6.0% for w_{10} , 6.8% for \hat{w}_{10} and 8.6% for w_{20} , to be compared with 3.7%, 6.5% and 9.3% for the w_{L+T}^N , $N=0,1,2$). Although the dominant errors (those from the K^* region of the us spectrum) are reasonably small, they are still large enough that the *relative* size of the residual statistical error grows very rapidly with the increase in the degree of cancellation. Thus, e.g., at $s_0 = m_\tau^2$, the statistical error represents 42%, 36%, 26%, 77%, 38% and 23% of the $ud-us$ spectral difference for the w_{10} , \hat{w}_{10} , w_{20} , w_{L+T}^0 , w_{L+T}^1 and w_{L+T}^2 sum rules, respectively.¹² The present experimental situation is, therefore, such that the errors in our final result for m_s are minimized by working with w_{20} , rather than w_{10} .

Working with the w_{20} sum rule in the window specified above we find, for our best fit,

$$m_s(1 \text{ GeV}^2) = 158.6 \pm 18.7 \pm 16.3 \pm 13.3 \text{ MeV}, \quad (20)$$

which is equivalent to

¹²Because of the high degree of cancellation, reducing s_0 , which increases the degree of suppression of the (already small) high- s us contributions, still has a non-trivial effect; e.g., the relative statistical error for the w_{20} sum rule is reduced from 26% to 19% when s_0 is lowered from m_τ^2 to 2.55 GeV^2 .




FIG. 2. The agreement between the OPE and hadronic sides of the FESR corresponding to the weight, $w_{20}(y)$ for $1.95 \text{ GeV}^2 \leq s_0 \leq m_\tau^2$. The solid line is the OPE side, using the values of m_s and C_8 obtained in the fitting procedure described in the text. The dashed line is the hadronic side, obtained using the ALEPH spectral data from which the longitudinal component has been subtracted as described in the text.

$$m_s(4 \text{ GeV}^2) = 115.1 \pm 13.6 \pm 11.8 \pm 9.7 \text{ MeV}, \quad (21)$$

where in both Eqs. (20) and (21) the first error is statistical, the second is due to the uncertainty on $|V_{us}|$, and the third theoretical. The theoretical error has been obtained by combining the following in quadrature [where we quote the numerical values corresponding to Eq. (20) to be specific]: $\pm 5.2 \text{ MeV}$, associated with the error on $\alpha_s(m_\tau^2)$; $\pm 3.6 \text{ MeV}$, associated with the uncertainty in $\langle \bar{s}s \rangle / \langle \bar{l}l \rangle$; $\pm 1.6 \text{ MeV}$, associated with the variation of m_s within the window $2.55 \text{ GeV}^2 \leq s_0 \leq m_\tau^2$; $\pm 0.6 \text{ MeV}$, associated with the uncertainty in the VSA-violating parameter, ρ ; and $\pm 11.6 \text{ MeV}$, associated with truncation of the $D=2$ series. The latter obviously remains the dominant source of theoretical error, despite the significant improvement produced by the use of the new weights. Figure 2 displays the quality of the match between

the OPE and spectral integral sides of the w_{20} sum rule corresponding to the fit above; the agreement in the previously established stability window, $s_0 > 2.55 \text{ GeV}^2$, is obviously excellent. The divergence of the OPE and spectral integral curves below $s_0 \sim 2.55 \text{ GeV}^2$ is precisely what one would expect based on the observation above that, for the w_{10} sum rule, $D > 10$ contributions, not included in the truncated OPE representation, begin to become important in this region.

The result of Eqs. (20) and (21) is in good agreement with the strange scalar channel results of Refs. [5] and [9], the strange pseudoscalar channel result of Ref. [8], and the recent hadronic τ decay analysis of Ref. [15], but, we believe, has significantly reduced theoretical and experimental errors. In particular, the statistical error has, at this point, been reduced almost to the level of that associated with the uncertainty in $|V_{us}|$.

Improvements in the accuracy of the experimental us spectral data, in particular in the K^* region, could lead to a significant improvement in the size of the statistical error. Such an improvement should be possible using BaBar data [41]. Reduced uncertainties in our knowledge of $|V_{us}|$ would also be helpful. On the theoretical side, while significant improvements in the accuracy of the spectral data would allow one to move from the w_{20} to the w_{10} sum rule, the decrease in the theoretical uncertainty that would result from this shift would be only $\sim 1.3 \text{ MeV}$. Far more likely to lead to a significant improvement in the size of the theoretical error would be a computation of the $\mathcal{O}(a^3)$ coefficient in the $D=2$ contribution to the flavor-breaking correlator difference, Π .

ACKNOWLEDGMENTS

The authors would like to thank A. Höcker and S. Chen for providing detailed information on the ALEPH nonstrange and strange spectral distributions, S. Chen for pointing out the normalization correction to the 1998 nonstrange data necessitated by the results of the 1999 strange data analysis, and G. Colangelo for his collaboration at an early stage of this work. K.M. acknowledges the ongoing support of the Natural Sciences and Engineering Research Council of Canada, and the hospitality of the Special Research Center for the Subatomic Structure of Matter at the University of Adelaide, where much of this work was performed, and J.K. the partial support of the Schweizerischer Nationalfonds and the EEC-TMR program, Contract No. CT 98-0169.

- [1] J. Bijnens, J. Prades, and E. de Rafael, Phys. Lett. B **348**, 226 (1995).
- [2] M. Jamin and M. Münz, Z. Phys. C **66**, 633 (1995); M. Jamin, Nucl. Phys. B (Proc. Suppl.) **64**, 250 (1998).
- [3] K. G. Chetyrkin, D. Pirjol, and K. Schilcher, Phys. Lett. B **404**, 337 (1997).
- [4] S. Narison, Phys. Lett. B **358**, 113 (1995).
- [5] P. Colangelo, F. De Fazio, G. Nardulli, and N. Paver, Phys. Lett. B **408**, 340 (1997).
- [6] T. Bhattacharya, R. Gupta, and K. Maltman, Phys. Rev. D **57**,

5455 (1998).

- [7] K. Maltman, Phys. Lett. B **428**, 179 (1998).
- [8] C. Dominguez, L. Pirovano, and K. Schilcher, Phys. Lett. B **425**, 193 (1998).
- [9] K. Maltman, Phys. Lett. B **462**, 195 (1999).
- [10] K. G. Chetyrkin and A. Kwiatkowski, Z. Phys. C **59**, 525 (1993); erratum hep-ph/9805232.
- [11] K. Maltman, Phys. Rev. D **58**, 093015 (1998).
- [12] A. Pich and J. Prades, J. High Energy Phys. **06**, 013 (1998).
- [13] K. G. Chetyrkin, J. H. Kuhn, and A. A. Pivovarov, Nucl. Phys.

- B533**, 473 (1998).
- [14] S. Chen, M. Davier, and A. Hocker, Report No. LAL-98-90, 1998.
- [15] A. Pich and J. Prades, *J. High Energy Phys.* **10**, 004 (1999).
- [16] ALEPH Collaboration, R. Barate *et al.*, *Eur. Phys. J. C* **11**, 599 (1999).
- [17] S. Narison, *Phys. Lett. B* **466**, 345 (1999).
- [18] S. Narison, *Nucl. Phys. B (Proc. Suppl.)* **86**, 242 (2000).
- [19] R. Gupta and T. Bhattacharya, *Phys. Rev. D* **55**, 7203 (1997); *Nucl. Phys. B (Proc. Suppl.)* **63**, 45 (1998); B. J. Gough *et al.*, *Phys. Rev. Lett.* **79**, 1622 (1997).
- [20] R. D. Kenway, *Nucl. Phys. B (Proc. Suppl.)* **73**, 16 (1999).
- [21] V. Lubicz, *Nucl. Phys. B (Proc. Suppl.)* **74**, 291 (1999).
- [22] T. Blum, A. Soni, and M. Wingate, *Phys. Rev. D* **60**, 114507 (1999); ALPHA/UKQCD Collaboration, J. Gardner, J. Heitger, R. Sommer, and H. Wettig, *Nucl. Phys.* **B571**, 237 (2000); JLQCD Collaboration, S. Aoki *et al.*, *Phys. Rev. Lett.* **82**, 4392 (1999); CP-PACS Collaboration, S. Aoki *et al.*, *ibid.* **84**, 238 (2000); W. Gökeler *et al.*, *Phys. Rev. D* **62**, 054504 (2000); D. Becirevic, V. Lubicz, G. Martinelli, and M. Testa, *Nucl. Phys. B (Proc. Suppl.)* **83**, 863 (2000); CP-PACS Collaboration, A. Ali Khan *et al.*, *ibid.* **83**, 176 (2000); hep-lat/0004010.
- [23] K. Maltman and C. E. Wolfe, *Phys. Rev. D* **59**, 096003 (1999).
- [24] ALEPH Collaboration, R. Barate *et al.*, *Z. Phys. C* **76**, 379 (1997); *Eur. Phys. J. C* **4**, 409 (1998).
- [25] E. Golowich and J. Kambor, *Phys. Rev. D* **53**, 2651 (1996).
- [26] S. Dürr and J. Kambor, *Phys. Rev. D* **61**, 114025 (2000).
- [27] Particle Data Group, C. Caso *et al.*, *Eur. Phys. J. C* **3**, 1 (1998).
- [28] Y. S. Tsai, *Phys. Rev. D* **4**, 2821 (1971); H. B. Thacker and J. J. Sakurai, *Phys. Lett.* **36B**, 103 (1971); F. J. Gilman and D. H. Miller, *Phys. Rev. D* **17**, 1846 (1978); F. J. Gilman and S. H. Rhie, *ibid.* **31**, 1066 (1985).
- [29] E. Braaten, *Phys. Rev. Lett.* **60**, 1606 (1988); S. Narison and A. Pich, *Phys. Lett. B* **211**, 183 (1988); E. Braaten, *Phys. Rev. D* **39**, 1458 (1989); S. Narison and A. Pich, *Phys. Lett. B* **304**, 359 (1993).
- [30] E. Braaten, S. Narison, and A. Pich, *Nucl. Phys.* **B373**, 581 (1992).
- [31] F. Le Diberder and A. Pich, *Phys. Lett. B* **286**, 147 (1992); **289**, 165 (1992).
- [32] A. Pich, in *Heavy Flavours II*, edited by A. J. Buras and M. Lindner (World Scientific, Singapore, 1997).
- [33] W. J. Marciano and A. Sirlin, *Phys. Rev. Lett.* **61**, 1815 (1988).
- [34] K. Maltman, *Phys. Lett. B* **440**, 367 (1998).
- [35] K. Maltman, *Phys. Lett. B* **462**, 14 (1999).
- [36] V. P. Spiridonov and K. G. Chetyrkin, *Yad. Fiz.* **47**, 818 (1988) [*Sov. J. Nucl. Phys.* **47**, 522 (1988)].
- [37] H. Leutwyler, *Phys. Lett. B* **374**, 163 (1996); **378**, 313 (1996); hep-ph/9609467.
- [38] S. Narison, *Phys. Lett. B* **361**, 121 (1995).
- [39] T. van Ritbergen, J. A. M. Vermaseren, and S. A. Larin, *Phys. Lett. B* **400**, 379 (1997).
- [40] K. G. Chetyrkin, *Phys. Lett. B* **404**, 161 (1997); T. Van Ritbergen, J. A. M. Vermaseren, and S. A. Larin, *ibid.* **405**, 327 (1997).
- [41] M. Roney (private communication).

