

# Prevention and Management of Asbestos-Related Diseases in Finland




## History of asbestos production and use in Finland

Geologically the Finnish ground is in many locations, particularly in Eastern Finland, rich in naturally occurring asbestos (anthophyllite) (Nikkarinen et al. 2001). The archaeological findings of asbestos use in Eastern Finland date back to 2500 BC. Asbestos was used for reinforcement and improvement of heat resistance of ceramic clay pots and other household items.

Industrial-scale asbestos mining was carried out in two mines in Eastern Finland. The larger one was located in Paakkila, Eastern Finland. Anthophyllite asbestos was first quarried in Paakkila by a Danish company for 1907–1910. The mining was continued in an industrial scale by Finnish companies since 1918 and it continued until 1975. The smaller Majasalmi mine was op-

erating only for 10 years 1943–1953. The industrial production of anthophyllite was totally 350 000 tons of which 120 000 tons was used in Finland (Nikkarinen et al. 2001; Palomäki and Halonen 1968). Finland also imported asbestos: chrysotile, amosite, and crocidolite. Asbestos products were manufactured at various facilities between 1923 and 1988. The main uses were in asbestos cement, insulation and roofing materials. The highest consumption of all asbestos was about 12 000 tons a year in the mid-seventies and total cumulative consumption has been estimated at the level of more than 300 000 tons, of which about 200 000 was estimated to be left in existing buildings and structures in 1989/1990. Mining in Paakkila was closed in 1975 for economic reasons, namely steeply falling consumption and the change in the owner company's production strategy towards man-made mineral fibres.



**Figure 1.** Consumption curve of asbestos in Finland in 1925–2008 and related health outcomes (Rantanen 2014a)

## National strategy & policies

### Policy

As typical in Nordic Countries, the actions for developing strategies, regulation and practical actions were carried out in collaboration between the Government (Ministry of Social Affairs and Health, the Occupational Safety and Health Administration (OSHA), and the Health Administration, MoSAH), the social partners (employers' and workers' federations), and expert institutions (mainly the Finnish Institute of Occupational Health, FIOH), social insurance institutions and NGOs, particularly the Organisation for Respiratory Health in Finland. The Finnish strategy combined soft regulation and hard law, as it was found that none of them alone was sufficient. Since Finland joined the European Union, EU regulations and their amendments concerning both the asbestos OSH regulations and the REACH regulations for chemicals have been continually transposed to Finnish regulations.


### Strategy

The Finnish national asbestos strategy was included as an element in the Health for All strategy. (HFA2000 1985) In March 1985, the Government of Finland presented the National Health for All 2000 strategy to the Parliament, which included activity line (target) No. 13: Stopping the use of hazardous chemicals (such as asbes-

tos and benzene) by 1995. In spring 1986, there was a public debate on the health risks caused by the improper dismantling of old asbestos insulation during the renovation of certain public buildings, such as schools. In April 1986, the Finnish Metal Workers' Union and the Construction Workers' Union called attention to the health problems associated with exposure to asbestos. The discussion led to actions by the Government and enhancement of activities related to asbestos by FIOH (See: Asbestos Programme).

The OSH law and related Government Decrees for the elimination of use of new asbestos and for the protection of workers' safety and health in demolition work, the law on Chemicals for banning asbestos and the tobacco law regulating tobacco smoke at workplaces represent hard legislation. The soft law used guidelines, requirements for the capacity and competence of actors dealing with asbestos, training programmes, information campaigns and numerous types of services, including health examinations of workers and advice for asbestos patients via an NGO phone service. Parallel to these preventive and diagnostic actions, social insurance provisions and practices were developed to better address the unique social security needs of workers with asbestos-related diseases (ARDs).

The strategy was shaped as follows (Figure 2):



**Figure 2.** Strategic outline for the Finnish Asbestos Programme (Rantanen 2014b)

## FIOH's National Asbestos Programme

### Background

Many industrialized countries paid increasing attention to the alleviation and prevention of health hazards caused by asbestos in the late 1970s and the 1980s. Extensive legislative, administrative and practical measures were introduced in, for example, the USA, Sweden, Norway and the then Federal Republic of Germany. The International Labour Organization (ILO) published numerous guides on safety in the use of asbestos, and in 1986 approved ILO Convention No. 162, which set technical and medical principles for the prevention of asbestos risks. Finland was one of the first countries to ratify the Convention in 1988. WHO experts have evaluated the effects of occupational and environmental exposure to asbestos in several contexts. (Huuskonen et al. 1993.) On the initiative of the International Commission on Occupational Health, ICOH, the 13th Joint ILO/WHO Committee on Occupational Health agreed upon a Joint Programme for Elimination of Asbestos-Related Diseases in 2003.

The Finnish Metal Workers' Union and the Construction Workers' Union called attention to the health problems associated with exposure to asbestos. In May 1986, the Ministry of Social Affairs and Health requested that FIOH prepare a proposal on how to tackle the problem. In October 1986, FIOH proposed the following:

1. Minimization of exposure to asbestos
2. Determination of the number of people exposed to asbestos, and identification of the asbestos-exposed individuals
3. Assessment of the health risks caused by exposure to asbestos
4. Development of diagnostics for asbestos diseases in Finland.

The National Asbestos Programme initiated by FIOH in 1987–1992 constituted a key instrument for the implementation of the National Strategies. In addition to the four tasks included in the FIOH proposal for the Ministry, the programme also contained a research section and a great deal of training supported by a wide element of awareness raising and information dissemination. A special initiative was the establishment of the Government Asbestos Committee for updating and completing regulations (Rantanen 1988; Huuskonen et al. 1994; Asbestos Committee 1989).

Objectives of the Asbestos Programme were:

- Identification of asbestos-exposed groups and screening for asbestos-related diseases
- Identification of health effects of asbestos exposure among the population
- Supporting the work of the Government Asbestos Committee in updating a regulatory basis
- Proposal to prohibit the manufacture, import, trade and use of new asbestos (see Asbestos Committee)
- Numerous actions for minimizing asbestos exposure, particularly in demolition work (see Asbestos Committee)
- Training and information for workers, employers, decision-makers and the public at large
- Research activities
- International activities.

### Research

During the Asbestos Programme, a number of scientific research projects related to asbestos diseases were conducted. The topics of the studies included the role of asbestos as an etiological factor in lung cancer and mesothelioma, diagnostic methods for asbestos-related diseases, the occurrence of pulmonary changes caused by exposure to asbestos among different population groups, and the impact of asbestos exposure on decisions to stop smoking. (Huuskonen et al. 1994; Huuskonen & Rantanen 2006; Huuskonen 2009).

Between 1990 and 1992, FIOH carried out broad-based screening of asbestos-related diseases which showed that 22% of circa 19 000 persons who had worked in housing construction and shipbuilding had pulmonary fibrosis and changes in their pleura (Koskinen et al. 1998).

The National Health Survey 'Mini-Suomi project' showed that, as a result of work-related exposure to asbestos, 6.8% of men and 2% of women in the population aged over 30 had bilateral pleural plaques. Radiological findings in the lung caused by asbestos are detectable in over 100 000 Finns. This incidence, which is very high by international standards, is a result of the extensive use of anthophyllite asbestos in building materials, which began in the 1920s (Zitting et al. 1995).

According to epidemiological studies, about 10% of the male population who smoke die of lung cancer, while the figure for insulation workers with high asbestos exposure who also smoke is about 25%. About half of those who have been highly exposed and fallen ill with asbestosis have died of lung cancer. Lung cancer incidence starts to increase 10 years after the beginning of exposure to asbestos, and is at its highest after 40 years (Meurman et al. 1994; Karjalainen 1994; Oksa 1998; Henderson et al. 2004).

The number of mesotheliomas in the general population and worker groups reflects how common exposure to asbestos has been. Even short exposures are sufficient to increase the occurrence of mesotheliomas. All types of asbestos, including anthophyllite and chrysotile have been associated with mesothelioma. Mesothelioma occurrence is increasing with increasing time since first exposure. Latency times of between 15 and 20 years have been reported, but latencies as long as 40 years have also been reported. (Tuomi 1991; Karjalainen et al. 1993; Karjalainen et al. 1994a; Karjalainen et al. 1994b; Karjalainen et al. 1996; Karjalainen et al. 1997; Takahashi et al. 1999; Tossavainen 2004; Hodgson et al. 2005; IARC 1987; IARC 2012).


In addition to well-known asbestos-related diseases, a new type of asbestos-related disease, retroperitoneal fibrosis, was described by Uibu and co-workers (2004).

### Training

Training related to asbestos dismantling work, asbestos surveys, and the diagnostics of asbestos-related diseases was organized during the Programme. In addition, numerous other training sessions and courses on asbestos were organized as part of the FIOH training programme. Asbestos as a topic was included in the training curriculum of occupational health physicians and occupational health nurses specializing at FIOH, and in the basic medical training of the University of Helsinki. During the programme, some 20 experts, including staff from the six Regional Institutes of Occupational Health, received training in the Dust Laboratory of FIOH on the microscopic asbestos analytics of material, dust, air and lung samples. OSH inspectors were specially trained to detect and advise on measures concerning the management of asbestos at the workplace, particularly in the demolition work.

### Information

One of the important aims of the Asbestos Programme was to raise awareness among the public at large. This was mainly done through the media: the press, radio and TV. Hundreds of articles to raise public awareness were published during the Programme. Numerous small guides and leaflets were produced for workplaces to increase the knowledge of all actors. One of the practical information activities was



**Figure 3.** Information as a tool for elimination of hazards (Modified from Lehtinen 2013)

the production of good practice guidelines for asbestos work and, for example, information on substitutes for asbestos materials.

In the times of the Asbestos Committee, there were about one million dwellings in detached houses and about an equal number of blocks of flats. The houses constructed between the 1920s and the 1970s were most likely constructed with materials containing asbestos. Therefore, the Committee recommended that the Ministry of Environment organize an information campaign with the Associations of dwelling building owners for the identification, labelling, and if necessary, safe removal of existing asbestos from the dwelling buildings.

### Services

Numerous types of services were organized within the frames of the Asbestos Programme. The assessment of exposure was supported by training in interview practices for occupational health physicians and occupational health nurses and for the physicians in hospitals' pulmonology clinics. A standardized method of asbestos fibre analysis was organized in FIOH's Regional Institutes of Occupational Health, the occupational hygiene experts of which provided practical advice for protective and preventive measures at workplaces.

Diagnostic services were strengthened in occupational health services, University Hospital polyclinics and in FIOH's Department of Occupational Medicine.


A national mesothelioma panel was set up at FIOH in 1989. It can be consulted in cases of difficulties in differential diagnosis. There are some 70 incident cases of mesothelioma every year. Diagnostics is managed well in Finland these days, and only the most difficult cases are sent to the panel for confirmation (Anttila et al. 2011; Huuskonen et al. 2006).

All the university hospitals in Finland have set up expert groups in pneumoconiosis diseases that can be consulted in problematical cases. The Helsinki pneumoconiosis group operates at FIOH. Every year, the asbestos fibre contents in the lung tissue of about 300 samples are examined (320 in 2005) for occupational disease diagnostics. In forensic medicine studies of cause of death carried out because of a suspicion of an occupational disease, tests for asbestos in the lung tissue are generally requested. The method has been found well standardized; the regional differences were found negligible (Nordman et al. 2006; Huuskonen et al. 2006).

## Government Asbestos Committee

The Council of State established a multisectorial Asbestos Committee in August 1988, which comprised representatives of six Ministries and State Agencies. The Asbestos Committee, chaired by the Director General of FIOH, submitted its report in December 1989, with a total of 27 recommendations for action in six different ministerial jurisdictions.

The Asbestos Committee reviewed the use and occurrence of asbestos in buildings, chemicals, consumer products and materials used at work. It also evaluated the occupational and environmental exposures to asbestos in Finland, as well as the data on asbestos-related diseases. The Committee proposed several technical, legislative, administrative and educational measures for the prevention of human exposure to asbestos. The proposed actions included a ban on manufacture, import, trade or use of asbestos-containing products such as chemicals, vehicle parts, consumer appliances and building materials. It was proposed that the use of asbestos at work be prohibited and the occupational exposure limit be lowered to the level of 0.1 fibres/cm<sup>3</sup>. More effective handling of asbestos waste was to be organized and its control strengthened. As an urgent measure, surveillance of the occurrence of asbestos in all public buildings was also proposed, as well as voluntary initiation of



measures for the identification of asbestos in private buildings and houses. These voluntary measures were to be supported by appropriate information and guidance. (Report of the Asbestos Committee 1989).

## Early recognition of diseases

Asbestos-related diseases constitute a unique type of morbidity in several aspects: Cumulative dose with poor or no elimination and clearance, long latency period between exposure and disease outcome, several types of morbidity with different shapes of dose-response, for example, asbestosis, pleural plaques and numerous types of cancers (FIOH Asbestos Programme 1993). In all these aspects, early diagnosis is of the utmost value, but simultaneously, challenging. The Finnish strategy started early with conventional (administrative) health examination schedules and proceeded to the use of biomarkers and, the screening of exposed workers with conventional chest X-rays, and gradually moved to advanced imaging technologies (HRCT and LDCT) and screening after the 1998 International Consensus Report (Kusaka et al. 2005). The sensitivity of screening tools for early detection has substantially increased.

The Finnish Strategy paid special attention to reliable exposure interview methods by the health personnel, and where possible, the assessment of exposure on the basis of measured data and the use of HRCT. Today HRCT is routinely used in occupational clinical practice (Nordman et al. 2006). Without the screening project of the Asbestos Programme for registered asbestos-exposed workers, a high number of asbestos-related cancers had remained undiagnosed or the diagnosis had been delayed. (Koskinen et al. 2003; Vehmas 2012)

It was found that the sensitivity of the HRCT test is higher than that of a conventional thorax X-ray for diagnosing diffuse interstitial pulmonary fibrosis, emphysema and changes in pleura. Low-dose spiral computed tomography is superior in sensitivity for diagnosing lung cancer (Oksa et al. 1994, Lehtola et al. 1999; Huuskonen et al. 2001, Tiitola et al. 2002a; Tiitola et al. 2002b; Kusaka et al. 2005; Huuskonen et al. 2006; Vierikko et al. 2007).

FIOH has proposed screening of lung cancer among asbestos-exposed workers (FIOH 2011). There is increasing evidence suggesting that screening of high-risk groups for lung cancer with low-dose computed tomography can reduce mortality from lung cancer and all causes

(Humphrey et al. 2013; Wender et al. 2013); however, the harms associated with screening must be balanced with the benefits. These new opportunities for early diagnosis are expected to allow early surgical and other interventions, which may improve the prognosis of the patients.

## Sharing information at international level

As asbestos-related diseases are an important issue of both European and international health, FIOH participated actively in the related international activities.

### 1997 Helsinki Criteria Meeting

The progress in Finland in tackling the asbestos situation in the 1980s and early 1990s led to a decision at FIOH to organize a meeting of international renowned experts at the Hanasaari Cultural Centre in 1997, to discuss the prevention of diseases caused by exposure to asbestos. The result of the expert meeting, now known as Helsinki Criteria, was published in the Scandinavian Journal of Work, Environment and Health (Consensus Report 1997), and in a separate publication with the background papers: Asbestos, Asbestosis, and Cancer (Work and Health, Research Reports 14).

### Asbestos Symposium for the Countries of Central and Eastern Europe

In December 1997, a regional meeting of the Countries of Central and Eastern Europe was organized, in collaboration between the Hungarian Institute of Occupational Health, FIOH, and Collegium Ramazzini. Ten Central and Eastern European countries described their situation concerning asbestos exposure and their plans for the prevention of asbestos-related occupational diseases. The country reports of Bulgaria, Estonia, Hungary, Latvia, Poland, Romania, Russia, Slovakia, Slovenia and Ukraine, keynote presentations and the recommendations of the meeting have been published as a separate report. (Work and Health, Research Reports 19. Helsinki, 1998)

### 2000

As new knowledge and diagnostic methods were developing rapidly, FIOH organized another expert meeting in February 2000, which dealt with new advances in radiology and the screening of asbestos-related diseases. The result of the expert meeting was published in the Scandinavian Journal of Work, Environment and Health (Tossavainen 2000). The papers presented and

the recommendations of the meeting were published as a separate report. (Work and Health, Research Reports 36)

## 2002

In September 2002, a Regional Asbestos Symposium for Asian Countries was organized in collaboration between the University of Occupational and Environmental Health, Kitakyushu, Japan and FIOH, and supported by ILO and WHO-WPRO. The Proceedings containing the asbestos situation reports of Singapore, Republic of Korea, China, Japan, Vietnam, the Philippines, Malaysia, Thailand, Indonesia and Taiwan, and the presentations held at the Symposium were published as a joint publication by UOEH and FIOH in the University Series: Journal of UOEH in 2002.

## 2006

In April 2006, an Asbestos Symposium for Latin American Countries was organized in Sao Paulo, Brazil, by FUNDACENTRO in collaboration with ICOH, FIOH, ILO, WHO and Mount Sinai Hospital. A few country reports (Brazil, Cuba, Ecuador and Venezuela) were published in a Chilean Journal on Occupational Health (Ciencia & Trabajo 2008).

## Demolition work

The strategy proposed by the Asbestos Committee was based on the immediate removal of asbestos in buildings and industrial facilities, in which the release of fibres and thus the exposure of workers was probable. In structures in which asbestos existed, but did not release fibres it was recommended that removal be carried out in conjunction with normal renovation projects (typically in Finland every 20 to 25 years). As well as by Government Decrees 1380/1994, 318/2006 and 863/2010 on asbestos work, the protection of workers in asbestos demolition work was supported by guidelines for good protection practice (Riala 1991; Oksa 2013). These used the information on research concerning the efficiency of respiratory protection, which was in certain cases found to be ineffective. Thus the development of safety in demolition work has been warranted (Ylioinas et al. 2012). Demolition work was estimated to produce about 5 million tons of asbestos-contaminated construction waste. Waste handling was regulated by amendment to the Waste law (1072/1993) and by drawing up a special Government Decree on the handling of asbestos waste (centralized collection of asbestos waste and special procedures for disposal).





## Current situation and further challenges

The post-hoc evaluation shows, by several indicators, that the Finnish strategy has in many respects been successful (See Summary table), though not 100% perfect. The use of new asbestos practically ceased 24 years ago, but asbestos will continue to cause health problems for decades due to the consequences of exposures in the past and from potential failures in protection during demolition work. Asbestosis cases still need follow-up throughout the whole life span of the exposed and new disease cases will still occur when the latency periods expire. Screening programmes for the early detection of both fibrosis and cancers among the exposed are planned. The protection of workers in demolition work and in waste handling needs improving. FIOH experts propose a review of safety and the efficiency of current working methods, and the development of safer techniques for the removal of existing asbestos.

Globally, the health consequences of the worldwide use of asbestos are starting to be increasingly visible in countries that have sufficiently efficient systems for diagnosis, notification, registration and compensation. The growth rates have created concern about possible over-diagnosis, while concern has also been expressed by trade unions worried about both under-diagnosis and under-reporting or delayed diagnosis. From a purely methodological perspective, we also know that the bias is more likely to be an underestimation instead of overestimation of the risk. There are several reasons for this; first, even in the best registered countries such as the Nordic ones under-reporting is substantial due to insufficient coverage of exposure assessment, and the high probability that occupational etiology is not observed by the clinicians who do the major part of cancer diagnosis. In addition, the prevention of asbestos exposures and

the elimination of asbestos-related diseases are not given equally high priority in all parts of the world. (Rantanen & Henderson 1998)

Mesothelioma is used as a marker of asbestos-related carcinogenicity. The 56 well registered countries show strong correlation between the cumulative historical asbestos consumption and the cumulative numbers of mesothelioma. On the other hand, a high number of countries with known consumption of asbestos show no or low rates of mesothelioma demonstrating a substantial (20-25%) gap in the global recognition and reporting system. (Park et al. 2011).

More than 100,000 people worldwide die each year from mesothelioma, lung cancer and asbestosis. Although the epidemic of asbestos-related disease has plateaued or is expected to plateau in the industrialized world, little is known about the epidemic in developing countries. It is obvious that increased asbestos use by these countries will result in an increase in asbestos-related diseases in future (WHO 2006; ILO 2006; Stayner et al. 2013; ICOH 2013).

Still less than one fifth of the 185 Member Countries of ILO has ratified the ILO Convention No. 162 and less than one third has decided on the ban of asbestos.

The global ban of asbestos has been declared or supported by several bodies, including Collegium Ramazzini and ICOH. In January 2014, a total of 55 out of 185 ILO Member Countries had banned asbestos. Two million tons of asbestos is still produced by five countries. Dozens of millions of tons of asbestos still prevail in the existing buildings and industrial facilities. The global epidemic of ARDs can only be managed by the global ban of asbestos and the global regulations and their effective implementation in demolition work.

## Summary

Table 1. Evolution of Finnish asbestos policies, regulations and practices

Time period	Action
~2500 BC	Use of fibres in clay pots for mechanical strength and fire resistance
1917–1975	Mining of anthophyllite in Eastern Finland
1922–1992	Use of asbestos in construction and construction materials (Top 1965–75).
1940s to 1990s	<p>Historical perspectives of asbestos exposure and disease in Finland</p> <p>Health research on asbestos miners and industrial workers, including detection of lung cancer (Noro 1945)</p> <p>Asbestosis radiology (Wegelius 1947)</p> <p>Asbestos bodies and pleural changes (Meurman 1966)</p> <p>An elevated mortality and increased lung cancer rates among anthophyllite miners (Nurminen 1972; Meurman et al. 1974, Meurman et al. 1979).</p> <p>Mesothelioma mortality (Nurminen 1975)</p> <p>Mortality of asbestosis patients (Huuskonen 1979)</p> <p>Morbidity and mortality of asbestos workers (Huuskonen 1979; Huuskonen et al. 1980)</p> <p>Mesothelioma studies (Tuomi 1991)</p> <p>Lung cancer studies (Karjalainen 1994)</p> <p>Follow-up of anthophyllite miners (Meurman et al. 1994)</p>
1975	Closure of asbestos mine for economic reasons (due to rapidly declining consumption and development of new materials because of growing information on health and environment hazards)
1979	<p>Ratification of ILO Occupational Cancer Convention No. 139 (1974), listing asbestos as registered carcinogen and registering asbestos-exposed workers</p> <p>Saalo et al. 2011. Registry of workers exposed to carcinogens in their work. FIOH: Annual Reports of the ASA Registry, 1979–</p>
1985	Health for All (HFA) National Programme for Finland (1985), Target line 13: Stopping the use of asbestos by the year 1995
1987–1992	<p>FIOH's National Asbestos Programme to support the achievement of the HFA Target, including:</p> <ul style="list-style-type: none"> <li>• Identification of asbestos-exposed people and screening of their health</li> <li>• Identification of health effects of asbestos exposure among the population</li> <li>• Initiative for establishing the Government Asbestos Committee</li> <li>• Proposal to prohibit manufacture, import, trade and use of asbestos (see Asbestos Committee)</li> <li>• Numerous actions for minimizing exposure, particularly in demolition work (see Asbestos Committee)</li> <li>• Training and information for workers, employers, decision-makers and the public at large</li> <li>• Research reports</li> <li>• Evaluation of impact: Policy &amp; regulations, consumption &amp; exposures, prevention &amp; protection, compensation for diseases, training, education, information, research knowledge (Final Report: Finnish Institute of Occupational Health 1993).</li> </ul>

1988–1989	Government Asbestos Committee: 27 proposals for asbestos management in 6 jurisdictions, including the total ban of asbestos, stringent regulation regarding demolition and waste handling work and surveys of the health of exposed workers (Government Committee Reports 1989:66)
1987	Government Decree 886/1987 on asbestos work.
1987–1988	Ratification of ILO Asbestos Convention No. 162 (1986)
1989	Decision by the National Board of Labour Protection 205/1989 on health examinations of workers in asbestos work
1989	Establishment of a National Mesothelioma Panel
1992–1993	Government Decree 852/1992 on the prohibition of manufacture, import, trade and use of all main types of asbestos, including chrysotile. Waste Law 1072/1993
1994	Government Decree 1380/1994 on asbestos work for the prevention and minimizing of asbestos exposure and health hazards at work, including employers' responsibilities for recognition of asbestos, registration, prevention and protection, training and education, licensing, health surveillance and occupational exposure limit, OEL, to 0.25 f/cm <sup>3</sup>
1997–2005	<p>International Expert Meeting: Asbestos, Asbestosis, and Cancer in Espoo, Finland 20–22 January 1997, publishing a Consensus Report: Asbestos, Asbestosis, and Cancer: the Helsinki criteria for diagnosis and attribution Scand J Work Environ Health 1997;23:311–6 (Meeting Report: Asbestos, Asbestosis, and Cancer. Work and Health. Research Reports 14. Finnish Institute of Occupational Health, Helsinki 1997. 98p.</p> <p>New Advances in Radiology and Screening of Asbestos-Related Diseases. Scandinavian Journal of Work, Environment and Health 2000;26(5):449–54. Work and Health. Research Reports 36. Finnish Institute of Occupational Health, Helsinki 2000. 89 p.</p> <p>FIOH contribution to the World Health Organization (1998) "Chrysotile Asbestos" and to the World Trade Organization (2001) ruled on the ongoing dispute concerning the banning of asbestos.</p> <p>Finnish contribution to international HRCT classification: Y. Kusaka, K.G. Hering, J.E. Parker (Eds.). International Classification of HRCT for Occupational and Environmental Respiratory Diseases (Springer 2005).</p>
1997–2006	<p>Regional Asbestos Meetings:</p> <p>Proceedings of the Asbestos Symposium for the Countries of Central and Eastern Europe. Work and Health, Research Reports 19. Finnish Institute of Occupational Health, Helsinki 1998. 112p.</p> <p>Proceedings of the Asbestos Symposium for Asian Countries. FIOH, UOEH 2002;24(Suppl. 2):1–120.</p> <p>Asbestos Symposium for Latin American Countries, 25–27 April 2006, Fundacentro, Sao Paulo, Brazil. Ciencia &amp; Trabajo 2008;27(10). <a href="http://www.cienciaytrabajo.cl">www.cienciaytrabajo.cl</a></p>
2004	Government Decree 975/2004, on the prohibition of manufacture, import, trade or use of products (with a few exceptions), containing any type of asbestos, including chrysotile


2006–2010	Government Decrees 318/2006 and 863/2010 on asbestos work. Setting OEL to 0.1f/cm <sup>3</sup> and requiring proven competence from supervisors and workers in asbestos (demolition) work
2007–2010	Guidelines for diagnostics of asbestos-related diseases (Huuskonen 2005; Nordman et al. 2006; Huuskonen et al. 2010). Research on asbestos exposures in demolition work & related guidelines
2006–2009	Guidelines and articles on exposure and diagnostics of ARDs for occupational health services (Riala 1991; Nordman et al. 2006; Huuskonen et al. 2006; Huuskonen et al. 2009; Oksa 2013)
2009	EU Commission Decree No. 552/2009 for the Harmonization of national regulations on chemicals with EU Regulation 1907/2006 called REACH Regulation containing prohibition of asbestos use
2006–2012	Research on increased risk in demolition work in spite of careful protection (Ylioinas et al. 2012) Guidelines for protection of workers in construction work (including demolition of asbestos) (FIOH data package 2011, Ekman 2011)
2000–2013	<p>About 80% reduction in the number of registered asbestos-exposed workers from the year 2001 to 2011 and about 50% reduction in the number of registered asbestosis cases from the highest values in the mid-90s. The numbers of lung cancer and mesothelioma show an almost flat trend.</p> <p>The Ministry of Social Affairs and Health Working Group on Occupational Cancer (2013): Compensation criteria for asbestos diseases in principle unchanged, but clarified. Compensation for mesothelioma even in low exposure, lung cancer and larynx cancer in heavy exposure (25 fibre-years), compensation for ovarian cancer in heavy exposure and individual judgement.</p>
2014	International Conference on Monitoring and Surveillance of Asbestos-Related Diseases, 10–13 February 2014, Espoo, Finland

## Literature

- Ammattisyöpätyöryhmän muistio 2013. Työterveyslaitos, Helsinki 2013. (Report of the Working Group on Occupational Cancer).
- Anttila S, Wolff H, Kettunen E, Böhling T, Knuutila A. Mesoteliomadiagnostiikan osuvuus ja ammattisyöpän toteamisen käytännöt Suomessa. Loppuraportti 31.8.2011. Työterveyslaitos, Helsinki 2011.
- Asbestikomitean mietintö. (Report of the Asbestos Committee). Komiteamietintö 1989:66. Ministry of Labour, Helsinki 1989. 76 p. + 9 Annexes.
- Asbestos, Asbestosis, and Cancer. Proceedings of an International Expert Meeting. Work and Health. Research Reports 14. Finnish Institute of Occupational Health, Helsinki 1997. 79 p.
- Ciencia & Trabajo 2008;10(27). [www.cienciaytrabajo.cl](http://www.cienciaytrabajo.cl)
- Consensus Report. Asbestos, asbestosis, and cancer: the Helsinki criteria for diagnosis and attribution. Scand J Work Environment Health 1997;23(4):311–6.
- Ekman A. Toimiva asbestipurku. Työturvallisuuskeskus, Helsinki 2011.
- FIOH: Annual Reports of the ASA Registry, 1979– . Finnish Institute of Occupational Health, Helsinki.
- FIOH: Finnish Institute of Occupational Health proposes screening for lung cancer among asbestos-exposed workers. Press release 73/2011. 08.12.2011, Helsinki.
- FIOH: Työterveyslaitos. Asbestituotteet (updated 1.12.2011). [www.ttl.fi/fi/toimialat/rakennus/turvapakki/vaaralliset\\_aineet/eristeaineet/asbestituotteet/Sivut/default.aspx](http://www.ttl.fi/fi/toimialat/rakennus/turvapakki/vaaralliset_aineet/eristeaineet/asbestituotteet/Sivut/default.aspx)
- Government Decree 886/1987 on asbestos work.
- Government Decree 975/2004, on the prohibition of manufacture, import, trade or use of products (with a few exceptions), containing any type of asbestos, including chrysotile
- Health for All 2000 Strategy. Sosiaali- ja terveystieteiden ministeriö. (Ministry of Social Affairs and Health). Helsinki 1985.
- Henderson DW, Rodelsperger K, Woitowitz HJ, Leigh J. After Helsinki: a multidisciplinary review of the relationship between asbestos exposure and lung cancer, with emphasis on studies published during 1997–2004. Pathology 2004;36(6):517–50.
- Hodgson JT, McElvenny DM, Darnton AJ, Price MJ, Peto J. The expected burden of mesothelioma mortality in Great Britain from 2002–2050. British Journal of Cancer 2005;92:587–92.
- Humphrey LL, Deffebach M, Pappas M, Baumann C, Artis K, Mitchell JP, Zakher B, Fu R, Slatore CG. Screening for lung cancer with low-dose computed tomography: a systematic review to update the U.S. Preventive services task force recommendation. Ann Intern Med 2013;Sept 17, 159(6):411–20.
- Huuskonen MS. Asbestosis in Finland: clinical and immunological findings and mortality among asbestosis patients. Doctoral thesis. University of Helsinki, Finland 1979, 69p.
- Huuskonen MS. Asbestos-related diseases. In: Evidence-Based Medicine Guidelines. Kunnamo I, editor in chief. Helsinki, The Finnish Medical Society DUODECIM Medical Publications Ltd and John Wiley & Sons, Ltd. 2005:226–8.
- Huuskonen MS. Chapter 3: Prevention and Deletion of Asbestos Diseases in Finland. In: Soto A and Salazar G. Asbestosis: Risks, Environment and Impact. Nova Biomedical Books. New York. Nova Science Publishers Inc. 2009:57–70.
- Huuskonen MS, Ahlman K, Tossavainen A. Asbestos disease in Finland. J Occ Med 1980;22(11):751–7.
- Huuskonen MS, Jahkola A, Oksa P. Asbestisairaudet. Julkaisussa. Kunnamo I (päätoim.) Lääkärin käsikirja (on line) Helsinki, Kustannus Oy Duodecim 7.6.2010.
- Huuskonen M, Karjalainen A, Koskinen K, Rinne J-P, Tossavainen A, Rantanen J. Asbestos Program 1987–92. Final Report. 44 p. + Appendices. Institute of Occupational Health, Helsinki 1993.
- Huuskonen M, Karjalainen A, Koskinen K, Rinne J-P, Tossavainen A, Rantanen J. Asbestos Program 1987–92. Final Report. 40 p. + Appendices. Institute of Occupational Health, Helsinki 1994. Second Edition.
- Huuskonen MS, Oksa P, Vehmas T, Anttila S, Tossavainen A. Asbestisairaudet Suomessa. Suom Lääkäril 2006;39:3961–6.
- Huuskonen MS, Rantanen J. Finnish Institute of Occupational Health (FIOH): Prevention and Detection of Asbestos-Related Diseases 1987–2005. Am J Ind Med 2006;49(3):215–20.
- Huuskonen O, Kivisaari L, Zitting A, Taskinen K, Tossavainen A, Vehmas T. High-resolution computed tomography classification of lung fibrosis for patients with asbestos-related disease. Scand J Work Environ Health 2001;27(2):106–12.
- IARC. Asbestos. IARC Monograph on Evaluation of Carcinogenic Risks to Humans. Overall Evaluations of Carcinogenicity: An Updating of IARC Monographs, Volumes 1–42. Supplement 7, IARC, Lyon 1987.
- IARC. A Review of Human Carcinogens volume 100 C. Arsenic, metals, fibres, and dusts. Asbestos (chrysotile, amosite, crocidolite, tremolite, actinolite, and anthophyllite). 2012:219–303. <http://monographs.iarc.fr/ENG/Monographs/vol100C/mono100C.pdf>
- International Commission on Occupational Health. ICOH Statement on Global Asbestos Ban and the Elimination of Asbestos-related Diseases 2013. [www.ichweb.org](http://www.ichweb.org)
- International Labour Organization. Resolution concerning asbestos, 2006.

- Karjalainen A. Occupational asbestos exposure, pulmonary fiber burden and lung cancer in the Finnish population. Helsinki: Finnish Institute of Occupational Health, 1994.
- Karjalainen A, Anttila S, Heikkilä L, Kyyronen P, Vainio H. Lobe of origin of lung cancer among asbestos-exposed patients with or without diffuse interstitial fibrosis. *Scand J Work Environ Health* 1993;19(2):102-7.
- Karjalainen A, Anttila S, Vanhala E, Vainio H. Asbestos exposure and the risk of lung cancer in a general urban population. *Scand J Work Environ Health* 1994a;20(4):243-50.
- Karjalainen A, Meurman L, Pukkala E. Four cases of mesothelioma in miners. *Occup Environ Med* 1994b;51:212-5.
- Karjalainen A, Nurminen M, Vanhala E, Vainio H, Anttila S. Pulmonary asbestos bodies and asbestos fibers as indicators of exposure. *Scand J Work Environ Health* 1996;22(1):34-8.
- Karjalainen A, Pukkala E, Mattson K, Tammilehto L, Vainio H. Trends in mesothelioma incidence and occupational mesotheliomas in Finland in 1960-1995. *Scand J Work Environ Health* 1997;23(4):266-70.
- Koskinen K, Zitting A, Tossavainen A, Rinne JP, Roto P, Kivekäs J, Reijula K, Huuskonen MS. Radiographic abnormalities among Finnish construction, shipyard and asbestos industry workers. *Scand J Work Environ Health* 1998;24(2):109-17.
- Koskinen K, Pukkala E, Reijula K, Karjalainen A. Incidence of cancer among the participants of the Finnish Asbestos Screening Campaign. *Scand J Work Environ Health* 2003;29(1):64-70
- Kusaka Y, Hering KG, Parker JE (Eds.). *International Classification of HRCT for Occupational and Environmental Respiratory Diseases*. Tokyo: Springer-Verlag 2005.
- Lehtinen S. Information as a tool for elimination of hazards and management of ARDs. 2013. Modified in 2014 by Rantanen and Lehtinen.
- Lehtola H, Huuskonen MS, Zitting A, Kivisaari L, Vehmas T, Mattson K, Koskinen K, Reijula K, Tossavainen A, Kaleva S, Rantanen J. Asbestisairauksien varhaisdiagnostiikka ammattitautiseurantatutkimusten yhteydessä 1996-99. Loppuraportti Työsuojelurahastolle. FIOH, Helsinki 1999. 19 p. (in Finnish)
- Meurman L. Asbestos bodies and pleural plaques in a Finnish series of autopsy cases. *Acta Radiologica and Microbiologica Scandinavica* 1966;Suppl. 181:1-107.
- Meurman LO, Kiviluoto R, Hakama M. Mortality and morbidity among the working population of anthophyllite asbestos miners in Finland. *Br J Ind Med* 1974;31:105-12.
- Meurman LO, Kiviluoto R, Hakama M. Combined effect of asbestos exposure and tobacco smoking on Finnish anthophyllite miners and millers. *Ann NY Acad Sci* 1979;330:491-5.
- Meurman L, Pukkala E, Hakama M. Incidence of cancer among anthophyllite asbestos miners in Finland. *Occup Environ Med* 1994;51:421-5.
- National Board of Labour Protection 205/1989. Decision by the National Board of Labour Protection on health examinations of workers in asbestos work.
- Nikkarinen M, Aatos S, Teräsvuori E. Asbestin esiintyminen ja sen vaikutus ympäristöön Tuusniemellä, Outokummussa, Kaavilla ja Heinävedellä. Geologian tutkimuskeskus, Tutkimusraportti — Geological Survey of Finland, Report of Investigation 152, 2001.
- Nordman H, Oksa P, Karjalainen A, Koskinen H. Asbestisairauksien diagnostiikka ja seuranta. Helsinki: Työterveyslaitos, Työ ja ihminen -tutkimusraportti 28, 2006;7-15.
- Noro L. Ammattitaudit ja niiden ehkäisy. (Prevention of occupational diseases). Helsinki 1945. 198 p.
- Nurminen M. A study of the mortality of workers in an anthophyllite asbestos factory in Finland. *Work-Environment-Health* 1972;9:112-8.
- Nurminen M. The epidemiologic relationship between pleural mesothelioma and asbestos exposure. *Scandinavian Journal of Work Environment and Health* 1975;1:128-37.
- Oksa P. Asbestosis, its detection and predictors of progression and cancer. *People and Work Research Reports* 17. Helsinki: Finnish Institute of Occupational Health 1998.
- Oksa P. Asbestialistuminen ja asbestisairaudet. Asbesti-iltapäivä. Työterveyslaitos 31.10.2013, Helsinki.
- Oksa P, Suoranta H, Koskinen H, Zitting A, Nordman H. High resolution computed tomography in the early detection of asbestosis. *Occup Environ Health* 1994;65:299-304.
- Palomäki A, Halonen O. 1968. Paakkilan antofylliittiasbestilouhos. *Vuoriteollisuus* 1968;26(2):92-8.
- Park E-K, Takahashi K, Hoshuyama T, Cheng T-J, Delgermaa V, Le GV, Sorahan T. Global Magnitude of Reported and Unreported Mesothelioma. *Environ Health Perspect* 2011;119:514-8. <http://dx.doi.org/10.1289/ehp.1002845> [online 06 January 2011]
- Proceedings of the Asbestos Symposium for the Asian Countries. FIOH, UOEH 2002;24(Suppl. 2):1-120.
- Proceedings of the Asbestos Symposium for the Countries of Central and Eastern Europe. *Work and Health, Research Reports* 19. Finnish Institute of Occupational Health, Helsinki 1998. 112p.
- Rantanen J. Analysis of asbestos consumption and asbestos-related occupational diseases. 2014a.
- Rantanen J. Strategic outline for the Finnish Asbestos Programme. 2014b.
- Rantanen J. Preventive strategies for occupational cancer in Finland. In: *Primary Prevention of Cancer* edited by WJ Eylenbosch, N Can Larebeke and AM Depoerter. Raven Press, Ltd. New York, 1988:313-35.

- Rantanen J, Hendersson DW. The global asbestos epidemic – need for the diagnostic criteria for out-comes. In: *Advances in the prevention of Occupational Respiratory Diseases. Proceedings of the 9th International Conference on Occupational Respiratory Diseases*, Kyoto, Japan 13–16 October 1997. Chiyotani K, Hosoda Y, Aizawa Y, editors. Amsterdam, Lausanne, New York, Oxford, Shannon, Singapore, Tokyo. Elsevier, Excerpta Medica International Congress Series 1153. 1998:259–65.
- Regulation (ec) no 1907/2006 of the European Parliament and of the Council of 18 December 2006 concerning the Registration, Evaluation, Authorization and Restriction of Chemicals (REACH), establishing a European Chemicals Agency, amending Directive 1999/45/EC and repealing Council Regulation (EEC) No 793/93 and Commission Regulation (EC) No 1488/94 as well as Council Directive 76/769/EEC and Commission Directives 91/155/EEC, 93/67/EEC, 93/105/EC and 2000/21/EC. Official Journal of the European Union L 396/11
- Riala R. Altisteet työssä, nro 5. Asbesti. FIOH, Work Environment Fund. Helsinki 1991. 43 p.
- Saalo A, Soosaar A, Länsimäki E, Kauppinen T. ASA 2009 Syöpäsairauden vaaraa aiheuttaville aineille ja menetelmille ammatissaan altistuneiksi ilmoitetut Suomessa. Helsinki: Työterveyslaitos 2011;39.
- Stayner L, Welch LS, Lemen R. The worldwide Pandemic of Asbestos-Related Diseases. *Annu Rev Public Health* 2013;34:4.1–4.12.
- Takahashi K, Huuskonen MS, Tossavainen A, Higashi T, Okubo T, Rantanen J. Ecological relationship between mesothelioma incidence/mortality and asbestos consumption in ten western countries and Japan. *Journal of Occupational Health* 1999;41:8–11.
- Tiitola M, Kivisaari L, Huuskonen MS, Mattson K, Koskinen H, Lehtola H, Zitting A, Vehmas T. Computed tomography screening for lung cancer in asbestos-exposed workers. *Lung Cancer* 2002a;35(1):17–22.
- Tiitola M, Kivisaari L, Zitting A, Kaleva S, Tossavainen A, Huuskonen MS, Vehmas T. Computed tomography of asbestos-related pleural abnormalities. *Int Arch Occup Environ Health* 2002b;75:224–8.
- Tossavainen A. Consensus Report. International expert meeting on new advances in the radiology and screening of asbestos-related diseases. *Scand J Work Environ Health* 2000;26(5):449–54.
- Tossavainen A. Global use of asbestos and the incidence of mesothelioma. *Int J Occup Environ Health* 2004;10:22–5.
- Tossavainen A, Lehtinen S, Huuskonen M, Rantanen J (eds.). *Proceedings of an International Expert Meeting on New Advances in Radiology and Screening of Asbestos-Related Diseases*. 9–11 February 2000, Espoo Finland. Finnish Institute of Occupational Health, Helsinki, Finland. People and Work. Research Reports 36. Helsinki 2000. 89 p.
- Tuomi T. Asbestos exposure of Finnish mesothelioma patients. Electron microscopic analysis of mineral fibres in the lung tissue and bronchoalveolar lavage fluid. Helsinki: Kuopion yliopisto, Työterveyslaitos 1991.
- Työsuojeluhallituksen päätös hyväksyttävistä asbestipurkutyössä käytettävistä menetelmistä ja laitteista 14.12.1989/231, muutossäädös 10.2.1992/176.
- Uibu T, Oksa P, Auvinen A, Honkanen E, Saha H, Uitti J, Roto, P. Asbestos exposure as a risk factor for retroperitoneal fibrosis. *Lancet* 2004;363:1422–6.
- Vehmas T. CT screening for asbestos-exposed workers. Expertise from Finland. DGUV Dresden 26 Apr 2012.
- VNP 1380/1994 Valtioneuvoston päätös asbestityöstä. Annettu Helsingissä 21 päivänä joulukuuta 1994. Amendments 27.4.2006/318 ja 30.9.2010/863.
- Vierikko T, Järvenpää R, Autti T, Oksa P, Huuskonen MS, Kaleva S, Laurikka J, Kajander S, Paakkola, K, Saarelainen S, Salomaa ER, Tukiainen P, Uitti J, Vanhanen M, Vehmas T. Chest CT screening of workers with asbestos exposure: lung tumors, benign nodules and incidental findings. *European Respiratory* 2007;27:78–84.
- Waste Law 1072/1993.
- Wegelius C, Changes in the lungs in 126 cases of asbestosis observed in Finland. *Acta Radiologica* 1947;28:139–52.
- Wender R et al. American Cancer Society Lung Cancer Screening Guidelines. *Ca Cancer J Clin* 2013;63:106–17. <http://onlinelibrary.wiley.com/doi/10.3322/caac.21172/pdf>.
- World Health Organization. Elimination of asbestos-related diseases. WHO/SDE/OEH/06.03. September 2006.
- Ylioinas P, Kropsu P, Oksa P. Asbestisairaudet eivät ole loppumassa – asbestipurkajapotilaan tapaus. *Suomen lääkärilehti* 2012;67(8):601–5. (No end in sight for asbestos-related illnesses – the case of an asbestos abatement worker).
- Zitting A, Karjalainen A, Impivaara O, Tossavainen A, Kuusela T, Mäki J, Huuskonen MS. Radiographic small lung opacities and pleural abnormalities as a consequence of asbestos exposure in an adult population. *Scand J Work Environ Health* 1995;21(6):470–7.
- This leaflet was written by  
 Jorma Rantanen  
 Suvi Lehtinen  
 Matti Huuskonen  
 Panu Oksa  
 Antti Tossavainen  
 Timo Tuomi  
 Harri Vainio
- Editing support by  
 Mirkka Salmensaari
- 3.2.2014  
 Cover photo: Antophyllite 5000x by Jaakko Sääntti



**ISBN 978-952-261-396-7 (nid.)**  
**ISBN 978-952-261-397-4 (pdf)**

**Unigrafia**  
**Helsinki 2014**