

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Publications from USDA-ARS / UNL Faculty

U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska

3-2017

DEVELOPMENT AND FITNESS OF *Spodoptera cosmioides* (WALKER) (LEPIDOPTERA: NOCTUIDAE) ON *Phaseolus vulgaris* L. GENOTYPES

M. M. Freitas
Universidade Estadual Paulista, freitasmm@hotmail.com

C. A. Freitas
Universidade Estadual Paulista, carloscaf77@gmail.com

L. Nogueira
Universidade Estadual Paulista

P. H. S. Barcelos
Universidade Estadual Paulista

A. L. Boiça Júnior
Universidade Estadual Paulista, aboicajr@fcav.unesp.br

Follow this and additional works at: <https://digitalcommons.unl.edu/usdaarsfacpub>

Freitas, M. M.; Freitas, C. A.; Nogueira, L.; Barcelos, P. H. S.; and Boiça Júnior, A. L., "DEVELOPMENT AND FITNESS OF *Spodoptera cosmioides* (WALKER) (LEPIDOPTERA: NOCTUIDAE) ON *Phaseolus vulgaris* L. GENOTYPES" (2017). *Publications from USDA-ARS / UNL Faculty*. 1747.
<https://digitalcommons.unl.edu/usdaarsfacpub/1747>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

DEVELOPMENT AND FITNESS OF *Spodoptera cosmioides* (WALKER) (LEPIDOPTERA: NOCTUIDAE) ON *Phaseolus vulgaris* L. GENOTYPES

M. M. Freitas¹, C. A. Freitas¹, L. Nogueira¹, P. H. S. Barcelos¹, A. L. Boiça Júnior¹

¹Universidade Estadual Paulista, Faculdade de Ciências Agrárias e Veterinárias
(UNESP/FCAV), Departamento de Fitossanidade, Laboratório de Resistência de Plantas à
Insetos, Via de Acesso Prof. Paulo Donato Castellane s/nº, CEP 14884-900, Jaboticabal, SP,
Brazil E-mail: freitasmm@hotmail.com

INTRODUCTION

The black armyworm *Spodoptera cosmioides* (Walker, 1858) (Lepidoptera: Noctuidae), is widely distributed in South America, except for southern Argentina, Chile, and regions located to the west of the Andes in Peru (SILVAIN, LALANNE -CASSOU, 1997). In Brazil, this species has occurred in several crops, including soybean, cotton and bean, causing injuries on leaves and pods that result in substantial losses in the plant yield (SPECHT et al., 2004).

Thus, Host-Plant Resistance becomes an alternative method as a strategy in the control of this pest, since resistance features expressed by some plants can cause changes in the behavior, and / or biology of phytophagous insects, or provide greater support capacity to their attack (BOIÇA JÚNIOR et al., 2013). Therefore, the aim of this study was to evaluate the development and fitness of *S. cosmioides* larvae fed on bean genotypes.

MATERIAL E METHODS

Bean genotypes used were BRS Pérola, IAC Harmonia, BRS Supremo, BRS Talisman and IPR Campos Gerais. The leaves of the genotypes (treatments) were individualized into 9-cm-diameter Petri dishes lined with deionized water moistened filter paper. In each plate, a newly hatched larvae of *S. cosmioides* was released. Experiment was carried out under a completely randomized design with 30 replications. Data on larval stage duration, larval survival and larval weight of *S. cosmioides* were recorded. In addition, the larval performance of *S. cosmioides* was observed by calculating the Fitness Index (FI), adapted from Boregas et al. (2013), using the formula: FI = larval survival (%) × pupal biomass (mg) / larval development period (days) / 10.

Larval development data were analyzed for residuals normality and variance homogeneity, and when necessary, were transformed to meet the assumptions of analysis of variance (ANOVA). Then, data were submitted to ANOVA (unidirectional ANOVA), and when significant, the means of treatments were compared by Tukey's test ($\alpha = 0.05$).

RESULTS AND DISCUSSION

There were significant differences on survival and larval weight at 10 days, and fitness index of *S. cosmioides* among bean genotypes evaluated. The highest survival was observed in larvae fed on the genotypes BRS Supremo, BRS Pérola and IPR Campos Gerais, in relation to the genotype BRS Talismã. There was total mortality of the larvae that fed on IAC Harmonia genotypes (Figure 1). Similar results obtained by Morando (2015) and Santos (2015) showed that the IAC Harmonia genotype presented resistance features to *Chrysodeixis includens* and *Spodoptera frugiperda* (Lepidoptera: Noctuidae), respectively.




Figure 1. Survival curve (\pm SE) of *Spodoptera cosmioides* larvae fed with bean genotypes. Different lowercase letters indicate significant difference by the Tukey test ($P < 0.0001$; $F = 18,32$; $SE = 9,16$). *Data not analyzed due to total insect mortality.




Figure 2. Average larval weight (\pm SE) of *Spodoptera cosmioides* fed with bean genotypes at 10 and 15 days. Bars topped with different lowercase letters indicate significant difference by the Tukey test ($P = 0.0006$; $F = 5,58$; $SE = 5,28$). ns: No significant difference by the Tukey test ($P = 0.8210$; $F = 0,38$; $SE = 54,23$).




Figure 3. Fitness index of *Spodoptera cosmioides* larvae fed with bean genotypes.

Regarding larval weight (Figure 2), there were significant differences only at 10 days after larvae hatching. Genotypes BRS Supremo, BRS Pérola and IPR Campos Gerais, provided greater *S. cosmioides* weight in relation to genotypes IAC Harmonia and BRS Talisman. The fitness index that demonstrates the host adequacy level for the evaluated population, was considered low (<1.0) for *S. cosmioides* in all bean genotypes used in this experiment (Figure 3). The results found herein demonstrate that bean genotypes evaluated show resistance features in the antibiosis category and provided low fitness level to *S. cosmioides*.

REFERENCES

- BOIÇA JÚNIOR, A. L.; SOUZA, B. H. S.; LOPES, G. S.; COSTA, E. N.; MORAES, R. F. O.; EDUARDO, W. I. Atualidades em resistência de plantas a insetos. In: BUSOLI, A. C.; ALENCAR, J. R. D. C. C.; FRAGA, D. F.; SOUZA, L. A.; SOUZA, B. H. S.; GRIGOLLI, J. F. J. (Eds.). Tópicos em entomologia agrícola – VI. Jaboticabal: Gráfica Multipress, 2013b. p. 207-224.
- BOREGAS, K.G.B.; MENDES, S.M.; WAQUIL, J.M.; FERNANDES, G.A.W. Estádio de adaptação de *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) em hospedeiros alternativos. Bragantia, v.72, p.61-70, 2013.
- MORANDO, R. Resistência de genótipos de feijoeiro a *Chrysodeixis Includens* (Walker) (Lepidoptera: Noctuidae). 2015. 81 f. Dissertação (Mestrado em Agronomia – Área de Concentração: Proteção de Plantas), Faculdade de Ciências Agronômicas, Universidade Estadual Paulista, Botucatu, 2015.
- SANTOS, J. D. V. Não preferência para alimentação e antibiose em genótipos de feijoeiro a *Spodoptera frugiperda* (J. E. Smith, 1797) (Lepidoptera: Noctuidae). 2015. 84 f. Dissertação (Mestrado em Agronomia – Área de Concentração: Entomologia Agrícola) – Faculdade de Ciências Agrárias e Veterinárias, Universidade Estadual Paulista, Jaboticabal, 2015.
- SILVAIN, J. F.; LALANNE-CASSOU, B. Distinction entre *Spodoptera latifascia* (Walk.) et *Spodoptera cosmioides* (Walk.), bona species (Lepidoptera: Noctuidae). Revue Française d'Entomologie, v. 19, p. 95-97, 1997.
- SPECHT, A.; SILVA, E.; LINK, D. Noctuideos (Lepidoptera, Noctuidae) do Museu Entomológico Ceslau Biezanko, Departamento de Fitossanidade, Faculdade de Agronomia "Eliseu Maciel", Universidade Federal de Pelotas, RS. Revista Brasileira de Agrociências, v. 4, p. 389-409, 2004.