

1942

Results of the Cooperative Uniform Soybean Tests

Albert H. Probst

Bureau of Plant Industry: Division of forage Crops and Diseases

Denver I. Allen

U.S. Bureau of Plant Industry: Division of forage Crops and Diseases

Charles R. Weber

U.S. Bureau of Plant Industry: Division of forage Crops and Diseases

Leonard F. Williams

U.S. Bureau of Plant Industry: Division of forage Crops and Diseases

Jackson L. Cartter

U.S. Bureau of Plant Industry: Division of forage Crops and Diseases

Follow this and additional works at: <https://docs.lib.purdue.edu/ars>

Recommended Citation

Probst, Albert H.; Allen, Denver I.; Weber, Charles R.; Williams, Leonard F.; and Cartter, Jackson L., "Results of the Cooperative Uniform Soybean Tests" (1942). *Uniform Soybean Tests Northern Region*. Paper 4.
<https://docs.lib.purdue.edu/ars/4>

**Agronomy Reading Room
S-506A Turner Hall
University of Illinois**

N. L. Burlison

UNITED STATES DEPARTMENT OF AGRICULTURE
AGRICULTURAL RESEARCH ADMINISTRATION

BUREAU OF PLANT INDUSTRY,
SOILS, AND AGRICULTURAL ENGINEERING
cooperating with the
STATE AGRICULTURAL EXPERIMENT STATIONS
of the
NORTH CENTRAL REGION

RESULTS OF THE COOPERATIVE UNIFORM
SOYBEAN TESTS

1942

U. S. REGIONAL SOYBEAN LABORATORY
Urbana, Illinois
February, 1943

RESULTS OF THE COOPERATIVE UNIFORM SOYBEAN TESTS

1942

by

Albert H. Probst, Denver I. Allen, Charles R. Weber,
Leonard F. Williams, and Jackson L. Cartter 1/

CONTENTS

Introduction.....	1
Cooperation.....	2
Location of Uniform Tests.....	3
Map of North Central Region.....	4
Methods.....	5
Uniform Test, Group I.....	6
Uniform Test, Group II.....	14
Uniform Test, Group III.....	30
Uniform Test, Group IV.....	45

INTRODUCTION

One of the main objectives of the U. S. Regional Soybean Laboratory 2/ is to develop improved varieties and strains of soybeans for commercial and industrial utilization. In order that new strains developed through the cooperative breeding work could be evaluated more rapidly and accurately, four uniform tests were established and designated as Uniform Test Groups I, II, III, and IV.

Group I was designed to include varieties for the northernmost parts of the North Central states and contained material of approximately the maturity of Mandarin. Likewise, Group II was designed to include varieties adapted to the northern parts of the "soybean belt" immediately south of the region for Group I, and contained material of approximately the maturity of Richland. Group III was established to include varieties adapted to the central "soybean belt" area, namely, southern Ohio, central

1/ Assistant Agronomist, Agent, Agent, Associate Agronomist, and Agronomist, respectively, Division of Forage Crops and Diseases, Bureau of Plant Industry, Soils and Agricultural Engineering, Agricultural Research Administration, United States Department of Agriculture.

2/ A cooperative organization participated in by the Bureau of Plant Industry, Soils, and Agricultural Engineering of the U. S. Department of Agriculture and the Agricultural Experiment Stations of the North Central states of Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin.

Indiana, central Illinois, southern Iowa, and northern Missouri, and contained material of approximately the maturity of Illini. Group IV contained material of approximately the maturity of Boone and Gibson, which are adapted to Missouri, southern Indiana, and southern Illinois.

This report includes the detailed results of all the Uniform Tests for the 1942 season, and also two- and three-year summaries of agronomic and chemical information for the different strains grown during the 1940-42 period.

COOPERATING AGENCIES AND PERSONNEL
for the
NORTH CENTRAL REGION

Bureau of Plant Industry:

Division of Forage Crops and Diseases: Jackson L. Cartter, Leonard F. Williams, Albert H. Probst, Denver I. Allen, Charles R. Weber, David Heusinkveld, Lewis C. Saboe, Morris A. Arneson, Floyd I. Collins, Orland A. Krober, Leon E. Wener, Donald C. Foster, and Samuel R. Aldrich.

Illinois Agricultural Experiment Station:

Agronomy Department: W. L. Burlison and C. H. Woodworth

Iowa Agricultural Experiment Station:

Farm Crops Department: H. D. Hughes

Missouri Agricultural Experiment Station:

Field Crops Department: W. C. Etheridge and B. L. King

Nebraska Agricultural Experiment Station:

Agronomy Department: T. A. Kiesselbach

North Dakota Agricultural Experiment Station:

Agronomy Department: T. E. Stoa

Ohio Agricultural Experiment Station:

Agronomy Department: R. D. Lewis

Purdue Agricultural Experiment Station:

Agronomy Department: G. H. Cutler


Minnesota Agricultural Experiment Station

Agronomy Department: A. C. Army

LOCATION OF COOPERATIVE NURSERIES

Location	Cooperator	Uniform Test Group I	Uniform Test Group II	Uniform Test Group III	Uniform Test Group IV
Strongsville, Ohio	NE.Br.Ohio Agr.Exp.Sta.	x			
Wooster, Ohio	Ohio Agr. Exp. Sta.	x	x		
Holgate, Ohio	NW.Br.Ohio Agr.Exp.Sta.		x	x	
Columbus, Ohio	Ohio Agr. Exp. Sta.			x	
Dearborn, Michigan*	Ford Motor Co.		x		
La Grange, Ind.	E. Frey & E. Bender	x	x		
Bluffton, Ind.	Gerald Bayless		x		
Wanatah, Ind.	Purdue Agr. Exp. Sta.		x		
Lafayette, Ind.	Purdue Agr. Exp. Sta.		x	x	
Greenfield, Ind.	Raymond Roney			x	
North Vernon, Ind.	Charles Robbins			x	x
Wheatland, Ind.	B. F. Carr				x
Evansville, Ind.	Leo Hirsch				x
Mt. Morris, Ill.	Earl Kump		x		
Dwight, Ill.	Frank Roeder		x	x	
Urbana, Ill.	Ill. Agr. Exp. Sta.		x	x	x
Clayton, Ill.	Russell S. Davis			x	x
Stonington, Ill.	Frank Garwood & Sons			x	x
Freeburg, Ill.	Loren Wilderman			x	x
Paris, Mo.	N.E. Mo. Agr. Exp. Sta.		x	x	
St. Joseph, Mo.*	G. C. McClain		x	x	
Columbia, Mo.	Mo. Agr. Exp. Sta.			x	x
Carrollton, Mo.*	Roy Monier			x	x
Elsberry, Mo.	Mo. Rice Exp. Field				x
Sikeston, Mo.	S. E. Mo. Agr. Exp. Field				x
Ames, Iowa	Iowa Agr. Exp. Sta.		x	x	
Kanawha, Iowa	North Ia. Agr. Exp. Assn.		x		
Hudson, Iowa	George M. Strayer		x		
Cherokee, Iowa	Cherokee State Hospital		x		
Waseca, Minn.	Br.Sta.Minn.Agr.Exp.Sta.		x		
Lincoln, Nebr.	Nebraska Agr. Exp. Sta.			x	
Fargo, N. Dakota	N. Dakota Agr. Exp. Sta.	x			
Torrington, Wyo.*	Wyoming Agr. Exp. Sta.	x			

*No yield data included in report at these locations, samples were submitted for chemical analyses.



Map of the North Central states showing location of the cooperative Uniform Tests.

METHODS

All Uniform Tests have been planted in replicated row-plots, using either a lattice square design with three replications or a restricted randomized block design with four replications. Row widths used at the different test locations have varied from twenty-one to forty-two inches depending upon the width in common use or the equipment available for handling the crop. Seeding rates have also varied with locations, the most prevalent rates being 150 to 200 viable seeds per row. Rates within this range have given satisfactory stands throughout the region under normal soil and weather conditions at planting time.

Yields were taken on individual replications after the seed had been dried to a uniform moisture content basis.

Chemical composition was determined for each strain in a Uniform Test on composite samples prepared by combining equal weights of seed from each replication at each location included in that particular Group Test. The location composites were prepared by combining equal weights of seed of each of the strains in a Group Test at an individual location. Percentage composition of the seed is expressed on a moisture-free basis. Seed size for each strain was also determined on the variety composite and was recorded as weight (in grams) per 100 seeds.

Lodging notes were recorded on a scale of 1 to 5 according to the following criteria:

- 1 Almost all plants erect.
- 2 Either all plants leaning slightly, or a few plants down.
- 3 Either all plants leaning moderately, or 25% to 50% of the plants down.
- 4 Either all plants leaning considerably, or 50% to 80% of the plants down.
- 5 All plants down badly.

Height was determined as the average length of plants in a plot from the ground to the top extremity at time of maturity.

Maturity was taken as the date when the leaves had dropped, the pods were ripe, and the stems were fairly dry. Maturity in all summaries is expressed as days earlier (-) or later (+) than a standard or reference variety. Reference varieties used for the different Uniform Tests are as follows: Group I, Mandarin; Group II, Illini; Group III, Illini; and Group IV, Gibson.

Seed Quality was rated from 1 to 5 according to the following scale:

1. Very good
2. Good
3. Fair
4. Poor
5. Very poor

The factors considered in estimating seed quality were: development of seed; wrinkling; damage; and color for the variety.

Uniform Test, Group I

The Group I Test was established in 1942 and was composed of sixteen named varieties, two selections from hybrids, and two U. S. D. ... foreign introductions. The origin of these varieties and strains is as follows:

Variety or Strain	Source or Originating Agency	Origin
Kabott	Central Exp.Farm,Ottawa	Selection from a Manchurian strain
Pagoda	Central Exp.Farm,Ottawa	Selection from(Manitoba Br.x Mandarin)
Cayuga	N.Y.Agr.Exp.Sta.,Ithaca	PI 65393 ^{1/}
Ontario	N.Y.Agr.Exp.Sta.,Ithaca	PI 65344
Goldsoy	Ontario Agr.College	Selection from O.A.C. 211
O.A.C. 211	Ontario Agr.College	Selection from Habaro
C39	Purdue Agr.Exp.Sta.	Selection from a natural hybrid
C40	Purdue Agr.Exp.Sta.	Selection from a natural hybrid
Earlyana	Purdue Agr.Exp.Sta.	C28 (Sel.from a natural hybrid)
Richland	Purdue Agr.Exp.Sta.	PI 70502-2
Minsoy	Minnesota Agr.Exp.Sta.	PI 27890
Manchu 831	S.Dakota Agr.Exp.Sta.	Selection from Manchu
PI. 79610	Minnesota Agr.Exp.Sta.	Foreign Plant Introduction
PI. 68666	U.S.Dept.of Agriculture	Foreign Plant Introduction
McR.Mandarin ^{2/}	Agr.Exp.Farm,Winnipeg	Selection from Mandarin
Wis.Man.3 Sel.	Wisconsin Agr.Exp.Sta.	Selection from Wisconsin Manchu 3
Wis.Manchu 606	Wisconsin Agr.Exp.Sta.	Selection from Manchu
Habaro	-	PI 20405
Mandarin	-	PI 36653
Sioux	-	PI 81021

^{1/}Division of Plant Exploration and Introduction, Bureau of Plant Industry, U. S. D. A.

^{2/}McRostie Mandarin

In order to meet a demand for material earlier than that in the Group II test, a Group I test was organized in 1942 to include strains too early for Group II. Richland and Earlyana were included as strains for comparison with the Group II test.

As this test is newly organized, its composition and the areas to which it is adapted are not so definitely settled. Probably some revision will be necessary for 1943. The test was grown at two locations in Ohio, one each in Indiana and North Dakota, and at four locations in Wyoming. Three of the locations in Wyoming were discarded due to early frosts. The other location in Wyoming and that at Fargo, North Dakota were less severely frosted and were included in the chemical analyses. Since the Fargo results differed so much from those of Ohio and Indiana, they were not included in the means for yield. Table I presents a summary of the mean agronomic and chemical data for these strains. Table 2 includes the individual yield data and yield ranks for four locations.

It should be noted that C40, C39, and Earlyana (C28) are very closely related strains. These data are of particular interest since Earlyana is being released by the Purdue Agricultural Experiment Station as an early variety for northern Indiana. Most of the difference in the mean yields of these strains is due to the exceptionally high yield of C40 at Strongsville. The summary of yield rank in Table 2 indicates that there is some interaction between varieties and fields but the comparison of mean yields with mean maturity ratings in Table 1 shows a strong tendency for earlier maturity to be associated with lower yields.

Lodging and height summaries are presented in Table 3. Richland continues to be outstanding in lodging resistance but is equalled in this test by McRostie Mandarin, Ontario, and Pagoda, and exceeded by Sioux. Sioux, however, is so very early and short that it is really not comparable. Sioux also had the poorest seed quality as is indicated in Table 4, a summary of individual maturity and seed quality notes. P.I. 68666 and Cayuga had the highest average ranking in this respect. Sioux is also outstanding in percentage of protein and reached a high of 57.2% at LaGrange, Indiana as shown in Table 5, a summary of individual seed size and chemical analyses. This high protein content is accompanied by an unusually low oil content. Most of the iodine numbers of the oils are relatively high. This might be expected since these are all northern locations.

Table 1. Summary of agronomic and chemical data for the strains in the Uniform Test, Group I, 1942.

Strain Number of Tests	Yield Bu./A.	Lodg- ing	Height Inches	Matu- rity ¹	Seed Quality	Percent-		Iodine Number of Oil	
						Seed age of Protein	Seed age of Oil		
	3	3	3	3	3	5 ²	5 ²	5 ²	
C40	35.2	1.8	32	+5.3	2.3	14.9	42.1	19.9	134
P.I.68666	34.7	2.2	25	+2.3	1.8	12.1	39.2	20.5	134
Manchu 831	34.4	1.8	27	+6.3	2.5	13.9	41.2	19.3	137
Habaro	34.2	1.5	24	+0.3	2.5	16.8	41.4	19.5	133
W.Man.3 Sel.	33.4	1.5	31	+3.7	2.3	16.0	39.5	20.6	135
Wis.Man.606	32.8	1.8	28	+7.3	2.0	15.7	40.4	20.3	135
C39	31.8	2.0	32	+4.3	2.2	14.4	41.7	20.0	135
Earlyana	31.7	2.0	33	+5.3	2.2	14.2	42.1	20.0	134
O.A.C. 211	31.7	1.8	23	+1.3	2.5	16.7	42.4	19.1	134
Richland	31.6	1.3	27	+10.7	2.2	14.0	37.7	20.4	130
McR.Mand. ²	31.6	1.3	22	-3.0	2.7	18.7	42.7	19.4	132
Mandarin	27.4	1.8	25	0.0	2.8	16.6	43.3	18.6	135
P.I. 79610	25.7	1.5	22	+0.3	2.4	15.7	42.6	19.5	134
Goldsoy	24.9	2.0	22	-4.0	3.3	17.8	42.3	18.7	136
Ontario	24.3	1.3	21	-0.7	2.5	18.2	40.0	21.1	132
Cayuga	24.2	1.8	25	-1.0	1.8	13.2	41.8	18.3	131
Kabott	20.5	1.7	19	-6.3	2.8	18.8	45.0	17.5	134
Minsoy	18.6	2.0	16	-11.0	2.8	11.9	41.2	19.1	129
Pagoda	16.1	1.3	21	-14.3	2.8	13.9	42.5	13.7	130
Sioux	9.3	1.0	11	-7.0	4.5	15.2	52.2	13.8	133

Bu.Nec.for Sig. 5.5
(5% level)

- ¹ Days earlier (-) or later (+) than Mandarin. Mandarin required 108 days to mature.
- ² McRostie Mandarin.
- ³ Mean of individual tests, composition on dry basis.

Table 2. Summary of yields in bushels per acre, and yield rank for strains in the Uniform Test, Group I, 1942.

Variety or Strain	Yield in Bu./A.					Yield Rank			
	Mean of 3 Tests ¹	Strongsville Ohio	Strongsville Wooster Ohio	La-Grange Ind.	Far-go N.Dak.	Strongsville Ohio	Strongsville Wooster Ohio	La-Grange Ind.	Far-go N.Dak.
C40	35.2	50.5	29.1	25.9	20.7	1	11	4	14
P.I.68666	34.7	42.8	34.0	27.3	16.9	4	1	1	17
Manchu 831	34.4	42.8	33.6	26.7	23.6	4	3	3	12
Habaro	34.2	46.0	31.2	25.5	30.3	2	7	7	5
W.Man.3 Sel.	33.4	43.2	31.0	25.9	24.1	3	8	4	9
Wis.Man.606	32.8	42.1	31.4	24.8	23.9	6	5	8	11
C39	31.8	39.1	30.7	25.6	19.6	11	9	6	15
Earlyana	31.7	39.8	27.9	27.2	18.4	10	13	2	16
O.A.C. 211	31.7	41.6	32.4	21.0	21.1	7	4	10	13
Richland	31.6	40.5	31.4	22.8	11.9	9	5	9	20
McR.Mandarin ²	31.6	40.9	33.8	20.1	39.8	8	2	12	2
Mandarin	27.4	33.2	28.6	20.6	34.5	15	12	11	3
P.I. 79610	25.7	33.3	25.9	17.6	25.5	14	15	14	7
Goldsoy	24.9	31.2	30.3	13.2	44.4	16	10	15	1
Ontario	24.3	35.1	20.2	17.8	26.9	12	19	13	6
Cayuga	24.2	34.2	26.3	11.8	24.0	13	14	16	10
Kabott	20.5	27.0	24.5	9.8	31.8	17	16	18	4
Minsoy	18.6	24.4	21.3	10.2	24.8	18	18	17	8
Pagoda	16.1	19.0	21.5	7.6	15.7	19	17	19	18
Sioux	9.3	13.8	12.6	1.6	12.7	20	20	20	19
Mean	27.7	36.0	27.9	19.2	24.5				
Coef.of Var.(%)	15.3	17.0	11.7	12.3	20.3				
Bu.Nec.for Sig. (5% level)	5.5	8.7	4.6	3.4	6.9				

¹ Fargo not included in mean because of severe frost damage on some varieties.

² McRostie Mandarin.

Table 3. Summary of lodging and height of strains in the Uniform Test, Group I, 1942.

Strain	Lodging				Height in Inches			
	Mean	Strong's-	La-		Mean	Strong's-	La-	
	of 3 Tests	ville Ohio	Wooster Ohio	Grange Ind.	of 3 Tests	ville Ohio	Wooster Ohio	Grange Ind.
C4C	1.8	3.5	1.0	1.0	32	40	30	26
PI. 68666	2.2	4.5	1.0	1.0	25	30	24	20
Manchu 831	1.8	3.5	1.0	1.0	27	30	27	23
Habaro	1.5	2.5	1.0	1.0	24	30	23	20
W.Man.3 Sel.	1.5	2.5	1.0	1.0	31	38	29	25
Wis.Man.606	1.8	3.5	1.0	1.0	28	34	26	24
C39	2.0	3.5	1.5	1.0	32	40	29	27
Earlyana	2.0	4.0	1.0	1.0	33	45	29	25
O.A.C. 211	1.8	3.5	1.0	1.0	23	28	22	18
Richland	1.3	2.0	1.0	1.0	27	33	24	25
McR.Mandarin	1.3	2.0	1.0	1.0	22	26	21	19
Mandarin	1.8	3.5	1.0	1.0	25	26	24	24
PI. 79610	1.5	2.5	1.0	1.0	22	26	21	18
Goldsoy	2.0	4.0	1.0	1.0	22	27	21	18
Ontario	1.3	2.0	1.0	1.0	21	26	19	17
Cayuga	1.8	3.5	1.0	1.0	25	32	23	20
Kabott	1.7	3.0	1.0	1.0	19	26	17	14
Minsoy	2.0	4.0	1.0	1.0	16	20	14	13
Pagoda	1.3	2.0	1.0	1.0	21	24	21	17
Sioux	1.0	1.0	1.0	1.0	11	12	11	9

Table 4. Summary of maturity and seed quality of strains in the Uniform Test, Group I, 1942.

Strain	Maturity				Seed Quality			
	Mean of 3 Tests	Strongs-ville Ohio	Wooster Ohio	La-Grange Ind.	Mean of 3 Tests	Strongs-ville Ohio	Wooster Ohio	La-Grange Ind.
C40	+5.3	+9	+12	-5	2.3	3.0	2.5	1.5
PL 68666	+2.3	+7	+12	-12	1.8	2.0	2.0	1.5
Manchu 831	+6.3	+7	+16	-4	2.5	3.0	3.0	1.5
Habaro	+0.3	+5	+12	-16	2.5	3.0	2.5	2.0
W,Man.3 Sel.	+3.7	+3	+12	-4	2.3	2.0	3.0	2.0
Wis.Man. 606	+7.3	+5	+16	+1	2.0	2.0	2.5	1.5
C39	+4.3	+7	+10	-4	2.2	2.0	2.5	2.0
Earlyana	+5.3	+10	+10	-4	2.2	3.0	2.0	1.5
O.A.C. 211	+1.3	+5	+10	-11	2.5	3.0	2.0	2.5
Richland	+10.7	+10	+18	+4	2.2	2.0	2.5	2.0
McR. Mandarin	-3.0	-1	-2	-6	2.7	3.0	2.0	3.0
Mandarin	0.0	0	0	0	2.8	4.0	2.0	2.5
PL 79610	+0.3	+1	+4	-4	2.4	2.8	2.0	2.5
Goldsoy	-4.0	-3	-3	-6	3.3	4.0	3.0	3.0
Ontario	-0.7	0	+4	-6	2.5	3.0	2.0	2.5
Cayuga	-1.0	+2	-5	0	1.8	2.0	2.0	1.5
Kabott	-6.3	-5	-5	-9	2.8	3.0	2.0	3.5
Minsoy	-11.0	-4	-7	-22	2.8	3.0	2.0	3.5
Pagoda	-14.3	-3	-7	-33	2.8	3.0	3.0	2.5
Sioux	-7.0	-8	-11	-2	4.5	5.0	4.0	4.5
Mandarin Matured		9/15	9/5	9/21				
Date of Planting		5/29	6/4	5/20				

Table 5. Summary of seed size and chemical data for the strains in the Uniform Test, Group I, 1942

Variety or Strain	Grams per 100 seeds										Percentage of Protein																																																																																																																																																																																																																																																																					
	Mean of 5 Tests					Strongsville					Mean of 5 Tests					Strongsville																																																																																																																																																																																																																																																																
	Ohio	La-Ind.	Wooster Ohio	Fargo N.Dak.	Torrington Wyo.	Ohio	La-Ind.	Wooster Ohio	Fargo N.Dak.	Torrington Wyo.	Ohio	La-Ind.	Wooster Ohio	Fargo N.Dak.	Torrington Wyo.	Ohio	La-Ind.	Wooster Ohio	Fargo N.Dak.	Torrington Wyo.																																																																																																																																																																																																																																																												
C40	14.9	17.6	14.9	18.3	10.9	12.9	42.1	41.8	44.9	45.8	40.2	38.0	PI. 68666	12.1	16.1	14.4	15.3	7.7	7.2	39.2	39.4	43.4	42.0	38.5	32.6	Manchu 831	13.9	17.3	15.0	15.5	10.9	11.0	41.2	42.9	44.7	45.1	40.2	33.3	Habaro	16.8	22.2	17.9	19.1	14.0	10.9	41.4	43.3	45.9	45.9	42.3	29.4	W.Man. 3 sel.	16.0	19.4	14.9	18.9	12.5	14.0	39.5	40.3	42.7	43.3	38.4	33.0	Wis.Man. 606	15.7	17.6	16.1	18.2	12.9	13.7	40.4	40.7	42.3	44.5	38.9	35.7	C39	14.4	17.5	15.1	17.3	10.0	12.0	41.7	41.4	45.2	44.5	39.4	37.9	Earlyana	14.2	17.4	14.3	17.2	10.2	12.0	42.1	42.9	44.9	45.4	40.2	37.0	O. A. C. 211	16.7	21.7	17.2	19.5	14.5	10.6	42.4	43.5	46.2	47.1	43.1	32.3	Richland	14.0	18.2	16.8	18.7	8.6	7.6	37.7	39.8	41.7	41.6	37.1	28.4	McR. Mand. ¹	18.7	20.9	16.2	22.2	18.5	15.9	42.7	44.6	45.3	47.6	41.9	34.2	Mandarin	16.6	18.4	14.8	19.7	17.0	13.0	43.3	45.9	45.0	47.4	42.3	56.0	PI. 79610	15.7	17.3	13.4	18.3	14.8	14.7	42.6	41.9	44.9	45.7	40.4	40.1	Goldsoy	17.8	19.5	17.6	18.3	19.0	14.7	42.3	44.5	46.0	46.3	42.5	32.3	Ontario	18.2	20.0	17.7	21.4	17.0	14.9	40.0	41.6	42.4	45.3	39.1	31.8	Cayuga	13.2	15.8	11.4	15.7	12.6	10.6	41.8	44.8	44.1	46.0	41.9	32.0	Kabott	18.8	20.2	17.2	21.7	18.6	16.4	45.0	45.3	47.3	48.4	44.1	39.7	Minsoy	11.9	13.1	11.0	14.8	10.7	10.0	41.2	44.1	44.1	46.4	41.3	30.0	Pegoda	13.9	15.1	13.3	15.8	12.9	12.2	42.5	43.2	44.6	45.3	45.4	34.0	Sioux	15.2	17.4	14.8	14.3	15.6	14.0	52.2	49.2	53.4	57.2	51.0	50.1	Mean							43.1	45.0	46.0	41.4	41.4	34.9

¹ Mc Rostic Mandarin

Table 5. (continued)

Variety or Strain	Percentage of Oil				Lardine Number of Oil									
	Mean of 5 Tests	StrongS- ville Ohio	Wooster Ohio	La- Grange Ind.	Mean of 5 Tests	StrongS- ville Ohio	Wooster Ohio	La- Grange Ind.	Tarring- ton Wyo.	Mean of 5 Tests	StrongS- ville Ohio	Wooster Ohio	La- Grange Ind.	Tarring- ton Wyo.
C40	19.9	20.7	19.5	19.5	134	134	131	133	135	134	134	131	133	137
PI. 68666	20.5	21.9	20.5	20.6	134	134	130	129	139	134	133	130	129	139
Manchu 831	19.3	19.6	18.8	18.0	137	137	135	134	138	137	137	135	134	141
Habaro	19.5	19.3	18.9	18.5	133	133	130	130	135	133	133	130	130	137
Wis. Jan. 3 Sel.	20.6	20.7	20.7	19.6	135	134	134	132	138	135	134	134	132	139
Wis. Jan. 606	20.3	20.4	20.5	19.0	135	135	133	132	138	135	135	133	132	137
C39	20.0	21.1	19.9	19.4	135	134	131	133	137	135	134	131	133	138
Earlyana	20.0	20.3	19.9	19.5	134	133	132	132	135	134	133	132	132	138
O. A. C. 211	19.1	19.3	18.8	17.7	134	134	131	132	136	134	134	131	132	139
Richland	20.4	20.7	20.4	20.5	130	129	126	126	136	130	129	126	126	135
Mc. R. Mand. ¹	19.4	18.6	18.5	17.9	132	133	131	129	136	132	133	131	129	131
Mandarin	18.6	17.7	18.2	17.8	135	137	133	131	139	135	137	133	131	135
PI. 79610	19.5	20.1	18.8	19.3	134	136	131	130	137	134	136	131	130	136
Goldsoy	18.7	17.8	17.6	17.6	136	137	132	133	139	136	137	132	133	137
Ontario	21.1	20.4	21.2	19.5	132	134	129	128	136	132	134	129	128	134
Cayuga	18.3	17.4	17.3	17.5	131	131	129	125	136	131	131	129	125	135
Kabott	17.5	17.7	17.0	16.7	134	136	132	131	136	134	136	132	131	134
Ilinsoy	19.1	17.8	17.9	18.0	129	131	127	122	134	129	131	127	122	132
Pagoda	18.7	19.0	18.5	18.4	130	129	127	125	136	130	129	127	125	133
Sioux	13.8	15.1	13.1	12.3	133	135	132	127	135	133	135	132	127	136
Mean		19.3	18.8	18.4	134	134	131	130	137	134	134	131	130	136

McRostie Mandarin

Uniform Test, Group II

The Group II Test in 1942 was composed of eleven named varieties, six U. S. D. A. foreign introductions, and three selections from hybrids. The origin of these strains is as follows:

Variety or Strain	Source or Originating Agency	Origin
Dunfield	Purdue Agr. Exp. Sta.	P.I. 36846 <u>1/</u>
Earlyana	Purdue Agr. Exp. Sta.	C28 (natural hybrid)
Mandell	Purdue Agr. Exp. Sta.	Selection from Manchu
Richland	Purdue Agr. Exp. Sta.	P.I. 70502-2
Mukden	Iowa Agr. Exp. Sta.	P.I. 50523-Q
Mingo	Ohio Agr. Exp. Sta.	Selection from Manchu
Seneca	N.Y. Agr. Exp. Sta., Ithaca	F.C. 05654G <u>2/</u>
Illini	Illinois Agr. Exp. Sta.	Selection from A. I.
L6-12	Illinois Agr. Exp. Sta.	Selection from (Mandarin x Manchu)
L6-685	Illinois Agr. Exp. Sta.	Selection from (Mandarin x Manchu)
L6-700	Illinois Agr. Exp. Sta.	Selection from (Mandarin x Manchu)
P.I. 68474	U.S. Dept. of Agriculture	Foreign Plant Introduction
P.I. 70478	U.S. Dept. of Agriculture	Foreign Plant Introduction
P.I. 91109	U.S. Dept. of Agriculture	Foreign Plant Introduction
P.I. 91161	U.S. Dept. of Agriculture	Foreign Plant Introduction
P.I. 92592	U.S. Dept. of Agriculture	Foreign Plant Introduction
P.I. 92717	U.S. Dept. of Agriculture	Foreign Plant Introduction
Wis. Manchu 3	Wisconsin Agr. Exp. Sta.	Selection from Manchu
Wis. Man. 3 Sel.	Wisconsin Agr. Exp. Sta.	Selection from Wisconsin Manchu 3
Wis. Man. 606	Wisconsin Agr. Exp. Sta.	Selection from Manchu

1/ Division of Plant Exploration and Introduction, Bureau of Plant Industry, U. S. D. A.

2/ Division of Storage Crops and Diseases, Bureau of Plant Industry, U. S. D. A.

The varieties and strains included in the Uniform Test, Group II, in 1942 varied widely in yield among the various locations and show a range of ten bushels between the means of the thirteen tests. There was, likewise, a wide range in maturity of the strains at the individual stations and a range of twelve days between the variety means of the ten stations at which maturity data were obtained. The wide variation in maturity appears to reflect itself upon the yield, and in the main the later maturing strains have been the higher both in 1942 and in the summary of three years, 1940-42, as shown in Table 13.

Two strains of hybrid origin, L6-685 and L6-12, were significantly higher in yield in 1942 than any of the commercial varieties in the test. Of these two strains, L6-685 is somewhat more promising when the whole array of agronomic and chemical features are considered. The three-year summary, Table 13, shows the outstanding performance of this strain in comparison to the commercial varieties and other strains. It is most worthy to note that L6-685 has yielded, on the average, six bushels more than the combined average of Lingo, Illini, Lukden, Mandell, and Dunfield which are of approximately the same maturity and represent the most commonly grown varieties of comparable maturity in the central area of the "soybean belt". In addition to the exceptional yielding ability of L6-685, it likewise has a higher oil content and a higher iodine number than any of the commercial varieties listed above. A comparison of the chemical analyses of L6-685 and L6-12 with several commercial varieties is shown in Table 17 for 1942 and in Table 16 for the 1941-42 summary.

Earlyana, the earliest variety in the test, was also grown in the Group I test and is being released by the Purdue Agricultural Experiment Station. It is recommended especially for northern Indiana where conditions require an earlier and taller growing variety than Richland. Earlyana compares favorably with Richland in yield, oil content, and seed quality. It grows taller and is usually about four days earlier than Richland, thus meeting the demands of farmers who prefer early harvesting of soybeans in order to seed wheat soon after the Hessian fly-free date. The taller growth also makes it more desirable on soils of low fertility. This new variety is not as lodging resistant as Richland and may be expected to lodge more excessively on highly fertile soils.

In addition to L6-685 and the named varieties, several strains merit consideration as possible breeding material. Notable among these are P.I. 60474 which is quite favorable in yield, oil content, and maturity, but is rather short; and P.I. 70478 which is high in oil content and has about the same maturity as Richland.

Wisconsin Manchu 3 Selection was selected from Wisconsin Manchu 3 for the purpose of purification of the latter variety. The 1942 data indicate that the selection varies considerably in agronomic characteristics from the parent variety. The most notable variations, as shown in Table 6, are maturity, seed quality, and yield.

Early frost probably affected the yields in 1942, particularly of the later maturing strains. This is shown by the fact that at several locations the earlier varieties were comparatively higher in yield than is usually obtained.

In a number of cases, due to early frost, maturity records represent not actual maturity, but the best estimates based on the stage of development of the plant at the time of the frost.

Table 6. Summary of agronomic and chemical data for the strains in the Uniform Test, Group II, 1942.

Strain	Yield	Lodg- ing	Height Inches	Matu- rity ¹	Seed Qual- ity	Seed Size	Percent-		Iodine Number of Oil
	in Bu./A.						age of Protein	age of Oil	
Number of Tests	13	15	12	10	10	16 ²	16 ²	16 ²	16 ²
L6-685	40.4	2.6	40	-1.6	1.6	14.7	39.9	22.1	136
L6-12	38.1	2.8	40	-1.4	1.6	14.0	41.4	21.3	137
L6-700	35.6	3.3	39	-1.2	1.6	14.5	41.6	21.5	137
P.I.92717	35.3	2.8	37	-5.2	1.8	14.3	41.5	20.9	136
Mingo	33.9	3.1	39	-1.7	2.2	15.4	42.2	20.8	134
P.I.68474	33.7	3.2	32	-4.7	1.6	14.7	38.8	22.2	135
Illini	33.4	3.3	43	0.0	1.9	13.9	40.7	21.0	135
P.I.91161	33.0	3.3	35	-4.8	2.0	16.7	38.5	22.2	132
P.I.92592	32.9	3.0	32	-4.7	1.8	16.9	40.2	22.3	134
Dunfield	32.8	3.1	39	-1.7	1.5	15.2	39.5	21.7	129
P.I.91109	32.7	3.0	31	-6.5	2.1	15.2	41.9	20.3	129
Seneca	32.5	2.8	38	-5.8	2.8	15.4	40.2	21.7	134
Mandell	32.3	2.6	40	-3.0	2.1	15.5	43.6	19.9	136
P.I.70478	32.1	3.0	31	-6.9	1.8	14.9	39.0	22.1	133
Mukden	31.8	2.6	40	-5.3	1.8	15.0	43.1	20.2	131
Wis.Man. 3	31.7	3.2	38	-6.4	2.6	16.9	41.9	20.8	135
Wis.Man.606	31.5	3.1	35	-6.7	2.1	16.7	41.9	20.7	135
Earlyana	31.4	2.7	36	-11.8	1.8	14.7	42.6	20.7	133
Richland	31.0	2.0	33	-7.5	2.0	15.9	40.3	21.1	129
W.Man.3 Sel.	30.4	3.0	37	-10.9	2.0	17.0	41.3	20.9	135

Bu.Nec.for Sig. 2.5
(5% level)

- ¹ Days earlier (-) or later (+) than Illini. Illini required 130 days to mature.
- ² Composite sample of 16 locations, composition on dry basis.

Table 7. Summary of rank for yield, arranged in order of mean yields, for the Uniform Test, Group II, 1942

Strain	Anes Iowa		Ur-bena Ill.		Bluff-ton Ind.		Lafayette Ind.		Dwight Ill.		Cherokee Iowa		Wanatah Ind.		Kana-wha Iowa		La-Grange Ind.		Wooster Ohio		Holgate Ohio		Hudson Iowa		Waseca Minn.		Paris I.O.		Mt. Morris Ill.																																										
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20																															
L6-685	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20																															
L6-12	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20																																
L6-700	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20																																	
P.I. 92717	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20																	
Mingo	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20																		
P.I. 68474	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20																	
Illini	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20						
P.I. 91161	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
P.I. 92592	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20								
Dunfield	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20														
P.I. 91109	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20																
Seneca	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20															
Mandell	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20												
P.I. 70478	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20					
Mukden	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20													
Wis.Man.3	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20																			
Wis.Man.606	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20			
Earlyana	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20							
Richland	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
W.Lan.3 Sel.	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20			

Table 8. Summary of yields in bushels per acre for the strains in the Uniform Test, Group II, 1942.

Strain	Mean of 13 Tests ¹	Ames Iowa	Ur- bana Ill.	Bluff- ton Ind.	Lafay- ette Ind.	Dwight Ill.	Choro- koc Iowa	Wana- tah Ind.
L6-685	40.4	52.3	57.3	54.8	50.0	45.8	37.4	40.6
L6-12	38.1	44.0	53.4	53.9	45.9	44.9	40.8	41.4
L6-700	35.6	40.8	51.3	48.2	46.4	38.8	39.2	43.4
P.I. 92717	35.3	44.1	45.8	45.6	42.5	40.5	34.9	38.4
Mingo	33.9	46.3	47.9	42.4	43.2	36.2	36.0	33.7
P.I. 68474	33.7	46.5	45.3	41.7	39.0	42.8	35.7	40.5
Illini	33.4	42.0	40.5	49.2	42.6	31.4	41.0	35.2
P.I. 91161	33.0	40.2	42.1	36.3	41.7	38.6	35.1	36.1
P.I. 92592	32.9	41.7	48.2	42.2	38.3	39.6	34.4	31.1
Dunfield	32.8	47.3	46.2	41.2	41.2	34.9	29.7	33.2
P.I. 91109	32.7	46.6	38.2	35.4	37.2	37.8	35.5	34.4
Seneca	32.5	47.2	38.1	42.0	37.6	40.6	36.1	33.0
Mandell	32.3	48.3	41.3	39.7	40.2	35.6	34.8	30.0
P.I. 70478	32.1	42.9	40.1	38.2	37.0	36.1	34.9	32.0
Mukden	31.8	47.9	45.3	38.7	41.4	35.1	33.7	30.8
Wis.Man.3	31.7	45.0	39.4	36.7	37.7	33.6	36.7	29.2
Wis.Man.606	31.5	43.3	38.9	33.6	39.0	30.8	34.1	34.2
Earlyana	31.4	41.8	37.4	35.1	35.4	30.6	40.8	35.8
Richland	31.0	44.4	37.9	35.6	34.0	38.9	31.6	31.2
W.Man.3 Sol.	30.4	43.3	36.6	31.9	33.2	30.5	35.0	31.4
Mean	33.3	44.8	43.5	41.1	40.2	37.1	35.9	34.8
Coef. of Var.	11.7	9.4	10.1	14.0	8.8	11.5	9.6	10.6
Bu. Nec. for	2.5	6.0	6.2	8.1	5.0	6.0	4.8	5.2
Sig.(5% level)								

¹ Waseca and Mt. Morris not included in the mean.

Table 3 (con't)

Strain	Kana- wha Iowa	La- Grango Ind.	Woost- er Ohio	Hol- gate Ohio	Hud- son Iowa	Wasoca Minn. ²	Paris Mo.	Mt. Morris Ill. ³
L6-685	32.2	36.7	30.8	36.6	32.4	24.2	18.1	13.2
L6-12	30.6	35.0	28.0	32.1	30.2	18.3	15.5	11.1
L6-700	24.5	31.2	30.5	31.9	21.2	12.4	15.5	10.7
P.I. 92717	31.0	28.2	30.8	32.9	27.2	25.0	15.5	13.2
Mingo	27.3	32.0	30.4	26.2	24.0	9.3	15.2	11.7
P.I. 68474	29.9	29.2	25.1	28.2	20.4	15.3	15.7	11.9
Illini	29.7	30.9	26.3	24.0	28.9	12.0	12.9	10.8
P.I. 91161	34.8	27.7	28.7	26.2	26.7	19.6	15.0	13.5
P.I. 92592	30.3	29.0	29.3	23.5	20.4	17.4	18.0	14.8
Dunfield	30.1	26.5	28.1	23.0	30.2	12.2	14.7	11.6
P.I. 91109	35.6	24.4	28.3	25.0	31.1	21.1	15.9	15.4
Seneca	31.6	27.1	26.9	21.1	27.9	13.0	15.0	13.1
Mandoll	29.0	26.8	24.7	27.3	27.6	21.2	13.9	12.9
P.I. 70478	36.2	28.9	28.3	29.5	22.9	12.4	10.6	13.8
Mukdon	31.1	27.4	27.4	17.2	25.8	25.6	12.1	12.0
Wis. Man.3	34.5	25.5	28.2	26.3	25.9	32.2	13.0	13.5
Wis. Man.606	33.8	26.8	29.3	27.9	24.0	16.0	13.6	18.0
Earlyana	35.1	25.2	27.4	28.3	24.7	26.0	10.7	16.9
Richland	34.2	22.4	26.9	23.7	27.8	27.4	14.6	13.2
W. Man.3 Sel.	35.8	25.6	28.7	26.2	21.8	17.3	13.1	14.0
Mean	31.8	28.4	28.2	27.0	26.1	19.0	14.3	13.3
Coef. of Var.	6.3	11.0	10.2	19.0	17.2	-	12.6	15.0
Bu. Nec. for Sig.(5% level)	2.8	4.4	4.1	7.2	6.4	-	2.5	2.8

² Only one replication at Wasoca

³ Mt. Morris planted quite late, some varieties severely frosted.

Table 9. Summary of lodging notes for the strains in the Uniform Test, Group II, 1942

Strain	Hoan of 15 Tests	Ames Iowa	Ur- bana Ill.	Bluff- ton Ind.	Lafay- ette Ind.	Dwight Ill.	Chero- keo Iowa	Wana- tah Ind.
L6-685	2.6	3.0	2.0	2.2	2.0	2.5	4.0	3.6
L6-12	2.8	3.3	2.0	2.9	2.1	2.0	3.5	3.6
L6-700	3.3	4.3	3.0	3.2	2.9	3.0	4.8	4.0
P.I. 92717	2.8	3.0	3.0	2.9	2.7	2.0	4.0	4.2
Mingo	3.1	3.0	3.0	3.1	2.4	4.0	4.0	3.9
P.I. 68474	3.2	3.0	4.0	2.7	3.2	3.0	5.0	4.4
Illini	3.3	4.5	3.0	3.0	2.4	4.0	4.5	4.1
P.I. 91161	3.3	3.5	3.5	3.1	2.7	3.0	5.0	4.2
P.I. 92592	3.0	3.0	4.0	3.0	3.1	2.0	4.3	4.4
Dunfield	3.1	3.3	3.0	3.5	2.2	3.0	4.0	4.1
P.I. 91109	3.0	3.5	4.0	2.9	2.5	2.5	4.5	4.4
Seneca	2.8	3.0	2.5	2.6	2.7	2.5	3.8	3.5
Mandell	2.6	3.5	2.0	3.0	2.6	2.0	3.3	3.9
P.I. 70478	3.0	3.5	3.5	2.7	2.6	3.0	4.5	4.2
Mukden	2.6	2.8	2.0	2.9	2.9	2.5	3.8	3.7
Wis.Man.3	3.2	3.3	4.0	2.7	3.6	4.0	4.5	4.1
Wis.Man.606	3.1	3.3	3.5	2.7	3.2	3.0	3.8	4.1
Earlyana	2.7	3.8	2.0	2.6	3.0	2.5	3.8	4.0
Richland	2.0	1.8	2.0	2.0	1.5	2.0	2.8	3.6
W.Man.3 Sel.	3.0	3.3	3.0	2.4	2.7	3.0	3.8	4.1

Table 9 (continued)

Strain	Kana- wha Iowa	La- Grange Ind.	Woost- er Ohio	Hol- gato Ohio	Hud- son Iowa	Wasceca Minn.	Paris Mo.	Mt. Morris Ill.
L6-685	2.8	1.1	1.0	2.3	3.0	5.0	1.5	2.5
L6-12	3.0	1.5	2.0	2.5	3.5	5.0	1.5	3.0
L6-700	3.5	1.5	1.0	2.3	4.3	5.0	2.0	4.0
P.I. 92717	2.3	1.1	1.0	3.5	3.5	5.0	1.5	2.5
Mingo	3.3	1.4	1.0	2.8	3.5	5.0	2.5	3.0
P.I. 68474	3.3	1.6	1.0	2.3	4.8	5.0	1.0	4.0
Illini	4.3	1.6	1.0	2.5	4.8	5.0	2.5	3.0
P.I. 91161	3.8	1.4	1.0	3.5	4.3	5.0	2.0	3.5
P.I. 92592	3.3	1.0	1.0	2.8	3.3	5.0	1.5	3.0
Dunfield	3.0	1.5	1.0	3.8	3.5	5.0	2.5	3.0
P.I. 91109	2.5	1.0	1.0	3.3	4.3	4.0	1.5	3.0
Seneca	2.8	1.0	1.0	2.8	3.8	5.0	1.5	3.0
Mandell	2.3	1.1	1.0	3.0	3.3	4.0	1.5	2.5
P.I. 70478	2.3	1.0	1.0	3.5	4.3	5.0	1.0	3.0
Mukden	2.5	1.1	1.0	2.8	3.5	4.0	1.0	2.0
Wis.Man.3	3.3	1.4	1.0	2.8	4.3	3.0	2.0	3.5
Wis.Man.606	2.8	1.1	1.0	3.8	5.8	5.0	2.0	3.0
Earlyana	2.8	1.0	1.0	2.8	3.8	4.0	1.0	2.5
Richland	1.3	1.0	1.0	2.5	2.5	3.0	1.0	2.0
W.Man.3 Sel.	3.0	1.0	1.0	3.8	3.5	4.0	3.0	4.0

Table 10. Summary of plant height for the strains in the Uniform Test, Group II, 1942

Strain	Mean of 13 Tests	Ur-bana		Bluff-ton		Lafayette		Cherokee		La-Grange		Hoost-er		Hudson		Paris Mo.
		Iowa	Ill.	Iowa	Ill.	Iowa	Ind.	Iowa	Ill.	Iowa	Ind.	Ohio	Ohio	Iowa	Iowa	
L6-685	40	48	39	38	42	48	41	43	32	28	53	43	30			
L6-12	40	48	39	37	41	47	41	43	34	25	51	43	30			
L6-700	39	46	39	34	41	46	40	42	28	28	51	43	30			
P.I. 92717	37	44	36	32	39	44	40	42	29	26	47	40	27			
Mingo	39	45	37	32	40	44	42	43	31	26	51	41	31			
P.I. 68474	32	37	32	24	35	37	34	34	25	20	49	35	23			
Illini	43	51	47	35	49	50	43	47	35	25	55	46	31			
P.I. 91161	35	42	32	29	38	41	33	40	27	24	45	39	27			
P.I. 92582	32	38	31	29	34	38	33	35	25	22	46	28	23			
Dunfield	39	46	40	35	41	45	41	42	33	25	50	42	31			
P.I. 91109	31	39	31	27	34	38	33	34	23	22	40	32	21			
Seneca	38	49	39	24	42	49	41	45	25	23	43	45	26			
Mandell	40	46	40	34	44	44	42	44	33	24	49	44	30			
P.I. 70478	31	37	36	22	34	38	32	33	23	21	44	33	22			
Mukden	40	49	44	32	44	49	40	46	28	25	49	48	28			
Wis.Man.3	38	45	40	30	43	45	42	41	25	24	44	43	29			
Wis.Man.606	35	44	37	27	39	43	41	36	26	20	44	37	28			
Earlyana	36	44	37	27	38	40	40	40	27	23	45	43	24			
Richland	33	40	32	25	34	40	33	40	23	22	40	36	25			
Wis.Man.3 Sel.	37	45	40	27	43	44	38	42	25	26	47	41	28			

Table 11. Summary of maturity notes for the strains in the Uniform Test, Group II, 1942.

Strain	Mean of 10 Tests	Ames	Ur-	Bluff-	Lafay-	Chero-	Kana-	La-	Woost-	Waseca	Paris
		Iowa	bana Ill.	ton Ind.	ette Ind.	keo Iowa	wha Iowa	Grange Ind.	er Ohio	Minn.	Mo.
I6-685	-1.6	-6	+4	0	-3	-3	-3	-2	0	-4	+1
I6-12	-1.4	-5	+2	0	-2	-3	-3	-2	+1	-4	+2
I6-700	-1.2	-4	+4	-1	-3	-2	-2	-2	0	-4	+2
P.I. 92717	-5.2	-9	0	-5	-6	-8	-8	-4	-6	-4	-2
Mingo	-1.7	-7	+8	-1	-4	-5	-5	-5	0	0	+2
P.I. 68474	-4.7	-6	-1	-6	-8	-3	-5	-7	-6	-4	-3
Illini	0.0	0	0	0	0	0	0	0	0	0	0
P.I. 91161	-4.8	-8	+2	-6	-7	-5	-4	-6	-7	-4	-3
P.I. 92592	-4.7	-6	0	-5	-7	-3	-5	-9	-6	-4	-4
Dunfield	-1.7	-2	-2	-2	-5	-2	-2	-5	0	0	+3
P.I. 91109	-6.5	-8	0	-6	-8	-6	-6	-6	-9	-9	-7
Seneca	-5.8	-8	-1	-10	-7	-4	-4	-7	-11	0	-6
Mandell	-3.0	-8	+2	+1	-2	-5	-5	-2	-1	-9	-1
P.I. 70478	-6.9	-8	+4	-8	-15	-5	-5	-15	-7	-4	-6
Mukden	-5.3	-7	-1	-4	-5	-7	-7	-7	-6	-4	-5
Wis.Man.3	-6.4	-13	+3	-6	-6	-9	-5	-6	-7	-14	-1
Wis.Man.606	-6.7	-11	-2	-9	-8	-8	-6	-7	-11	-4	-1
Earlyana	-11.8	-14	-6	-14	-21	-8	-8	-16	-15	-9	-7
Richland	-7.5	-12	-3	-8	-8	-8	-4	-9	-8	-9	-6
W.Man.3 Sel.	-10.9	-13	-6	-9	-20	-9	-6	-16	-14	-9	-7
Illini Matured		10/2	9/18	10/4	10/7	10/3	10/3	10/5	10/2	10/14	9/15
Date of Planting		5/16	5/21	5/21	5/25	5/27	5/26	5/20	6/4	5/23	5/29

Table 12. Summary of seed quality notes for the strains in the Uniform Test, Group II, 1942.

Strain	Mean of 10 Tests	Ur-	Bluff-	Wana-	La-	Woost-	Hol-	Paris	Mt.
		Ill.	ton Ind.	tah Ind.	Grange Ind.	er Ohio	gate Ohio	Mo.	Morris Ill.
L6-685	1.6	1.0	1.0	2.5	1.0	2.0	1.0	1.0	3.0
L6-12	1.6	1.0	1.0	2.5	1.0	2.0	1.0	1.0	3.0
L6-700	1.6	1.0	1.5	2.0	1.5	2.0	1.5	1.0	3.0
P.I. 92717	1.8	2.0	1.0	2.5	2.0	1.5	1.0	2.5	3.0
Mingo	2.2	2.0	1.5	3.0	1.5	2.0	2.0	2.5	3.0
P.I. 68474	1.6	1.0	1.0	2.5	2.0	2.5	1.0	1.5	2.0
Tillini	1.9	1.0	2.0	2.5	1.5	2.5	2.0	1.5	2.0
P.I. 91161	2.0	1.0	2.0	3.5	2.0	2.5	2.0	1.5	2.0
P.I. 92592	1.8	1.0	2.5	2.0	2.5	3.0	1.5	1.0	2.0
Dunfield	1.5	1.0	1.5	2.5	1.0	2.0	1.5	2.0	1.0
P.I. 91109	2.1	2.0	1.5	3.0	2.0	3.0	1.5	2.0	2.0
Seneca	2.8	2.0	3.0	4.0	3.0	5.0	3.0	1.0	2.0
Mandell	2.1	2.0	2.5	2.5	2.5	2.0	2.0	2.0	2.0
P.I. 70478	1.8	1.0	2.0	2.5	1.0	3.0	1.5	2.0	2.0
Mukden	1.8	1.0	2.0	2.0	2.5	3.0	2.0	1.0	2.0
Wis.Man.3	2.6	2.0	2.0	4.0	2.0	3.0	2.0	2.0	3.0
Wis.Man.606	2.1	2.0	2.0	3.5	2.5	2.0	2.0	2.0	2.0
Earlyna	1.8	2.0	2.0	2.5	1.5	1.5	2.0	2.0	1.0
Richland	2.0	2.0	2.0	2.5	2.0	1.5	1.5	2.0	2.0
Wis.Man.3 Sel.	2.0	2.0	2.0	3.5	1.5	3.0	1.5	1.5	2.0

Table 13. Three-year summary of mean agronomic and chemical data for the strains in the Uniform Test, Group II, 1940-1942.

Strain	Mean Yield Bu/A	Lodging Inches	Height Inches	Maturity ¹	Seed Quality	Seed Size	Per-cent Protein	Per-cent Oil	Iodine Number of Oil
No. of Tests	31	27	22	27	28	35*	35*	35*	35*
L6-685	34.2	2.3	36	-0.8	1.5	15.0	41.4	21.4	135
L6-12	33.9	2.4	36	-0.7	1.5	14.5	42.4	20.8	135
P.I. 92717	29.9	2.3	33	-4.1	1.8	14.1	43.2	20.0	134
Mingo	29.8	2.9	35	-1.0	2.1	15.7	43.6	20.0	132
Illini	29.0	2.8	38	0.0	1.6	14.2	41.7	20.3	133
P.I. 68474	28.8	2.7	29	-6.2	1.6	14.6	40.5	21.3	132
Mandell	28.3	2.2	36	-1.7	1.9	14.9	45.1	19.1	133
Dunfield	27.9	2.7	34	-0.4	1.6	16.4	41.0	20.8	127
P.I. 91161	27.9	2.6	32	-4.5	2.1	16.1	40.4	21.1	128
P.I. 70478	26.8	2.3	28	-5.8	1.7	14.5	40.1	21.4	128
Mukden	26.1	2.1	36	-4.4	1.8	14.5	44.8	19.5	128
Richland	26.1	1.5	31	-5.8	1.9	16.2	41.7	20.2	126
Earlyana	25.1	2.3	33	-9.6	2.1	14.4	44.4	20.0	130
Wis.Man.3	25.0	2.7	35	-4.5	2.5	16.6	43.6	19.9	131
Wis.Man.606	24.4	2.3	31	-8.3	2.4	16.1	43.6	20.0	132

¹ Days earlier (-) or later than (+) Illini. Illini required 130 days to mature (3 year average of all locations).

* Mean of composite samples by years, composition on dry basis. Sixteen tests in 1942, 10 tests in 1941, and 9 tests in 1940.

Table 14. Analysis of variance for yield of seed for the Uniform Test, Group II, 1942.

Source of Variation	Degrees of Freedom	Mean Squares
Replications	39	107.52**
Locations	12	5,808.54**
Varieties	19	308.00**
Varieties x Locations	228	42.68**
Error	741	15.18

**Highly significant

Table 15. Three-year summary of yield in bushels per acre and yield rank for the strains in the Uniform Test, Group II, 1940-1942.

Strain	Mean of 31 Tests	Lafayette Ind.	Bluffton Ind.	Wahatah Ind.	Urbana Ill.	Mazon or Dwight Ill.	Clayton Ill.*	Stonington Ill.*	Kanawha Iowa	Holgate or Van Wert, Ohio
				<u>Yield, Bu/A</u>						
L6-685	34.2	44.7	37.3	26.5	47.8	38.6	34.0	28.3	27.3	28.8
L6-12	33.9	41.4	36.9	27.9	46.5	39.2	32.9	30.8	28.0	29.1
P.I. 92717	29.9	37.3	31.6	23.8	40.3	33.2	27.3	25.1	27.2	27.1
Mingo	29.8	37.0	30.7	23.0	41.2	31.2	31.1	25.8	26.2	24.6
Illini	29.0	35.7	33.5	24.2	35.9	32.2	27.0	25.7	26.5	22.1
P.I. 68474	28.8	34.7	29.4	25.4	36.5	34.9	27.6	24.9	24.4	26.0
Mandell	28.3	34.2	29.8	19.5	37.8	33.0	28.7	24.9	26.4	22.6
Dunfield	27.9	32.6	28.5	22.4	37.7	31.8	30.9	25.7	25.1	20.5
P.I. 91161	27.9	35.1	27.1	22.2	36.1	32.6	25.7	24.0	26.9	23.7
P.I. 70478	26.8	32.6	27.9	21.9	34.2	28.8	24.3	24.4	25.2	24.7
Mukden	26.1	35.0	27.4	20.8	34.5	30.7	23.1	20.5	24.3	18.0
Richland	26.1	30.9	26.5	20.4	35.1	29.9	27.2	20.6	25.6	20.1
Earlyana	25.1	30.5	25.9	22.0	29.7	27.9	21.2	19.4	27.4	20.8
Wis.Man.3	25.0	30.1	26.0	17.4	31.4	27.6	23.7	20.7	25.5	21.2
Wis.Man.606	24.4	30.0	23.6	19.7	32.2	25.0	21.7	18.1	24.7	22.9
Mean Yield	28.2	34.8	29.5	22.5	37.1	31.8	27.1	23.9	26.0	23.5
				<u>Yield Rank</u>						
L6-685		1	1	2	1	2	1	2	3	2
L6-12		2	2	1	2	1	2	1	1	1
P.I. 92717		3	4	5	4	4	7	6	4	3
Mingo		4	5	6	3	9	3	3	8	6
Illini		5	3	4	9	7	9	4	6	10
P.I. 68474		8	7	3	7	3	6	7	14	4
Mandell		9	6	14	5	5	5	7	7	9
Dunfield		11	8	7	6	8	4	5	12	13
P.I. 91161		6	11	8	8	6	10	10	5	7
P.I. 70478		10	9	10	12	12	11	9	11	5
Mukden		7	10	11	11	10	13	13	15	15
Richland		12	12	12	10	11	8	12	9	14
Earlyana		13	14	9	15	13	15	14	2	12
Wis.Man.3		14	13	15	14	14	12	11	10	11
Wis.Man.606		15	15	13	13	15	14	15	13	8

* Two-year data only, 1940-1941.

Table 16. Two-year summary of chemical composition of two improved strains in comparison with three commercial varieties grown at twelve locations, 1941-1942.

Variety or Strain	Mean of 24 Tests	Colum- bus Ohio	Hol- gate Ohio	Bluff- ton Ind.	Wana- tah Ind.	La- fayette Ind.	North Vernon Ind.	Green- field Ind.	Dwight Ill.	Ur- bana Ill.	Clay- ton Ill.	Stoning- ton Ill.	kana- wha Iowa
L6-685	41.5	42.8	40.4	41.6	43.4	41.0	44.6	41.0	40.5	37.9	42.3	41.5	41.9
L6-12	42.4	42.5	41.7	42.3	44.2	42.0	45.8	42.0	48.8	38.7	43.2	42.7	41.8
Mingo	-	-	42.2	44.0	46.0	44.0	-	-	42.6	40.6	-	-	43.9
Illini	41.7	42.4	39.8	41.9	44.4	41.4	46.0	40.2	41.6	38.1	42.3	40.2	42.3
Dunfield	40.6	41.7	39.6	41.0	44.0	40.5	43.4	38.4	39.9	37.9	40.7	40.1	39.9
Percentage of Oil													
L6-685	21.7	20.7	21.7	21.2	20.7	22.2	20.9	21.7	22.2	23.0	21.9	22.5	21.5
L6-12	21.1	20.5	21.1	21.0	20.5	21.6	20.2	21.3	21.5	22.6	21.1	21.7	20.6
Mingo	-	-	21.1	20.3	19.5	20.2	-	-	21.1	21.6	-	-	20.1
Illini	20.6	19.8	21.1	20.4	19.5	20.9	19.4	21.0	21.1	21.9	20.6	21.5	20.1
Dunfield	21.5	20.8	21.3	20.7	20.3	21.6	20.8	22.3	22.1	23.0	22.3	22.5	21.1
Iodine Number of Oil													
L6-685	134	134	135	132	134	135	132	133	134	134	134	133	136
L6-12	134	135	135	133	134	134	133	133	136	134	134	134	136
Mingo	-	-	131	131	130	132	-	-	133	131	-	-	134
Illini	132	134	134	131	132	132	131	132	133	132	133	131	134
Dunfield	126	128	129	123	126	124	124	126	127	124	126	125	131

Table 17. Summary of chemical composition of two improved strains and three commercial varieties grown at twenty-six locations in 1942.

Variety or Strain	Mean of 26 Tests	Colum- bus Ohio		Hol- gate Ohio		Dear- born Mich.		Bluf- fton Ind.		Wana- tah Ind.		La- Grange Ind.		La- fayette Ind.		Green- field Ind.		North Mt. Morris Ill.		Dwight Ill.
		41.4	41.1	39.4	41.3	42.0	42.1	40.2	40.5	44.7	33.0	40.7								
L6-685	40.1	41.4	41.1	39.4	41.3	42.0	42.1	40.2	40.5	44.7	33.0	40.7								
L6-12	41.5	43.1	41.8	41.4	42.3	42.5	42.7	41.5	40.8	45.5	34.6	43.9								
Mingo	-	44.4	-	41.3	43.7	44.6	42.1	42.9	-	-	35.9	43.1								
Illini	40.7	43.0	41.5	38.9	41.2	43.2	42.2	40.6	38.0	46.6	35.9	43.1								
Dunfield	39.5	42.3	40.4	38.6	40.5	41.7	40.3	39.5	35.5	43.2	33.6	39.8								
<u>Percentage of Protein</u>																				
L6-685	22.2	21.2	21.1	21.9	20.2	21.5	21.6	21.2	22.2	21.4	23.4	22.5								
L6-12	21.4	20.9	20.8	20.8	19.4	21.1	21.3	20.9	21.6	21.4	22.0	21.5								
Mingo	-	20.1	-	21.1	18.8	20.4	20.4	20.5	-	-	21.7	21.4								
Illini	21.0	19.7	19.9	20.9	18.4	20.6	20.2	21.0	21.4	19.2	21.4	21.4								
Dunfield	21.9	20.9	20.8	21.2	19.2	20.9	21.2	21.7	23.0	20.9	21.7	22.3								
<u>Percentage of Oil</u>																				
L6-685	135	136	137	137	138	136	135	136	135	135	138	135								
L6-12	136	136	137	138	139	136	136	137	137	136	139	137								
Mingo	-	132	-	134	133	135	134	135	-	-	136	134								
Illini	134	134	136	136	136	133	134	134	134	131	137	134								
Dunfield	130	128	131	132	130	129	128	127	130	127	133	129								

Table 17. (continued)

Variety or Strain	Ur- bana Ill.	Clay- ton Ill.	Stoning- ton Ill.	Free- burg Ill.	Paris Mo.	St. Joseph Mo.	Car- rollton Mo.	Column- bia Mo.	Iowa- iana- wha Iowa		Chero- kee Iowa		Ames Iowa	Lincoln Neb.
									Minn.	Iowa	Iowa	Iowa		
	Percentage of Protein													
L6-635	36.9	44.0	40.3	41.0	40.4	27.2	29.3	40.6	41.8	40.8	39.6	33.6	41.7	41.3
L6-12	37.2	44.8	42.0	42.3	39.9	41.1	31.6	42.3	43.1	42.5	41.5	39.7	43.4	41.8
Mingo	38.8	-	-	-	40.7	40.6	-	-	42.1	43.8	42.7	40.5	44.4	-
Illini	36.0	42.9	41.0	40.1	40.0	37.3	30.4	44.6	41.1	42.0	41.4	38.4	42.7	40.7
Dunfield	36.5	42.1	39.8	38.6	38.3	38.5	31.6	41.7	41.5	40.3	39.2	38.0	41.3	40.6
	Percentage of Oil													
L6-685	23.1	21.5	22.7	22.4	22.7	24.1	25.9	23.1	20.5	22.0	22.2	22.8	21.4	22.6
L6-12	22.8	20.7	21.8	21.1	22.3	22.3	25.0	21.8	20.0	21.1	21.2	21.9	20.4	21.8
Mingo	22.2	-	-	-	21.2	21.6	-	-	19.3	20.3	20.9	21.4	20.1	-
Illini	22.4	20.5	21.4	21.5	21.8	23.1	24.8	20.2	18.9	20.7	21.1	21.3	20.4	22.3
Dunfield	23.2	21.9	22.5	22.8	23.0	22.8	24.9	21.7	19.5	21.9	22.3	22.6	21.4	22.8
	Iodine Number of Oil													
L6-685	135	135	133	133	134	132	135	134	137	136	137	137	136	134
L6-12	135	136	135	134	135	134	136	134	137	136	137	139	136	133
Mingo	134	-	-	133	133	133	-	-	135	133	135	136	134	-
Illini	133	135	132	130	132	132	127	130	137	136	137	137	134	134
Dunfield	126	129	130	127	129	130	132	129	133	132	134	133	130	128

Uniform Test, Group III

The Group III Test in 1942 was composed of six named varieties, 23 selections from hybrids, and one selection obtained as a rogue. The origin of these varieties and strains is as follows:

Variety or Strain	Source or Originating Agency	Origin
C2	Purdue Agr. Exp. Sta.	Selection from X231 (Dunfield x Midwest)
C56	Purdue Agr. Exp. Sta.	Selection from X331 (Illini x Mandell)
C60	Purdue Agr. Exp. Sta.	Selection from X331 (Illini x Mandell)
C66	Purdue Agr. Exp. Sta.	Selection from X831 (Dunfield x Manchu)
C72	Purdue Agr. Exp. Sta.	Selection from X831 (Dunfield x Manchu)
Dunfield	Purdue Agr. Exp. Sta.	P.I. 36846 1/
Patoka	Purdue Agr. Exp. Sta.	Selection from P.I. 70218-2
L4-12	Illinois Agr. Exp. Sta.	Selection from mixed hybrid population
L4-42	Illinois Agr. Exp. Sta.	Selection from mixed hybrid population
L4-45	Illinois Agr. Exp. Sta.	Selection from mixed hybrid population
L6-12	Illinois Agr. Exp. Sta.	Selection from (Mandarin x Manchu)
L6-685	Illinois Agr. Exp. Sta.	Selection from (Mandarin x Manchu)
L6-690	Illinois Agr. Exp. Sta.	Selection from (Mandarin x Manchu)
L6-700	Illinois Agr. Exp. Sta.	Selection from (Mandarin x Manchu)
L7-1087	Illinois Agr. Exp. Sta.	Selection from X157 (Illini x T48)
L7-1111	Illinois Agr. Exp. Sta.	Selection from X157 (Illini x T57)
L7-1280	Illinois Agr. Exp. Sta.	Unknown
L7-1355	Illinois Agr. Exp. Sta.	Rogue from a plot of P.I.81041
Chief	Illinois Agr. Exp. Sta.	Selection from (Illini x T95)
Illini	Illinois Agr. Exp. Sta.	Selection from A. K.
S32-3	Missouri Agr. Exp. Sta.	Selection from (P.I.37062 x Illini)
S32-8	Missouri Agr. Exp. Sta.	Selection from (P.I.37062 x Illini)
Scioto	Ohio Agr. Exp. Sta.	Selection from Manchu
McClave	Charles McClave	Farmer's selection

1/ Division of Plant Exploration and Introduction, Bureau of Plant Industry,
U. S. D. A.

The Uniform Test, Group III was enlarged in 1942 to include additional promising strains. Among the additional strains entered which are of particular interest because of their previous outstanding performance in Group II were L6-685 and L6-12. Their entry in this group afforded an opportunity to test them still more widely than in Group II alone. In this test, as in Group II, L6-685 was higher in yield than any other variety or strain, and averaged 8.5 bushels per acre above the mean of Illini, Scioto, and Dunfield which are commonly grown varieties of similar maturity. The summary of the 1942 agronomic and chemical data is presented in Table 13.

McClave was the latest maturing variety in the test and was likewise the lowest in yield both in 1942, as shown in Table 13, and for the two-year period, 1941-42, as shown in Table 27. In addition to being low in yield of seed, it is also lower in oil content than any other entry in the test and over 2% lower than Illini. The two-year data, Table 25, also shows McClave to be the lowest in oil content in the test and 2.4% lower than Illini.

Comparisons of varieties grown in this test two and three years are shown in Tables 25 to 26 inclusive. In these periods L6-690 and C2 have been higher in yield than the commercial varieties but are about as low as, or lower than, the lowest of the commercial varieties in oil content.

As in the other tests, there was frost damage, particularly to the later maturing strains and varieties, at some locations. Because of frost damage, maturity data in some cases represents the best estimates of the investigator based upon the stage of development at frost.

Table 18. Summary of agronomic and chemical data for the strains in the Uniform Test, Group III, 1942.

Strain	Mean	Lodg-		Ma-	Seed	Seed	Percent-	Percent-	Iodine
	Yield	ing	Height	tur-	Qual-		age of	age of	
No. of	Bu/A			ity ¹	ity	Size	Protein	Oil	of Oil
Tests	13	14	13	9	14	15 ²	15 ²	15 ²	15 ²
L6-685	35.2	2.4	39	-1.1	1.7	15.0	39.7	22.4	135
L6-12	31.9	2.7	39	-0.7	1.8	14.1	40.1	21.9	136
I4-45	31.8	2.7	37	-2.3	1.9	14.3	42.7	21.0	134
I4-12	31.7	2.7	37	-0.6	1.7	13.9	40.8	21.9	136
I4-42	31.4	2.6	44	+0.8	1.7	13.0	43.0	20.6	138
C56	31.3	2.9	40	-0.4	2.2	13.9	42.0	20.5	135
L7-1280	31.0	2.9	49	+3.9	1.9	13.0	40.2	21.5	135
L6-700	30.9	3.0	39	-0.3	1.7	14.6	44.8	21.5	134
L6-690	30.7	2.7	43	+2.5	2.0	11.8	42.4	20.8	137
L7-1087	30.1	3.1	44	+4.0	1.9	12.1	43.0	20.5	134
L7-1111	30.1	2.6	41	+1.1	2.0	12.5	41.1	20.7	136
Chief	29.6	3.0	47	+6.1	2.1	12.0	39.8	21.5	133
Patoka	29.4	2.6	36	+6.6	2.1	16.8	41.6	21.8	133
C2	29.2	3.1	43	+6.0	2.3	12.7	40.6	21.5	132
C60	29.1	2.6	38	-1.3	1.7	17.3	41.9	21.1	128
C66	28.5	2.1	34	-3.6	2.0	14.8	39.9	22.2	128
Dunfield	27.9	2.9	37	-1.3	1.8	14.2	37.9	22.2	130
C72	27.8	3.1	41	+1.9	2.1	15.4	39.6	21.8	131
L7-1355	27.7	2.8	50	+6.7	2.6	13.5	39.4	22.1	130
Scioto	26.2	3.8	40	+3.6	2.2	13.6	39.6	22.5	156
Illini	26.0	3.3	42	0.0	1.9	12.8	40.4	21.1	134
S32-3	25.3	2.8	44	+0.2	1.6	12.1	41.4	21.1	136
S32-8	23.0	2.8	47	+5.4	2.4	12.0	41.3	21.5	134
McClave	21.5	2.7	34	+9.2	3.1	9.5	42.3	18.9	136

Bu.Nec. 2.7
for Sig. (5% level)

¹ Days earlier (-) or later (+) than Illini. Illini required 122 days to mature.

² Composite sample of 15 locations, composition on dry basis.

Table 20. Summary of yields in bushels per acre for the strains in the Uniform Test, Group III, 1942.

Strain	Mean of 13 Tests ¹	Ur-bana Ill.	Columbus Ohio	Lafayette Ind.	Greenfield Ind.	Ames Iowa	Dwight Ill.	Stonington Ill.
L6-685	35.2	54.8	48.2	41.2	36.8	41.3	38.6	36.4
L6-12	31.9	47.4	44.6	39.2	40.6	31.6	34.7	32.6
L4-45	31.8	51.3	44.4	39.1	30.1	42.6	37.9	36.2
L4-12	31.7	47.0	46.6	38.2	34.6	33.4	37.0	31.6
L4-42	31.4	49.1	39.8	32.7	36.0	32.0	30.8	35.5
C56	31.3	49.7	42.4	38.6	33.9	33.8	33.1	35.6
L7-1280	31.0	49.5	35.3	38.2	33.7	27.9	30.9	31.8
L6-700	30.9	47.5	34.2	40.1	36.3	33.3	34.4	30.9
L6-690	30.7	44.1	43.3	38.0	37.9	31.9	30.5	35.0
L7-1087	30.1	45.7	42.1	34.5	36.6	30.5	28.8	30.6
L7-1111	30.1	48.2	42.5	39.0	35.2	37.1	38.3	32.3
Chief	29.6	46.5	36.2	36.7	34.0	27.0	28.1	31.5
Patoka	29.4	48.4	42.4	36.3	30.4	24.0	27.5	26.1
C2	29.2	53.1	37.8	40.4	30.0	28.8	27.9	26.4
C60	29.1	40.1	41.1	40.6	36.0	38.5	32.8	28.1
C66	28.5	43.5	38.0	36.1	29.4	40.4	32.4	30.1
Dunfield	27.9	48.3	28.0	32.1	26.4	38.0	35.5	32.9
C72	27.8	39.6	32.2	41.0	32.0	29.0	28.2	27.5
L7-1355	27.7	44.6	27.3	40.8	32.2	23.5	24.5	27.5
Scioto	26.2	42.5	27.2	31.2	29.7	28.7	26.3	26.1
Illini	26.0	37.8	26.1	35.1	28.1	33.8	30.9	32.1
S32-3	25.3	36.9	28.8	33.9	29.2	29.8	26.5	32.2
S32-8	23.0	38.9	26.0	35.6	26.6	19.7	20.5	23.1
McClave ²	21.5	31.4	28.0	22.2	28.1	18.6	19.1	17.9
Mean	29.1	45.2	36.8	36.7	32.6	31.7	30.6	30.4
Coef. of Var. (%)	12.4	9.9	14.8	9.7	12.8	14.6	8.4	12.6
Bu. Nec. for Sig. (5% level)	2.7	6.3	7.7	5.0	5.9	6.5	3.6	5.4

¹ Yields at Ames not included in mean because of severe frost damage to some varieties.

² McClave not harvested at Lincoln because of excessive shattering.

Table 20. (continued)

Strain	North Vernon Ind.	Free- burg Ill.	Clay- ton Ill.	Hol- gate Ohio	Colum- bia Mo.	Lin- coln Neb.	Paris Mo.
L6-685	29.0	35.4	34.7	39.8	21.9	23.3	17.1
L6-12	28.9	28.1	31.9	33.1	22.4	19.1	11.9
L4-45	22.4	30.4	29.6	34.4	18.9	25.4	13.7
L4-12	31.2	26.0	29.3	34.5	21.2	21.6	14.0
L4-42	29.1	35.4	25.6	32.1	23.5	23.9	14.4
C56	24.6	31.2	27.0	32.2	19.6	22.9	15.7
L7-1280	30.8	36.3	30.8	24.4	25.5	20.8	14.9
L6-700	36.0	29.0	25.3	31.0	23.8	20.1	12.7
L6-690	28.6	30.8	28.3	26.5	21.3	21.2	13.3
L7-1087	30.8	26.2	26.6	30.3	22.3	21.1	13.6
L7-1111	29.1	24.0	24.3	28.3	15.3	20.1	14.1
Chief	34.0	32.7	29.1	22.4	22.0	18.9	13.0
Patoka	40.7	37.6	24.9	18.1	21.2	15.8	13.2
C2	31.6	32.7	29.0	22.6	21.8	14.7	12.3
C60	31.1	28.3	27.0	24.9	17.9	19.2	11.7
C66	29.4	26.0	21.2	29.5	20.4	23.1	11.4
Dunfield	24.4	26.6	30.7	23.9	17.7	23.0	13.1
C72	29.8	26.1	24.0	23.6	22.3	20.5	14.3
L7-1355	36.2	33.3	29.3	18.3	23.4	13.4	11.7
Scioto	26.0	25.3	24.6	26.0	21.2	19.3	15.4
Illini	24.4	22.5	26.1	24.8	14.4	22.6	13.0
S32-3	22.2	25.2	26.8	17.7	20.0	17.3	11.7
S32-8	24.8	25.6	23.3	9.3	21.0	10.4	14.0
McClave	26.6	25.5	17.9	19.0	12.3	-- ^a	9.9
Mean	29.3	29.3	26.9	26.1	20.4	19.9	13.3
Coef. of Var. (%)	11.6	11.4	9.2	17.9	11.2	14.7	15.7
Bu. Nec. for Sig. (5% level)	4.8	4.7	3.5	6.6	3.4	4.1	2.9

^a Not harvested because of excessive shattering.

Table 21. Summary of lodging notes for the strains in the Uniform Test, Group III, 1942.

Strain	Mean of 14 Tests ¹	Urbana Ill.	Columbus Ohio	Lafayette Ind.	Greenfield Ind.	Ames Iowa	Dwight Ill.	Stonington Ill.
L6-685	2.4	2.0	3.8	2.1	1.4	3.0	3.0	2.0
L6-12	2.7	3.0	4.8	2.4	2.2	4.0	3.0	2.0
L6-45	2.7	3.0	3.3	2.7	2.0	4.0	2.5	3.0
L6-12	2.7	3.0	4.3	2.6	2.2	4.0	3.0	2.0
L6-42	2.6	3.0	5.0	3.0	1.9	3.0	3.0	2.0
C56	2.9	3.0	4.0	2.9	1.9	4.0	4.0	3.0
L7-1280	2.9	3.0	3.5	2.7	2.5	3.0	4.0	2.5
L6-700	3.0	3.0	5.0	3.0	2.2	5.0	4.0	3.0
L6-300	2.7	3.0	4.0	2.9	2.5	3.0	3.5	2.0
L7-1037	3.1	3.5	4.5	3.0	2.2	4.0	4.0	3.0
L7-1111	2.6	2.0	3.5	2.2	2.0	3.0	3.0	3.0
Chief	3.0	3.0	4.0	3.0	2.6	4.0	4.0	3.0
Patoka	2.6	3.0	4.3	2.7	1.6	4.0	4.0	4.0
C2	3.1	3.0	4.8	2.5	2.4	4.0	4.0	4.0
C60	2.6	3.0	4.5	2.9	1.9	3.0	3.0	2.5
C66	2.1	3.0	3.3	2.9	1.4	3.0	3.0	1.0
Dunfield	2.9	2.5	4.8	3.1	1.4	4.0	4.0	2.5
C72	3.1	3.0	5.0	3.7	1.5	5.0	5.0	4.0
L7-1355	2.8	3.0	4.8	2.6	1.9	4.0	4.0	3.0
Scioto	3.8	5.0	5.0	3.7	3.1	4.0	4.0	4.0
Illini	3.3	3.0	4.8	2.9	1.7	5.0	4.0	3.0
S32-3	2.8	3.0	5.0	2.5	1.4	4.0	4.0	2.0
S32-8	2.8	3.0	4.8	2.9	1.9	3.0	4.5	3.0
McClave	2.7	3.0	4.8	2.6	2.1	3.0	2.5	2.5

¹ Carrollton not included in mean since no varieties lodged.

Table 21. (continued)

Strain	North Vernon Ind.	Free- burg Ill.	Clay- ton Ill.	Hol- gate Ohio	Colum- bia Mo.	Paris Mo.	St. Joseph Mo.	Car- rollton Mo.
L6-685	2.6	3.0	3.0	2.8	1.5	1.0	2.0	1.0
L6-12	2.1	3.5	3.0	2.5	2.0	1.0	2.0	1.0
L4-45	3.0	3.5	3.0	2.5	2.5	1.5	2.0	1.0
L4-12	2.7	3.0	2.5	2.5	2.5	1.0	2.0	1.0
L4-42	2.4	2.5	2.0	2.5	2.5	1.5	1.5	1.0
C56	3.1	3.0	3.0	2.8	2.5	1.5	2.0	1.0
L7-1280	2.7	3.0	3.0	2.2	3.0	2.5	2.5	1.0
L6-700	1.9	3.0	3.0	2.5	3.0	1.5	2.0	1.0
L6-690	2.5	2.5	3.0	2.2	3.5	1.5	1.5	1.0
L7-1087	2.9	3.0	3.0	2.0	3.5	1.5	3.0	1.0
L7-1111	2.2	3.0	3.0	2.2	3.5	1.5	3.0	1.0
Chief	4.0	3.5	3.0	2.2	2.5	1.5	2.0	1.0
Patoka	1.0	2.5	4.0	1.8	1.0	1.0	2.0	1.0
C2	2.7	3.0	3.0	2.0	3.5	2.0	2.0	1.0
C60	1.9	2.5	3.0	2.0	1.5	1.5	3.0	1.0
C66	1.5	2.0	2.0	2.8	1.5	1.0	1.5	1.0
Dunfield	2.0	3.5	2.0	2.5	4.0	1.5	3.0	1.0
C72	2.4	3.0	4.0	2.5	2.0	1.5	2.5	1.0
L7-1355	2.0	3.0	3.0	1.5	2.5	1.5	2.0	1.0
Scioto	4.0	4.0	4.0	2.8	4.0	2.5	3.5	1.0
Illini	2.9	3.0	3.0	2.8	4.0	2.5	3.0	1.0
S32-3	2.5	3.0	2.0	2.0	3.0	1.5	3.0	1.0
S32-8	2.2	3.0	2.0	2.2	2.5	2.0	2.5	1.0
McClave	1.0	2.0	3.5	2.5	4.0	2.0	3.0	1.0

Table 22. Summary of plant height for the strains in the Uniform Test, Group III, 1942

Strain	Mean of 13 Tests	Ur- Colum- Lafayette- Green-		North Hol- Colum- Lin-		Paris Joseph rollton		St. Mo.	Car- Mo.					
		bana Ill.	bus Ohio	ette Ind.	field Ind.	Ames Iowa	Dwight Ill.			Vernon Ind.	gate Ohio	coln Ncb.		
L6-685	39	44	60	44	43	48	43	28	34	35	23	31	36	21
L6-12	39	44	52	46	41	49	46	32	39	36	21	24	38	23
L6-45	37	43	52	44	36	49	43	25	49	33	22	25	30	24
L6-12	37	43	52	42	41	49	41	50	55	36	22	26	30	20
L4-42	44	51	59	48	46	56	47	36	37	39	27	31	42	28
056	40	48	54	46	42	54	45	37	53	34	23	30	30	21
L7-1280	49	58	68	55	51	59	55	38	67	42	28	37	48	28
L6-700	39	44	52	45	42	46	45	31	56	35	22	26	36	25
L6-690	43	43	65	51	46	56	51	33	53	38	26	31	40	29
L7-1087	44	55	60	49	50	54	49	34	55	37	24	30	42	27
L7-1111	41	47	53	47	45	53	47	29	55	32	22	28	44	28
Chief	47	55	69	54	51	59	53	40	63	40	26	31	42	30
Patoka	36	40	48	39	37	48	38	30	51	35	19	29	30	22
C2	43	49	60	49	48	54	48	34	56	40	24	28	42	28
C60	38	42	50	45	40	48	44	33	49	33	22	25	32	25
C66	34	41	46	41	36	36	40	22	47	30	19	24	30	24
Dunfield	37	45	50	45	38	46	44	28	50	32	22	27	36	22
C72	41	45	61	45	46	49	46	34	53	33	20	30	32	25
L7-1355	50	59	68	56	52	60	56	45	74	42	27	36	44	28
Scioto	40	47	62	45	45	49	44	33	54	34	22	30	34	22
Illini	42	50	66	48	44	54	47	32	62	31	24	30	40	23
S32-3	44	54	62	48	47	54	48	34	66	36	26	27	38	26
S32-8	47	52	72	53	51	55	53	38	66	40	29	36	42	29
McClave	34	36	51	45	33	42	34	22	52	27	20	26	30	21

Table 23. Summary of maturity notes for the strains in the Uniform Test, Group III, 1942*

Strain	Mean of 9 Tests	Ur-bana Ill.	Colum-bus Ohio	Lafay-ette Ind.	Green-field Ind.	North Vernon Ind.	Colum-bia Mo.	Paris Mo.	St. Joseph Mo.	Car-rollton Mo.
L6-685	-1.1	-3	-1	-3	-2	-1	+3	-1	-2	0
L6-12	-0.7	-2	+1	-3	-2	+1	+3	-2	-2	0
L4-45	-2.3	-2	-4	-3	-3	0	+1	-4	-4	-2
L4-12	-0.6	-1	-1	-3	-2	+1	+3	-2	-2	+2
L4-42	+0.8	+1	-1	+2	-1	+3	+2	+3	-2	0
C56	-0.4	0	-3	0	-1	+2	+1	-1	-2	-2
L7-1280	+3.9	+1	+3	+5	+5	+7	+6	+4	+2	+2
L6-700	-0.3	-1	-3	+3	-2	+2	+2	-2	-2	0
L6-690	+2.5	+10	0	+3	+1	+2	+5	+2	0	+2
L7-1087	+4.0	+1	0	+4	+2	+6	+7	+6	+4	+6
L7-1111	+1.1	+2	-1	-1	0	+2	+2	+4	0	+2
Chief	+6.1	+10	+5	+7	+5	+6	+10	+4	+4	+4
Patoka	+6.6	+10	+4	+9	+7	+7	+10	+8	+2	+2
C2	+6.0	+10	+4	+4	+4	+4	+11	+7	+6	+4
C60	-1.3	0	+1	-3	-2	+1	+1	-2	-4	-4
C66	-3.6	-2	-6	-6	-4	+1	-1	-6	-6	-2
Dunfield	-1.3	-3	-2	-5	-1	-1	+1	+1	-2	0
C72	+1.9	+1	+1	+1	+1	+2	+7	+2	0	+2
L7-1355	+6.7	+10	+7	+7	+6	+8	+6	+6	+4	+6
Scioto	+3.6	+10	+4	+2	+2	0	+7	+3	+2	+2
Illini	0.0	0	0	0	0	0	0	0	0	0
S32-3	+0.2	+1	0	+4	+1	+4	0	-2	-4	-2
S32-8	+5.4	+10	+6	+8	+3	+4	+7	+5	+4	+2
McClave	+9.2	+10	+4	+10	+9	+11	+15	+12	+6	+6
Date Illini matured		9/29	10/9	10/9	10/6	9/20	9/15	9/18	9/28	9/26
Date Planted		5/23	5/29	5/25	5/22	5/29	6/3	5/29	6/3	5/29

* Days earlier (-) or later (+) than Illini.

Table 24. Summary of seed quality notes for the strains in the Uniform Test, Group III, 1942.

Strain	Mean of 14 Tests	Ur- bana Ill.	Colum- bus Ohio	Lafay- ette Ind.	Green- field Ind.	Dwight Ill.	Ston- ington Ill.	North Vernon Ind.
L6-685	1.7	1.0	3.0	1.5	1.5	2.0	1.0	3.0
L6-12	1.8	1.0	2.0	1.5	2.0	2.0	1.0	3.0
L4-45	1.9	2.0	3.0	1.5	2.0	1.0	2.0	3.0
L4-12	1.7	1.0	2.0	1.5	1.5	2.0	1.0	3.0
L4-42	1.7	1.0	2.0	1.5	1.5	2.0	2.0	1.5
C56	2.2	2.0	2.5	1.5	1.5	3.0	2.0	3.0
L7-1280	1.9	1.0	1.5	1.5	1.5	3.0	2.0	2.0
L6-700	1.7	1.0	2.5	1.0	1.0	2.0	2.0	1.5
L6-690	2.0	1.0	2.0	1.5	1.5	3.0	3.0	2.5
L7-1087	1.9	1.0	2.0	1.0	1.0	3.0	2.0	1.5
L7-1111	2.0	1.0	3.0	1.0	1.0	2.0	3.0	3.0
Chief	2.1	1.0	2.0	1.0	1.5	3.	3.0	2.0
Patoka	2.1	1.0	2.0	1.0	1.5	3.0	4.0	1.5
C2	2.3	2.0	1.5	1.0	1.0	3.0	5.0	1.5
C60	1.7	1.0	2.0	1.5	1.0	3.0	2.0	2.5
C66	2.0	1.0	2.0	2.0	1.5	3.0	1.0	3.0
Dunfield	1.8	1.0	2.0	1.5	1.5	1.0	1.0	3.5
C72	2.1	1.0	1.5	1.0	1.0	3.0	4.0	2.0
L7-1355	2.6	2.0	2.0	1.5	1.5	5.0	5.0	2.0
Scioto	2.2	2.0	2.0	1.5	1.5	3.0	4.0	1.5
Illini	1.9	1.0	2.0	1.5	2.0	2.0	2.0	2.5
S32-3	1.6	1.0	1.5	1.0	1.5	2.0	1.0	2.0
S32-8	2.4	2.0	2.0	1.5	1.0	3.0	5.0	1.5
McClave	3.1	2.0	4.0	1.5	1.5	3.0	5.0	1.5

Table 24 (continued)

Strain	Free- burg Ill.	Clay- ton Ill.	Hol- gate Ohio	Colum- bia Mo.	Paris Mo.	St. Joseph Mo.	Car- rollton Mo.
L6-685	1.0	1.0	1.0	1.0	1.0	2.0	3.5
L6-12	1.0	1.0	1.5	1.0	1.0	3.0	4.0
L4-45	1.0	1.0	1.5	2.0	1.0	2.0	4.0
L4-12	1.0	2.0	1.5	1.0	1.0	2.5	2.5
L4-42	1.0	2.0	1.0	1.0	1.0	3.0	3.0
C56	1.0	2.0	3.0	1.5	1.0	3.5	3.0
L7-1280	1.0	2.0	2.5	1.0	1.0	3.5	2.5
L6-700	1.0	2.0	2.0	1.0	1.0	2.5	3.0
L6-690	1.0	1.0	2.0	1.0	1.5	3.5	4.0
L7-1087	1.0	1.0	2.0	2.0	2.0	3.5	4.0
L7-1111	1.0	2.0	2.5	1.5	1.0	3.5	3.0
Chief	1.0	2.0	1.5	1.0	1.5	4.5	4.0
Patoka	1.0	2.0	2.0	1.0	2.0	4.0	3.0
C2	2.0	2.0	2.5	2.0	2.0	4.5	2.5
C60	1.0	2.0	2.0	1.0	1.0	2.0	2.0
C66	2.0	1.0	2.5	2.0	2.5	1.5	3.5
Dunfield	1.0	1.0	2.0	1.5	2.0	3.0	3.0
C72	1.0	2.0	3.0	2.0	1.0	4.0	3.0
L7-1355	2.0	2.0	3.0	1.0	2.0	3.0	4.0
Scioto	1.0	4.0	2.0	1.0	1.0	3.5	3.0
Illini	1.0	1.0	2.0	2.0	1.5	3.0	2.5
S32-3	1.0	1.0	3.5	1.0	1.0	2.5	2.5
S32-8	2.0	2.0	4.0	1.0	1.0	3.0	2.0
McClave	2.0	4.0	3.0	3.0	3.0	3.0	3.0

Table 25. Two-year summary of mean agronomic and chemical data for the strains in the Uniform Test, Group III, 1941-1942.

Strain	Mean Yield Bu/A	Lodging	Height Inches	Maturity ¹	Seed Quality	Seed Size	Percentage Protein	Percentage Oil	Iodine No. of Oil
No. of Tests	25	20	17	18	25	27 ²	27 ²		27 ²
C2	28.4	3.0	40	+6.5	2.3	13.5	41.2	21.0	131
L6-690	28.3	2.4	40	+2.2	1.8	12.1	42.9	20.7	136
L7-1355	27.7	2.6	41	+3.8	1.8	12.6	42.9	20.3	133
Patoka	27.0	2.0	33	+6.3	2.3	17.2	42.7	21.6	132
Chief	26.8	2.6	44	+5.2	2.2	12.5	40.1	21.2	132
L7-1355	26.8	2.5	46	+6.4	2.4	14.2	40.4	21.6	128
Scioto	25.5	3.5	36	+3.0	2.3	14.2	40.8	22.0	135
Dunfield	24.4	2.4	34	-1.2	2.0	15.0	39.2	21.9	126
Illini	23.8	2.6	38	0	2.2	13.2	41.0	20.8	132
McClave	21.5	2.3	30	+11.1	2.8	10.4	43.2	18.4	135

Table 26. Three-year summary of mean agronomic and chemical data for the strains in the Uniform Test, Group III, 1940-1942.

Strain	Mean Yield Bu/A	Lodging	Height Inches	Maturity ²	Seed Quality	Seed Size	Percentage Protein	Percentage Oil	Iodine No. of oil
No. of Tests	35	27	21	27	35	37 ²	37 ²	37 ²	37 ²
L6-690	27.7	2.3	36	+2.8	1.7	12.2	42.9	20.5	135
C2	27.0	2.7	36	+6.5	2.0	13.8	41.7	20.6	130
L7-1355	26.7	2.2	42	+5.7	2.1	14.6	40.7	21.2	127
Patoka	26.4	1.7	31	+6.6	2.0	17.3	43.1	21.1	130
Chief	26.1	2.5	39	+4.4	1.8	12.7	40.5	21.0	132
Scioto	25.1	3.3	33	+2.7	2.1	14.4	41.5	21.5	134
Dunfield	23.4	2.1	32	-1.4	1.8	15.3	39.9	21.5	127
Illini	23.3	2.2	33	0	2.0	13.3	41.4	20.6	131

¹ Days earlier (-) or later (+) than Illini. Illini required 121 days to mature.

² Mean of composite samples by years, composition on dry basis, 15 tests in 1942, 12 tests in 1941, and 10 tests in 1940.

Table 27. Two-year summary of yield in bushels per acre and yield rank for the strains in the Uniform Test, Group III. 1941-1942.

Strain	Mean Yield 25 Tests	Fortville		North Vernon Ind.	Urbana Ill.	Dwight Ill.	Clayton Ill.	Ston- ington Ill.	Free- burg Ill.	Colum- bia Ill.
		Colum- bus Ohio	or Greenville Ind.							
C2	28.4	28.6	27.9	24.2	50.8	37.6	35.1	27.2	30.1	21.7
L6-690	28.3	31.8	30.2	20.4	44.4	38.5	30.7	32.6	27.4	21.4
L7-1087	27.7	30.6	30.6	21.8	45.4	35.2	30.8	30.1	27.2	21.9
Patoka	27.6	32.2	26.0	29.4	47.7	33.8	29.5	26.6	32.4	20.2
Chief	26.8	27.4	29.1	23.1	45.2	36.0	20.9	29.2	26.2	20.2
L7-1355	26.8	24.6	27.6	23.7	45.0	35.2	31.0	28.0	29.9	22.2
Scioto	25.5	23.6	28.4	21.4	39.2	33.6	30.2	27.0	24.6	19.8
Dunfield	24.4	22.9	22.2	18.5	41.2	35.3	30.8	29.3	23.4	16.4
Illini	23.8	22.7	26.0	18.3	36.2	37.1	26.8	29.6	18.4	14.8
McClave	21.5	23.2	23.2	20.6	32.0	27.4	25.6	20.0	25.6	11.9
					<u>Yield Bu/A</u>					
C2		4	5	2	<u>Yield Rank</u>	2	1	7	2	3
L6-690		2	2	8	1	1	6	1	4	4
L7-1087		3	1	5	3	6	4	2	5	2
Patoka		1	7	1	2	8	8	9	1	5
Chief		5	3	4	4	4	3	5	6	5
L7-1355		6	6	3	5	6	2	6	3	1
Scioto		7	4	6	8	9	7	8	8	7
Dunfield		9	10	9	7	5	4	4	9	8
Illini		10	7	10	9	3	9	3	10	9
McClave		8	9	7	10	10	10	10	9	10

Table 28. Three-year summary of yield in bushels per acre and yield rank for the strains in the Uniform Test, Group III, 1940-1942.

Strain	Mean Yield 35 Tests	Colum- bus Ohio	North Vernon Ind.	Yield in Bu/A		Clay- ton Ill.	Ston- ington Ill.	Colum- bia Mo.
				Ur- bana Ill.	Dwight or Mazon Ill.			
L6-690	27.7	27.5	21.6	43.1	33.1	31.4	31.5	21.6
C2	27.0	25.5	24.2	44.5	33.0	33.7	25.9	21.6
L7-1355	26.7	23.4	25.0	42.4	31.5	31.1	27.4	22.9
Patoka	26.4	27.9	28.0	42.5	29.6	28.4	26.1	20.9
Chief	26.1	25.3	23.5	40.8	31.7	30.2	28.1	21.3
Scioto	25.1	21.9	25.7	37.6	29.8	29.3	26.2	21.2
Dunfield	23.4	20.2	18.8	36.3	29.9	29.6	27.2	19.1
Illini	23.3	21.9	20.5	33.3	32.0	27.0	27.9	14.7
				Yield Rank				
L6-690		2	6	2	1	2	1	2
C2		3	3	1	2	1	8	2
L7-1355		5	2	4	5	3	4	1
Patoka		1	1	3	8	7	7	6
Chief		4	5	5	4	4	2	4
Scioto		6	4	6	7	6	6	5
Dunfield		8	8	7	6	5	5	7
Illini		6	7	8	3	8	3	8

Table 29. Analysis of variance for yield of seed for the Uniform Test, Group III, 1942.

Source of Variation	Degrees of Freedom	Mean Squares
Replications	39	1,344.68**
Locations	12	6,563.72**
Varieties	23	503.29**
Varieties x Locations	276	50.95**
Error	897	13.05

** Highly significant

Uniform Test, Group IV

The Group IV Test in 1942 was composed of six named varieties, seventeen selections from hybrids, and one selection obtained as a rogue. The origin of these varieties and strains is as follows:

Variety or Strain	Source or Originating Agency	Origin
C2	Purdue Agr. Exp. Sta.	Selection from X231 (Dunfield x Midwest)
C6	Purdue Agr. Exp. Sta.	Selection from X331 (Illini x Landell)
C146	Purdue Agr. Exp. Sta.	Selection from X231 (Dunfield x Midwest)
C148	Purdue Agr. Exp. Sta.	Selection from X231 (Dunfield x Midwest)
C149	Purdue Agr. Exp. Sta.	Selection from X231 (Dunfield x Midwest)
C153	Purdue Agr. Exp. Sta.	Selection from X231 (Dunfield x Midwest)
C154	Purdue Agr. Exp. Sta.	Selection from X231 (Dunfield x Midwest)
C155	Purdue Agr. Exp. Sta.	Selection from X231 (Dunfield x Midwest)
C160	Purdue Agr. Exp. Sta.	Selection from X331 (Illini x Mandell)
C175	Purdue Agr. Exp. Sta.	Selection from X731 (Dunfield x Man.Sel.31)
C178	Purdue Agr. Exp. Sta.	Selection from X831 (Dunfield x Man.Sel.21)
Gibson	Purdue Agr. Exp. Sta.	Selection from X531 (Midwest x Dunfield)
Patoka	Purdue Agr. Exp. Sta.	Selection from P.I. 70218-2
L7-923	Illinois Agr. Exp. Sta.	Selection from mixed hybrid population
EV-1160	Illinois Agr. Exp. Sta.	Selection from LX157 (Illini x T148)
Chief	Illinois Agr. Exp. Sta.	Selection from (Illini x Manchu)
Macoupin	Elmor Hulcher	Selection from commercial lot
Morse	U.S.Dept.of Agriculture	P.I. 19186
S32-11	Missouri Agr. Exp. Sta.	Selection from (P.I.37062 x Illini)
S49-5	Missouri Agr. Exp. Sta.	Selection from (Virginia x P.I.54610-3)
S49-12	Missouri Agr. Exp. Sta.	Selection from (Virginia x P.I.54610-3)
S49-18	Missouri Agr. Exp. Sta.	Selection from (Virginia x P.I.54610-3)
S100	Missouri Agr. Exp. Sta.	Rogue from a plot of Illini.
Boone	Missouri Agr. Exp. Sta.	Selection from P.I.54563-3

All of the higher yielding strains or varieties in this group in 1942 were of hybrid origin. Likewise, the earlier strains, as a rule, were among the highest in yield. This is probably due to the fact that yields of the later maturing strains were lowered by the extremely early frost at most locations in relation to the amount of frost damage sustained, and would support the contention that too much weight should not be placed on one year's results alone.

A remarkable amount of local adaptation is also evident in the 1942 results. Varieties or strains best at one location, or even one state, were often poorest at another. This, too, may be due in part to the frost damage to some varieties. On the other hand, it would seem to show that there is considerable differential varietal response in soybeans to soil and climatic conditions encountered at the various locations.

Many of the strains in this test show practically no improvement in yield or chemical composition when compared to the more recently released varieties in the two- and three-year summaries and will therefore be eliminated from the test.

Table 30. Summary of agronomic and chemical data for the strains in the Uniform Test, Group IV, 1942.

Variety or Strain	Mean Yield Bu./A.	Lodging	Height in Inches	Maturity ¹	Seed Quality	Seed Size	Percent- age of Protein	Percent- age of Oil	Iodine Number of Oil
Number of Tests	10	10	9	7	11	11 ²	11 ²	11 ²	11 ²
Patoka	29.1	2.2	35	-5.5	2.1	16.4	41.8	21.9	132
C2	28.9	3.2	43	-3.4	2.2	12.8	40.8	21.6	131
C155	28.3	2.8	45	-0.6	1.8	14.5	40.4	22.1	130
Chief	28.2	2.8	47	-4.3	2.0	12.1	39.5	21.8	131
C175	27.1	3.2	45	-1.6	2.3	13.9	39.8	21.7	130
C146	27.0	2.6	43	-0.1	2.1	23.2	38.5	22.4	127
C149	26.9	2.7	43	-1.3	1.9	12.7	39.7	22.4	132
C6	26.6	2.7	43	-4.6	2.0	12.7	42.1	21.0	132
Gibson	26.1	3.4	39	0.0	1.8	12.9	39.2	21.7	133
C160	25.6	2.7	41	-4.0	1.9	13.5	40.9	21.6	132
L7-923	25.5	2.7	46	-1.4	2.2	11.6	38.3	21.9	131
L7-1160	25.3	2.7	45	-1.1	2.2	13.3	40.4	21.8	131
C178	25.0	2.7	43	+1.9	2.5	14.3	40.5	20.9	132
C154	24.8	3.1	40	-3.6	2.3	13.7	39.2	22.4	131
C153	24.6	3.2	43	-0.9	2.4	13.5	40.1	22.0	130
Maccupin	24.2	3.1	43	-3.6	2.3	14.3	39.4	22.4	131
S49-18	23.9	3.0	42	-1.4	2.0	11.7	39.6	21.4	134
Morso	23.8	3.6	42	-1.4	2.5	15.7	40.6	21.2	134
Boone	23.2	2.7	42	-0.1	2.0	12.6	40.0	22.1	131
S49-5	23.0	3.2	46	+0.3	2.1	12.1	37.9	21.8	133
C148	22.7	3.4	44	-2.6	2.2	13.8	39.5	21.9	128
S32-11	22.3	2.9	43	-1.0	1.4	10.4	39.5	21.7	135
S100	22.2	3.0	45	+5.6	2.5	10.9	40.3	20.1	132
S49-12	22.0	2.6	41	-5.9	1.6	11.5	41.5	20.9	135

Bu.Nec. 3.0
for Sig. (5% level)

¹ Days earlier (-) or later (+) than Gibson. Gibson required 130 days to mature.

² Composite sample of eleven tests, composition on dry basis.

Table 31. Summary of yields in bushels per acre for the strains in the Uniform Test, Group IV, 1942.

Strain	Mean of 10 Tests	Evans-	Ur-	North	Free-	Clay-	Wheat-	Els-	Ston-	Sikes-	Colum-
		ville Ind.	bana Ill.	Vernon Ind.	burg Ill.	ton Ill.	land Ind.	berry Mo.	ington Ill.	ton Mo.	bia Mo.
Patoka	29.1	41.2	45.4	34.1	36.8	25.8	26.0	20.2	25.4	16.9	20.3
C2	28.9	47.4	49.2	31.6	32.0	25.4	22.2	21.7	22.0	17.3	24.0
C155	28.3	45.4	46.7	28.4	32.7	25.8	21.4	24.5	20.4	21.4	19.4
Chief	28.2	41.5	36.6	31.1	31.5	27.9	28.8	26.0	26.8	19.0	15.4
C175	27.1	43.4	43.0	30.0	29.2	22.8	24.8	22.6	20.3	18.5	22.3
C146	27.0	44.6	49.8	28.0	27.8	23.4	19.1	20.7	16.5	23.4	20.6
C149	26.9	40.6	40.4	29.7	29.8	23.9	20.8	26.5	19.8	18.2	21.0
C6	26.6	39.8	44.5	29.2	25.8	22.3	26.7	23.7	22.7	18.2	18.0
Gibson	26.1	44.0	42.9	27.0	24.4	24.3	25.4	18.7	19.1	21.2	18.1
C160	25.6	36.7	39.9	29.4	27.8	24.0	22.2	21.5	22.6	18.7	16.4
L7-923	25.5	43.0	37.8	29.6	27.5	20.3	19.4	22.6	16.1	21.9	18.6
L7-116C	25.3	37.2	38.1	24.0	24.7	21.4	22.4	24.2	19.6	20.1	23.5
C178	25.0	40.6	38.9	29.8	27.6	20.0	20.6	19.0	17.8	23.9	14.7
C154	24.8	40.6	31.0	29.6	22.7	22.5	24.6	17.7	22.2	18.1	20.3
C153	24.6	41.6	31.0	32.2	26.9	20.0	23.2	18.2	16.5	21.7	16.0
Macopin	24.2	38.2	28.1	25.5	26.0	24.5	22.8	18.5	22.4	17.8	18.5
S49-18	23.9	33.0	31.2	26.0	26.3	23.5	20.6	24.9	16.2	19.4	19.4
Morse	23.8	31.1	30.6	25.9	26.6	20.2	22.6	24.3	21.1	17.7	18.9
Boone	23.2	32.2	36.2	22.1	22.0	21.6	20.2	20.0	17.9	19.9	22.8
S49-5	23.0	31.4	29.5	23.0	22.4	22.7	21.6	23.8	18.8	18.3	18.1
C148	22.7	39.5	28.1	24.7	20.6	22.3	19.8	19.6	17.6	16.5	18.6
S32-11	22.3	33.1	31.4	19.2	21.5	20.3	20.8	20.2	20.6	20.0	18.8
S100	22.2	41.5	31.3	23.8	22.4	18.2	19.1	17.1	10.0	24.9	15.5
S49-12	22.0	29.9	30.3	18.3	21.8	23.4	23.9	13.2	27.1	16.4	18.6
Mean	25.3	39.1	37.2	27.2	26.5	22.8	22.5	21.3	20.0	19.5	19.1
Coef. of Var.	13.7	9.9	16.3	12.9	15.3	8.4	11.2	18.4	14.7	8.2	16.0
Du. Nec. for Sig.	3.0	5.3	10.0	5.0	5.7	2.7	4.1	5.5	4.2	2.2	4.3

Table 32. Yield rank for the strains in the Uniform Test, Group IV, 1942.

Strain	Evans- ville Ind.	Ur- bana Ill.	North Vernon Ind.	Free- burg Ill.	Clay- ton Ill.	Wheat- land Ind.	Els- berry Mo.	Ston- ington Ill.	Sikes- ton Mo.	Colum- bia Mo.
Patoka	10	4	1	1	2	3	14	3	22	7
C2	1	2	3	3	4	12	11	8	21	1
C155	2	3	12	2	2	15	5	11	6	9
Chief	8	13	4	4	1	1	2	2	12	23
C175	5	6	5	6	12	5	10	12	14	4
C146	3	1	13	7	10	23	13	20	5	6
C149	11	8	7	5	8	16	1	13	16	5
C6	14	5	11	15	15	2	8	4	17	19
Gibson	4	7	14	17	6	4	19	15	7	18
C160	18	9	10	7	7	12	12	5	13	20
L7-923	6	12	8	10	19	22	9	23	4	15
L7-1160	17	11	19	16	18	11	7	14	8	2
C178	11	10	6	9	23	18	18	18	2	24
C154	11	18	8	18	14	6	22	7	18	8
C153	7	18	2	11	23	8	21	20	5	21
Macoupin	16	23	17	14	5	9	20	6	19	16
S49-18	20	17	15	13	9	18	4	22	11	9
Morse	23	20	16	12	22	10	6	9	20	11
Boone	21	14	22	21	17	20	16	17	10	3
S49-5	22	22	21	19	13	14	3	16	15	17
C148	15	24	18	24	15	21	17	19	23	14
S32-11	19	15	23	23	19	16	15	10	9	12
S100	8	16	20	19	24	23	23	24	1	22
S49-12	24	21	24	22	10	7	24	1	24	13

Table 33. Summary of lodging notes for the strains in the Uniform Test, Group IV, 1942.

Strain	Mean of 10 Tests ¹		Evansville Ind.		Ur-bana Ill.		North Vernon Ind.		Freeburg Ill.		Clayton Ill.		Wheatland Mo.		Elsberry Mo.		Stonington Ill.		Sikes-ton Mo.		Columbia Mo.		Carrollton Mo.	
Patoka	2.2	2.2	3.0	3.0	1.0	1.0	2.0	3.0	3.0	1.0	1.0	3.5	3.5	1.0	1.0	3.5	3.5	3.5	3.5	1.5	1.5	1.0	1.0	
C2	3.2	3.0	3.0	4.0	3.1	3.1	3.0	4.0	4.0	1.5	1.5	4.0	4.0	4.0	4.0	4.0	5.0	5.0	4.0	4.0	4.0	1.0	1.0	
C155	2.8	2.6	3.0	3.5	3.2	3.2	3.0	3.5	3.5	1.0	1.5	3.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.5	3.5	1.0	1.0	
Chief	2.8	2.4	3.0	3.5	3.1	3.1	3.0	3.5	3.5	1.3	1.5	3.0	4.0	4.0	4.0	4.0	5.0	5.0	4.0	3.0	3.0	1.0	1.0	
C175	3.2	3.0	3.0	3.5	3.0	3.0	3.5	3.5	3.5	2.0	2.0	3.5	4.0	4.0	4.0	4.0	5.0	5.0	4.0	3.0	3.0	1.0	1.0	
C146	2.6	2.6	3.0	3.0	1.7	1.7	3.0	3.0	3.0	1.2	2.0	3.0	3.0	4.0	4.0	4.0	4.0	4.0	3.0	3.0	3.0	1.0	1.0	
C149	2.7	2.9	3.0	3.0	2.9	2.9	4.0	3.0	3.0	1.2	1.0	3.0	3.5	3.0	3.0	3.5	3.5	3.0	3.5	2.5	2.5	1.0	1.0	
C6	2.7	3.2	3.0	3.0	2.7	2.7	3.5	3.0	3.0	1.0	1.0	3.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	2.5	2.5	1.0	1.0	
Gibson	3.4	2.9	4.0	4.0	2.6	2.6	3.0	4.0	4.0	1.7	3.5	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	1.0	1.0	
C160	2.7	2.9	3.0	3.0	2.2	2.2	3.0	3.0	3.0	1.0	1.5	3.0	3.0	3.0	3.0	3.0	5.0	5.0	4.0	2.0	2.0	1.0	1.0	
L7-923	2.7	2.6	3.0	3.0	2.6	2.6	3.0	3.0	3.0	1.7	1.0	3.0	3.0	3.0	3.0	4.0	4.0	4.0	4.0	3.5	3.5	1.0	1.0	
L7-1160	2.7	2.7	3.0	3.0	3.1	3.1	3.0	3.0	3.0	1.6	1.0	3.0	3.0	3.0	3.0	4.0	4.0	4.0	4.0	3.0	3.0	1.0	1.0	
C178	2.7	3.0	3.0	3.0	2.5	2.5	3.0	3.0	3.0	2.0	1.0	3.0	3.0	3.0	3.0	3.5	3.5	3.0	3.0	3.0	3.0	1.0	1.0	
C154	3.1	3.2	4.0	4.0	2.1	2.1	4.0	4.0	4.0	1.2	1.0	4.0	4.0	4.0	4.0	4.0	5.0	5.0	4.0	2.5	2.5	1.0	1.0	
C153	3.2	2.9	4.0	4.0	2.5	2.5	4.0	2.5	2.5	1.7	2.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	1.0	1.0	
Macoupin	3.1	3.1	3.0	4.0	2.9	2.9	3.0	4.0	4.0	1.0	2.0	3.0	3.0	3.0	3.0	3.0	5.0	5.0	4.0	4.0	4.0	1.0	1.0	
S49-18	3.0	3.2	3.0	3.5	3.2	3.2	3.0	3.5	3.5	2.0	2.0	3.0	3.0	3.0	3.0	4.0	4.0	4.0	4.0	3.0	3.0	1.0	1.0	
Morse	3.6	3.1	4.0	4.0	3.5	3.5	3.0	4.0	4.0	2.0	3.0	4.0	4.0	4.0	4.0	4.0	5.0	5.0	4.0	4.0	4.0	1.0	1.0	
Boone	2.7	3.4	3.0	3.0	2.2	2.2	3.0	3.0	3.0	1.2	1.0	3.0	3.0	3.0	3.0	4.0	4.0	4.0	4.0	3.5	3.5	1.0	1.0	
S49-5	3.2	3.1	4.0	4.0	3.5	3.5	3.5	3.0	3.0	2.0	1.5	3.0	3.0	3.0	3.0	4.0	4.0	4.0	4.0	4.0	4.0	1.0	1.0	
C148	3.4	3.5	4.0	4.0	3.0	3.0	4.0	4.0	4.0	1.2	2.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	1.0	1.0	
S32-11	2.9	3.1	3.0	3.0	3.0	3.0	3.0	3.0	3.0	2.0	2.0	3.0	3.0	3.0	3.0	4.0	4.0	4.0	4.0	3.0	3.0	1.0	1.0	
S49-12	2.6	3.5	3.0	3.0	2.9	2.9	3.0	2.0	2.0	1.0	1.5	2.0	2.0	2.0	2.0	3.0	3.0	3.0	3.0	2.0	2.0	1.0	1.0	
S100	3.0	3.0	4.0	4.0	3.4	3.4	4.0	3.0	3.0	2.0	1.5	4.0	4.0	4.0	4.0	4.0	3.0	3.0	3.0	2.0	2.0	1.0	1.0	

¹Carrollton not included in the mean since no varieties lodged.

Table 34. Summary of plant height for the strains in the Uniform Test, Group IV, 1942

Strain	Mean of 9 Tests	Evans- ville Ind.	Ur- bana Ill.	North Vernon Ind.	Free- burg Ill.	Clay- ton Ill.	Wheat- land Ind.	Els- berry Mo.	Colum- bia Mo.	Car- rollton Mo.
Patoka	35	41	45	30	40	43	33	29	33	23
C2	43	49	54	38	50	53	42	34	38	31
C155	45	56	54	38	47	54	46	42	41	29
Chief	47	58	58	37	54	57	48	38	39	32
C175	45	57	55	36	50	54	43	35	41	33
C146	43	52	51	37	48	51	42	39	38	30
C149	43	52	52	36	50	52	42	35	40	31
C6	43	47	53	35	50	53	43	35	42	32
Gibson	39	49	45	33	42	45	40	35	33	25
C160	41	50	50	33	51	50	41	36	36	26
L7-923	46	58	57	39	48	56	47	40	41	32
L7-1160	45	53	51	38	54	50	44	37	42	37
C178	43	51	51	39	51	50	43	36	39	31
C154	40	50	48	33	51	48	40	33	36	25
C153	43	50	50	41	47	50	44	37	38	29
Macoupin	43	53	52	30	48	51	41	35	39	36
S49-18	42	54	50	34	50	50	40	37	36	25
Morse	42	51	50	32	46	49	41	37	40	29
Boone	42	51	50	31	47	50	40	36	40	30
S49-5	46	58	53	37	51	53	45	38	43	32
C148	44	53	54	39	50	53	43	36	36	33
S32-11	43	55	54	37	48	53	40	34	38	26
S100	45	56	56	38	48	55	44	36	40	34
S49-12	41	51	48	32	45	49	41	38	36	29

Table 35. Summary of maturity notes for the strains in the Uniform Test, Group IV, 1942.*

Strain	Mean of 7 Tests	Evans- ville Ind.	North Vernon Ind.	Wheat- land Ill.	Els- berry Mo.	Sikes- ton Mo.	Colum- bia Mo.	Car- rollton Mo.
Patoka	-3.3	-5	-4	-4	-3	-8	0	+1
C2	-3.4	-8	-7	-6	-3	-8	+5	+3
C155	-0.6	-1	-2	-5	-2	-2	+6	+2
Chief	-4.3	-5	-4	-6	-6	-10	0	+1
C175	-1.6	-1	-3	-1	-2	-10	+1	+5
C146	-0.1	-1	-2	-1	0	-2	+5	0
C149	-1.3	-1	-3	-1	-3	-2	0	+1
C6	-4.6	-9	-7	-8	-3	-6	0	+1
Gibson	0.0	0	0	0	0	0	0	0
C160	-4.0	-4	-5	-7	-5	-6	-1	0
L7-923	-1.4	-4	-5	0	-1	-8	+4	+4
L7-1160	-1.1	-1	-3	-3	-2	-4	+3	+2
C178	+1.9	0	-2	+2	0	0	+6	+7
C154	-3.6	-4	-2	-6	-5	-6	-2	0
C153	-0.9	-4	-2	-3	-2	-4	+6	+3
Macoupin	-3.6	-7	-6	-5	-4	-8	+3	+2
S49-18	-1.4	+1	-6	-1	-6	0	0	+2
Horse	-1.4	-2	-5	-3	-5	0	+3	+2
Boone	-0.1	+3	-3	-2	-2	0	+2	+1
S49-5	+0.3	+4	-1	+1	-7	-2	+5	+2
C148	-2.6	-4	-3	-4	-5	-6	+4	0
S32-11	-1.0	+5	-4	0	-7	0	-1	0
S100	+5.6	+8	+6	+6	+2	+2	+6	+9
S49-12	-5.9	-2	-6	-5	-10	-10	-8	0
Date Gibson Matured		10/4	9/30	10/6	9/26	9/28	9/26	9/28
Date Planted		5/14	5/29	5/28	5/13	5/11	6/3	5/29

*Days earlier (-) or later (+) than Gibson.

Table 36. Summary of seed quality notes for the strains in Uniform Test, Group IV, 1942.

Strain	Mean of 11 Tests		Evansville Ind.		North Vernon Ind.		Freeburg Ill.		Clayton Ill.		Wheatland Ind.		Elsberry Mo.		Stonington Ill.		Sikes-ton Mo.		Columbia Mo.		Carrollton Mo.	
Patoka	2.1	3.0	1.0	1.5	1.5	1.0	2.0	2.0	2.0	2.5	1.0	4.0	3.0	1.0	3.0	1.0	3.5					
C2	2.2	2.0	1.0	1.5	1.5	1.0	3.0	3.0	3.0	1.5	1.0	5.0	3.0	2.0	3.0	2.0	3.0					
Cl55	1.8	2.0	1.0	2.0	2.0	1.0	1.0	2.0	1.0	2.0	1.0	3.0	2.0	1.0	2.0	1.0	4.0					
Chief	2.0	2.5	1.0	1.5	1.5	1.0	2.0	2.0	2.0	2.0	1.0	4.0	2.0	1.0	2.0	1.0	4.0					
Cl75	2.3	2.0	1.0	2.0	2.0	1.0	2.0	2.0	2.0	3.0	1.5	4.0	3.0	1.0	3.0	1.0	5.0					
Cl46	2.1	1.5	1.0	2.0	2.0	1.0	2.0	2.0	2.0	3.0	1.0	5.0	3.0	1.0	3.0	1.0	2.5					
Cl49	1.9	2.0	1.0	1.5	1.5	1.0	2.0	2.0	2.0	1.5	1.0	5.0	2.0	1.0	2.0	1.0	3.0					
C6	2.0	1.5	1.0	1.0	1.0	1.0	3.0	3.0	3.0	1.5	1.0	5.0	3.0	1.5	3.0	1.5	3.0					
Gibson	1.8	2.5	1.0	2.0	2.0	1.0	2.0	2.0	2.0	1.5	1.0	3.0	2.5	1.0	2.5	1.0	2.5					
Cl60	1.9	1.5	1.0	1.5	1.5	2.0	1.0	1.0	1.0	2.0	1.0	4.0	2.0	1.0	2.0	1.0	3.5					
L7-923	2.2	1.5	1.0	1.5	1.5	1.0	3.0	3.0	3.0	2.0	2.0	4.0	2.0	2.0	2.0	2.0	4.5					
L7-1160	2.2	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	1.0	5.0	2.0	1.0	2.0	1.0	3.5					
Cl78	2.5	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	3.0	3.0	3.0	2.0	2.0	2.0	2.0	4.0					
Cl54	2.3	1.5	1.0	1.5	1.5	1.0	2.0	2.0	2.0	3.0	1.0	3.0	3.0	1.0	3.0	1.0	2.5					
Cl53	2.4	1.5	1.0	2.0	2.0	1.0	3.0	3.0	3.0	3.0	1.0	5.0	2.5	2.5	2.5	2.5	4.0					
Macoupin	2.3	2.0	1.0	2.0	2.0	1.0	2.0	2.0	2.0	2.5	2.0	4.0	3.0	1.0	3.0	1.0	4.5					
S49-18	2.0	2.5	1.0	1.5	1.5	1.0	2.0	2.0	2.0	2.5	1.0	4.0	3.0	1.0	3.0	1.0	3.0					
Morse	2.5	3.5	1.0	2.5	2.5	1.0	2.0	2.0	2.0	3.0	1.0	3.0	3.0	2.0	3.0	2.0	5.0					
Boone	2.0	2.0	1.0	2.0	2.0	1.0	2.0	2.0	2.0	1.5	1.0	4.0	3.0	1.0	3.0	1.0	3.0					
S49-5	2.1	3.0	1.0	2.0	2.0	1.0	2.0	2.0	2.0	2.0	1.0	4.0	2.5	1.5	2.5	1.5	3.5					
Cl48	2.2	2.0	1.0	2.5	2.5	1.0	2.0	2.0	2.0	2.5	1.0	4.0	3.5	1.5	3.5	1.5	5.0					
S32-11	1.4	1.5	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	3.0	1.5	1.0	1.5	1.0	2.5					
Sl00	2.5	2.0	2.0	2.0	2.0	2.0	3.0	3.0	3.0	1.5	3.0	5.0	1.5	1.5	1.5	1.5	4.0					
S49-12	1.6	1.5	1.0	1.5	1.5	1.0	1.0	1.0	1.0	2.0	1.0	2.0	3.0	1.0	3.0	1.0	2.5					

Table 37. Two-year summary of mean agronomic and chemical data for the strains in the Uniform Test, Group IV, 1941-1942.

Strain	Mean		Height		Seed		Percent-	Percent-	Iodina-
	Yield	Lodg-	in	Matu-	Qual-	Seed	age of	age of	Number
	Bu./A.	ing	Inches	urity ¹	ity	Size	Protein	Oil	of Oil
Number of Tests	21	17	15	15	22	22 ²	22 ²	22 ²	22 ²
C155	26.3	2.6	43	-0.3	2.3	15.3	41.9	21.3	128
C146	26.1	2.4	40	+0.5	2.4	14.0	40.1	21.7	126
Patoka	26.0	1.8	32	-3.2	2.7	16.8	43.1	21.4	131
C149	25.2	2.6	40	-1.2	2.1	13.4	41.7	21.8	130
L7-923	25.1	2.8	44	-0.7	2.2	12.1	39.7	21.5	131
C175	24.8	2.9	42	-2.3	2.6	14.4	41.2	21.2	129
C6	24.6	2.6	41	-3.3	2.4	13.3	43.6	20.3	131
C178	24.6	2.4	39	+1.5	2.7	14.6	42.2	20.3	132
Gibson	24.5	3.1	36	0	2.1	13.3	40.4	21.2	132
C153	24.3	3.0	40	-0.5	2.3	14.4	42.0	21.5	128
Chief	24.1	2.5	44	-3.2	2.6	12.6	40.5	21.4	131
C160	23.6	2.4	38	-2.5	2.1	13.6	42.1	21.1	131
L7-1160	23.1	2.6	43	-0.6	2.6	13.9	41.6	21.2	130
C154	23.0	2.6	37	-3.3	2.5	13.9	40.1	21.8	130
Macoupin	22.2	2.8	40	-2.3	2.6	14.8	41.1	22.0	129
Morse	22.1	3.4	40	-1.7	2.8	16.8	41.4	21.0	132
Boone	22.0	2.8	39	+0.5	2.3	13.4	41.5	21.5	130
C148	21.8	2.8	40	-2.3	2.5	13.9	40.8	21.4	126

¹ Days earlier (-) or later (+) than Gibson. Gibson required 127 days to mature.

² Mean of composite samples, composition on dry basis. Eleven tests in 1942 and eleven tests in 1941.

Table 38. Three-year summary of mean agronomic and chemical data for the strains in the Uniform Test, Group IV, 1940-1942.

Variety or Strain	Mean		Height		Seed		Percent-	Percent-	Iodine
	Yield	Lodg-	in	Matu-	Qual-	Seed	age of	age of	Number
	Bu./A.	ing	Inches	urity ¹	ity	Size	Protein	Oil	of Oil
Number of Tests	30	25	24	24	27	31 ²	31 ²	31 ²	31 ²
C149	25.2	2.6	39	-0.8	1.9	13.6	41.6	21.5	129
Patoka	25.1	1.8	31	-3.0	2.4	16.9	43.2	21.2	130
Gibson	24.8	2.9	36	0.0	2.1	13.4	40.1	21.1	132
Chief	24.3	2.6	43	-3.0	2.3	12.7	40.6	21.3	130
C178	23.7	2.5	38	+1.3	2.6	14.6	42.1	20.2	132
Boone	22.0	2.7	39	+1.1	2.2	13.6	40.9	21.5	130
Macoupin	22.0	2.7	40	-1.1	2.4	14.9	40.4	21.8	129

¹ Days earlier (-) or later (+) than Gibson. Gibson required 126 days to mature.

² Mean of composite samples, composition on dry basis. Eleven tests in 1942, eleven tests in 1941, and nine tests in 1940.

Table 39. Two-year summary of yield in bushels per acre and yield rank for the strains in the Uniform Test, Group IV, 1941-1942

Strain	Mean Yield 21 Tests	Evans- ville Ind.	Wheat- land Ind.	Ur- bana Ill.	Ston- ington Ill.	Clay- ton Ill.	Free- burg Ill.	Colum- bia Mo.	Sikes- ton Mo.
<u>Yield in bushels per acre</u>									
C155	26.3	41.6	17.6	44.3	23.7	29.5	28.7	22.4	18.2
C146	26.1	40.2	17.1	48.0	21.7	29.5	28.9	23.0	20.8
Patoka	26.0	42.3	20.4	43.7	27.0	28.3	32.7	19.0	14.6
C149	25.2	38.8	16.9	39.7	23.0	29.1	28.8	21.9	16.0
L7-923	25.1	42.0	17.2	41.8	22.1	26.8	26.9	20.3	18.6
C175	24.8	37.8	18.7	42.7	23.4	28.9	26.5	22.4	15.3
C6	24.6	38.6	19.9	43.4	24.6	28.0	24.7	19.6	16.0
C178	24.6	39.9	17.7	40.3	21.5	25.5	26.8	19.4	21.0
Gibson	24.5	41.3	18.7	41.5	22.8	26.5	23.1	19.6	18.4
C153	24.3	40.7	19.3	35.4	21.4	27.9	25.7	19.3	19.2
Chief	24.1	38.3	20.3	36.7	25.5	27.9	24.5	17.5	13.1
C160	23.6	35.9	17.1	41.1	23.2	26.7	26.5	17.0	15.6
L7-1160	23.1	33.8	16.4	39.2	22.3	27.2	21.5	20.9	17.7
C154	23.0	36.4	17.6	36.3	24.1	25.9	24.5	19.2	13.2
Macoupin	22.2	32.2	17.4	32.5	23.9	27.3	24.1	18.8	14.4
Morse	22.1	29.7	17.7	32.6	22.0	25.1	24.4	19.0	14.0
Boone	22.0	32.7	17.3	34.4	20.0	25.9	20.8	22.0	16.8
C148	21.8	37.5	15.5	34.1	20.9	25.3	22.5	20.1	12.2
<u>Yield Rank</u>									
C155		3	9	2	6	1	4	2	6
C146		6	14	1	14	1	2	1	2
Patoka		1	1	3	1	5	1	14	13
C149		8	16	9	9	3	3	5	9
L7-923		2	13	6	12	11	5	7	4
C175		11	5	5	7	4	7	2	12
C6		9	3	4	3	6	10	9	9
C178		7	7	10	15	16	6	11	1
Gibson		4	5	7	10	13	15	9	5
C153		5	4	14	16	7	9	12	3
Chief		10	2	12	2	7	11	17	17
C160		14	14	8	8	12	7	18	11
L7-1160		15	17	11	11	10	17	6	7
C154		13	9	13	4	14	11	13	16
Macoupin		17	11	18	5	9	14	16	14
Morse		18	7	17	13	18	13	14	15
Boone		16	12	15	18	14	18	4	8
C148		12	18	16	17	17	16	8	18

Table 40. Three-year summary of yield in bushels per acre and yield rank for the strains in the Uniform Test, Group IV, 1940-1942.

Strain.	Mean Yield 31 Tests	Evans- ville Ind.	Ur- bana Ill.	Ston- ington Ill.	Clay- ton Ill.	Els- berry Mo. ¹	Col- umbia Mo.	Sikes- ton Mo.
<u>Yield in bushels per acre.</u>								
C149	25.2	36.2	38.5	23.5	28.9	22.1	23.2	16.7
Patoka	25.1	34.3	40.1	25.9	28.5	19.0	20.0	16.0
Gibson	24.8	37.4	39.9	22.7	26.0	18.4	20.3	18.5
Chief	24.3	33.1	36.7	26.0	28.0	24.3	18.7	15.3
C178	23.7	36.2	35.2	20.4	24.0	19.7	20.6	19.6
Boone	22.0	30.7	33.1	19.6	25.2	18.1	22.5	16.9
Macoupin	22.0	28.3	32.1	23.8	26.7	17.9	19.8	14.4
<u>Yield Rank</u>								
C149		2	3	4	1	2	1	4
Patoka		4	1	2	2	4	5	5
Gibson		1	2	5	5	5	4	2
Chief		5	4	1	3	1	7	6
C178		2	5	6	7	3	3	1
Boone		6	6	7	6	6	2	3
Macoupin		7	7	3	4	7	6	7

¹Two years data only, 1940 and 1942.

Table 41. Analysis of variance for yield of seed for the Uniform Test, Group IV, 1942.

Source of Variation	Degrees of Freedom	Mean Squares
Replications	28	8,212.09**
Locations	9	4,578.05**
Varieties	23	178.31**
Varieties x Locations	207	42.54**
Error	642	12.03

**Highly significant

Acknowledgement of Seed Sources

Aside from the cooperating agencies, the following have very generously supplied seed for use in these tests. Their assistance is greatly appreciated.

<u>Source</u>	<u>Varieties</u>
McFayden Seed Co. Winnipeg, Man.	Pagoda Sioux Kabbot
Dr. G. P. McRostie Central Exp. Sta. Ottawa, Ontario	Goldsoy McRostie Mandarin
Professor R. G. Wiggins N. Y. Agr. Exp. Sta.	Ontario Cayuga
Professor B. D. Leith Wisconsin Agr. Exp. Sta.	Wis. Manchusel.
Mr. H. R. Bellin Wapeton, N. D.	Minsoy Habaro

Effect of Location on Composition

Average chemical composition of a strain of soybeans in the area of its adaptation is of importance to the plant breeder. The chemical composition to be expected in each locality is of importance also. The most desirable way to obtain this information would be by analyzing separately, seed of each strain at each location, thus also obtaining information on the interaction or failure of the strains to maintain their same rank or order among the locations where the nurseries were grown. Since this would entail the analysis of many samples, the compositing of strains appeared more feasible. By making up a field composite at each location composed of equal weights of seed of each variety or strain in a given Group, information on the average chemical composition of soybean seed at that location in that season may be obtained. This gives no information on the interaction between varieties and locations, however, from previous work (U. S. D. A. Bulletin 787) it may be assumed that for protein percent, oil percent and iodine number of oil, the varieties x locations interaction is non-significant. The average composition of soybean seed at each location will be found in Table 42.

Table 42. Chemical composition of soybean seed grown at each of the Uniform Test locations, and the location means for 1940-42. (composite of all strains grown in each respective Group Test, composition on dry basis)

Location	1940			1941			1942			Three-year mean		
	Percent-Protein	Percent-Iodine	Percent-Iodine	Percent-Protein	Percent-Iodine	Percent-Iodine	Percent-Protein	Percent-Iodine	Percent-Protein	Percent-Iodine	Percent-Iodine	
	age of	Number of	Number of	age of	Number of	Number of	age of	Number of	age of	Number of	Number of	
Wooster, Ohio	43.1	133	131	45.0	18.8	151	43.1	19.3	134	41.5	20.5	132
Strongsville, Ohio	45.9	129	125	43.1	19.3	132	46.0	18.4	130	43.6	19.4	129
Holgate, Ohio	45.9	131	128	43.1	19.3	132	41.4	19.2	137	43.6	19.9	131
LaGrange, Ind.	47.3	134	126	34.9	20.5	136	41.4	19.2	137	45.8	18.5	131
Bluffton, Ind.	43.1	128	129	45.0	20.5	131	41.4	19.2	137	40.8	21.3	130
Lafayette, Ind.	47.6	131	131	45.0	20.5	131	34.9	20.5	136	43.6	19.6	132
Wanatah, Ind.	42.8	130	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132
Urbana, Ill.	44.3	128	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132
Dwight, Ill.	44.3	128	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132
Stonington, Ill.	44.3	128	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132
London Mills, Ill.	44.3	128	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132
Clayton, Ill.	44.3	128	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132
Mt. Morris, Ill.	44.3	128	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132
Hudson, Iowa	44.3	128	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132
Kanawha, Iowa	44.3	128	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132
Ames, Iowa	44.3	128	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132
Cherokee, Iowa	44.3	128	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132
Waseca, Minn. ¹	44.3	128	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132
Dearborn, Mich.	44.3	128	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132
Paris, Mo.	44.3	128	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132
St. Joseph, Mo.	44.3	128	129	45.0	20.5	131	45.0	20.5	131	43.6	19.6	132

Group I (composite of 25 strains in 1940 and 1941 and 20 strains in 1942)

Group II (composite of 20 strains)

Group III (composite of 16 strains in 1940 and 1941, and 24 strains in 1942)

Columbus, Ohio	44.3	18.2	133	45.3	19.0	131	42.3	20.0	136	44.0	19.1	133
Holgate, Ohio	--	--	--	--	--	--	41.0	19.7	137	--	--	--
N. Vernon, Ind.	41.9	20.4	132	46.4	19.1	128	45.0	21.8	133	44.4	20.4	131
Greenfield, Ind. ²	--	--	--	42.2	20.3	129	40.6	21.2	135	--	--	--
Lafayette, Ind.	--	--	--	--	--	--	42.1	20.4	136	--	--	--
Urbana, Ill.	43.5	19.8	130	41.6	21.0	131	39.4	21.7	135	41.5	20.8	132
Dwight, Ill.	45.1	18.2	132	43.1	21.0	132	42.1	21.5	134	43.4	20.2	133
F. Greenwood, Ill.	43.6	20.2	128	41.4	21.6	127	--	--	--	--	--	--
Stonington, Ill.	40.9	20.8	131	43.7	20.4	129	42.4	21.4	133	42.3	20.9	131
London Mills, Ill.	40.4	21.2	132	43.2	20.1	130	--	--	--	--	--	--
Freeburg, Ill.	--	--	--	47.7	19.0	127	41.8	21.4	133	--	--	--
Clayton, Ill.	43.1	20.2	130	42.3	20.8	130	43.6	20.8	135	43.0	20.6	132
Paris, Mo.	43.8	20.0	127	--	--	--	40.1	22.2	134	--	--	--
Columbia, Mo.	42.3	20.1	131	41.8	20.5	128	43.0	20.8	131	42.4	20.5	130
Carrollton, Mo.	--	--	--	33.6	23.2	129	32.2	24.7	134	--	--	--
St. Joseph, Mo.	--	--	--	--	--	--	36.4	23.0	133	--	--	--
Ames, Iowa	--	--	--	--	--	--	43.4	21.0	135	--	--	--
Lincoln, Nebr. ¹	--	--	--	--	--	--	42.4	21.3	132	--	--	--

Group IV (composite of 22 strains in 1940 and 24 strains in 1941 and 1942)

Wheatland, Ind.	--	--	--	45.9	19.4	123	38.0	22.2	130	--	--	--
Evansville, Ind.	40.4	21.4	128	41.6	21.7	128	40.7	21.6	131	40.9	21.6	129
Mt. Vernon, Ind.	43.5	19.1	128	46.4	17.9	126	--	--	--	--	--	--
F. Vernon, Ind.	--	--	--	--	--	--	44.3	19.9	131	--	--	--
Urbana, Ill.	41.5	20.1	129	40.8	21.1	131	39.4	21.5	135	40.6	20.9	132
Edgewood, Ill.	41.8	20.4	127	40.9	22.1	126	--	--	--	--	--	--
Stonington, Ill.	39.6	21.6	131	41.8	21.7	129	40.9	22.2	131	40.8	21.8	130
Freeburg, Ill.	--	--	--	46.6	19.4	128	40.8	21.4	131	--	--	--
Clayton, Ill.	40.2	20.9	128	40.6	21.6	128	42.3	20.8	135	41.0	21.1	130
Sikeston, Mo.	41.1	20.3	127	46.6	18.7	123	40.1	21.8	127	42.6	20.3	126
Elsberry, Mo.	39.1	21.8	130	--	--	--	39.2	22.4	129	--	--	--
Columbia, Mo.	41.3	20.6	130	40.0	21.2	129	41.3	21.4	131	40.9	21.1	130
Carrollton, Mo.	--	--	--	33.4	23.5	130	33.3	24.0	132	--	--	--

¹ Composite of 19 strains at Waseca, and composite of 23 strains at Lincoln.

² Test located at Fortville near Greenfield in 1941.

