

PUBLISHED VERSION

Crocker, Roland Murley; Giunti, Carlo; Mortlock, Daniel J.
[Neutrino interferometry in curved spacetime](#) Physical Review, 2004; 69(6):063008

©2004 American Physical Society

<http://link.aps.org/doi/10.1103/PhysRevD.69.063008>

PERMISSIONS

<http://publish.aps.org/authors/transfer-of-copyright-agreement>

“The author(s), and in the case of a Work Made For Hire, as defined in the U.S. Copyright Act, 17 U.S.C.

§101, the employer named [below], shall have the following rights (the “Author Rights”):

[...]

3. The right to use all or part of the Article, including the APS-prepared version without revision or modification, on the author(s)' web home page or employer's website and to make copies of all or part of the Article, including the APS-prepared version without revision or modification, for the author(s)' and/or the employer's use for educational or research purposes.”

15th May 2013

<http://hdl.handle.net/2440/37091>

Neutrino interferometry in curved spacetime

Roland M. Crocker*

*School of Physics, The University of Melbourne, 3010 Victoria, Australia*Carlo Giunti[†]*INFN, Sezione di Torino, and Dipartimento di Fisica Teorica, Università di Torino, Via P. Giuria 1, I-10125 Torino, Italy*Daniel J. Mortlock[‡]*Institute of Astronomy, Madingley Road, Cambridge CB3 0HA, United Kingdom*

(Received 25 September 2003; published 25 March 2004)

Gravitational lensing introduces the possibility of multiple (macroscopic) paths from an astrophysical neutrino source to a detector. Such a multiplicity of paths can allow for quantum mechanical interference to take place that is qualitatively different from neutrino oscillations in flat space. After an illustrative example clarifying some underappreciated subtleties of the phase calculation, we derive the form of the quantum mechanical phase for a neutrino mass eigenstate propagating nonradially through a Schwarzschild metric. We subsequently determine the form of the interference pattern seen at a detector. We show that the neutrino signal from a supernova could exhibit the interference effects we discuss *were* it lensed by an object in a suitable mass range. We finally conclude, however, that—given current neutrino detector technology—the probability of such lensing occurring for a (neutrino-detectable) supernova is tiny in the immediate future.

DOI: 10.1103/PhysRevD.69.063008

PACS number(s): 98.62.Sb, 14.60.Pq, 95.30.Sf

I. INTRODUCTION

Spacetime curvature allows, in general, for there to be more than one macroscopic path from a particle source to a detector. This means that there is a quantum mechanical interference phenomenon that may occur—at least in principle—with gravitationally lensed, astrophysical neutrinos that is qualitatively different from “conventional” neutrino oscillation. The possibility for this different type of interference arises because—with, generically, each path from source to detector having a different length—a phase difference may develop at the detector due to affine *path* difference(s). This is to be contrasted with flat spacetime neutrino oscillations which arise because different mass eigenstates generically have different *phase* velocities. One might expect, in fact, gravitationally induced neutrino interference (GINI) to exhibit a phenomenology partially analogous to that produced by Young’s double slit experiment, *viz.*, regular patterns of maxima and minima across a detected energy spectrum. As we show below, for ultrarelativistic neutrinos, each maximum and minimum at some particular energy is characterized by, respectively, enhancement and depletion of *all* neutrino species (not relative depletion of one species with respect to another, which characterizes flat space neutrino oscillations).

Below, we shall provide the theoretical underpinning to all the contentions made above. We also sketch a proof-of-principle that this interference effect could actually be seen in the neutrinos detected from a supernova *given* a suitable

lens. There are other situations where the GINI effect might, in principle, also be evident. Reluctantly, however, we conclude that pragmatic considerations mean that GINI effects will be very difficult to see in these cases.

II. SURVEY

Particle interferometry experiments enjoy a venerable lineage and—apart from their intrinsic interest—have often found utility in the measurement of intrinsically small quantities. The idea that the effects of *gravity*—the epitome of weakness as far as particle physics is concerned—on the phase of particles might become manifest in interferometry dates back to the seminal, theoretical work of Overhauser and Colella [1]. It was these researchers, themselves, together with Werner [2], who were the first to experimentally confirm the effect they were predicting (in what has come to be labeled a COW experiment after the initials of these researchers; see Ref. [3] for a review).

Another interesting idea involving gravitational effects on interferometry of neutral particles—though, to the authors’ knowledge, without yet having received experimental confirmation—is the idea that gravitational microlensing of *light* might realize a *de facto* Young double slit arrangement. There is extensive literature devoted to this idea (see Refs. [4–12]), which has been labeled “femtolensing” because of the natural angular scales involved for cosmologically distant sources and lenses [7]. Femtolensing is somewhat more closely analogous to the idea we present (indeed, as we show below, the analogy becomes exact in the massless neutrino limit) than COW-type experiments. This is because in femtolensing, gravity not only affects the phase of the propagating photons, but it is also itself responsible for the “bending” of these particles so that diverging particle beams (or, more precisely, wave packets) can be brought back together to

*Electronic address: r.crocker@physics.unimelb.edu.au

[†]Electronic address: giunti@to.infn.it; URL: <http://www.to.infn.it/~giunti>[‡]Electronic address: mortlock@ast.cam.ac.uk

interfere. Furthermore, while the interfering particles are relativistic in both the femtolensing and GINI cases, they are nonrelativistic in COW experiments.

As far as sources go, light from GRBs has received particular attention in the context of femtolensing [4,7,8]. While we would, of course, also require astrophysical objects as the sources for a GINI “experiment,” the sources best able to offer a chance for the detection of this effect are probably supernovae. A Galactic supernova would generate excellent statistics in existing solar (and other) neutrino detectors (thousands of events—see below). And with a much larger, generation of neutrino detectors on the drawing board—some having as one of their chief design goals the detection of neutrinos from supernovae occurring almost anywhere in our Local Group—prospects for the detection of the effect we predict can only improve with time [13].

By way of a pedagogical detour, please note the following: we believe the “time-delay” nomenclature is misleading in the context of either femtolensing or GINI effects. It is much better, we contend, to think in terms of path difference(s). The idea of a well-defined time delay belongs to classical physics. The time delay is—in the frame of some observer—the time elapsed between the arrival of two signals. These should have their origins in the “same” (macroscopic) event at a source, but then travel down different classical geodesics from source to detector. Now, from the viewpoint of quantum mechanics, there is a limit in which the classical description just given makes sense and is useful. This limit is that in which the size of the wave packets describing the signaling particles is small in comparison to the affine path length differences between the different classical trajectories under consideration. This limit will usually be satisfied in observationally interesting cases of gravitational lensing. But this limit must *not* be satisfied if femtolensing or GINI effects are to be observed. Indeed, we require the opposite situation to pertain, viz., an affine path length difference of the order of or smaller than the wave packet size. This is required so that wave packets created in the same (microscopic) event can overlap at the detector—with interference effects being the result. In this sense, there is *no* time delay because the wave packets have to be overlapping at the detector position at the same (observer) time, i.e., overlap must be satisfied at the spacetime location of the observation event. Note further that, to paraphrase Dirac, each photon—or neutrino—only interferes with itself. So it is the wave packet of the single particle that results from a single (microscopic) event—like the decay of an unstable parent particle—that, in simple terms, splits to travel down all the classical geodesics from source to detector, only to interfere when recombined there. The idea we are describing, then, does not require some weird (and impossible) analog of a “neutrino laser”; it works at the level of individual, particle wave packets.

Another strand that will be peripherally drawn into this paper is the behavior—at a classical level—of neutrinos in a curved spacetime background (i.e., gravitational lensing of neutrinos treated as ultrarelativistic, classical particles). This topic became of immediate interest with the detection of neutrinos from SN 1987A [14,15]. Timing information from the

nearly simultaneous detection of these neutrinos and the supernova’s photon signal and from the time and energy spread of the neutrino burst alone has been investigated in many papers as an empirical limit on the neutrino mass scale (see, e.g., Refs. [16,17] and Refs. [18–20] for reviews and the seminal references concerning this idea) and also as a probe of the equivalence principle over intergalactic distance scales [21–23]. More speculatively but germane to this work, the apparently bimodal distribution of SN 1987A neutrinos observed by the then-operating Kamiokande solar neutrino observatory was given an explanation in terms of an intervening gravitational lens (in the $5 \times 10^5 M_{\odot}$ mass range: [24]). Further, the idea that astrophysical neutrino “beams” might be gravitationally focused by massive objects like the Sun has been investigated and it has been found that such focusing can amplify an intrinsic neutrino signal by many orders of magnitude (see [25] and [26] for more recent work).

An early and important work treating the *quantum mechanical* aspects of neutrino propagation through a curved metric is that of Brill and Wheeler [27]. Their work is particularly important for its elucidation of the formalism that allows one to treat (massless) spinor fields under the influence of gravitational effects (i.e., the extension of the Dirac equation to curved spacetime).

As presaged above, in this paper we shall be particularly concerned with the *phase* of neutrinos (more particularly, neutrino mass eigenstates) in curved spacetime. The seminal work treating the phase of quantum mechanical particles in curved spacetime is that of Stodolsky [28]. In this work the author argued that the phase of a *spinless* particle in an arbitrary metric is identical with the particle’s classical action (divided by \hbar). Later work conducted on neutrino oscillations in curved spacetime [29–31] has often—though not always [32,33]—implicitly assumed the correctness of Stodolsky’s contention (that the phase is given by the classical action) for spinor fields as well and taken this as its starting point. Somewhat ironically—as we set out in detail below—recent researches [34] have revealed that the equality of classical action and phase holds for spin half particles, but *not* for spin zero or one particles, or, at least not in an unqualified sense. In any case, that Stodolsky’s contention holds for spinors means a considerable simplification for our calculations as we can avoid directly treating the covariant Dirac equation.

A full review of the literature (see [29–41]) on neutrino phase in the presence of gravity is beyond the scope of this work. Suffice it to say that most work here to date has been concerned with the calculation of neutrino phase in radial propagation of neutrinos through stationary, spherically symmetric, spacetimes. There is active controversy in this context as to at what order in neutrino mass (m_{ν}^2 or m_{ν}^4) gravitational corrections show up in the phase [30,33,35,36]. The answer to this hangs critically on how energy and distance, in particular, are defined [36]. We shall have to treat such issues carefully, but all the subtleties of this debate need not particularly concern us. This is because we are *primarily* interested in gravity not for its effect on phase *per se*, but for its ability to generate multiple macroscopic paths from source to detector. And it is what might actually be measured

at the detector that concerns us. Detectors count the neutrinos—registered in terms of flavor and (local) energy—that interact within their volume. From these one can infer interference patterns, but one does not, of course, have any direct experimental access to the phase difference(s) (a point that does sometimes seem to be forgotten).

In regard to interference phenomenology, note the following: whereas interference patterns with flat space neutrino oscillations take the form of variations in neutrino flavor *ratios* across energy, with GINI, because there will be constructive and destructive interference between the multiple allowed routes, one (also) expects to see, in general, maxima and minima (distributed across energy) in the counts of *all* neutrino flavors. These maxima and minima will be present irrespective of what measure of distance, say, we settle on (though, of course, they may be undetectably small in amplitude—but that is a separate issue). To put this in a different way, flat space neutrino oscillations modify the *relative* abundances of neutrinos expressed as a function of energy, whereas GINI effects can modify *absolute* abundances.

Interestingly, of all the papers devoted to the topic of the gravitationally affected neutrino phase, to the authors' knowledge, only one [29] has previously examined GINI, which, to reiterate, is the idea that neutrino *oscillations* in the presence of gravitational lensing—or, to be strict, gravitational focusing—might present interesting phenomenology. (This is the analog of the femtolensing described above that involved light.) To examine this idea, the authors of Ref. [29] were obliged to develop a formalism to deal with nonradial propagation of neutrinos around a lensing mass, and we shall adopt much of this formalism in the current work. Unfortunately, Ref. [29] contains an incorrect result which it is one of the major aims of this paper to point out. Moreover, other works which have considered the gravitationally affected neutrino phase contain results—and commentary thereon—which, if not strictly incorrect, can be misleading if one does not realize the restricted nature of their tenability. In brief, most authors have failed to consider the possibility of multiple paths. Any result which suggests the vanishing of a neutrino phase difference in the massless limit [see, e.g., Eq. (13) of Ref. [30] or Eq. (4.7) of [33]] should be interpreted with extreme caution [42].

Essentially, the incorrect result in Ref. [29], as alluded to above, is then that the phase for a neutrino mass eigenstate, j , propagating nonradially through a Schwarzschild metric is purely proportional to its mass, m_j , squared [see Eq. (58) of [29] and also Eq. (25) of [40]]. This result is incorrect [43]: in the massless limit, the neutrino phase in curved spacetime should reduce (modulo spin-dependent corrections which vanish for radial trajectories [27] and are negligible except in extreme, gravitational environments [34]) to the result for photons. And the photon phase is not zero (otherwise the interference fringes—in space or energy—predicted by the femtolensing literature would not be produced), even though, of course, the classical action is zero along null geodesics. Indeed, the photon phase is essentially proportional to energy. As we show below, furthermore, this $\propto E$ term is the leading order term for the neutrino phase as well.

What has gone wrong when one's analysis misses the $\propto E$ term in the neutrino phase is that one has tried to simultaneously employ two incompatible notions: the fundamentally wave or quantum mechanical idea of phase with the particle notion of trajectory so that x is given in terms of t or vice versa. Even in the simpler case of flat space oscillations, the introduction (often implicitly) of the idea of a trajectory—or, more particularly, a group velocity—into the calculation of neutrino phase leads to error (in particular, the recurring bugbear that the conventional formula for the neutrino oscillation length is wrong by a factor of two: see [44]). In the calculations set out below, we show the reader how the error of introducing a trajectory can be avoided. Furthermore, our method allows calculations to be performed along the actual (classical) paths [not trajectories; i.e., we have $r(\phi)$, say, rather than $r(t), \phi(t)$] of the neutrino mass eigenstates, rather than taking the approach of calculation along the null geodesic employed in Ref. [29]. On the other hand, our approach also circumvents the obligation to introduce extra phase shifts “by hand.” This artificial device becomes necessary when one offsets either the emission times or positions of the different mass eigenstates with respect to each other so that they arrive at the same spacetime point (see [30] for an example of this).

The plan of this paper is the following. In Sec. III we describe, for illustrative purposes, interference of neutrino plane waves propagating through flat space along both different (classical) paths and having, in general, different phase velocities. Then in Sec. IV we describe the calculation of the neutrino mass eigenstate phase in a Schwarzschild metric, correcting an erroneous result that has existed in the literature for some time. We then set out, in Sec. V the calculation of the analog of the survival and oscillation probabilities in flat space for neutrinos that have been gravitationally lensed by a point mass. Section VI describes some generic limiting factors to the size of the GINI effect. In Sec. VII we examine—at a heuristic level—questions of coherence that can effect the visibility of the GINI effect for neutrinos from supernovae. We give a proof-of-principle that the effect should be detectable. In Sec. VIII we describe some limitations of our method—which stem particularly from the assumption of exclusively classical paths—and set out improvements to be made in further work. Finally, in an Appendix, we set out a wave packet treatment of the neutrino beam splitter toy model treated in Sec. III in terms of plane waves. We derive results here pertaining to the analog of the coherence length in conventional neutrino oscillations.

III. NEUTRINO BEAM SPLITTER

By way of an illustrative introduction to this topic we consider a toy model of interference effects that can arise when there are both multiple paths from a source to a detector, in the manner of Young's double slit experiment, and different phase velocities for the propagating particles, in the manner of neutrino oscillations. Of course, interference requires that our experimental apparatus be unable to distinguish between the propagating particles (just as we likewise require for interference that the apparatus be unable to identify *which* path any single particle has propagated down). We




FIG. 1. The geometry of either a double slit interference experiment or a two-image gravitational lens. The bold lines are the two neutrino paths, + and -, between the source, A , and the observer, B , and the geometry is defined both in terms of physical variables (r_A , r_B , b_{\pm} and s) and astronomical/lensing variables (D_d , D_s , D_{ds} , θ_{\pm} and β). Also included is the deflection angle, $\alpha_{\pm}(b)$, parametrized as a function of the physical impact parameter. It is implicit in this diagram that the deflector is rotationally symmetric, and that the paths are thus confined to the plane of the page defined by the source, deflector and observer.

therefore require the propagating and detected particles to be different objects. In this context, let us consider, for the sake of definiteness and relevance, a *gedanken experiment* involving an imaginary (flat space) neutrino beam splitter in the geometry illustrated in Fig. 1. We take it that the neutrinos' paths can be approximated as two straight-line segments along which momentum is constant in magnitude. We need only treat, therefore, one spatial dimension: $\int \mathbf{p}_j \cdot d\mathbf{x} = \int p_j dx = p_j |x|$, where $|x|$ is the total distance along the two line segments. We expect that the qualitative behavior of this device shall illustrate many of the features expected to emerge from interference of gravitationally lensed neutrinos. Note that for reasons of clarity we present only a plane wave treatment here, leaving a full wave packet calculation for an Appendix. We stress, however, that wave packet considerations—which allow, in particular, for a proper treatment of decoherence effects—are, in general, important and must certainly be considered when one is dealing with neutrinos that have propagated over long distances (see Sec. VI).

Let us write the ket associated with the neutrino flavor eigenstate α that has (in a loose sense) propagated from the source spacetime position $A = (x_A, t_A)$ to detection position $B = (x_B, t_B)$ as

$$|\nu_{\alpha}; A, B\rangle = N \sum_p \sqrt{I_p} \sum_j U_{\alpha j} \times \exp[-i\Phi_j^p(E_j; L_p^{AB}, T^{AB})] | \nu_j \rangle, \quad (1)$$

where $T^{AB} \equiv t_B - t_A$, L_p^{AB} is the distance from source position x_A to detector position x_B along one of a finite number of paths labeled by p , $\Phi_j^p(E_j; L_p^{AB}, T^{AB}) \equiv E_j T^{AB} - p_j(E_j) L_p^{AB}$, with $p_j(E_j)$ denoting the momentum of mass eigenstate j with energy E_j , and, finally, U is a unitary matrix relating the neutrino flavor eigenstates to the neutrino mass eigenstates. We have included the $\sqrt{I_p}$ factor to account for the fact that, in general, we should allow for a path-dependence to the amplitude, a situation where, for paths p and q , I_p

$\neq I_q$ is the analog of Young's slit experiment in which the slits are not equally illuminated [thus reducing the visibility—see Eq. (92)—of the resulting interference fringes]. The Schwarzschild lens scenario explored in Secs. IV and V presents a situation analogous to this: light or neutrinos propagating down the two classical paths from source to observer will not be, in general, equally magnified. Note that we choose throughout this paper—unless otherwise indicated—to work in units such that $\hbar = c = 1 \neq G$. The mass eigenstates are assumed to be on mass shell: $p_j(E_j) \equiv (E_j^2 - m_j^2)^{1/2}$. The amplitude for a neutrino created as type α at the spacetime position A to be detected as type β at the spacetime position, B , of the detection event is then

$$\langle \nu_{\beta} | \nu_{\alpha}; A, B \rangle = N \sum_p \sqrt{I_p} \sum_j U_{\alpha j} U_{\beta j}^* \times \exp[-i\Phi_j^p(E_j; L_p^{AB}, T^{AB})]. \quad (2)$$

Assuming a stationary source, we can get rid of the unwanted dependence on time by averaging over T^{AB} in the above to determine a time-averaged oscillation probability analog at the detector position x_B [45–47]. This maneuver gives us that

$$|\langle \nu_{\beta} | \nu_{\alpha}; x_A, x_B \rangle|^2 \propto \int dT |\langle \nu_{\beta} | \nu_{\alpha}; A, B \rangle|^2 \propto |N|^2 \sum_{pq} \sqrt{I_p I_q} \sum_{jk} U_{\alpha j} U_{\beta j}^* U_{\beta k} U_{\alpha k}^* \times \exp\{i[p_j(E) L_p^{AB} - p_k(E) L_q^{AB}]\}, \quad (3)$$

where we have $E_j = E_k \equiv E$ because of the $\delta(E_j - E_k)$ that arises from the integration over time.

We find, then, after a simple calculation that the oscillation probability analog becomes

$$\begin{aligned}
& |\langle \nu_\beta | \nu_\alpha ; x_A, x_B \rangle|^2 \\
& \simeq \frac{1}{\sum_{rs} \sqrt{I_r I_s}} \sum_{pq} \sqrt{I_p I_q} \sum_{jk} U_{\alpha j} U_{\beta j}^* U_{\beta k} U_{\alpha k}^* \\
& \times \exp[-i\Delta\Phi_{jk}^{pq}], \tag{4}
\end{aligned}$$

where the phase difference is given by

$$\Delta\Phi_{jk}^{pq} \equiv -E(L_p^{AB} - L_q^{AB}) + \left(\frac{m_j^2 L_p^{AB} - m_k^2 L_q^{AB}}{2E} \right). \tag{5}$$

Note that in Eq. (4) the normalization has been determined by requiring that $|\langle \nu_\beta | \nu_\alpha ; x_A, x_B \rangle|^2 \leq 1$. The presence of the \leq sign (as opposed to a simple equality) comes about as the interference between states propagating down different paths can result in minima at which the total neutrino detection probability is zero, in which case the usual unity normalization is impossible (so now $|\langle \nu_\beta | \nu_\alpha ; x_A, x_B \rangle|^2$ no longer has a direct interpretation as a probability). This is qualitatively different behavior from that seen in neutrino oscillations where, given maximal mixing between ν_α and ν_β , an experiment might be conducted in a position where only neutrinos of type α , say, are to be found or, alternatively, only of type β , but never in a position where none can be found in principle.

On the other hand the behavior explained above—involving interference minima and maxima—is obviously analogous to what one would expect in a double slit experiment or similar. In fact, we shall show below that the phenomenology of neutrino interference—when there is more than one path from source to detector—is a convolution of the two types of interference outlined above. This means that, in general, one cannot simply recast $|\langle \nu_\beta | \nu_\alpha ; x_A, x_B \rangle|^2$ in terms of a conditional probability, separating out the overall interference pattern (with its nulls, etc.) from the conditional probability that any *detected* neutrino has certain properties. To understand this point, imagine setting all mixing angles to zero so that the mass eigenstates and weak eigenstates are identical. The point now is that the interference patterns for the various detected weak/mass eigenstates are still different: the phase difference (which now stems purely

from the path difference) is dependent on the mass of the neutrino species involved. In other words, $|\langle \nu_\beta | \nu_\alpha ; x_A, x_B \rangle|^2$ will *not*, in general, factorize into a conditional probability multiplied by an interference envelope because the putative interference envelope is different for different mass eigenstates. As we show below, however, in the ultrarelativistic limit, the cross term (that mixes different path indices with different mass eigenstate indices) always turns out to be small with respect to the other terms. In this limit, then, factorization is a good approximation.

We now consider two particular, illustrative cases of the neutrino beamsplitter *gedanken experiment* for which we calculate relevant phase differences and oscillation probability analogs.

A. Two-path neutrino beam splitter

A particularly perspicuous example is given by the two-path example of the above equations. For this we specify a reference length, L , to which the two paths, of lengths L_+ and L_- , are related by

$$L_\pm \equiv L \pm \frac{\Delta L}{2}. \tag{6}$$

This means that there are four phase difference types as labeled by the path indices $p, q \in \{+, -\}$, viz.,

$$\begin{aligned}
\text{(i)} \quad \Delta\Phi_{jk}^{++} &= + \frac{\delta m_{jk}^2 L + \Delta L/2}{2E} \\
\text{(ii)} \quad \Delta\Phi_{jk}^{--} &= + \frac{\delta m_{jk}^2 L - \Delta L/2}{2E} \\
\text{(iii)} \quad \Delta\Phi_{jk}^{+-} &= -E \left(1 - \frac{m_j^2 + m_k^2}{4E^2} \right) \Delta L + \frac{\delta m_{jk}^2 L}{2E} \\
\text{(iv)} \quad \Delta\Phi_{jk}^{-+} &= +E \left(1 - \frac{m_j^2 + m_k^2}{4E^2} \right) \Delta L + \frac{\delta m_{jk}^2 L}{2E}. \tag{7}
\end{aligned}$$

In this case, then, Eq. (4) becomes

$$\begin{aligned}
|\langle \nu_\beta | \nu_\alpha ; x_A, x_B \rangle|^2 & \simeq \frac{1}{I_+ + I_- + 2\sqrt{I_+ I_-}} \sum_{jk} U_{\alpha j} U_{\beta j}^* U_{\beta k} U_{\alpha k}^* \exp\left(-i\frac{\delta m_{jk}^2 L}{2E}\right) \left\{ I_+ \exp\left(+i\frac{\delta m_{jk}^2 \Delta L}{4E}\right) + I_- \exp\left(-i\frac{\delta m_{jk}^2 \Delta L}{4E}\right) \right. \\
& \left. + 2\sqrt{I_+ I_-} \cos\left[E\left(1 - \frac{m_j^2 + m_k^2}{4E^2}\right)\right] \right\}. \tag{8}
\end{aligned}$$

This is an interesting result. It shows that the interference factorizes into a conventional, flat space oscillation term and an interference “envelope” in curly brackets (as might be

expected from Young’s slit type experiment). If we now further particularize to the case where $I_+ = I_-$, this envelope term becomes

$$\begin{aligned} & \cos\left(\frac{\delta m_{jk}^2 \Delta L}{4E}\right) + \cos\left[E\left(1 - \frac{m_j^2 + m_k^2}{4E^2}\right)\right] \\ &= 2 \cos\left[E\left(1 - \frac{m_j^2}{2E^2}\right)\frac{\Delta L}{2}\right] \cos\left[E\left(1 - \frac{m_k^2}{2E^2}\right)\frac{\Delta L}{2}\right], \quad (9) \end{aligned}$$

which obviously reduces to the expected photon interference term $\propto \cos^2$ in the massless limit. Later (in Sec. V) we shall see that this sort of factorization property also arises, under a different set of assumptions, with gravitational lensing of astrophysical neutrinos.

B. Double slit experiment geometry

Having considered a hypothetical neutrino plane wave beam-splitter (Sec. III A), it is now possible to combine the rigorous, if simplistic, results derived above with heuristic arguments to investigate a more realistic double slit neutrino interference experiment. The choice of a double slit experiment is particularly relevant not only because of its links with more familiar interference phenomena, but also because a point-mass gravitational lens admits two (significant) paths from source to observer. Thus the results obtained in this section should provide a useful guide to the qualitative behavior of GINI in curved spacetime considered in Sec. IV.

The relevant geometry is illustrated in Fig. 1, showing the two classical paths (+ and -) from source to detector, with the important introduction of a more physically motivated set of lengths than the reference length, L , used previously. The entire experiment is taken to be planar, with the coordinate system origin on the line defined by the two slits. (This is an arbitrary decision at present, but will coincide with the position of the deflector in Sec. IV.) The slits are defined by their positions, b_+ and b_- (which will correspond to image positions in Sec. IV E); the source position is given by both its radial coordinate, r_A , and its perpendicular offset, s ; the observer position is defined by its radial coordinate, r_B , alone. There are several other plausible ways in which this geometry could be defined, but all derived results become equivalent under the assumption that $|b_{\pm}| \ll r_{\{A,B\}}$, as applied throughout. Note also that the observer and source are interchangeable.

As discussed in Sec. III, all flat-space interference phenomena can be treated in terms of path lengths. The lengths of the two paths illustrated in Fig. 1 are

$$\begin{aligned} L_{\pm} &= (r_B^2 + b_{\pm}^2)^{1/2} + [r_A^2 - s^2 + (b_{\pm} - s)^2]^{1/2} \\ &\simeq (r_A + r_B) \left[1 + \frac{1}{2r_A r_B} \left(b_{\pm}^2 - \frac{2r_B}{r_A + r_B} s b_{\pm} \right) \right], \quad (10) \end{aligned}$$

where, in most cases, b_- and b_+ are of opposite sign, and the second line explicitly utilizes the fact that $|b_{\pm}| \ll r_{\{A,B\}}$. The path difference is thus

$$\begin{aligned} \Delta L_{+-} &= L_+ - L_- \\ &\simeq \frac{r_A + r_B}{2r_A r_B} \left[b_+^2 - b_-^2 - \frac{2r_B}{r_A + r_B} s(b_+ - b_-) \right] \quad (11) \end{aligned}$$

and the average path length is

$$L \simeq (r_A + r_B) \left\{ 1 + \frac{1}{4r_A r_B} \left[b_+^2 + b_-^2 - \frac{2r_B}{r_A + r_B} s(b_+ + b_-) \right] \right\}. \quad (12)$$

C. Schwarzschild slit geometry

In a laboratory-based double slit experiment the two slit positions can be chosen arbitrarily, but in the case of gravitational lensing the impact parameters of the beams are determined by a combination deflector and source parameters (Sec. IV). Given the Schwarzschild metric around a point-mass (Sec. IV E), the assumption of small deflection angles implies that the source position and impact parameters are related by [cf. Eq. (75)]

$$b_+^2 - b_-^2 = \Delta b_{+-}^2 \simeq \frac{r_B}{r_A + r_B} s(b_+ - b_-). \quad (13)$$

Applying this result in the more general context of the double slit experiment, Eq. (13) can be rewritten as

$$\Delta L_{+-} = L_+ - L_- \simeq -\frac{r_A + r_B}{2r_A r_B} (b_+^2 - b_-^2). \quad (14)$$

Similarly, the mass-length expression that appears in, e.g., Eq. (5) can be simplified to

$$\begin{aligned} & m_j^2 L_+ - m_k^2 L_- \\ & \simeq (r_A + r_B) \left[\delta m_{jk}^2 - \frac{1}{2r_A r_B} (m_j^2 b_+^2 - m_k^2 b_-^2) \right]. \quad (15) \end{aligned}$$

Substituting these expressions in Eq. (5) then gives the phase difference between mass eigenstates j and k traveling down paths + and - as

$$\begin{aligned} \Delta \Phi_{jk}^{+-} & \simeq +E(r_A + r_B) \frac{\Delta b_{+-}^2}{2r_A r_B} + \frac{\delta m_{jk}^2}{2E} (r_A + r_B) \\ & \quad - \frac{r_A + r_B}{2E} \frac{(m_j^2 b_+^2 - m_k^2 b_-^2)}{2r_A r_B}. \quad (16) \end{aligned}$$

This expression includes contributions from both different phase velocities and different path lengths, and can be understood further by considering the special cases in which (i) different mass neutrinos travel down the same path or (ii) the same mass eigenstate travels down different paths.

(i) From Eq. (5) for the general case of the phase difference between the *samemass* eigenstate propagating along *different* paths we find

$$\Delta\Phi_{jj}^{pq} \equiv - \left[E - \frac{m_j^2}{2E} \right] (L_p - L_q) \quad (17)$$

so that if we further particularize, as above, to paths passing through slits at b_+ and b_- we find

$$\Delta\Phi_{jj}^{+-} \simeq \left[E - \frac{m_j^2}{2E} \right] \frac{(r_a + r_b)\Delta b_{+-}^2}{2r_a r_b}. \quad (18)$$

Notice in the above the similarity to the phase difference for an ordinary Youngs double slit type experiment using photons, namely,

$$\begin{aligned} \Delta\Phi_{\gamma}^{pq} &= -\bar{E}v(L_p - L_q) \\ &= - \left(\bar{E} - \frac{m^2}{2\bar{E}} \right) (L_p - L_q), \end{aligned} \quad (19)$$

where v is the (phase) velocity of the interfering particle (which we assume to be relativistic). We shall see below that the analog of this phase—essentially proportional to energy \times path difference—has been missed in the existing literature on neutrino oscillations in curved space. This has led to an incomplete result suggesting that the phase difference vanishes in the massless limit even when there is more than one path from source to detector.

(ii) Again for the general case, for the phase difference between *different* mass eigenstates propagating along the *same* path (i.e., the analog of the usual phase difference encountered in neutrino oscillation experiments), we find from Eq. (5)

$$\begin{aligned} \Delta\Phi_{jk}^{xx} &\equiv + \left(\frac{\delta m_{jk}^2 L_x}{2E} \right) \\ &\simeq \frac{\delta m_{jk}^2}{2E} (L_a + L_b) \left(1 + \frac{b_x^2}{2L_a L_b} \right), \end{aligned} \quad (20)$$

where $x \in \{p, q\}$.

It is worth keeping the above expressions in mind when considering the results for the neutrino phase difference in curved spacetime presented in Sec. IV E. As will be seen, the results obtained in this more complex physical situation are analogous to those derived above, e.g., compare Eq. (18) with Eq. (58) and Eq. (20) with Eq. (59).

IV. THE PHASE OF A NEUTRINO MASS EIGENSTATE IN CURVED SPACETIME

A neutrino beam splitter is in the realm of fantasy—except for the interesting case of gravitational lensing of neutrinos: a gravitational field can bring to focus diverging neutrino beams, and therefore provide for multiple (classical) paths from a source to a detector. In the remainder of this paper we explore whether any interesting, quantum mechanical interference effects can arise in this sort of situation.

We shall be concerned below, therefore, with deriving an expression for the neutrino oscillation phase in curved spacetime, in particular a Schwarzschild metric (this providing the

simplest case in which gravitational lensing is possible). Here we shall follow the development laid out in [29] and [32] but, importantly, we shall also employ the prescription set out in Sec. III that allows for the removal of time from consideration in the oscillation “probability” by integration over $T \equiv t_B - t_A$ where $A = (r_A, t_A)$ and $B = (r_B, t_B)$ are the emission and detection events respectively [45]. Note that we are assuming the semiclassical limit in which gravity is not quantized and its effects can be described completely by a nonflat metric, $g_{\mu\nu} \neq \eta_{\mu\nu}$.

The procedure we follow is to start with the generalization of the equation for a mass eigenstate’s phase in flat spacetime to curved spacetime first arrived at by Stodolsky [28]:

$$\Phi_k(B, A) = \int_A^B p_{\mu}^{(k)} dx^{\mu}, \quad (21)$$

where

$$p_{\mu}^{(k)} = m_k g_{\mu\nu} \frac{dx^{\nu}}{ds}, \quad (22)$$

is the canonically conjugate momentum to the coordinate x^{μ} . Actually, as pointed out by Alsing *et al.* in Ref. [34], Stodolsky’s expression for the phase is missing in general, small correction terms that arise from quantum mechanical modifications to the classical action. These vary according to the spin of the particle under consideration. Completely fortuitously, the would-be correction terms are identically zero in the case of spin half particles in a *static* metric (whereas for particles with, e.g., spin zero or one they are nonzero) so the Stodolsky expression happens to be exact for the Schwarzschild metric and many other cases of interest. Note in passing that this restriction to a static metric means that this technology cannot—as it stands—treat, e.g., particle phases in a cosmological context.

We now introduce the metric of the Schwarzschild spacetime. This may be written in radial coordinates, $x^{\mu} = (t, r, \vartheta, \varphi)$, as

$$ds^2 = B(r)dt^2 - B(r)^{-1}dr^2 - r^2d\vartheta^2 - r^2\sin^2\vartheta d\varphi^2, \quad (23)$$

where

$$B(r) \equiv \left(1 - \frac{2GM}{r} \right), \quad (24)$$

and G is the Newtonian constant and M is the mass of the source of the gravitational field, i.e., the lensing mass. Given the isotropy of the gravitational field, the motion of the neutrino mass eigenstate will be confined to a plane which we take to be the equatorial one, $\vartheta = \pi/2$ and $d\vartheta = 0$.

The relevant components of the canonical momentum, Eq. (22), are then [29]

$$p_t^{(k)} = m_k B(r) \frac{dt}{ds}, \quad (25)$$

$$p_r^{(k)} = -m_k B^{-1}(r) \frac{dr}{ds}, \quad (26)$$

and

$$p_\varphi^{(k)} = -m_k r^2 \frac{d\varphi}{ds}. \quad (27)$$

These are all interrelated through the mass-shell condition [29],

$$\begin{aligned} m_k^2 &= g^{\mu\nu} p_\mu^{(k)} p_\nu^{(k)} \\ &= \frac{1}{B(r)} (p_t^{(k)})^2 - B(r) (p_r^{(k)})^2 - \frac{(p_\varphi^{(k)})^2}{r^2}. \end{aligned} \quad (28)$$

Given that the components of the metric are independent of the coordinates t and φ , the momenta associated with these quantities, $p_t^{(k)}$ and $p_\varphi^{(k)}$, shall be conserved along the classical geodesic traced out by ν_k . We define these constants of motion as $E_k \equiv p_t^{(k)}$ and $J_k \equiv -p_\varphi^{(k)}$. These two are, respectively, the energy and angular momentum seen by an observer at $r \rightarrow \infty$ for the k th mass eigenstate [29]. They are *not* identical with the energy and angular momentum that would be measured for ν_k at some definite, finite position r . In general, however, one may relate these quantities using the transformation law that relates a local reference frame $\{x^{\hat{\alpha}}\} = \{\hat{t}, \hat{r}, \hat{\vartheta}, \hat{\varphi}\}$ to the frame $\{x^\mu\} = \{t, r, \vartheta, \varphi\}$ [48]:

$$x^{\hat{\alpha}} = L_{\mu}^{\hat{\alpha}} x^\mu, \quad g_{\mu\nu} = L_{\mu}^{\hat{\alpha}} L_{\nu}^{\hat{\beta}} \eta_{\hat{\alpha}\hat{\beta}}, \quad (29)$$

where the $L_{\mu}^{\hat{\alpha}}$'s are the coefficients of the transformation between the two bases:

$$L_{\hat{t}}^t = \sqrt{|g_{tt}|}, \quad L_{\hat{r}}^r = \sqrt{|g_{rr}|}, \quad L_{\hat{\vartheta}}^{\vartheta} = \sqrt{|g_{\vartheta\vartheta}|}, \quad L_{\hat{\varphi}}^{\varphi} = \sqrt{|g_{\varphi\varphi}|}. \quad (30)$$

We therefore follow the prescription set out in [45] to rid ourselves of the unwanted time parameter: we assume a stationary source and integrate the interference term, $\exp[-i\Delta\Phi_{kj}^{pq}]$, over the unknown emission time t_A (or, equivalently, the transmission time $T \equiv t_B - t_A$). This results in a very useful $\delta(E_j - E_k)$.

Note here that though the energies of different mass eigenstates are different [49]—so that the $\delta(E_k - E_j)$ arising from the time integration would seem to imply no interference—in fact, in a correct treatment, massive neutrinos are described by wave packets, not plane waves as here. This means that, though the average energies of different mass eigenstate wave packets are, in general, different, each massive neutrino wave packet has an energy spread and the

So we have, in particular, that the local energy is given by [32]

$$E_k^{(loc)}(r) = |g_{tt}|^{-1/2} E_k = B(r)^{-1/2} E_k. \quad (31)$$

A. Calculating the phase difference

Given the above definitions, we now have that

$$\begin{aligned} \Phi_j^p(B, A) &= \int_A^B p_\mu^{(j)} dx^\mu \\ &= \int_A^B [E_j dt - p_j(r) dr_p - J_j d\varphi_p], \end{aligned} \quad (32)$$

where we have implicitly defined $p_j(r) \equiv -p_r^{(j)}$. Note that we have explicitly introduced the path index p which allows for the possibility of multiple paths from source to detector. Again, however, the integration over t is independent of the path as the end points of this integration are defined by the emission event and detection events. In fact, as discussed above, E_j is conserved over classical paths, so that if mass eigenstate j is assumed to travel down such a path, we can calculate the phase it accumulated after leaving the source to be

$$\begin{aligned} \Phi_j^p(B, A) &= \int_{t_A}^{t_B} E_j dt - \int_{r_A}^{r_B} \left[p_j(r) + J_j \left(\frac{d\varphi}{dr} \right)_j^p \right] dr_p \\ &= E_j (t_B - t_A) - \int_{r_A}^{r_B} \left[p_j(r) + J_j \left(\frac{d\varphi}{dr} \right)_j^p \right] dr_p. \end{aligned} \quad (33)$$

Of course, the quantity that governs the oscillation phenomenology is the phase difference $\Delta\Phi_{kj}^{pq}$ where, generically, interference can be between different mass eigenstates and/or different paths (cf. discussion in Sec. III). As things stand this quantity would be parametrized in terms of both t and r :

$$\Delta\Phi_{jk}^{pq}(r_B, t_B, r_A, t_A) = (E_j - E_k)(t_B - t_A) - \left\{ \int_{r_A}^{r_B} \left[p_j(r) + J_j \left(\frac{d\varphi}{dr} \right)_j^p \right] dr_p - \int_{r_A}^{r_B} \left[p_k(r) + J_k \left(\frac{d\varphi}{dr} \right)_k^q \right] dr_q \right\}. \quad (34)$$

detection process *can* pick up the same energy component for different massive neutrinos (see Refs. [45,50]). If the energy spread of the wave packets is small there is a suppression factor that, formally, can only be calculated only with a wave packet treatment (cf. Sec. III), but can also be assessed at the heuristic level (cf. Sec. VI).

Let us see how all the above works in practice.

B. Radial propagation

We consider first the simple case of radial propagation, in which case there is a single classical path from source to detector. Along this path, the angular momentum vanishes and we have

$$\Delta\Phi_{jk}(r_B, t_B, r_A, t_A) = (E_j - E_k)(t_B - t_A) - \int_{r_A}^{r_B} [p_j(r) - p_k(r)] dr. \quad (35)$$

We can determine $p_j(r) - p_k(r)$ from the mass-shell relation, Eq. (28) [29],

$$p_k(r) = \pm \frac{1}{B(r)} \sqrt{E_k^2 - B(r)m_k^2}, \quad (36)$$

where the $+$ sign refers to neutrinos propagating outwards from the gravitational well and the $-$ sign refers to neutrinos propagating inwards. We can further simplify this relation by employing the binomial expansion which, as in the flat space case, holds for relativistic particles,

$$\sqrt{E_k^2 - B(r)m_k^2} \simeq E_k - B(r) \frac{m_k^2}{2E_k}, \quad (37)$$

where E_0 is the energy at infinity for a neutrino mass eigenstate in the massless limit [see [29] for a detailed account of the region of applicability of Eq. (37)]. We therefore have that

$$p_j(r) - p_k(r) \simeq \pm \frac{1}{B(r)} (E_j - E_k) \mp \left(\frac{m_j^2}{2E_j} - \frac{m_k^2}{2E_k} \right). \quad (38)$$

The phase difference then becomes

$$\Delta\Phi_{jk}(r_B, t_B, r_A, t_A) \simeq (E_j - E_k) \left[(t_B - t_A) \mp \int_{r_A}^{r_B} \frac{dr}{B(r)} \right] + \left(\frac{m_j^2}{2E_j} - \frac{m_k^2}{2E_k} \right) |r_B - r_A|. \quad (39)$$

Given the oscillation ‘‘probability’’ shall be, following our previously established procedure (cf. Sec. III), integrated over $T \equiv t_B - t_A$, the *relevant* phase difference can be seen to be

$$\begin{aligned} \Delta\Phi_{jk}(r_B, r_A) &\simeq \frac{\delta m_{jk}^2}{2E_\nu} |r_B - r_A| \\ &\simeq \frac{\delta m_{jk}^2}{2E_0} |r_B - r_A|, \end{aligned} \quad (40)$$

where $E_\nu = E_j = E_k$ and E_0 is the energy at infinity for a massless particle and, as in flat space, the following relation holds [29]:

$$E_k \simeq E_0 + \mathcal{O}\left(\frac{m_k^2}{2E_0}\right). \quad (41)$$

To digress a little, note that the result presented in Eq. (40), arrived at previously [29,30,32,51], must be interpreted with some care: in Eq. (40) one must keep in mind that the radial distance $|r_B - r_A|$ is a *coordinate distance*, and not the proper distance the various mass eigenstates experience [except in

the flat space case to which Eq. (40) clearly reduces in the limit of a vanishing lensing mass] and that E_0 does not represent a locally detected energy. Following [29], however, we can convert the phase difference so that it appears in terms of these parameters. The proper distance is given by [cf. Eq. (30)]

$$L_{\text{prop}} \equiv \int_{r_A}^{r_B} \sqrt{g_{rr}} dr \quad (42)$$

$$\simeq r_B - r_A + GM \ln \frac{r_B}{r_A}, \quad (43)$$

where in the second line we have assumed the weak field limit holds. This demonstrates that, in a gravitational field, the length relevant to the calculation of phases, $|r_B - r_A|$, is actually shorter than the distances experienced by the propagating particles, L_{prop} . Substituting Eqs. (31) and (42) into Eq. (40) we determine that (cf. [29])

$$\Delta\Phi_{jk}(r_B, r_A) \simeq \left(\frac{\Delta m_{jk}^2 L_{\text{prop}}}{2E_0^{(loc)}(r_B)} \right) \left[1 - GM \left(\frac{1}{L_{\text{prop}}} \ln \frac{r_B}{r_A} - \frac{1}{r_B} \right) \right]. \quad (44)$$

C. Nonradial propagation

We turn now to the more interesting case presented by nonradial propagation. Here there will be, generically, more than one path for the mass eigenstates to take from source to detector and we have the possibility, therefore, of interference between particles on these different paths.

The phase difference we must calculate is given by Eq. (34). To proceed with this calculation we must determine a value for

$$p_j(r) + J_j \left(\frac{d\phi}{dr} \right)_j^p.$$

Firstly recall that J_j is constant along the classical path taken by ν_j . Now, using the fact [see [52], Eq. (101.5)] that

$$\begin{aligned} \left(\frac{d\phi}{dr} \right)_j^p &= \pm \frac{J_j^p}{r^2 \sqrt{E_j^2 - \left(m_j^2 + \frac{(J_j^p)^2}{r^2} \right) B(r)}} \\ &= \pm \frac{b_p \left(1 - \frac{m_j^2}{2E_j} \right)}{r^2 \sqrt{1 - \frac{B(r)}{E_j^2} \left[m_j^2 + \frac{E_j^2 b_p^2}{r^2} \left(1 - \frac{m_j^2}{E_j^2} \right) \right]}}, \end{aligned} \quad (45)$$

and, given that from the mass-shell relation we have [see Eq. (49) of [29]]

$$B(r)p_j(r) \simeq \pm \sqrt{1-B(r)} \frac{b_p^2}{r^2} \times \left[1 - \frac{B(r)(1-b_p^2/r^2)}{1-B(r)b_p^2/r^2} \frac{m_j^2}{2E_j^2} \right], \quad (46)$$

where b_p is the impact parameter for path p , one may determine that

$$p_j(r) + J_j^p \left(\frac{d\phi}{dr} \right)_j^p \simeq \pm \frac{E_j}{B(r) \sqrt{1-B(r)} \frac{b_p^2}{r^2}} \mp \frac{m_j^2}{2E_j^2} \frac{1 + [1-2B(r)] \frac{b_p^2}{r^2}}{\left(1 - B(r) \frac{b_p^2}{r^2} \right)^{3/2}}. \quad (47)$$

In the above we have also employed the fact that the angular momentum of mass eigenstate j (traveling along path p) at

infinity is given in terms of ν_j 's energy at infinity, E_j , the impact parameter along the path being considered, b_p , and ν_j 's velocity at infinity, $v_j^{(\infty)}$ [29],

$$J_j^p = E_j b_p v_j^{(\infty)} \simeq E_j b_p \left(1 - \frac{m_j^2}{2E_j^2} \right). \quad (48)$$

We can further evaluate Eq. (47) by replacing the path-dependent impact parameter, b_p , with the minimal radial coordinate for the same path, r_0^p . The relation between these two is found by noting that at the position of closest approach the rate of change of the coordinate r with respect to the angle ϕ vanishes [29]. For the massive case [Eq. (45)], this implies that

$$b_p^2 \simeq \frac{1 + \frac{m_j^2}{E_j^2} \frac{2GM}{r_0^p}}{B(r_0^p)}. \quad (49)$$

Employing Eq. (49), taking the weak field limit, and also expanding to $\mathcal{O}(m_j^2/E_j^2)$ we find that

$$p_j(r) + J_j^p \left(\frac{d\phi}{dr} \right)_j^p \simeq \pm E_j \left[\frac{r}{\sqrt{r^2 - (r_0^p)^2}} + GM \frac{2r - 3r_0^p}{\sqrt{r^2 - (r_0^p)^2}(r + r_0^p)} \right] \mp \frac{m_j^2}{E_j} \left[\frac{r}{\sqrt{r^2 - (r_0^p)^2}} - GM \frac{r_0^p}{\sqrt{r^2 - (r_0^p)^2}(r + r_0^p)} \right]. \quad (50)$$

With this result in hand, we can complete the calculation of Eq. (33), the phase accumulated by mass eigenstate in nonradial propagation from spacetime position $A=(r_A, t_A)$ to $B=(r_B, t_B)$, where either r_A or r_B is the minimal radial coordinate encountered over the journey (i.e., the path is either nonradially inwards *or* outwards but not both). After an elementary integration we find that

$$\Phi_j^p(B, A) \simeq E_j(t_B - t_A) \mp E_j \left[\sqrt{r_B^2 - (r_0^p)^2} - \sqrt{r_A^2 - (r_0^p)^2} + 2GM \ln \left(\frac{r_B + \sqrt{r_B^2 - (r_0^p)^2}}{r_A + \sqrt{r_A^2 - (r_0^p)^2}} \right) + GM \left(\sqrt{\frac{r_B - r_0^p}{r_B + r_0^p}} - \sqrt{\frac{r_A - r_0^p}{r_A + r_0^p}} \right) \right] \pm \frac{m_j^2}{2E_j} \left[\sqrt{r_B^2 - (r_0^p)^2} - \sqrt{r_A^2 - (r_0^p)^2} - GM \left(\sqrt{\frac{r_B - r_0^p}{r_B + r_0^p}} - \sqrt{\frac{r_A - r_0^p}{r_A + r_0^p}} \right) \right], \quad (51)$$

where the upper signs pertain if dr is positive (outward propagation) and the lower if dr is negative.

D. Neutrino lensing

Finally let us consider the case of gravitational lensing of neutrinos. In this case the neutrinos propagate nonradially along classical paths labeled by index p , from radial position r_A inwards to a path-dependent minimal radial coordinate r_0^p and outwards again to a detector situated at radial coordinate r_B . As presaged above, in this situation there will be (at least potentially) interference not only between different mass eigenstates propagating down the same classical path, but also between mass eigenstates propagating down different paths (p and q , say). Taking into account the sign of the momentum along these two legs, we find, following the developments above, that the relevant phase is given by

$$\Phi_j^p(B, A) \simeq E_j(t_B - t_A) - E_j \left[\sqrt{r_A^2 - (r_0^p)^2} + 2GM \ln \left(\frac{r_A + \sqrt{r_A^2 - (r_0^p)^2}}{r_0^p} \right) + GM \left(\sqrt{\frac{r_A - r_0^p}{r_A + r_0^p}} \right) + (r_A \rightarrow r_B) \right] + \frac{m_j^2}{2E_j} \left[\sqrt{r_A^2 - (r_0^p)^2} - GM \left(\sqrt{\frac{r_A - r_0^p}{r_A + r_0^p}} \right) + (r_A \rightarrow r_B) \right], \quad (52)$$

where $(r_A \rightarrow r_B)$ mean add another term of the same form but with r_A replaced with r_B .

Before proceeding any further with the calculation, it behooves us here to establish the plausibility of Eq. (52) by showing its relation to results known from some simpler cases. In the $M \rightarrow 0$ limit this equation becomes

$$\Phi_j^p(B,A) \approx E_j(t_B - t_A) - E_j \left(1 - \frac{m_j^2}{2E_j^2} \right) \left[\sqrt{r_A^2 - (r_0^p)^2} + \sqrt{r_B^2 - (r_0^p)^2} \right]. \quad (53)$$

Now refer back to Fig. 1, and take the coordinate origin on the diagram to denote the position of a lensing point mass. In the massless case, the two classical paths reduce to the single “undeflected” path denoted by the dashed line in the diagram. Denote the minimal radial coordinate along this path by r_0 (which intersects the dashed line at right angles). Clearly, then, the geometrical length of the path from source to detector is $\sqrt{r_A^2 - r_0^2} + \sqrt{r_B^2 - r_0^2}$. Now, given we know that in flat space the phase of mass eigenstate k is given by Eq. (21) with Minkowski metric, viz.,

$$\Phi_j = E_j(t_B - t_A) - \mathbf{p}_j \cdot (\mathbf{x}_B - \mathbf{x}_A), \quad (54)$$

then, for the $M \rightarrow 0$ case illustrated in Fig. 1 this becomes

$$\Phi_j \approx E_j(t_B - t_A) - E_j \left(1 - \frac{m_j^2}{2E_j^2} \right) (\sqrt{r_A^2 - r_0^2} + \sqrt{r_B^2 - r_0^2}). \quad (55)$$

With Eq. (55) we have then independently established the plausibility of Eq. (53), once one takes into account the fact that, in the massless lens case, all classical paths converge on the same undeflected path (as mentioned above) so that in this limit $r_0^p = r_0^q \equiv r_0 \forall p, q$.

The other limit of interest is to take $m_j \rightarrow 0$ in Eq. (52). In doing this—and then setting the temporal and spatial contributions to the phase equal as appropriate for a null geodesic—we find that we have rederived the Shapiro time delay [see, e.g., Eq. (8.7.4) of Ref. [53]].

Continuing with our main calculation, we can rewrite Eq. (52) in terms of b_p by inverting Eq. (49). If we also expand to $\mathcal{O}(b_p^2/r_{B \leftrightarrow A}^2)$, we find that

$$\begin{aligned} \Phi_j^p(B,A) \approx E_j(t_B - t_A) - E_j(r_A + r_B) & \left\{ 1 - \frac{b_p^2}{2r_A r_B} \right. \\ & + \frac{2GM}{r_A + r_B} \left[1 + \ln \left(\frac{4r_A r_B}{b_p^2} \right) \right] \left. \right\} \\ & + \frac{m_j^2}{2E_j} (r_A + r_B) \left(1 - \frac{b_p^2}{2r_A r_B} - \frac{2GM}{r_A + r_B} \right). \end{aligned} \quad (56)$$

We can now find the phase difference, which allows for interference between different paths and/or different mass eigenstates, by the usual integration over T (so that we have $E_\nu = E_j = E_k$):

$$\begin{aligned} \Delta\Phi_{jk}^{pq}(r_B, r_A) \approx & + E_\nu(r_A + r_B) \left(\frac{\Delta b_{pq}^2}{2r_A r_B} + \frac{4GM}{r_A + r_B} \ln \left| \frac{b_p}{b_q} \right| \right) \\ & + \frac{\delta m_{jk}^2}{2E_\nu} (r_A + r_B) \left(1 - \frac{2GM}{r_A + r_B} \right) \\ & - \frac{r_A + r_B}{2E_\nu} \left(\frac{m_j^2 b_p^2 - m_k^2 b_q^2}{2r_A r_B} \right), \end{aligned} \quad (57)$$

where $\Delta b_{pq}^2 \equiv b_p^2 - b_q^2$ and, in our notation, $\Delta\Phi_{jk}^{pq}(r_B)$ denotes the phase difference between mass eigenstate j traveling down path p and mass eigenstate k traveling down path q [54]. Equation (57) is one of the major results of this paper. Note that the presence of the $\propto E_\nu$ term in this equation—missed in Ref. [29]—ensures that the phase difference behaves properly in the massless limit (i.e., does *not* vanish). In passing, also note that the above equation satisfies the discrete symmetry of swapping B and A , as it should: the same result must be obtained for the phase difference (in a static spacetime) if we swap the positions of source and observer.

Also recall that, excluding the case of perfect alignment of source, lens, and observer, there are only two possible classical paths from source to observer for the Schwarzschild case. These we label by $+$ (this path having an impact parameter somewhat greater than the impact parameter for an undeflected ray) and $-$ (this path having an impact parameter on the “opposite” side of the lens to the undeflected ray). We require, therefore, that $p, q \in \{+, -\}$ and in the particular case in which we are considering interference between the same mass eigenstates propagating down different paths, Eq. (57) becomes

$$\begin{aligned} \Delta\Phi_{jj}^{+-}(r_B, r_A) \approx & + E_\nu(r_A + r_B) \left(\frac{\Delta b_{+-}^2}{2r_A r_B} + \frac{4GM}{r_A + r_B} \ln \left| \frac{b_+}{b_-} \right| \right) \\ & - \frac{m_j^2}{2E_\nu} (r_A + r_B) \frac{\Delta b_{+-}^2}{2r_A r_B}. \end{aligned} \quad (58)$$

Alternatively, in the case of different mass eigenstates traveling down the same path (i.e., “ordinary” neutrino oscillations, but in curved space), Eq. (57) becomes

$$\begin{aligned} \Delta\Phi_{jk}^{pp}(r_B, r_A) \approx & \frac{\delta m_{jk}^2}{2E_\nu} (r_A + r_B) \\ & \times \left(1 - \frac{b_p^2}{2r_A r_B} - \frac{2GM}{r_A + r_B} \right). \end{aligned} \quad (59)$$

Note that Eqs. (57), (58), and (59) give us that

$$\begin{aligned} \Delta\Phi_{jk}^{pq}(r_B, r_A) &= \Delta\Phi_{jj}^{pq}(r_B, r_A) + \Delta\Phi_{jk}^{pp}(r_B, r_A) \\ &+ \mathcal{O}\left[\frac{\delta m_{jk}^2}{2E_\nu}(r_A + r_B)\frac{\Delta b_{pq}^2}{2r_A r_B}\right]. \end{aligned} \quad (60)$$

This correction term will be small with respect to other terms (given our assumptions of ultrarelativistic neutrinos and undeflected impact parameters small with respect to the overall distances between source-lens and lens-observer). In fact, the third term of Eq. (57) can be expected to be suppressed with respect to the first term by $\mathcal{O}(m^2/E_\nu^2)$ and with respect to the second term by $\mathcal{O}[b^2/(r_A r_B)]$. The consequence of this is that the phase may be written

$$\Delta\Phi_{jk}^{pq} \approx \Delta\Phi_{jk} + \Delta\Phi^{pq}, \quad (61)$$

satisfying what we label ‘‘separability,’’ where

$$\begin{aligned} \Delta\Phi^{pq} \equiv E_\nu(r_A + r_B) &\left(\frac{\Delta b_{pq}^2}{2r_A r_B} + \frac{4GM}{r_A + r_B} \ln\left|\frac{b_p}{b_q}\right|\right) \\ &- \frac{\bar{m}^2}{2E_\nu}(r_A + r_B)\frac{\Delta b_{pq}^2}{2r_A r_B} \end{aligned} \quad (62)$$

and

$$\Delta\Phi_{jk} \equiv \frac{\delta m_{jk}^2}{2E_\nu}(r_A + r_B) \left(1 - \frac{\bar{b}^2}{2r_A r_B} - \frac{2GM}{r_A + r_B}\right), \quad (63)$$

where

$$\bar{m} \equiv \frac{1}{N_\nu} \sum_j^{N_\nu} m_j \quad \text{and} \quad \bar{b} \equiv \frac{1}{N_{\text{path}}} \sum_j^{N_{\text{path}}} b_p, \quad (64)$$

with N_ν the number of neutrino mass eigenstates and N_{path} the number of classical paths from source to detector (two in the case of the Schwarzschild metric). What Eq. (61) says in words is that the phase difference that develops between source and detector is due to two effects that can be considered separately: (i) a phase difference—independent of which the mass eigenstate is under consideration—that develops because of the different lengths of the paths involved and (ii) the phase difference that develops because the different mass eigenstates travel with different phase velocities. This situation is analogous to two runners who run along two very similar—though not identical—paths, with similar—though not identical—velocities: to first order, the difference in the finishing times between the two depends on terms proportional to the difference in lengths of the two courses, ΔL , and the difference in the runners’ velocities, Δv , but not, by definition, on terms $\propto \Delta L \Delta v$.

E. Phase difference in terms of conventional lensing parameters

To facilitate interpretation of the above results in an astrophysical context—and, eventually, to introduce an evolving cosmological model—it is useful to reexpress the phase

difference in the language of standard gravitational lensing theory (despite the fact that the particles being lensed are not photons).

1. The lens equation

The classically allowed neutrino paths in the presence of a deflector can be derived by reconsidering the geometry shown in Fig. 1. Under the assumption that $|b_\pm| \ll r_{\{A,B\}}$, the source offset, s , can be related to the impact parameter, b , by the lens equation,

$$s \approx \frac{r_B}{r_A + r_B} b + r_A \alpha(b), \quad (65)$$

where $\alpha(b)$ is the deflection angle of the lens as a function of impact parameter.

It is standard practice to reexpress the lens equation in terms of angular variables. This entails replacing the source offset and impact parameters with angles (on the sky of the observer) and radial coordinates with line-of-sight distances. These conversions are summarized graphically in Fig. 1, which leads to the following replacements: $r_A \rightarrow (D_{ds}^2 + s^2)^{1/2} \approx D_{ds}$, where D_{ds} is the distance from deflector to source; $r_B \rightarrow D_d$, where D_d is the distance from observer to deflector; and $r_A + r_B \rightarrow D_s$, where D_s is the distance from observer to source. The notation employed for the distance measures is suggestive of these being the angular diameter distances used to relate angles and lengths in an evolving cosmological model, and they fulfill an analogous role here. It is most important to note, however, that they are not true angular diameter distances and the following results are only quantitatively valid on scales sufficiently small that the expansion of the Universe can be ignored (e.g., the Milky Way or the Local Group). These results will be extended to an evolving cosmology in [55].

The above caveats notwithstanding, the angular position (relative to the deflector) of an image with impact parameter b is now simply

$$\theta \approx \frac{b}{r_B} \approx \frac{b}{D_d}, \quad (66)$$

and the position of the source can be given in terms of an unobservable angular parameter β as

$$\beta \approx \frac{s}{r_A + r_B} \approx \frac{s}{D_s}. \quad (67)$$

Inserting these definitions into Eq. (65), the lens equation becomes

$$\beta \approx \theta + \frac{D_{ds}}{D_s} \alpha(D_d \theta). \quad (68)$$

The position(s) of the images formed by a source in a given position can then be found for a given choice of deflector model.

2. The Schwarzschild lens

In a Schwarzschild metric, the total angular deflection of a particle of mass m impinging on a point-mass M with undeflected impact parameter b is (see, e.g., [56])

$$\alpha_{\text{gen}}(b) = -\frac{4GM}{b} \frac{1}{2} \left(1 + \frac{1}{v_\infty^2} \right), \quad (69)$$

where v_∞ is the particle's speed at an infinite distance from the mass and it has been assumed that $b \gg 2GM = R_S$, the deflector's Schwarzschild radius. For an ultrarelativistic particle, this becomes

$$\alpha_{\text{rel}}(b) \approx -\frac{4GM}{b} \left(1 + \frac{m^2}{2E^2} \right), \quad (70)$$

where E is its coordinate energy (equal to the energy measured at an infinite distance from the mass).

For astrophysical neutrinos, however, $m^2/(2E^2) \ll 1$ and so it is an excellent approximation to assume they travel along classical photon paths, for which

$$\alpha_{\text{light}}(b) = -\frac{4GM}{b}. \quad (71)$$

Previously we have been rigorous in taking the classical paths of massive particles from source to observer but, as will be seen below, this assumption is entirely self-consistent when dealing with weak-field gravitational effects and ultrarelativistic particles. Note also that a corollary of this approximation is that the different mass eigenstates are assumed to travel down identical paths (whereas in reality the heavier eigenstates will fall marginally deeper into the deflector's potential well).

Applying the above deflection law to Eq. (68) gives the point-mass lens equation as

$$\beta = \theta - \frac{\theta_E^2}{\theta}, \quad (72)$$

where

$$\theta_E = \sqrt{4GM \frac{D_{ds}}{D_d D_s}} \quad (73)$$

is the Einstein radius of the lens. This is the angular radius of the circular image that would be formed in the case of perfect source-deflector-observer alignment (i.e., $\beta = 0$) and thus depends on distance factors as well as the lens mass. Solving the lens equation then gives the image positions as

$$\theta_\pm \approx \frac{1}{2} (\beta \pm \sqrt{\beta^2 + 4\theta_E^2}). \quad (74)$$

This also implies the useful Schwarzschild-specific result that

$$\theta_+^2 - \theta_-^2 = \Delta\theta_{+-}^2 \approx \beta(\theta_+ - \theta_-). \quad (75)$$

Having found a relationship between the angular position of a neutrino source and its images, the expression for the phase difference given in Eq. (57) can be recast in a form containing only line-of-sight distances and angular variables. This yields the Schwarzschild-specific result that

$$\begin{aligned} \Delta\Phi_{jk}^{pq} \approx & E_\nu \frac{D_d D_s}{D_{ds}} \left[\frac{\Delta\theta_{pq}^2}{2} + \theta_E^2 \ln \left(\left| \frac{\theta_p}{\theta_q} \right| \right) \right] \\ & + \frac{\delta m_{jk}^2}{2E_\nu} D_s \left(1 - \frac{D_d}{D_{ds}} \theta_E^2 \right) \\ & - \frac{1}{2E_\nu} \frac{D_d D_s}{D_{ds}} \frac{m_j^2 \theta_p^2 - m_k^2 \theta_q^2}{2}. \end{aligned} \quad (76)$$

The second term in this equation is simply the phase difference that develops between mass eigenstates j and k traveling along the same path for distance D_s , with a small correction for the presence of the deflector. The first term encodes the path difference along the trajectories p and q , with separate contributions from the geometrical effect ($\propto \Delta\theta_{pq}^2$) and the reduced coordinate velocity close to the deflector [$\propto \ln(|\theta_p/\theta_q|)$]. The final cross term is the leading order contribution from different eigenstates traveling down different paths. From the discussion in the previous section, this term will be small in general.

Given that interference effects can only ever be important when the detector cannot resolve different image positions (i.e., it cannot know down which path the neutrino has traveled), having the phase difference in terms of θ_p and θ_q is not as useful as expressing it as a function of the (angular) source position, β .

For the Schwarzschild lens the conversion from θ to β is given in Eq. (72), and substituting this into Eq. (76) then gives (for mass eigenstate j down path + and mass eigenstate k down path -)

$$\begin{aligned} \Delta\Phi_{jk}^{+-} \approx & E_\nu \frac{D_d D_s}{D_{ds}} \left[\frac{\beta \sqrt{\beta^2 + 4\theta_E^2}}{2} + \theta_E^2 \ln \left(\left| \frac{\beta + \sqrt{\beta^2 + 4\theta_E^2}}{\beta - \sqrt{\beta^2 + 4\theta_E^2}} \right| \right) \right] \\ & + \frac{\delta m_{jk}^2}{2E_\nu} D_s \left(1 - \frac{D_d}{D_{ds}} \theta_E^2 \right) \\ & - \frac{1}{2E_\nu} \frac{D_d D_s}{D_{ds}} \frac{1}{4} [\delta m_{jk}^2 (\beta^2 + 2\theta_E^2) \\ & + (m_j^2 + m_k^2) \beta \sqrt{\beta^2 + 4\theta_E^2}]. \end{aligned} \quad (77)$$

Thus the phase difference is expressed in terms of essentially independent astronomical variables: the line-of-sight distances between observer, deflector and source, the mass of the deflector (encoded uniquely in θ_E once the distances have been chosen) and the perpendicular source offset, β .

Most of the important results obtained towards the end of Sec. IV D can be recast similarly in terms of standard lensing variables, either in terms of the unobservable image positions or the source position. Assuming separability (see Sec.

IV D), for instance, the contribution to the $\Delta\Phi$ due to path difference effects alone [Eq. (62)] can be written as

$$\Delta\Phi^{pq} \simeq E_\nu \frac{D_d D_s}{D_{ds}} \left[\frac{\Delta\theta_{pq}^2}{2} + \theta_E^2 \ln \left(\left| \frac{\theta_p}{\theta_q} \right| \right) \right] - \frac{\bar{m}^2}{2E_\nu} \frac{D_d D_s}{D_{ds}} \frac{\Delta\theta_{pq}^2}{2}, \quad (78)$$

which for $p = +$ and $q = -$ becomes

$$\begin{aligned} \Delta\Phi^{+-} \simeq E_\nu \frac{D_d D_s}{D_{ds}} & \left[\frac{\beta \sqrt{\beta^2 + 4\theta_E^2}}{2} \right. \\ & \left. + \theta_E^2 \ln \left(\left| \frac{\beta + \sqrt{\beta^2 + 4\theta_E^2}}{\beta - \sqrt{\beta^2 + 4\theta_E^2}} \right| \right) \right] \\ & - \frac{\bar{m}^2}{2E_\nu} \frac{D_d D_s}{D_{ds}} \frac{\beta \sqrt{\beta^2 + 4\theta_E^2}}{2}. \end{aligned} \quad (79)$$

Similarly, the contribution due solely to the different phase velocities of two mass eigenstates traveling down the same path [Eq. (63)] can be expressed as

$$\Delta\Phi_{jk} \simeq \frac{\delta m_{jk}^2}{2E_\nu} D_s \left[1 - \frac{\theta_E^2}{8} (\beta^2 + 4\theta_E^2) \right]. \quad (80)$$

V. THE OSCILLATION “PROBABILITY”

With the above results, we can now calculate the analog, in curved space, of the flat-space neutrino oscillation probability:

$$|\langle \nu_\beta | \nu_\alpha, D_s \rangle|^2 \propto \int dT |\langle \nu_\beta | \nu_\alpha; A, B \rangle|^2 \quad (81)$$

so that we can write

$$\begin{aligned} |\langle \nu_\beta | \nu_\alpha, D_s \rangle|^2 = & |N|^2 \sum_{pq} \sqrt{I_p I_q} \sum_{jk} U_{\alpha j} U_{\beta j}^* U_{\beta k} U_{\alpha k}^* \\ & \times \exp[-i\Delta\Phi_{jk}^{pq}(D_s)], \end{aligned} \quad (82)$$

where I_p and I_q account for the fact that different paths may be differentially magnified by a lens. We remind the reader that $|\langle \nu_\beta | \nu_\alpha, D_s \rangle|^2$ is no longer strictly a probability—see Sec. III.

In the case that the “separability” defined by Eq. (61) is satisfied, if, for the moment, we are interested only in determining the (energy) spacing of the interference maxima and minima, we need only consider a plane-wave-like calculation (and can therefore set to one side the coherence length effects and so on that emerge from a wave packet calculation). So, following considerations similar to those that lead to Eq. (A14) in the Appendix we can calculate that

$$\begin{aligned} |\langle \nu_\beta | \nu_\alpha, D_s \rangle|^2 = & |N|^2 \left[N_{\text{path}} I + 2 \sum_{p,q < p} \sqrt{I_p I_q} \cos(\Delta\Phi^{pq}) \right] \\ & \times \left[\delta_{\alpha\beta} - 4 \operatorname{Re} \left\{ \sum_{j,k < j} U_{\alpha j} U_{\beta j}^* U_{\alpha k}^* U_{\beta k} \left[\sin^2 \left(\frac{\Delta\Phi_{jk}}{2} \right) + \frac{i}{2} \sin(\Delta\Phi_{jk}) \right] \right\} \right], \end{aligned} \quad (83)$$

where N_{path} is again the number of classical paths from source to detector (two in the case of a Schwarzschild metric), $I \equiv \sum_p I_p / N_{\text{path}}$, and the normalization, $|N|^2$ is again given (cf. Sec. III) by requiring that $\sum_\beta |\langle \nu_\beta | \nu_\alpha, D_s \rangle|^2 \leq 1$, i.e.,

$$\max\{\langle \nu_\alpha, B | \nu_\alpha, B \rangle\} = 1. \quad (84)$$

This means that

$$|N|^2 = \frac{1}{N_{\text{path}} I + 2 \sum_{p,q < p} \sqrt{I_p I_q}}. \quad (85)$$

Equation (83) establishes the contention made above that interference effects that emerge with gravitationally lensed neutrinos are a combination of Young’s double slit type interference {the $[\dots \cos(\Delta\Phi^{pq}) \dots]$ envelope term} and flat space oscillations [the $(\delta_{\alpha\beta} - \dots)$ term]. Further, assuming the separability requirement is satisfied, we can see how

$|\langle \nu_\beta | \nu_\alpha, D_s \rangle|^2$ factorizes into an interference pattern and a conditional probability. This is a repeat of the behavior seen in Sec. III A.

One should also note that in the particular case of the Schwarzschild lens under consideration in the last section, the two (assuming nonperfect alignment) classical paths from source to detector, denoted by the subscripts $+$ and $-$, experience magnifications given by [see Eq. (2.24) of Ref. [56]]

$$I_\pm = \frac{1}{4} \left(\frac{\beta}{\sqrt{\beta^2 + 4\theta_E^2}} + \frac{\sqrt{\beta^2 + 4\theta_E^2}}{\beta} \pm 2 \right). \quad (86)$$

This gives us that

$$|N|^2 = \frac{1}{I_+ + I_- + 2\sqrt{I_+ I_-}} = \frac{\beta}{\sqrt{\beta^2 + 4\theta_E^2}}. \quad (87)$$

Equation (83) then becomes

$$|\langle \nu_\beta | \nu_\alpha, D_s \rangle|^2 \simeq \frac{1}{\beta^2 + 4\theta_E^2} \{ \beta^2 + 2\theta_E^2 [1 + 2\cos(\Delta\Phi^{+-})] \} \\ \times \left[\delta_{\alpha\beta} - 4 \operatorname{Re} \left\{ \sum_{j,k < j} U_{\alpha j} U_{\beta j}^* U_{\alpha k}^* U_{\beta k} \left[\sin^2 \left(\frac{\Delta\Phi_{jk}}{2} \right) + \frac{i}{2} \sin(\Delta\Phi_{jk}) \right] \right\} \right], \quad (88)$$

with $\Delta\Phi^{+-}$ and $\Delta\Phi_{jk}$ given by Eqs. (79) and (80) respectively (where, again, care should be taken not to confuse β as a label on the neutrino flavor with β as the source angular position).

Another result of interest is that for the *magnification*, μ_{ν_β} , which is defined to be the ratio of the flux of neutrinos of type β actually received (from the source at Y and given the lensing mass is where it is) to the flux of neutrinos of the same type that would be received with the lens absent (but with the source in the same position),

$$\mu_{\nu_\beta} = \frac{F(D_s, E_\nu) \sum_\alpha |\langle \nu_\beta | \nu_\alpha, D_s \rangle|_{\text{lens}}^2 \times P_\alpha}{F(D_s, E_\nu) \sum_\alpha |\langle \nu_\beta | \nu_\alpha, D_s \rangle|_{\text{no lens}}^2 \times P_\alpha}, \quad (89)$$

where $F(D_s, E_\nu)$ denotes the flux of neutrinos of all types that would be received, at an energy of E_ν and factoring in geometrical effects, in the absence of the lens. Also, P_α denotes the probability that a neutrino generated by the source under consideration is of type α . Now, for the Schwarzschild lens, assuming mass degeneracy, the path difference and phase velocity contributions to $|\langle \nu_\beta | \nu_\alpha, D_s \rangle|^2$ factorize into an interference pattern and a conditional probability, as mentioned above [Eq. (83)]. This has the effect that the magnification is independent of the neutrino flavor under consideration,

$$\mu_{\nu_\beta} \simeq \frac{\sum_\alpha |N|^2 [I_+ + I_- + 2\sqrt{I_+ I_-} \cos(\Delta\Phi^{+-})] P(\alpha \rightarrow \beta)_{\text{flat}} \times P_\alpha}{\sum_{\alpha'} |N|^2 P(\alpha' \rightarrow \beta)_{\text{flat}} \times P_{\alpha'}} = [I_+ + I_- + 2\sqrt{I_+ I_-} \cos(\Delta\Phi^{+-})] \\ = \frac{1}{\beta\sqrt{\beta^2 + 4\theta_E^2}} \{ \beta^2 + 2\theta_E^2 [1 + \cos(\Delta\Phi^{+-})] \}, \quad (90)$$

where $P(\alpha \rightarrow \beta)_{\text{flat}}$, the flat space neutrino oscillation probability, is given by

$$P(\alpha \rightarrow \beta)_{\text{flat}} = \left[\delta_{\alpha\beta} - 4 \operatorname{Re} \left\{ \sum_{j,k < j} U_{\alpha j} U_{\beta j}^* U_{\alpha k}^* U_{\beta k} \right. \right. \\ \left. \left. \times \left[\sin^2 \left(\frac{\Delta\Phi_{jk}}{2} \right) + \frac{i}{2} \sin(\Delta\Phi_{jk}) \right] \right\} \right]. \quad (91)$$

The result for the magnification is as expected given what is known about the photon case [see Eq. (9) of Ref. [4]].

Finally for this section, we determine, for future reference, the fringe *visibility*, $\mathcal{V}(\beta)$,

$$\mathcal{V}(\beta) \equiv \frac{\mu_{\nu_\beta}^{\max} - \mu_{\nu_\beta}^{\min}}{\mu_{\nu_\beta}^{\max} + \mu_{\nu_\beta}^{\min}} \simeq \frac{2\theta_E^2}{\beta^2 + 2\theta_E^2}. \quad (92)$$

VI. PHENOMENOLOGY: HEURISTIC CONSIDERATIONS

Above we have presented the calculation of the phase and the consequent phase difference, oscillation probability

analog— $|\langle \nu_\beta | \nu_\alpha, D_s \rangle|^2$ —which determines the form of the oscillation pattern seen at a detector, and magnification factor, all for the Schwarzschild lens. We now turn briefly to the question of the phenomenological consequences of all these theoretical developments. We shall deal with the issues presented here at greater length in another work [55]. There are a number of factors which broadly determine the visibility of GINI effects [57]:

(i) *Suitability of potentially lensed sources*. The first consideration must be, what qualifies as a suitable source? We require sources that produce a neutrino signal that might be both gravitationally lensed and of sufficient intensity.

(ii) *Geometrical optics limit*. Because our theoretical evaluation for the neutrino phase difference has been performed within the geometrical optics limit (where only the classical paths from source to detector need be considered in determining the form of the interference pattern), we require that this limit holds in the experimental situation under consideration. This translates to the requirement that the de Broglie wavelength of the neutrino mass eigenstates is not larger than the Schwarzschild radius of the lens (the latter quantity setting the scale of the path difference: see below) [4,6,11,58].

(iii) *Detector energy resolution.* Even if there exists an interference pattern to be mapped out—and sufficient events to achieve this—a separate question is whether the smearing of this pattern caused by the finite energy resolution of any real neutrino detector is so large as to completely wash it out.

(iv) *Just-so condition for lensing mass.* Points (ii) and (iii) imply a range for a “just-right” lensing mass (given the energy scale of the neutrinos is already set)—not too large and not too small—inside which GINI effects may become evident. This can be roughly determined by the following considerations: for a (point mass) lensing system to produce images of similar brightness (so that interference effects might be seen), we require that source to be sufficiently well aligned with the line from the observer through the lens (i.e., $\beta \lesssim \theta_E$). Granted this, the scale of the path difference is then set by the Schwarzschild radius (see, e.g., [56], p. 240),

$$R_S \equiv 2GM \approx 3 \times 10^{-12} \left(\frac{M}{10^{-17} M_\odot} \right) \text{cm}, \quad (93)$$

of the lens (and—very broadly—can be considered as independent of the distance to the detector). Once one has settled on a generic astrophysical source which emits neutrinos in some characteristic energy range, the lensing mass range is determined. This is because we require

$$E_\nu \times R_S \approx 1 \quad (94)$$

at an energy either within or not too far below that detectable by the particular detector technology under consideration (see Sec. VI C below). We label this constraint on the lensing mass range the just-so condition.

(v) *Wave packet spreading and decoherence.* By analogy with the considerations set out in Sec. III, we expect that the full expression for the oscillation probability analog include exponential decay factors that account for coherence loss effects. These essentially factor in the interference attenuation which occurs when the different neutrino wave packets, traveling with different group velocities and/or along paths of different affine length, overlap significantly less than completely at the detector. See Sec. VI A for more detail on this issue.

(vi) *Finite source size effects.* Our derivation of the phase difference has assumed a stationary point source (and detector). Of course, this is at variance with Heisenberg uncertainty requirements. But more significantly, any real, *macroscopic* source (the region giving birth to all the neutrinos that are identified as having come from a particular astrophysical object) will be of finite—indeed macroscopic—size. This can, like detector energy resolution issues, tend to wash away the interference pattern because the path difference is now different for the various neutrinos that come from different parts of the “same” object. More concisely, an effective source angular extent of the order of—or larger than—the angular extent of the Einstein radius means that the visibility of the interference fringes is reduced [7]. If the source size is denoted by r_{source} , then this translates to the requirement that

$$r_{\text{source}} < \sqrt{4GM \frac{D_s D_{ds}}{D_d}}. \quad (95)$$

Equation (95) is not a sufficient condition to guarantee a pointlike source, however; as energy—and, therefore, phase along any particular path—increases, there will come a point where (while the lens-induced path difference might still generate the greatest component of the phase difference for neutrinos from all parts of the lens) the phase difference for neutrinos emerging from one part of the source will be noticeably different from that for neutrinos generated from a different part of the source. At this point the interference pattern will, again, become smeared out. That this does *not* occur bounds the energy,

$$E_\nu \lesssim \frac{2D_s D_{ds}}{D_d r_{\text{source}}^2}. \quad (96)$$

(vii) *Finite detector size effects.* Much of the discussion immediately above carries through, all things being equal, to considerations stemming from finite detector size. Explicitly, finite detector size effects can tend to wash away the interference pattern because the path difference (at any particular energy) will be nonconstant across the volume of the detector. One must determine whether this is a significant effect.

(viii) *Finite lens size effects.* We have calculated the neutrino phase difference in a Schwarzschild metric, i.e., assuming the lens to be effectively pointlike. This assumption will hold, at least roughly, if the Einstein radius of the lensing system [θ_E , as defined in Eq. (73)] is larger than the scale of the physical dimension of the lens.

(ix) *Source-lens-alignment probability.* In order to see interference fringes we require that the *visibility* [defined in Eq. (92)] be sufficiently good. This requires a sufficient degree of alignment between source, lens, and detector (i.e., a small Y or η). One can then ask, given the lensing mass scale, as determined by point (iv) above, and the expected distance to a source (of the chosen, generic type), how likely is it that there is a lens within a certain distance of the line from the source to the detector?

(x) *Time scale of lens crossing.* Further to the point immediately above, one must consider over what time scale the lens will cross the “beam” from source to detector and, therefore, how temporally stable—and, indeed, how long-lasting—any interference pattern will be.

(xi) *Intrinsic source spectrum.* In order to confidently identify interference effects one must be able to rule out the possibility of the intrinsic spectrum of the source mimicking these effects. Moreover, even given a well-understood source spectrum, a separate question is whether there is a measurable neutrino flux over a sufficient energy extent that a number of interference fringes might be seen at a detector.

A. More detail on decoherence effects in curved spacetime

We only attempt a heuristic treatment here [59]. Ignoring detector effects (see below), coherence requires that there be a significant overlap between mass eigenstates at a detector. As explained in the Appendix, the various mass eigenstates

may have traveled with both different group velocities and along different paths. Let us take a source located on a source plane at D_s and neutrino mass eigenstates with an effective width of σ_x . Then, by analogy with the second exponential damping term in Eq. (A14) of the Appendix and given the scale of the path difference is given by R_S , interference between mass eigenstate j traveling down one macroscopic path through a Schwarzschild spacetime and k the other roughly requires

$$\left(R_S \mp \frac{|\delta m_{jk}^2|}{2E_\nu^2} D_s \right)^2 \lesssim 8\sigma_x^2. \quad (97)$$

Here the upper sign refers to the case when the lighter mass eigenstate traverses the longer path, the heavier along the shorter path, and the lower sign refers to the opposite case (there are now four broad cases depending on this sign and the relative sizes of R_S and $2\sqrt{2}\sigma_x$).

Note that if we wish to consider interference between different mass eigenstates traveling down the *same* path—i.e., the direct analog of flat space neutrino oscillations—we take $R_S \rightarrow 0$ in Eq. (97) so that we require

$$D_s \lesssim \frac{2E_\nu^2}{|\delta m_{jk}^2|} 2\sqrt{2}\sigma_x, \quad (98)$$

(then the equality in the above is satisfied for $D_s \simeq L_{\text{coh}}$, where L_{coh} is the coherence length), whereas if we wish to consider interference between the same mass eigenstate traveling down different paths, then from Eq. (97) we require

$$R_S \lesssim 2\sqrt{2}\sigma_x. \quad (99)$$

Below we shall determine some plausible numbers to put in these relations (for a number of different neutrino sources), but first we recall some considerations behind the determination of σ_x .

B. Determining the size of the wave packet

At a heuristic level—adequate to the order of magnitude calculations we will make—the neutrino wave packet size (in position space) is given by the size, d , of the region to which the neutrino parent particle is localized [60–62]:

$$\sigma_x \simeq d. \quad (100)$$

In turn, d is related to t_{eff} , the effective time available for the coherent emission, by the parent of a neutrino wave train,

$$d \simeq t_{\text{eff}}. \quad (101)$$

In free space the coherent emission time corresponds to the decay time of the parent particle, τ , but if the parent particle is in a dense and hot medium and undergoing collisions with its neighbors on a time scale, $t_{\text{collision}}$, smaller than τ , then $t_{\text{eff}} \simeq t_{\text{collision}}$ [63,64]. This effect corresponds to the collision or pressure broadening of atomic spectral lines. In summary, we shall take

$$\sigma_x \simeq \gamma t_{\text{eff}} \quad \text{with} \quad t_{\text{eff}} = \min\{t_{\text{collision}}, \tau\}, \quad (102)$$

where we have explicitly introduced a Lorentz boost, γ , to allow for any bulk motion of the source with respect to the detector frame. This factor can, of course, be large for astrophysical sources.

C. Determining energy ranges for GINI phenomenology

There are two energy ranges that must be considered in our analysis:

(i) *Extrinsic energy range.* Forgetting GINI effects for the moment, one energy range—which we label extrinsic—is delimited by the minimum and maximum energies, E_{min} and E_{max} , at which the generic source under consideration can be seen in neutrinos by a particular detector technology. These limiting energies are determined by either detector or intrinsic source spectrum considerations (whichever is the more severe). The extrinsic energy range is defined by

$$E_{\text{min}} \leq E_\nu \leq E_{\text{max}}. \quad (103)$$

(ii) *Intrinsic energy range* that is given by the following considerations: the lower bound on this range is given by the critical energy, E_{crit} , at which the pertinent phase difference is equal to one (below this value our treatment of the phase breaks down). The scale of this energy is given by requiring

$$E_{\text{crit}} \simeq \frac{\hbar c}{R_S}. \quad (104)$$

Note that the relation is not exact because the right-hand side (RHS) does not account for the effect of the source alignment parameter, Y , on the phase difference. The upper bound on this range, E_{washout} , is determined by the energy at which detector energy resolution issues mean that one interference fringe can no longer be resolved from another. Washout occurs generically because, although interference fringes are distributed at equal energy intervals, the absolute uncertainty in neutrino energy determined by a detector can be expected to be an increasing function of energy.

VII. SUITABLE SOURCE-LENS-DETECTOR CONFIGURATIONS FOR GINI

We can think of four scenarios for source-lens-detector configurations that *might* exhibit GINI effects (there may well be more):

(i) Sun-moon-solar neutrino detector (i.e., in a solar eclipse).

(ii) Cosmological neutrino source—intervening lensing object—large scale water/ice Čerenkov neutrino detector or airshower array.

(iii) Artificial neutrino beam on one side of Earth aimed through the center of Earth to a detector on the opposite side of the planet.

(iv) Galactic (i.e., Milky Way) core collapse supernova (types II, Ib and Ic)—intervening lensing object—solar neutrino detector.

Unfortunately, scenarios (i)–(iii) fail one or more of the

heuristic tests we have set out above and we must, reluctantly, dismiss them. Scenario (iv), however, holds out some promise and it is to this that we now briefly turn (see [55] for more detail on all the scenarios mentioned), though we alert the reader from the beginning that scenario is unlikely to be realized *at present* because of the low probability of supernovae at (neutrino-)detectable distances being lensed by objects in a suitable mass range.

A. Core collapse supernovae as sources for GINI observation

1. Core collapse supernovae as neutrino sources: general considerations

Let us take the characteristic scale of the distance to a galactic core collapse supernova to be $10 \text{ kpc} \approx 3 \times 10^{22} \text{ cm}$, the approximate distance to the galactic center. A core collapse SN observed today at the fiducial 10 kpc would produce around 10^4 and 10^3 events in SuperKamiokande and the Sudbury Neutrino Observatory, respectively [65]. Over the medium term, prospects for SN neutrino detection may become even better than at present with the construction of the next generation of 1 Mt underground, water Čerenkov detectors [66–68]. For a supernova at 10 kpc , a 1 Mt device should detect $\mathcal{O}(10^5)$ events [68].

2. Natural scale for lensing mass required for GINI effects with SN neutrinos

Writing

$$\begin{aligned} \Delta\Phi^{+-} &\sim E_\nu R_S \\ &\approx 1.5 \times 10^{17} \left(\frac{E_\nu}{10 \text{ MeV}} \right) \left(\frac{M}{M_\odot} \right), \end{aligned} \quad (105)$$

we can determine that the *smallest* lensing mass that might produce a phenomenological effect (that we can treat using our formalism) is, very roughly, $10^{-17} M_\odot \approx 3 \times 10^{16} \text{ g}$. This is in the cometary mass range. A more detailed calculation [55] demonstrates—for the specific case of SuperKamiokande—a sensitivity to the GINI effect with lensing masses in the range

$$10^{-18} M_\odot \lesssim M_{\text{lens}} \lesssim 10^{-16} M_\odot. \quad (106)$$

This range is both conservative and fairly sensitive to the SuperK energy thresholds and energy resolution.

3. Coherence of supernova neutrinos

A neutrino wave packet leaving the neutrinosphere of a nascent neutron star will have a size [61,69]

$$\sigma_x^{SN} \approx 10^{-9} \text{ cm}. \quad (107)$$

This is to be contrasted with the scale of the affine path difference for the lensing mass range under consideration [Eq. (106)] which is supplied by the range of the Schwarzschild radius, viz,

$$3 \times 10^{-13} \text{ cm} \gtrsim R_S \gtrsim 3 \times 10^{-11} \text{ cm}. \quad (108)$$

We do not, therefore, expect any significant damping of the interference amplitude by decoherence due to path difference effects.

There is, however, also decoherence due to group velocity difference to be considered, i.e., the direct analog of decoherence effects for conventional neutrino oscillations. The inequality to be satisfied is given by Eq. (98), the RHS of which translates to $\sim 3 \times 10^{13} \text{ cm}$ for 10 MeV neutrinos [70] much smaller than the fiducial scale of D_s , $\sim 3 \times 10^{22} \text{ cm}$. We can expect, therefore, to be beyond the flat space coherence length. This means that the neutrino signal will be characterized by flavor ratios that are constant across (measurable) energy. For supernova neutrinos, then, if a suitable lens were present, GINI would cause patterns of maxima and minima across energy in the detected neutrino spectra. Furthermore, the positions, in energy, of these maxima and minima would be essentially the same for all neutrino flavors. Interference effects would be, in principle, directly evident even in neutral current interaction data. On the other hand, we would not expect a noticeable change in the ratios between different neutrino species across energy. In other words, for supernova neutrinos, given a suitable lens, it is possible to see interference effects due to path difference effects but not due to phase velocity difference (i.e., flat space oscillation) effects.

4. Finite source effects with supernova neutrinos

Given a scale for the neutrinosphere, r_{SN} , of $\sim 10 \text{ km} = 10^6 \text{ cm}$, a calculation shows that the point source condition, Eq. (95), fails at the lower end of the phenomenologically interesting lensing mass range *assuming* $D_{ds} \approx D_d$. Furthermore, from Eq. (96) we find that in order that the phase uncertainty introduced by the finite size of the supernova neutrino source not be too large, we require that the neutrino energy be less than $\sim 1 \text{ MeV}$, a condition that, with 10 MeV neutrinos, we fail to meet by an order of magnitude, *again assuming* $D_{ds} \approx D_d$. We hasten to add, however, that we do not believe that either of these two is necessarily fatal. A numerical study is needed here, and this may well establish that GINI effects are visible even when the crude, heuristic inequalities above are violated [71], and moreover that $D_{ds} \approx D_d$ need not hold (over the galactic scales we are considering). Further, we might have $D_s > 10 \text{ kpc}$ (at the cost of a reduced event rate). In either case, point source conditions could easily be satisfied.

5. Finite lens size effects

For a lens in the mass range determined above, and both source and lens at galactic length scales, the Einstein length scale is

$$D_d \theta_E \approx 3 \sqrt{\left(\frac{M}{10^{-17} M_\odot} \right) \left(\frac{D_d}{D_s} \right) \left(\frac{D_{ds}}{5 \text{ kpc}} \right)} \text{ km}. \quad (109)$$

On the other hand, for a lens with a density of $\sim 1 \text{ gm cm}^{-3}$, the scale of the dimensions of the lensing object, l , is given by

$$l \approx 0.6 \left(\frac{M}{10^{-17} M_{\odot}} \right)^{1/3} \text{ km}, \quad (110)$$

meaning that the classical paths pass very close to the object, and, in some cases, one path might actually pass inside the object. Given the order of magnitude nature of the calculations we have performed here, however, this fact will not significantly impinge on the observability of the effect we predict. Certainly the neutrinos will not interact significantly with the material of the lens. Of course, if the lensing object is a black hole, taking the lens to be a point source is unproblematic.

6. Finite detector size effects

A quick calculation shows [55] that finite detector size/position resolution effects never become insurmountable over the whole range of possible lens positions.

7. Crossing time scales for supernova neutrinos

If the lens has a transverse velocity $v \approx 30 \text{ km s}^{-1}$, it will move across the Einstein ring in a time scale of $\sim 1 \text{ s}$ [7]. Given, then, that we expect a detectable neutrino signal will be received from a SN over a period of around 10 s, interference fringes will shift over the time of observation, but not so quickly that they cannot be observed.

8. Lensing probability

The above paragraphs detail the conditions that a source-deflector-observer alignment must satisfy in order for GINI to be measured, but implicit at all stages is that such an alignment has occurred. Unfortunately the chance of a suitable deflector lying sufficiently close to the line-of-sight to a source in the Local Group is not high.

The Galaxy's rotation curve places a strict upper bound on the total mass in its halo (e.g., [72]), which then implies a maximum possible alignment probability to, say, the Magellanic Clouds. Even if the halo consisted only of point-masses of suitable size, the simple calculation made by Paczynski [73] implies that the lensing optical depth, τ —essentially equal to the probability that any single source is lensed at a given time—is $\sim 10^{-6}$. This result has been corroborated experimentally by monitoring stars in both the galactic center and the Large Magellanic Cloud for period variations: both the MACHO [74] and OGLE [75] groups have found $\tau \approx 3 \times 10^{-6}$. It is important, then, to note that even if the halo is dominated by point-masses of $M \approx 10^{-17} M_{\odot}$ suitable for GINI, the alignment probability to any neutrino sources sufficiently close to be detected at all is only $\sim 10^{-6}$.

In the future, however, as detector technology improves, it may be possible to observe neutrinos from cosmologically distant sources at effective distances of up to Gpc. The lensing optical depth is thus increased, both because a given mass can act as a more efficient lens and because the chance of alignment increases proportionally with source distance. A simple calculation of these effects (e.g., [56]) implies that optical depths of close to unity are plausible; thus when neutrinos are detected from cosmologically distant sources,

GINI effects will *have* to be taken into account in the interpretation of any such data obtained.

VIII. EXTENSION OF THEORETICAL RESULTS

Besides treating the potential phenomenological effects of GINI at greater length in another work [55], there are, of course, a number of directions in which our theoretical treatment will be extended. Some issues we intend to deal with further in another publication [55] include the following.

(i) From consideration of interference of lensed *photons* in a Schwarzschild metric [see Eq. (7.8) of [56] and also see Eq. (9) of [4]], Eq. (88) is actually subtly in error: there is an extra $-\pi/2$ phase shift missing from the argument of the cos term (in other words, the interference envelope should actually go as $\beta^2 + 2\theta_E^2[1 + \sin(\Delta\Phi^{+-})]$, generating a central minimum for $Y=0$). This is present in the case of light—and will also be present in the case of neutrinos—because of the opposite parities of the two images produced by a Schwarzschild lens (i.e., the images—*were* they able to be distinguished—would be flipped with respect to each other [76]). The reason why our treatment has failed to pick this extra phase shift up is that we have artificially restricted the paths under consideration to only the classical ones. In other words, we have assumed the geometrical optics limit which is strictly only valid for phase differences of order unity and larger. A more complete treatment using the techniques of physical optics—involving integration over *all* paths through the lens plane (each such path being uniquely specified by its impact parameter)—would recover this phase [and, in fact, demonstrate that the full expressions for the oscillation probability and magnification involve confluent hypergeometric functions that only reduce to trigonometric functions in the large phase limit: cf. Eq. (7.11) of Ref. [56] or Eq. (6) of [4]]. Moreover, a more complete treatment would also demonstrate that the singularity at $\beta=0$ for Eq. (90) is not a real effect.

(ii) So far we have assumed a static metric. But the GINI effect, as noted, requires neutrinos from astrophysical sources that would probably need to be located at extragalactic or even cosmological distances for a decent chance that lensing occur (though it should be stressed that current detector technologies would not allow detection of neutrinos from supernovae beyond our Galaxy and its satellites [65]) and the introduction of cosmology into the formalism developed here would require the treatment of a nonstatic metric [77,78]. We note in passing that were GINI effects ever seen in neutrinos from cosmological sources, these effects would provide for a test of quantum mechanics over the very longest scales. We speculate, then, that GINI could be sensitive to the effects of spacetime foam (cf., say [79]). In principle, we also expect that a GINI pattern in cosmologically sourced neutrinos could be interrogated to determine the value of the Hubble constant H_0 [80,81]. Observation (or nonobservation) of GINI effects would also constitute a *de facto* probe of the distribution of dark matter objects within a certain well-defined (and interesting) mass range [7].

(iii) The Schwarzschild lens is an ideal case never precisely encountered in nature. For the realistic situation we

may need to account for shear [8], multiple lensing masses, etc. Treating these effects may very well demonstrate that the just-so mass range is considerably larger than the estimate given by Eq. (106) [8]. In any case, it is fairly easy to understand, at least at the heuristic level, how more general lenses might be treated: considering, say, Eq. (76) it can be seen that almost all the contributions to the phase difference are essentially geometrical. The only expression which contains information on the mass distribution of the deflector is the logarithmic term. This suggests that it may be possible to modify our results to arbitrary weak deflectors simply by inserting the appropriate lens potential (and replacing a Schwarzschild-specific result for $\Delta\theta_{pq}^2$).

(iv) In Sec. VI we added into the mix coherence loss considerations. Formally, these can only arise in a full wave packet treatment which we have not attempted for the curved spacetime case. We remind the reader, however, that our results for phase and phase-difference will continue to hold in any more detailed calculation because these are independent of wave packet considerations.

(v) Our treatment of effects due to the finite nature of any real source is very much at a heuristic level. Furthermore, finite detector effects can also be important, as has been remarked. In this regard, note, in passing, that logically connected to this concern is the consideration that the observation of GINI effects with two—or, preferably, more—widely separated detectors holds out some interesting possibilities [7]. One would expect here that the interference patterns seen by different detectors would be, in general, displaced in energy with respect to each other. The degree of this displacement will be related to the lensing mass and the geometry. So the displacement could probably be used to better constrain relevant parameters than observation with a single detector. Moreover, observation of fringes with more than one neutrino detector would certainly lend credence to the idea that these have their origins in GINI.

IX. CONCLUSION

In this paper we have explicitly calculated the phase for a neutrino mass eigenstate propagating through curved spacetime, in particular a Schwarzschild metric. With this expression in hand, we have shown how a novel interference effect—gravitationally induced neutrino interference (GINI)—may show up for gravitationally lensed, astrophysical neutrinos. These interference effects lead to a neutrino transition phenomenology qualitatively different from flat space neutrino oscillations. We have shown, further, that a result extant in the literature [29] for the phase difference with gravitational lensing must be in error. We have also derived the form of this phase difference when it is given in terms of conventional lensing parameters. We have derived the analog of the neutrino oscillation probability in flat space for the Schwarzschild metric. This quantity controls the phenomenology at a detector, in particular the pattern of maxima and minima (across energy) for neutrino wave packets which have propagated from source to detector along multiple paths. We have adduced heuristic arguments that establish that this interference pattern could be seen in the neutrino

signal from a supernova, *provided* a suitably lensed supernova event occurs. Current—and probably even next-generation—neutrino detector technologies would seem to mean, however, that the probability of such lensing occurring for a neutrino-detectable supernova is small. Still, for astrophysical neutrinos originating at cosmological distances, the lensing probability approaches 1 and some day the technology to detect large numbers of these from single sources may become available. We have mapped out a program for further research in this field.

In summary, the material presented in this paper serves as a proof-of-principle that the GINI effect is both real—in a theoretical sense—and, what is more, could lead, one day, to interesting phenomenological consequences for supernova neutrinos.

ACKNOWLEDGMENT

R.M.C. would like to sincerely thank Paul Alsing, Jesse Carlsson, Tim Garoni, Matt Garbutt, Sasha Ignatiev, Bruce McKellar, Andrew Melatos, Keith Nugent, Alicia Oshlack, David Paganin, Andrew Peele, Ray Protheroe, Georg Raffelt, Rob Scholten, Cath Trott, Rachel Webster, and Stuart Wyithe for enlightening discussions. He also thanks Nicole Bell and John Beacom for a useful correspondence. Finally, he particularly thanks German Kälberman, Ray Volkas, and Randall Wayth, who all devoted considerable time to setting this author straight on a number of subtle issues. D.J.M. is supported by PPARC.

APPENDIX: WAVE PACKET TREATMENT OF NEUTRINO BEAM SPLITTER

In this appendix we set forth a full (Gaussian) wave packet treatment of the neutrino beam splitter *gedanken experiment* (treated in terms of plane waves in Sec. III). Note that the results we derive for the exponential damping terms [in the equation for the oscillation probability analog—see Eq. (A14)] serve as a heuristic justification of the treatment of decoherence we present in Sec. VI.

We write the ket associated with the neutrino flavor eigenstate α that has propagated from the source spacetime position $A = (x_A, t_A)$ to detection position $B = (x_B, t_B)$ as

$$|\nu_\alpha, B\rangle = N \sum_p \sqrt{I_p} \sum_j U_{\alpha j} \times \int dE \exp[-i\Phi_j^p(E; L_p^{AB}, T^{AB})] A_j(E) |\nu_j\rangle, \quad (\text{A1})$$

where the various quantities are as explained in Sec. III. The amplitude for a neutrino created as type α to be detected as type β at the spacetime position, B , of the detection event is then

$$\begin{aligned} \langle \nu_\beta | \nu_\alpha ; A, B \rangle &= N \sum_p \sqrt{I_p} \sum_j U_{\alpha j} U_{\beta j}^* \\ &\times \int dE A_j(E) \exp[-i\Phi_j^p(E; L_p^{AB}, T^{AB})]. \end{aligned} \quad (\text{A2})$$

Again, we can get rid of the unwanted dependence on time by averaging over T^{AB} in the above to determine a time-averaged oscillation probability *analog* at the detector position x_B [45–47]. This gives us that

$$\begin{aligned} |\langle \nu_\beta | \nu_\alpha ; x_A, x_B \rangle|^2 &\propto \int dT^{AB} |\langle \nu_\beta | \nu_\alpha ; A, B \rangle|^2 \\ &\propto |N|^2 \sum_{pq} \sqrt{I_p I_q} \sum_{jk} U_{\alpha j} U_{\beta j}^* U_{\beta k} U_{\alpha k}^* \\ &\times \int dE A_j(E) A_k^*(E) \\ &\times \exp\{i[p_j(E) L_p^{AB} - p_k(E) L_q^{AB}]\}, \end{aligned} \quad (\text{A3})$$

where one integral over energy has disappeared because of the $\delta(E - E')$ that arises from the integration over time.

Assuming a Gaussian form for the wave packets leads to

$$A_j \propto \exp\left[-\frac{(E - \bar{E}_j)^2}{4(\sigma_E^{(j)})^2}\right] \simeq \exp\left[-\frac{(E - \bar{E}_j)^2}{4\sigma_E^2}\right], \quad (\text{A4})$$

where \bar{E}_j is the peak or average energy of mass eigenstate j and we employ the very good approximation [82] that the wave packet spread is the same for different mass eigenstates. This gives us that

$$A_j(E) A_k^*(E) \propto \exp\left[-\frac{2E^2 - 2E(\bar{E}_j + \bar{E}_k) + (\bar{E}_j^2 + \bar{E}_k^2)}{4\sigma_E^2}\right]. \quad (\text{A5})$$

Now defining the peak momentum of mass eigenstate j via $\bar{p}_j = (\bar{E}_j^2 - m_j^2)^{1/2}$, we can write

$$p_j(E) \equiv \bar{p}_j + \Delta p_j(E), \quad (\text{A6})$$

where

$$\begin{aligned} \Delta p_j(E) &\equiv p_j(E) - \bar{p}_j \\ &= \sqrt{E^2 - m_j^2} - \sqrt{\bar{E}_j^2 - m_j^2} \\ &\simeq E - \frac{m_j^2}{2E} - \bar{E}_j + \frac{m_j^2}{2\bar{E}_j} \\ &= (E - \bar{E}_j) \left(1 - \frac{m_j^2}{2E\bar{E}_j}\right). \end{aligned} \quad (\text{A7})$$

Employing the group velocity of mass eigenstate j , viz.

$$v_j = \bar{p}_j / \bar{E}_j = \frac{\sqrt{\bar{E}_j^2 - m_j^2}}{\bar{E}_j} \simeq 1 - \frac{m_j^2}{2\bar{E}_j^2}, \quad (\text{A8})$$

we determine that $\Delta p_j(E) \simeq (E - \bar{E}_j)v_j$ [83], so that

$$p_j(E) \simeq \bar{p}_j + (E - \bar{E}_j)v_j. \quad (\text{A9})$$

We find, then, after a simple calculation that the oscillation probability analog becomes

$$\begin{aligned} |\langle \nu_\beta | \nu_\alpha ; x_A, x_B \rangle|^2 &\simeq \frac{1}{\sum_{rs} \sqrt{I_r I_s}} \sum_{pq} \sqrt{I_p I_q} \sum_{jk} U_{\alpha j} U_{\beta j}^* U_{\beta k} U_{\alpha k}^* \exp\left\{-i\left[(v_j L_p^{AB} + v_k L_q^{AB}) \frac{\bar{E}_j - \bar{E}_k}{2} - (p_j L_p^{AB} - p_k L_q^{AB})\right]\right\} \\ &\times \exp\left[-\sigma_x^2 \frac{(\bar{E}_j - \bar{E}_k)^2}{2}\right] \exp\left[-\frac{(v_j L_p^{AB} - v_k L_q^{AB})^2}{8\sigma_x^2}\right]. \end{aligned} \quad (\text{A10})$$

Note that in the above calculation, though we have been employing the group velocity, this does not—and should not be seen to—enter into the phase in any fundamental way [44]. Indeed, the phase can be calculated entirely with plane waves (see Sec. III) and, therefore, totally without reference to wave packet notions like group velocity—though the exponential damping factors above critically depend on these. Note also that, again, the normalization has been determined by requiring that $|\langle \nu_\beta | \nu_\alpha ; x_A, x_B \rangle|^2 \leq 1$.

The two damping factors in Eq. (4) can be traced back to

considerations following from (i) source localization and (ii) the requirement for overlap of wave packets at the detector's position. Observe that the second damping term accounts for an interesting possibility: having a heavier—and slower—mass eigenstate travel down the shorter path and the lighter mass eigenstate down the longer path will tend to restore coherence.

As a further particularization of the expression for neutrino oscillation “probability” with an imaginary beamsplitter, we take the expressions for the energies, momenta, and

velocities of the various wave functions given in terms of expansions around the energy in massless limit, E_0 , viz.

$$\bar{E}_i \approx E_0 + (1 - \xi) \frac{m_i^2}{2E_0}, \quad (\text{A11})$$

$$\bar{p}_i \approx E_0 - \xi \frac{m_i^2}{2E_0}, \quad (\text{A12})$$

where ξ is a dimensionless parameter of order unity determined by kinematical considerations [44]. The group velocity will then be

$$v_i = \frac{\bar{p}_i}{\bar{E}_i} \approx 1 - \frac{m_i^2}{2E_0^2}. \quad (\text{A13})$$

Employing the above, we find

$$\begin{aligned} |\langle \nu_\beta | \nu_\alpha ; x_A, x_B \rangle|^2 \approx & \frac{1}{\sum_{rs} \sqrt{I_r I_s}} \sum_{pq} \sqrt{I_p I_q} \sum_{jk} U_{\alpha j} U_{\beta j}^* U_{\beta k} U_{\alpha k}^* \exp[-i\Delta\Phi_{jk}^{pq}] \\ & \times \exp\left[-\frac{\sigma_x^2}{2} \left(\xi \frac{\delta m_{jk}^2}{2E_0}\right)^2\right] \exp\left[-\frac{1}{2} \left(\frac{L_p^{AB} - L_q^{AB}}{2\sigma_x} - \frac{1}{2E_0^2} \frac{m_j^2 L_p^{AB} - m_k^2 L_q^{AB}}{2\sigma_x}\right)^2\right], \end{aligned} \quad (\text{A14})$$

where the phase difference is given by

$$\Delta\Phi_{jk}^{pq} \equiv -\left(E_0 + \xi \frac{m_j^2 + m_k^2}{4E_0}\right) (L_p^{AB} - L_q^{AB}) + \left(\frac{m_j^2 L_p^{AB} - m_k^2 L_q^{AB}}{2E_0}\right). \quad (\text{A15})$$

Note that we can take the plane wave limit of the above equation by setting $\xi \rightarrow 0$ and $\sigma_x \rightarrow \infty$. This allows us to recover Eq. (4).

-
- [1] A.W. Overhauser and R. Colella, Phys. Rev. Lett. **33**, 1237 (1974).
- [2] R. Colella, A.W. Overhauser, and S.A. Werner, Phys. Rev. Lett. **34**, 1472 (1975).
- [3] D.M. Greenberger and A.W. Overhauser, Rev. Mod. Phys. **51**, 43 (1979).
- [4] K.Z. Stanek, B. Paczynski, and J. Goodman, Astrophys. J. Lett. **413**, L7 (1993).
- [5] P. Schneider and J. Schmid-Burgk, Astron. Astrophys. **148**, 369 (1985).
- [6] J.B. Peterson and T. Falk, Astrophys. J. Lett. **374**, L5 (1991).
- [7] A. Gould, Astrophys. J. Lett. **386**, L5 (1992).
- [8] A. Ulmer and J. Goodman, Astrophys. J. **442**, 67 (1995).
- [9] A.V. Mandzhos, Sov. Astron. Lett. **7**, 213 (1981).
- [10] H.C. Ohanian, Astrophys. J. **271**, 551 (1983).
- [11] S. Deguchi and W.D. Watson, Astrophys. J. **307**, 30 (1986).
- [12] S. Deguchi and W.D. Watson, Phys. Rev. D **34**, 1708 (1986).
- [13] Note here that while there is some similarity between GINI and femtolensing effects, there are at least two effects that suggest that neutrino interference might be, at a pragmatic level, intrinsically more observable than femtolensing. (i) The typical length scale for the impact parameter in gravitational lensing is given by the Einstein radius of the lens. It may happen that the source-lens-observer geometry means that the Einstein radius of a lens is actually “inside” the lens body. There will be many situations, then, where the lensing object is optically thick to photons at the Einstein radius but transparent to neutrinos, meaning that interference effects are, in principle, observable in the former situation but not the latter. (ii) Interference effects can only show up when different lensed images are unresolved (i.e., one’s apparatus must *not* be able to determine which photon—or neutrino—belongs to which image). But, because of the very different, *intrinsic* angular resolutions of the microscopic processes involved in neutrino and photon detection, a clearly resolved, astrophysical light source may well be an *unresolved* source as far as neutrinos are concerned.
- [14] K. Hirata, T. Kajita, M. Koshiba, M. Nakahata, and Y. Oyama, Phys. Rev. Lett. **58**, 1490 (1987).
- [15] R.M. Bionta, G. Blewitt, C.B. Bratton, D. Caspere, and A. Ciocio, Phys. Rev. Lett. **58**, 1494 (1987).
- [16] J.N. Bahcall and S.L. Glashow, Nature (London) **326**, 476 (1987).
- [17] W.D. Arnett and J.L. Rosner, Phys. Rev. Lett. **58**, 1906 (1987).
- [18] S.M. Bilenky, C. Giunti, J.A. Grifols, and E. Massó, Phys. Rep. **379**, 69 (2003).
- [19] W. Hillebrandt and P. Hoflich, Rep. Prog. Phys. **52**, 1421 (1989).
- [20] J.F. Beacom, R.N. Boyd, and A. Mezzacappa, Phys. Rev. Lett. **85**, 3568 (2000).
- [21] M.J. Longo, Phys. Rev. Lett. **60**, 173 (1988).
- [22] L.M. Krauss and S. Tremaine, Phys. Rev. Lett. **60**, 176 (1988).
- [23] Data from SN 1987A neutrinos have been used to constrain other neutrino properties including neutrino mixing and mass hierarchy, neutrino lifetime, and neutrino magnetic moment: see Ref. [18] for a review.
- [24] J.D. Barrow and K. Subramanian, Nature (London) **327**, 375 (1987).

- [25] J.L. Gerver, Phys. Lett. A **127**, 301 (1988).
- [26] R. Escrivano, J.-M. Frère, D. Monderen, and V. Van Elewyck, Phys. Lett. B **512**, 8 (2001).
- [27] D.R. Brill and J.A. Wheeler, Rev. Mod. Phys. **29**, 465 (1957).
- [28] L. Stodolsky, Gen. Relativ. Gravit. **11**, 391 (1979).
- [29] N. Fornengo, C. Giunti, C.W. Kim, and J. Song, Phys. Rev. D **56**, 1895 (1997).
- [30] T. Bhattacharya, S. Habib, and E. Mottola, Phys. Rev. D **59**, 067301 (1999).
- [31] Y. Kojima, Mod. Phys. Lett. A **11**, 2965 (1996).
- [32] C.Y. Cardall and G.M. Fuller, Phys. Rev. D **55**, 7960 (1997).
- [33] K. Konno and M. Kasai, Prog. Theor. Phys. **100**, 1145 (1998).
- [34] P.M. Alsing, J.C. Evans, and K.K. Nandi, Gen. Relativ. Gravit. **33**, 1459 (2001).
- [35] D.V. Ahluwalia and C. Burgard, Gen. Relativ. Gravit. **28**, 1161 (1996).
- [36] J. Wudka, Phys. Rev. D **64**, 065009 (2001).
- [37] D.V. Ahluwalia and C. Burgard, Phys. Rev. D **57**, 4724 (1998).
- [38] Y. Grossman and H.J. Lipkin, Phys. Rev. D **55**, 2760 (1997).
- [39] C.M. Zhang and A. Beesham, Gen. Relativ. Gravit. **33**, 1011 (2001).
- [40] B. Linet and P. Teyssandier, “General relativity and quantum cosmology,” gr-qc/0206056.
- [41] C.M. Zhang and A. Beesham, Int. J. Mod. Phys. D **12**, 727 (2003).
- [42] As an example of this, take the contention on p. 1483 of Alsing *et al.* in Ref. [34] that the phase of photons propagating through a Schwarzschild metric vanishes to lowest order. This applies only to radially propagating photons and even then should really be thought of as a statement about the *action* along the classical, null-geodesic, rather than as a claim that the phase—and therefore a potentially measurable phase difference—actually is (close to) zero. Likewise, note that while the statement made in Ref. [38] that “. . . if one compares two experimental setups with and without gravity with the same curved distance in both cases there is no effect” is true, it implicitly assumes that there is only one path from source to detector that need be considered—and this does not hold in general in the presence of gravity.
- [43] Though it should be stressed here that the authors of Ref. [29] and many of the other papers we have mentioned *do* obtain the correct result for the phase *difference* between neutrino mass eigenstates traveling along the same macroscopic paths in curved spacetime (i.e., for flat space neutrino oscillations in the presence of a point mass) because, in such cases, any putative $\propto E$ term will vanish in subtracting one phase from the other (this term being the same for both phases).
- [44] C. Giunti, Phys. Scr. **67**, 29 (2003).
- [45] M. Beuthe, Phys. Rep. **375**, 105 (2003).
- [46] M. Beuthe, Phys. Rev. D **66**, 013003 (2002).
- [47] C. Giunti, “High energy physics—Phenomenology,” hep-ph/0302026.
- [48] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (W.H. Freeman, San Francisco, 1973).
- [49] C. Giunti, Mod. Phys. Lett. A **16**, 2363 (2001).
- [50] C. Giunti, “High energy physics—Phenomenology,” hep-ph/0205014.
- [51] T. Bhattacharya, S. Habib, and E. Mottola, gr-qc/9605074.
- [52] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields*, Course of Theoretical Physics, Pergamon International Library of Science, Technology, Engineering and Social Studies, 4th rev. ed. (Pergamon Press, Oxford, 1975).
- [53] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity* (Wiley, New York, 1972).
- [54] Note that the $M \rightarrow 0$ case of this equation can be rederived by, again, considering Fig. 1 while noting, in particular, that the Schwarzschild lens satisfies Eq. (13) for the two classical paths.
- [55] R. M. Crocker and D. Mortlock (unpublished).
- [56] P. Schneider, J. Ehlers, and E. E. Falco, *Gravitational Lenses* (Springer-Verlag, Berlin, 1992), Vol. XIV. (Also Astronomy and Astrophysics Library, 1992.)
- [57] Note that we will assume separability of the phase difference is satisfied in the following discussion.
- [58] This is analogous to the requirement that, for the “usual” equation (obtained in the geometrical optics limit) describing the intensity on a screen in Young’s slit type experiment to be correct, the wavelength of the interfering radiation must not be larger than the slit separation. Otherwise, the equation suggests a (nonphysical) violation of conservation of total intensity in the form of an interference maximum over the whole screen.
- [59] See the Appendix of Ref. [32] for a treatment of the direct analog of the coherence length of neutrino oscillations in flat space for curved spacetime, though note that the treatment presented here does not apply for multiple, macroscopic paths.
- [60] B. Kayser, Phys. Rev. D **24**, 110 (1981).
- [61] C. W. Kim and A. Pevsner, *Neutrinos in Physics and Astrophysics*, Contemporary Concepts in Physics, Vol. 8 (Harwood, Chur, Switzerland, 1993).
- [62] Note that we ignore here the contribution of the *detection* process to the effective wave packet size that can, in principle, act to restore coherence via broadening the effective wave packet width through accurate energy/momentum measurement: see Ref. [84] for more details and also Ref. [46] for a rigorous, quantum-field-theoretic treatment of neutrino coherence length.
- [63] S. Nussinov, Phys. Lett. **63B**, 201 (1976).
- [64] H. Anada and H. Nishimura, Phys. Rev. D **37**, 552 (1988).
- [65] J.F. Beacom and P. Vogel, Phys. Rev. D **60**, 033007 (1999).
- [66] Super-Kamiokande Collaboration, T. Nakaya *et al.*, presented at the XXth Int. Conf. on Neutrino Physics and Astrophysics, Munich (2002).
- [67] P. Antonioli *et al.*, Nucl. Instrum. Methods Phys. Res. A **433**, 104 (1999).
- [68] C.K. Jung, “High energy physics—Experimental,” hep-ex/0005046.
- [69] H. Anada and H. Nishimura, Phys. Rev. D **41**, 2379 (1990).
- [70] We take the *largest* possible value for this quantity by assuming the scale of the smallest experimentally determined δm^2 , i.e., the solar mass splitting at $O(10^{-5})$ eV².
- [71] Certainly, in their numerical study of femtolensing with a disk source, Peterson and Falk [6] found that, allowing for a realistic deviation from smoothness in the source function—which describes the intensity across the disk of the source—interference effects were visible with a source size significantly larger than the Einstein ring. We have not allowed for this (potential) effect for a supernova neutrino source.

- [72] T. Sakamoto, M. Chiba, and T.C. Beers, *Astron. Astrophys.* **397**, 899 (2003).
- [73] B. Paczynski, *Astrophys. J.* **304**, 1 (1986).
- [74] C. Alcock *et al.*, *Astrophys. J.* **541**, 734 (2000).
- [75] P.R. Wozniak, A. Udalski, M. Szymanski, M. Kubiak, G. Pietrzynski, I. Soszynski, and K. Zebrun, *Acta Astron.* **51**, 175 (2001).
- [76] An experimental analog of interference in such a situation would be Young's double slit apparatus where one beam of light is reflected in a mirror. This produces, of course, a central minimum.
- [77] D.J. Wagner and T.J. Weiler, *Mod. Phys. Lett. A* **12**, 2497 (1997).
- [78] M.R. Mbone, *Gen. Relativ. Gravit.* **34**, 1865 (2002).
- [79] R. Ragazzoni, M. Turatto, and W. Gaessler, *Astrophys. J. Lett.* **587**, L1 (2003).
- [80] A.S. Bolton and S. Burles, *Astrophys. J.* **592**, 17 (2003).
- [81] S. Refsdal, *Mon. Not. R. Astron. Soc.* **128**, 307 (1964).
- [82] C. Giunti, C.W. Kim, and U.W. Lee, *Phys. Rev. D* **44**, 3635 (1991).
- [83] Note that here we are ignoring the energy dependence of the velocities of the different plane wave components of each wave packet so that our calculation makes no account of dispersive effects which, in general, extend the coherence length; see [61].
- [84] C. Giunti and C.W. Kim, *Phys. Rev. D* **58**, 017301 (1998).