

## REFORMULATION OF GENERAL CHANCE CONSTRAINED PROBLEMS USING THE PENALTY FUNCTIONS

MARTIN BRANDA

**ABSTRACT.** We explore reformulation of nonlinear stochastic programs with several joint chance constraints by stochastic programs with suitably chosen penalty-type objectives. We show that the two problems are asymptotically equivalent. Simpler cases with one chance constraint and particular penalty functions were studied in [5, 9]. The obtained problems with penalties and with a fixed set of feasible solutions are simpler to solve and analyze than the chance constrained programs. We discuss solving both problems using Monte-Carlo simulation techniques for the cases when the set of feasible solution is finite or infinite bounded. The approach is applied to the financial optimization problem with Value at Risk constraint, transaction costs and integer allocations. We compare the ability to generate a feasible solution of the original chance constrained problem using the sample approximations of the chance constraints directly or via sample approximation of the penalty function objective.

### 1. INTRODUCTION

Stochastic programming treats problems where optimization and uncertainty appears together. Such problems arise in economy, finance, industry, agriculture and logistics, cf. [21].

In general, we consider the following program with a random factor

$$(1.1) \quad \min \{f(x) : x \in X, g_i(x, \omega) \leq 0, i = 1, \dots, k\},$$

where  $g_i, i = 0, \dots, k$ , are real functions on  $\mathbb{R}^n \times \mathbb{R}^{n'}$ ,  $X \subseteq \mathbb{R}^n$  and  $\omega \in \mathbb{R}^{n'}$  is a realization of a  $n'$ -dimensional random vector defined on the probability space  $(\Omega, \mathcal{F}, P)$ . However,  $\omega$  is unknown for us, hence a question is how to deal with the uncertain constraints. In [14], three suggestions how to deal with the stochastic constraints of the form  $g_i(x, \omega) = \omega_i - h_i(x) \leq 0$ ,  $i = 1, \dots, k$ , where  $\omega_i$  are random bounds with marginal distributions  $P_i$ , are introduced. First, the constraints can be incorporated into the objective function of the optimization problems as the penalty function

$$\sum_{i=1}^k N_i \int_{h_i(x)}^{\infty} [\omega_i - h_i(x)] P_i(d\omega).$$

with  $N_i > 0$  being constant. Next, the reliability type model with a chance or probabilistic constraint can be considered

$$P(h_i(x) \geq \omega_i, i = 1, \dots, k) \geq 1 - \varepsilon$$

for some level  $\varepsilon \in (0, 1)$ . Finally, the constraints involving the conditional expectations can be used

$$\mathbb{E}[\omega_i - h_i(x) | \omega_i - h_i(x) > 0] \leq l_i, i = 1, \dots, k$$

for some small levels  $l_i > 0$ .

Solving the chance constrained problems is not easy. In general, the feasible region is not convex even if the functions are convex and in many cases it is even not easy to check feasibility because it leads to computations of multivariate integrals. On the other hand, there are some special cases under which the convexity is preserved, e.g. the log-concave distributions [16], or it is relatively easy to check the feasibility of a point, e.g. for the normal distribution. There are several methods for numerical solving of particular chance constrained problems, you can see [17]. For the problems with discretely distributed random variables, p-efficient points can be used, cf. [15]. For continuously distributed random variables the methods based on supporting hyperplanes and reduced gradients are available. In the case that the underlying distribution is continuous or discrete with many realizations, the sample approximation techniques and mixed-integer programming reformulation can help us to solve the problem approximately, see [1, 12].

In this paper, we will study the relation between the nonlinear problems with several chance constraints and the penalty function problems. We will show that the model with chance constraints and the penalty type model are asymptotically equivalent under quite mild assumptions. In [9], the equivalence between the problem with one joint chance constraint and the problem with simple penalty function was shown. The approach was recently extended to a whole class of penalty functions in [5]. We propose further extension to multiple jointly chance constrained problems which cover the joint as well as the separate chance constrained problems as special cases.

The approach for solving nonlinear deterministic programs with several constraints using the penalty functions is well studied in literature. Algorithms and basic theory based on continuity and Karush-Kuhn-Tucker conditions are explained in [3, 11]. Theoretical analysis of the penalty function method is provided by [19]. The penalized objective function epi-converges to the objective function of the nonlinear problem with several constraints, which implies "stable" behaviour of optimal values and optimal solutions.

We will show that the penalty function approach can be helpful in numerical solution of stochastic optimization problems with chance constraints. The reformulation of chance constrained problems using the penalties was applied in insurance and water-management, cf. [8, 9]. We will draw our attention to the nonconvex case with a finite set of feasible solutions, which can appear in bounded integer programming, and with an infinite bounded set. We will extend the result on the rates of convergence for the sample approximations of the chance constrained problems and summarize the results for the problems with the expectation in the objective which cover the penalty function problems. The approach will be applied to the financial optimization problem with Value at Risk constraint, transaction costs and integer allocations. We compare the ability to generate a feasible solution of the original chance constrained problem using the sample approximations of the chance constraints directly or via sample approximation of the penalty function objective.

The paper is organized as follows. In section 2, we formulate the multiple jointly chance constrained problem and the problem with penalty type objective and we show that they are asymptotically equivalent. In section 3, Monte Carlo techniques

for solving the problems are discussed. Numerical study is included in section 4. In section 5, we will summarize our results.

## 2. REFORMULATION

Let  $g_{ji}(x, \omega), i = 0, \dots, k_j, j = 1, \dots, m$ , be real functions on  $\mathbb{R}^n \times \mathbb{R}^{n'}$  measurable in  $\omega$  for all  $x \in X$ . Then the multiple chance constrained problem can be formulated as follows:

$$(2.1) \quad \begin{aligned} \psi_\epsilon &= \min_{x \in X} f(x), \\ \text{s.t.} & \\ P(g_{11}(x, \omega) \leq 0, \dots, g_{1k_1}(x, \omega) \leq 0) &\geq 1 - \epsilon_1, \\ &\vdots \\ P(g_{m1}(x, \omega) \leq 0, \dots, g_{mk_m}(x, \omega) \leq 0) &\geq 1 - \epsilon_m, \end{aligned}$$

with an optimal solution  $x_\epsilon$ , where  $\epsilon = (\epsilon_1, \dots, \epsilon_m)$ , with the levels  $\epsilon_j \in (0, 1)$ . The formulation covers the joint ( $k_1 > 1$  and  $m = 1$ ) as well as the separate ( $k_j = 1$  and  $m > 1$ ) chance constrained problems as special cases.

In [9], asymptotic equivalence between the problem with one joint chance constraint and the problem with simple penalty function is shown. The approach by [9] can be extended to a whole class of penalty functions with desirable properties which was done in [5]. We propose further extension to the multiple jointly chance constrained problems (2.1).

Below, we will consider the penalty functions  $\vartheta_j : \mathbb{R}^{k_j} \rightarrow \mathbb{R}_+$ ,  $j = 1, \dots, m$ , which are continuous nondecreasing in their components, equal to 0 on  $\mathbb{R}_-^{k_j}$  and positive otherwise. Two special penalty functions are readily available:  $\vartheta^{1,o}(u) = \sum_{i=1}^k ([u_i]^+)^o, o > 0$ , where  $\vartheta^{1,1}(u) = \sum_{i=1}^k [u_i]^+$  was applied in [9], and  $\vartheta^2(u) = \max_{1 \leq i \leq k} [u_i]^+$  applied in [8]. Both functions preserve convexity,  $\vartheta^2$  is usually used for the joint chance constraints. Another penalty functions are also available:

$$\vartheta^3(u) = \min \left\{ t \geq 0 : u_i - t \leq 0, i = 1, \dots, k \right\},$$

and the ideal (perfect) penalty function, which is closely connected to the duality in nonlinear programming:

$$\vartheta^4(u) = \sup_{y \geq 0} \sum_{i=1}^k y_i u_i,$$

where  $y \in \mathbb{R}^k$ . For any nonpositive  $u$  it holds  $\vartheta^4(u) = 0$ , and  $\vartheta^4(u) = \infty$  otherwise.

We denote

$$p_j(x, \omega) = \vartheta_j(g_{j1}(x, \omega), \dots, g_{jk_j}(x, \omega)) : \mathbb{R}^n \times \mathbb{R}^{n'} \rightarrow \mathbb{R}$$

the penalized constraints. Our choice is appropriate, because it holds

$$(2.2) \quad P(g_{ji}(x, \omega) \leq 0, i = 1, \dots, k_j) \geq 1 - \epsilon_j \iff P(p_j(x, \omega) > 0) \leq \epsilon_j.$$

The corresponding penalty function problem can be formulated as follows:

$$(2.3) \quad \varphi_N = \min_{x \in X} \left[ f(x) + N \cdot \sum_{j=1}^m \mathbb{E}[p_j(x, \omega)] \right]$$

with  $N$  a positive parameter. We denote  $x_N$  an optimal solution of (2.3).

A rigorous proof of the relationship between the optimal values of (2.1) and those of (2.3) for a special additive penalty function and one chance constraint was given

by [9]. The following main theorem states the asymptotic equivalence of the models in generalized settings.

**Theorem 2.1.** *Consider the two problems (2.1) and (2.3) and assume:  $X \neq \emptyset$  is compact,  $f(x)$  is a continuous function,  $\vartheta_j : \mathbb{R}^{k_j} \rightarrow \mathbb{R}_+$ ,  $j = 1, \dots, m$ , are continuous functions, nondecreasing in their components, which are equal to 0 on  $\mathbb{R}_-^{k_j}$  and positive otherwise, denote*

$$p_j(x, \omega) = \vartheta_j(g_{j1}(x, \omega), \dots, g_{jk_j}(x, \omega)), \quad j = 1, \dots, m,$$

and assume

- (i)  $g_{ji}(\cdot, \omega)$ ,  $i = 1, \dots, k_j$ ,  $j = 1, \dots, m$ , are almost surely continuous;
- (ii) there exists a nonnegative random variable  $C(\omega)$  with  $\mathbb{E}[C^{1+\kappa}(\omega)] < \infty$  for some  $\kappa > 0$ , such that  $|p_j(x, \omega)| \leq C(\omega)$ ,  $j = 1, \dots, m$ , for all  $x \in X$ ;
- (iii)  $\mathbb{E}[p_j(x', \omega)] = 0$ ,  $j = 1, \dots, m$ , for some  $x' \in X$ ;
- (iv)  $P(g_{ji}(x, \omega) = 0) = 0$ ,  $i = 1, \dots, k_j$ ,  $j = 1, \dots, m$ , for all  $x \in X$ .

Denote  $\eta = \kappa/(2(1 + \kappa))$ , and for arbitrary  $N > 0$  and  $\epsilon \in (0, 1)^m$  put

$$\begin{aligned} \varepsilon_j(x) &= P(p_j(x, \omega) > 0), \quad j = 1, \dots, m, \\ \alpha_N(x) &= N \cdot \sum_{j=1}^m \mathbb{E}[p_j(x, \omega)], \\ \beta_\epsilon(x) &= \varepsilon_{max}^{-\eta} \sum_{j=1}^m \mathbb{E}[p_j(x, \omega)], \end{aligned}$$

where  $\varepsilon_{max}$  denotes maximum of the vector  $\epsilon = (\varepsilon_1, \dots, \varepsilon_m)$  and  $[1/N^{1/\eta}] = (1/N^{1/\eta}, \dots, 1/N^{1/\eta})$  is the vector of length  $m$ .

THEN for any prescribed  $\epsilon \in (0, 1)^m$  there always exists  $N$  large enough so that minimization (2.3) generates optimal solutions  $x_N$  which also satisfy the chance constraints (2.1) with the given  $\epsilon$ .

Moreover, bounds on the optimal value  $\psi_\epsilon$  of (2.1) based on the optimal value  $\varphi_N$  of (2.3) and vice versa can be constructed:

$$\begin{aligned} \varphi_{1/\varepsilon_{max}^\eta(x_N)} - \beta_{\epsilon(x_N)}(x_{\epsilon(x_N)}) &\leq \psi_{\epsilon(x_N)} \leq \varphi_N - \alpha_N(x_N), \\ \psi_{\epsilon(x_N)} + \alpha_N(x_N) &\leq \varphi_N \leq \psi_{[1/N^{1/\eta}]} + \beta_{[1/N^{1/\eta}]}(x_{[1/N^{1/\eta}]}), \end{aligned} \tag{2.4}$$

with

$$\lim_{N \rightarrow +\infty} \alpha_N(x_N) = \lim_{N \rightarrow +\infty} \varepsilon_j(x_N) = \lim_{\varepsilon_{max} \rightarrow 0_+} \beta_\epsilon(x_\epsilon) = 0$$

for any sequences of optimal solutions  $x_N$  and  $x_\epsilon$ .

PROOF. We denote

$$\delta_N = \sum_{j=1}^m \mathbb{E}[p_j(x_N, \omega)]$$

for some sequence  $x_N$  of optimal solutions of the problem (2.3). Our assumptions and general properties of the penalty function method, see [3, Theorem 9.2.2], ensure that for any sequence  $x_N$  of optimal solutions  $\delta_N \rightarrow 0_+$  and also  $\alpha_N(x_N) =$

$N\delta_N \rightarrow 0$  as  $N \rightarrow \infty$ . Then by Chebyshev inequality

$$\begin{aligned} & P\left(p_j(x_N, \omega) > 0\right) = \\ & = P\left(0 < p_j(x_N, \omega) \leq \sqrt{\delta_N}\right) + P\left(p_j(x_N, \omega) > \sqrt{\delta_N}\right) \\ & \leq G_j(x_N, \sqrt{\delta_N}) - G_j(x_N, 0) + \frac{1}{\sqrt{\delta_N}} \mathbb{E}[p_j(x_N, \omega)] \\ & \leq G_j(x_N, \sqrt{\delta_N}) - G_j(x_N, 0) + \sqrt{\delta_N} \rightarrow 0, \text{ as } N \rightarrow \infty, j = 1, \dots, m. \end{aligned}$$

Here for a fixed  $x$ ,  $G_j(x, \cdot)$  denotes the distribution function of  $p_j(x, \omega)$  defined by

$$G_j(x, y) = P\left(p_j(x, \omega) \leq y\right), j = 1, \dots, m.$$

Assumption (iii) implies that for every vector  $\epsilon > 0$  (with small components) there exists some  $x_\epsilon \in X$  such that

$$P(g_{ji}(x_\epsilon, \omega) \leq 0, i = 1, \dots, k_j) \geq 1 - \epsilon_j, j = 1, \dots, m.$$

Then for any  $\epsilon > 0$  the following relations hold

$$\begin{aligned} & \sum_{j=1}^m \mathbb{E}[p_j(x_\epsilon, \omega)] = \\ & = \sum_{j=1}^m \int_{\Omega} |p_j(x_\epsilon, \omega)| I_{(p_j(x_\epsilon, \omega) > 0)} P(d\omega) \\ & \leq \sum_{j=1}^m \int_{\Omega} C(\omega) I_{(p_j(x_\epsilon, \omega) > 0)} P(d\omega) \\ & \leq \left( \int_{\Omega} C^{1+\kappa}(\omega) P(d\omega) \right)^{1/(1+\kappa)} \cdot \sum_{j=1}^m \left( \int_{\Omega} I_{(p_j(x_\epsilon, \omega) > 0)} P(d\omega) \right)^{\kappa/(1+\kappa)} \\ & \leq c \cdot \sum_{j=1}^m P\left(p_j(x_\epsilon, \omega) > 0\right)^{\kappa/(1+\kappa)} \\ & \leq c \cdot m \cdot \varepsilon_{max}^{\kappa/(1+\kappa)}, \end{aligned}$$

where  $c := \left( \int_{\Omega} C^{1+\kappa}(\omega) P(d\omega) \right)^{1/(1+\kappa)}$ , which is finite due to the assumption (ii). Accordingly, for  $\varepsilon_{max} \rightarrow 0_+$

$$0 \leq \sum_{j=1}^m \mathbb{E}[p_j(x_\epsilon, \omega)] \leq c \cdot m \cdot \varepsilon_{max}^{\kappa/(1+\kappa)} \rightarrow 0,$$

and also  $\beta_\epsilon(x_\epsilon) \rightarrow 0$ . If we set

$$\varepsilon_j(x_N) = P\left(p_j(x_N, \omega) > 0\right), j = 1, \dots, m,$$

then the optimal solution  $x_N$  of the expected value problem is feasible for the chance constrained program with  $\epsilon(x_N) = (\varepsilon_1(x_N), \dots, \varepsilon_m(x_N))$ , because the following relations hold

$$\begin{aligned} & P(g_{ji}(x_N, \omega) \leq 0, i = 1, \dots, k_j) \geq 1 - \varepsilon_j(x_N) \\ & \iff P\left(p_j(x_N, \omega) > 0\right) \leq \varepsilon_j(x_N). \end{aligned}$$

Hence, we get the inequality

$$\begin{aligned}\varphi_N &= f(x_N) + N \cdot \sum_{j=1}^m \mathbb{E}[p_j(x_N, \omega)] \\ &\geq f(x_{\varepsilon(x_N)}) + N \cdot \sum_{j=1}^m \mathbb{E}[p_j(x_N, \omega)] \\ &= \psi_{\varepsilon(x_N)} + \alpha_N(x_N).\end{aligned}$$

Finally,

$$\begin{aligned}\psi_\varepsilon &= \left( \psi_\varepsilon + \varepsilon_{max}^{-\eta} \sum_{j=1}^m \mathbb{E}[p_j(x_\varepsilon, \omega)] \right) - \varepsilon_{max}^{-\eta} \sum_{j=1}^m \mathbb{E}[p_j(x_\varepsilon, \omega)] \\ &\geq \varphi_{\varepsilon_{max}^{-\eta}} - \varepsilon_{max}^{-\eta} \sum_{j=1}^m \mathbb{E}[p_j(x_\varepsilon, \omega)] \\ &= \varphi_{\varepsilon_{max}^{-\eta}} - \beta_\varepsilon(x_\varepsilon).\end{aligned}$$

This completes the proof.

Note that the theorem does not make any statement on the convergence of optimal solutions but it relates optimal values for certain values of the levels and the penalty parameter. We will investigate the behaviour of the optimal solutions in the numerical study.

*Remark.* The assumption (iii) can be very strong. The problem is that the overall feasible set may shrink with increasing levels to the empty set, which makes the approach less appropriate for probability measures with an unbounded support.

*Remark.* The assumption (iv) ensures that the probability function

$$P(g_{ji}(x, \omega) \leq 0, i = 1, \dots, k_j)$$

is continuous in the decision vector, which can be easily seen if we realize that the only point of discontinuity of the function is  $g_{ji}(x, \omega) = 0$ ,  $i = 1, \dots, k_j$  for any  $x$ .

The bounds (2.4) and the terms  $\alpha_N(x)$ ,  $\varepsilon(x)$  and  $\beta_\varepsilon(x)$  depend on the choice of the penalty function  $\vartheta$ . Notice, however, that when we want to evaluate one of the bounds in (2.4), we must be prepared to face some problems. We are able to compute  $\alpha_N(x_N)$ ,  $\varepsilon(x_N)$ , hence the upper bound for the optimal value  $\psi_{\varepsilon(x_N)}$  of the chance constrained program (2.1) with probability levels  $\varepsilon(x_N)$ . But we are not able to compute  $\beta_{\varepsilon(x_N)}(x_{\varepsilon(x_N)})$  without having the solution  $x_{\varepsilon(x_N)}$  which we do not want to find or even may not be able to find.

### 3. SAMPLE APPROXIMATIONS USING MONTE-CARLO TECHNIQUES

In this part, we will address the rates of convergence for the chance constrained problems and the problems with expectation type objectives which cover the penalty type objectives. Usually, the sample approximation of the chance constrained problems leads only to the feasible solutions of the original problem. Moreover, the sample reformulation results in a large mixed-integer optimization problem, see below. Hence, it may be interesting to investigate the ability to generate the feasible solutions of the original chance constrained problem using the penalty function

TABLE 1. Formulation and approximation schema

	1.		2.		3.	
	Stochastic prog. formulation		Sample approx. (SA)		Solution validation	
Program with a random factor	→	Chance constrained problem (CCP)	→	SA CCP	→	Reliability
	↘	↓ Penalty function problem (PFP)	→	SA PFP	→	Reliability

problems, where no additional integer variables are necessary. Our approach is summarized in Table 1.

For the case when the set of feasible solutions, the objective function and the constraints are convex, stronger results on the sample approximations are valid, cf. [6]. The results below generalize those of [1, 10, 13] for the case with several chance constraints and they are valid without assuming convexity of any parts of the problems. We will draw our attention to the case when the set of feasible solutions is finite, i.e.  $|X| < \infty$ , and to the bounded infinite  $X$ .

In this section, we will refer to the problem (2.1) as the original problem. We denote the probability functions using the equivalence (2.2)

$$(3.1) \quad q_j(x) = P(p_j(x, \omega) > 0).$$

Then the multiple chance constrained problem (2.1) can be rewritten as

$$(3.2) \quad \begin{aligned} \psi_\epsilon &= \min_{x \in X} f(x), \\ \text{s.t.} \quad & q_1(x) \leq \epsilon_1, \\ & \vdots \\ & q_m(x) \leq \epsilon_m, \end{aligned}$$

Let  $\omega^1, \dots, \omega^S$  be an independent Monte Carlo sample of the random vector  $\omega$ . Then, the sample version of the function  $q_j$  is defined to be

$$(3.3) \quad \hat{q}_j^S(x) = S^{-1} \sum_{s=1}^S I_{(0, \infty)}(p_j(x, \omega^s)).$$

Finally, the sample version of the multiple jointly chance constrained problem (3.2) is defined as

$$(3.4) \quad \begin{aligned} \hat{\psi}_\gamma^S &= \min_{x \in X} f(x), \\ \text{s.t.} \quad & \hat{q}_1^S(x) \leq \gamma_1, \\ & \vdots \\ & \hat{q}_m^S(x) \leq \gamma_m, \end{aligned}$$

where the levels  $\gamma_j$  are allowed to be different from the original levels  $\epsilon_j$ . Let the set  $X$  be compact and  $g_{ji}(\cdot, \omega^s)$  be continuous for all triplets  $(i, j, s)$ . The sample approximation of the chance constrained problem can be reformulated as a large





**3.2. Feasibility for the chance constrained problem.** We derive the rate of convergence of the probability that the set of feasible solutions of the sample approximated problem is contained in the feasibility set of the original problem.

3.2.1. *Finite  $|X|$ .* First, we will draw our attention to the case when the set of feasible solutions is finite, i.e.  $|X| < \infty$ , which appears in the bounded integer programs. We will assume that it holds  $\gamma_j < \varepsilon_j$  for all  $j$ , i.e. that the levels of the sample approximated problem are more restrictive.

We define the random variable  $Y_{sj} = I_{(p_j(x, \omega^s) \leq 0)}$ , i.e.  $Y_{sj} = 1$  if  $p_j(x, \omega^s) \leq 0$  and 0 otherwise. Let

$$\begin{aligned} X_{\gamma_j}^S &= \{x \in X : \frac{1}{S} \sum_{s=1}^S Y_{js} \geq 1 - \gamma_j\}, \\ X_{\varepsilon_j} &= \{x \in X : P(p_j(x, \omega) \leq 0) \geq 1 - \varepsilon_j\}, \\ X_{\gamma}^S &= \bigcap_{j=1}^m X_{\gamma_j}^S, \\ X_{\varepsilon} &= \bigcap_{j=1}^m X_{\varepsilon_j}. \end{aligned}$$

Then, for  $x \in X \setminus X_{\varepsilon_j}$  we obtain  $\mathbb{E}[Y_{js}] = P(p_j(x, \omega) \leq 0) < 1 - \varepsilon_j$ , which we can use to get an estimate for the probability

$$\begin{aligned} P(x \in X_{\gamma_j}^S) &= P\left(\frac{1}{S} \sum_{s=1}^S Y_{js} \geq 1 - \gamma_j\right) \\ &\leq P\left(\sum_{s=1}^S (Y_{js} - \mathbb{E}[Y_{js}]) \geq S(\varepsilon_j - \gamma_j)\right) \\ (3.8) \quad &\leq \exp\{-2S(\varepsilon_j - \gamma_j)^2\}, \end{aligned}$$

where we used Hoeffding's inequality, cf. [7]. We use this estimate to get an upper bound for the probability that there exists a feasible solution of the sample approximated problem which is infeasible for the original problem.

$$\begin{aligned} 1 - P(X_{\gamma}^S \subseteq X_{\varepsilon}) &= P(\exists_{j \in \{1, \dots, m\}} \exists_{x \in X_{\gamma}^S} : P(p_j(x, \omega) \leq 0) < 1 - \varepsilon_j) \\ &\leq \sum_{j=1}^m \sum_{x \in X \setminus X_{\varepsilon_j}} P(x \in X_{\gamma_j}^S) \\ &\leq |X \setminus X_{\varepsilon}| \sum_{j=1}^m \exp\{-2S(\varepsilon_j - \gamma_j)^2\} \\ &\leq m|X \setminus X_{\varepsilon}| \exp\left\{-2S \min_{j \in \{1, \dots, m\}} (\varepsilon_j - \gamma_j)^2\right\}. \end{aligned}$$

Using previous upper bound it is possible to estimate the sample size  $S$  such that the feasible solutions of the sample approximated problems are feasible for the original problem with a high probability  $1 - \delta$ , i.e.

$$(3.9) \quad S \geq \frac{1}{2 \min_{j \in \{1, \dots, m\}} (\gamma_j - \varepsilon_j)^2} \ln \frac{m|X \setminus X_{\varepsilon}|}{\delta}.$$

If we set  $m = 1$ , we get the same inequality as [10].

3.2.2. *Bounded  $|X|$ .* Below we will consider the case when the set of feasible solutions  $X$  is bounded but infinite in general. Again, let  $\gamma_j < \varepsilon_j$  for all  $j$ . However, we will need the following additional assumption which states Lipschitz continuity of the penalized constraints, i.e.

$$|p_j(x, \omega) - p_j(x', \omega)| \leq L_j \|x - x'\|, \quad \forall x, x' \in X, \quad \forall \omega \in \Omega, \quad \forall j,$$

for some  $L_j > 0$ . Let  $D = \sup\{\|x - x'\|_\infty : x, x' \in X\}$  be the diameter of  $X$ . In this case, it is necessary to consider the constraints which are satisfied strictly, i.e. with some deviation  $\tau$ :

$$\begin{aligned} X_{\gamma_j, \tau}^S &= \{x \in X : \frac{1}{S} \sum_{s=1}^S I_{(p_j(x, \omega^s) + \tau \leq 0)} \geq 1 - \gamma_j\}. \\ X_{\gamma, \tau}^S &= \bigcap_{j=1}^m X_{\gamma_j, \tau}^S. \end{aligned}$$

According to the proof of [10, Theorem 10], for  $\lambda_j \in (0, \varepsilon_j - \gamma_j)$  there exist finite sets  $Z_j^\tau \subseteq X$  with

$$|Z_j^\tau| \leq \lceil 1/\lambda_j \rceil \lceil 2L_j D/\tau \rceil^n$$

where  $\lceil \cdot \rceil$  denotes the upper integer part, and for any  $x \in X_{\gamma_j, \tau}^S$  and any  $j$  there exists  $z \in Z_j^\tau$  such that  $\|z - x\|_\infty \leq \tau/L_j$ . Using the finite sets  $Z_j^\tau$  we can define

$$\begin{aligned} Z_{\gamma_j}^{\tau, S} &= \{x \in Z_j^\tau : \frac{1}{S} \sum_{s=1}^S I_{(p_j(x, \omega^s) \leq 0)} \geq 1 - \gamma_j\}, \\ Z_{\varepsilon_j - \lambda_j}^\tau &= \{x \in Z_j^\tau : P(p_j(x, \omega) \leq 0) \geq 1 - \varepsilon_j + \lambda_j\}, \\ Z_\gamma^{\tau, S} &= \bigcap_{j=1}^m Z_{\gamma_j}^{\tau, S}, \\ Z_{\varepsilon - \lambda}^\tau &= \bigcap_{j=1}^m Z_{\varepsilon_j - \lambda_j}^\tau, \end{aligned}$$

Moreover, for all  $j$  it holds that  $Z_{\gamma_j}^{\tau, S} \subseteq Z_{\varepsilon_j - \lambda_j}^\tau$  implies  $X_{\gamma_j}^{\tau, S} \subseteq X_{\varepsilon_j}$ . For the previous finite sets, the inequality (3.8) is valid, i.e. we obtain

$$\begin{aligned} 1 - P(Z_\gamma^{\tau, S} \subseteq Z_{\varepsilon - \lambda}^\tau) &\leq m \left[ \frac{1}{\min_{j \in \{1, \dots, m\}} \lambda_j} \right] \left[ \frac{2L_{\max} D}{\tau} \right]^n \\ &\quad \exp \left\{ -2S \min_{j \in \{1, \dots, m\}} (\varepsilon_j - \gamma_j - \lambda_j)^2 \right\}, \end{aligned}$$

where  $L_{\max} = \max_j L_j$ . Since  $Z_\gamma^{\tau, S} \subseteq Z_{\varepsilon - \lambda}^\tau$  implies  $X_\gamma^{\tau, S} \subseteq X_\varepsilon$ , we get the inequality for the probabilities

$$P(X_{\gamma, \tau}^S \subseteq X_\varepsilon) \geq P(Z_\gamma^{\tau, S} \subseteq Z_{\varepsilon - \lambda}^\tau).$$

Using the bound it is possible to estimate the sample size  $S$  such that the feasible solutions of the sample approximated problems are feasible for the original problem

with a high probability  $1 - \delta$ , i.e.

$$S \geq \frac{1}{2 \min_{j \in \{1, \dots, m\}} (\varepsilon_j - \gamma_j - \lambda_j)^2} \left( \ln \frac{m}{\delta} + \ln \left[ \frac{1}{\min_{j \in \{1, \dots, m\}} \lambda_j} \right] + n \ln \left[ \frac{2L_{\max} D}{\tau} \right] \right).$$

If we choose  $\lambda_j = (\varepsilon_j - \gamma_j)/2$ , we obtain

$$S \geq \frac{2}{\min_{j \in \{1, \dots, m\}} (\varepsilon_j - \gamma_j)^2} \left( \ln \frac{m}{\delta} + \ln \left[ \frac{2}{\min_{j \in \{1, \dots, m\}} (\varepsilon_j - \gamma_j)} \right] + n \ln \left[ \frac{2L_{\max} D}{\tau} \right] \right).$$

Setting  $m = 1$  we obtain the same estimate as [10].

**3.3. Sample approximation for stochastic programs with expectation type objectives.** In this section we will review the main results of [20] on the sample average approximation (SAA) techniques for the expectation type stochastic programs with a finite or bounded set of feasible solutions.

3.3.1. *Finite*  $|X|$ . Let  $F(x, \omega)$  denote the objective function which is integrated over  $\omega$ , e.g. in the penalty approach

$$F(x, \omega) = f(x) + N \cdot \sum_{j=1}^m p_j(x, \omega),$$

and  $f(x) = \mathbb{E}[F(x, \omega)]$  be its expectation. Let  $\Phi_\zeta$  be the set of  $\zeta$ -optimal solutions. Let

$$Y(x, \omega) = F(u(x), \omega) - F(x, \omega),$$

where  $u$  is a function from  $X \setminus \Phi_\zeta$  into the set  $X$  such that

$$f(u(x)) \leq f(x) - \zeta^*, \quad \forall x \in X \setminus \Phi_\zeta$$

for some  $\zeta^* > \zeta$ . Denote

$$\nu(\hat{\zeta}, \zeta) = \min_{x \in X \setminus \Phi_\zeta} R(x, -\hat{\zeta}),$$

where  $R$  is the large deviations rate function of the random variable  $Y$  which is defined as the conjugate function to the logarithmic moment generating function, i.e.

$$R(x, \zeta) = \sup_{t \in \mathbb{R}} \left\{ t\zeta - \ln \mathbb{E}[e^{tY(x, \omega)}] \right\}.$$

Then, for the probability that the set of  $\hat{\zeta}$ -optimal solutions of the sample average approximated problem is included in the set of  $\zeta$ -optimal solutions, it holds

$$1 - P(\hat{\Phi}_\zeta^S \subseteq \Phi_\zeta) \leq |X| \exp\{-S\nu(\hat{\zeta}, \zeta)\}.$$

The function  $\nu$  can be further estimated as

$$\nu(\hat{\zeta}, \zeta) \geq \frac{(\zeta - \hat{\zeta})^2}{3\sigma_{\max}^2},$$

where

$$\sigma_{max}^2 = \max_{x \in X \setminus \Phi_\zeta} \text{Var}[F(u(x), \omega) - F(x, \omega)].$$

Then, the sample size  $S$ , which is necessary to generate  $\hat{\zeta}$ -optimal solutions which are also  $\zeta$ -optimal for the original problem with a high probability  $1 - \delta$ , can be estimated as, cf. [20],

$$S \geq \frac{3\sigma_{max}^2}{(\zeta - \hat{\zeta})^2} \ln \frac{|X|}{\delta},$$

where  $1 - \delta$  is the prescribed probability. It is necessary to mention that the term  $\sigma_{max}^2$  depends on the penalty parameter  $N$  in quadratic manner.

3.3.2. *Bounded  $|X|$ .* In the case that the set of feasible solutions  $X$  is bounded, not necessarily finite, and the function  $F(x, \omega)$  is Lipschitz continuous on  $X$  modulus  $L$  which does not depend on  $\omega$ , i.e.

$$|F(x, \omega) - F(x', \omega)| \leq L \|x - x'\|, \quad \forall x, x' \in X, \quad \forall \omega \in \Omega,$$

then we can get the following estimate for the sample size necessary to generate  $\hat{\zeta}$ -optimal solutions which are also  $\zeta$ -optimal for the original problem with a high probability  $1 - \delta$ , cf. [20],

$$S \geq \frac{12\sigma_{max}^2}{(\zeta - \hat{\zeta})^2} \left( n \ln \frac{2DL}{\zeta - \hat{\zeta}} - \ln \delta \right).$$

As can be easily seen, the estimate depends linearly on the dimension  $n$  of the decision variables  $x$ .

#### 4. MIXED-INTEGER VAR AND PENALTY FUNCTION PROBLEMS

In this section, we compare the penalty function approach with the chance constrained problems on a mixed-integer portfolio problem of a small investor. We consider 13 most liquid assets which are traded on the main market (SPAD) on Prague Stock Exchange. Weekly returns from the period 6th February 2009 to 10th February 2010 are used to estimate the means and the variance matrix. Suppose that the small investor trades assets on the "mini-SPAD" market. This market enables to trade "mini-lots" (standardized number of assets) with favoured transaction costs.

We denote  $Q_i$  the quotation of the "mini-lot" of security  $i$ ,  $f_i$  the fixed transaction costs (not depending on the investment amount),  $c_i$  the proportional transaction costs (depending on the investment amount),  $R_i$  the random return of the security  $i$ ,  $x_i$  the number of "mini-lots",  $y_i$  binary variables which indicate, whether the security  $i$  is bought or not. Then, the random loss function depending on our decisions and the random returns has the following form

$$-\sum_{i=1}^n (R_i - c_i) Q_i x_i + \sum_{i=1}^n f_i y_i.$$

The chance constrained portfolio problem can be formulated as follows

$$(4.1) \quad \min_{(r,x,y) \in \mathbb{R} \times X} r \\ P\left(-\sum_{i=1}^n (R_i - c_i)Q_i x_i + \sum_{i=1}^n f_i y_i \leq r\right) \geq 1 - \varepsilon,$$

which is in fact minimization of Value at Risk (VaR). Corresponding penalty function problem using the penalty  $\vartheta^{1,1}$  is

$$(4.2) \quad \min_{(r,x,y) \in \mathbb{R} \times X} r + N \cdot \mathbb{E}\left[-\sum_{i=1}^n (R_i - c_i)Q_i x_i + \sum_{i=1}^n f_i y_i - r\right]^+.$$

Setting  $N = 1/(1 - \varepsilon)$  we minimize Conditional Value at Risk (CVaR) exactly, see [18]. Similar problem with CVaR and transaction costs was considered by [2] and its stability was studied by [4].

The set of feasible solutions contains a budget constraint and the restrictions on the minimal and the maximal number of "mini-lots" which can be bought, i.e.

$$X = \{x \in \mathbb{N}^n \times \{0, 1\}^n \\ B_l \leq \sum_{i=1}^n (1 + c_i)Q_i x_i + \sum_{i=1}^n f_i y_i \leq B_u, \\ l_i y_i \leq x_i \leq u_i y_i, \quad i = 1, \dots, n\},$$

where  $B_l$  and  $B_u$  are the lower and the upper bound on the capital available for the portfolio investment,  $l_i > 0$  and  $u_i > 0$  are the lower and the upper number of units for each security  $i$ .

**4.1. Estimated sample sizes.** In our case, the cardinality of the integer part of the set of feasible solutions is bounded, i.e.  $|X| \leq 116^{13} \cdot 2^{13}$ . Moreover, if the support of the distribution of the returns is bounded, than the free variable  $t$  can be restricted to the closed interval which is bounded by the worst loss and by the best profit which can occur for our loss function considering the restrictions. Then we get the following estimate for the sample size which is necessary to generate a lower bound for the optimal value

$$S \geq \frac{2\varepsilon}{(\gamma - \varepsilon)^2} \ln \frac{1}{\delta},$$

and to generate a feasible solution

$$S \geq \frac{2}{(\varepsilon - \gamma)^2} \left( \ln \frac{1}{\delta} + 13 \ln 116 + 13 \ln 2 + \ln \left\lceil \frac{2}{(\varepsilon - \gamma)} \right\rceil + \ln \left\lceil \frac{2D}{\tau} \right\rceil \right),$$

which is based on the decomposition of the set of feasible solutions into the integer and real bounded part. In Tables 2 and 3, there are examples of the sample sizes for different combinations of the parameters  $\gamma, \varepsilon, \delta$  where we have chosen  $\tau = 10^{-6}$  and  $D = 2 \cdot 10^6$  which is the difference between the worst loss and the best profit. The sample size which is necessary to generate the lower bound for the optimal value of the original problem is quite low and will be covered partly by the following numerical experiment, see Table 2. However, the samples, which are necessary to ensure that the set of feasible solutions of the sample approximated problem is contained in the feasibility set of the original problem, are quite large and rapidly increase with decreasing level  $\varepsilon$ , see Table 3.

TABLE 2. Sample sizes - lower bound

$\varepsilon$	$\gamma$	$\delta$	$S$
0.1	0.2	0.01	93
0.05	0.1	0.01	185
0.01	0.02	0.01	9211
0.1	0.2	0.001	139
0.05	0.1	0.001	277
0.01	0.02	0.001	13816

TABLE 3. Sample sizes - feasibility

$\varepsilon$	$\gamma$	$\delta$	$S$
0.1	0.05	0.01	86496
0.05	0.025	0.01	348199
0.01	0.005	0.01	901792970
0.1	0.05	0.001	88338
0.05	0.025	0.001	355567
0.01	0.005	0.001	920213650

**4.2. Numerical comparison.** We generated 100 samples for each sample size  $S$ , i.e.  $100 \times S$  realizations, from the truncated normal distribution where the truncation points were set to  $-1$  for all random returns. We used the modelling system GAMS and the solver CPLEX to solve the sample approximations of the chance constrained problems (4.1) and the penalty function problems (4.2) for different sample sizes  $S$ , levels  $\gamma$  and penalty parameters  $N$ . Descriptive statistics for the results are contained in Tables 4, 5, 6. As we can see from Table 6, the "Penalty term"

$$N \cdot \mathbb{E} \left[ - \sum_{i=1}^n (R_i - c_i) Q_i x_i + \sum_{i=1}^n f_i y_i - r \right]^+$$

really decreases with increasing penalty parameter  $N$  and reduces violations of the constraint  $(R_i - c_i) Q_i x_i + \sum_{i=1}^n f_i y_i - r \leq 0$  for each sample size.

To verify the reliability of the obtained optimal solutions, we used the independent samples of 10 000 realizations from the truncated normal distribution which was used to model the random returns. The columns "Reliability" contain relative number of realizations for which the chance constraint is fulfilled. As can be easily seen, the reliability of the obtained solutions increases with increasing levels  $\gamma$  and penalty parameters  $N$  for each sample size  $S$ . Both problems are also able to generate comparable solutions for the same sample sizes, see Tables 4, 5. Furthermore, we can compare the descriptive statistics of the optimal values  $\hat{\psi}_\gamma^S$ ,  $\hat{\varphi}_N^S$  and the optimal solutions  $\hat{r}_N^S$  of the problems. We observe that the variability of the values increases with the sample size. Thus, we pay for the increasing reliability of the optimal solutions by decreasing reliability of the optimal values when we increase the size of the sample. Finally, we can compare the used sample sizes with theoretically estimated sizes in Tables 2 and 3, which can be now seen as very conservative.

TABLE 4. Chance constrained problems

$S$	$\gamma$	Reliability				$\hat{\psi}_\gamma^S$			
		min	max	mean	st.dev	min	max	mean	st.dev
100	0.1	0.8844	0.9967	0.9592	0.0255	29739.36	66854.82	41784.66	7525.69
100	0.05	0.9054	0.9869	0.9516	0.0189	29739.36	66854.82	41821.60	7465.46
100	0.01	0.8939	0.9941	0.9456	0.0250	29680.35	69513.05	42312.34	7612.11
250	0.1	0.9546	0.9968	0.9824	0.0098	37609.63	121252.72	52429.77	9887.54
250	0.05	0.9545	0.9950	0.9820	0.0086	37609.63	121252.72	52431.23	9884.16
250	0.01	0.9555	0.9950	0.9807	0.0115	38260.62	121972.21	52626.23	9909.60
500	0.1	0.9744	0.9982	0.9903	0.0043	45085.97	125638.34	67824.32	15849.91
500	0.05	0.9744	0.9982	0.9903	0.0043	45085.97	125638.34	67824.32	15849.91
500	0.01	0.9726	0.9982	0.9906	0.0043	45085.97	125638.34	67942.02	15757.14
750	0.1	0.9849	0.9994	0.9952	0.0033	48562.73	160984.79	74655.08	19435.11
750	0.05	0.9849	0.9994	0.9952	0.0033	48562.73	160984.79	74652.82	19436.71
750	0.01	0.9866	0.9994	0.9953	0.0032	48562.73	155469.46	74679.40	19187.28
1000	0.1	0.9870	1.0000	0.9966	0.0025	59129.41	187831.95	93390.26	28293.28
1000	0.05	0.9870	1.0000	0.9966	0.0025	59129.41	187831.95	93414.25	28269.13
1000	0.01	0.9870	1.0000	0.9966	0.0025	59129.41	187831.95	93384.85	28264.63

## 5. CONCLUSION

Reformulation of chance constrained programs by incorporating a suitably chosen penalty function into the objective helps to arrive at problems with expectation in the objective and a fixed set of feasible solutions. The obtained problems are much simpler to solve and analyze than the chance constrained programs. The recommended form of the penalty function follows the basic ideas of penalty methods and its suitable properties follow by generalization of the results from [5, 9].

The numerical study shows that not only the sample approximated chance constrained problems but also the penalty function problems are able to generate the solutions which are feasible for the original chance constrained problem with a high reliability.

**Acknowledgements** I would like to thank to Prof. RNDr. Jitka Dupačová, DrSc. for her valuable comments. I gratefully acknowledge the support of the Czech Science Foundation (GA CR 201/08/0486 and GD CR 402/09/H045).

TABLE 5. Penalty function problems

		Reliability				$\hat{r}_N^S$			
$S$	$N$	min	max	mean	st.dev	min	max	mean	st.dev
100	0	0.5504	0.5504	0.5504	0.0000	0.00	0.00	0.00	0.00
100	0.1	0.0000	0.0225	0.0030	0.0049	-9731888	-107661	-3400803	3404909
100	1	0.7622	0.9480	0.8770	0.0303	14479.93	40608.34	25672.46	3800.89
100	10	0.8967	0.9976	0.9581	0.0220	30739.36	67854.82	42827.32	7492.45
100	100	0.8967	0.9976	0.9581	0.0219	30739.36	67854.82	42902.79	7484.36
100	1000	0.8967	0.9976	0.9581	0.0218	30739.36	67854.82	42903.93	7474.20
250	0	0.5453	0.5453	0.5453	0.0000	0.00	0.00	0.00	0.00
250	0.1	0.0000	0.0105	0.0007	0.0018	-9840593.01	-193870	-5387627	3120485
250	1	0.8330	0.9290	0.8888	0.0199	20333.22	62991.61	27709.38	4866.67
250	10	0.9495	0.9950	0.9788	0.0101	36429.23	116137.42	49586.12	8798.91
250	100	0.9571	0.9973	0.9841	0.0089	39630.90	122252.72	53493.47	9862.21
250	1000	0.9571	0.9973	0.9840	0.0089	39630.90	122252.72	53458.34	9898.87
500	0	0.5408	0.5408	0.5408	0.0000	0.00	0.00	0.00	0.00
500	0.1	0.0000	0.0061	0.0004	0.0011	-9880574	-248703	-5721038	3324282
500	1	0.8716	0.9270	0.9016	0.0134	22916.95	54037.31	31671.51	5783.07
500	10	0.9723	0.9955	0.9871	0.0044	42674.84	100497.95	58776.94	12368.39
500	100	0.9813	0.9996	0.9935	0.0033	46085.97	126638.34	68995.38	15851.31
500	1000	0.9813	0.9995	0.9934	0.0033	46085.97	126638.34	68914.67	15748.83
750	0	0.5408	0.5408	0.5408	0.0000	0.00	0.00	0.00	0.00
750	0.1	0.0000	0.0032	0.0002	0.0006	-9912905	-281868	-6224877	3088217
750	1	0.8697	0.9330	0.8990	0.0108	23694.91	51361.54	31731.28	5614.38
750	10	0.9785	0.9950	0.9878	0.0036	43208.99	133243.07	60923.36	14886.03
750	100	0.9890	0.9995	0.9957	0.0026	49562.73	157103.91	75669.31	19379.62
750	1000	0.9890	0.9993	0.9956	0.0026	49562.73	157103.91	75541.31	19234.11
1000	0	0.5537	0.5537	0.5537	0.0000	0.00	0.00	0.00	0.00
1000	0.1	0.0000	0.0026	0.0002	0.0005	-9818182	-291063	-6513630	3051261
1000	1	0.8739	0.9253	0.8976	0.0097	25121.67	59977.76	35192.10	7145.00
1000	10	0.9753	0.9964	0.9886	0.0038	46083.49	134622.66	72959.07	19872.90
1000	100	0.9900	0.9999	0.9966	0.0023	59121.39	182075.76	94331.08	27977.78
1000	1000	0.9900	0.9999	0.9966	0.0023	59121.39	182561.86	94357.45	28209.17

## REFERENCES

- [1] S. Ahmed, A. Shapiro (2008). *Solving chance-constrained stochastic programs via sampling and integer programming*. In *Tutorials in Operations Research*, Z.-L. Chen and S. Raghavan (eds.), INFORMS.
- [2] E. Angelelli, R. Mansini, M.G. Speranza (2008). *A comparison of MAD and CVaR models with real features*. *Journal of Banking and Finance* 32, 1188-1197.
- [3] M.S. Bazara, H.D. Sherali, C.M. Shetty (1993). *Nonlinear programming: theory and algorithms*. Wiley, Singapore.
- [4] M. Branda (2008). *Stability Analysis of Mean-CVaR Investment Model with Transaction Costs and Integer Allocations*. *Proceedings of 10th International PhD Workshop on Systems and Control*, ed. Lenka Pavelková.
- [5] M. Branda, J. Dupačová (2008). *Approximations and contamination bounds for probabilistic programs*. SPEPS-13.
- [6] G. Calafiore, M.C. Campi (2005). *Uncertain convex programs: randomized solutions and confidence levels*. *Mathematical Programming, Ser. A* 102, 25-46.
- [7] A. DasGupta (2008). *Asymptotic theory of Statistics and Probability*. Springer, New York.
- [8] J. Dupačová, A. Gaivoronski, Z. Kos, T. Szantai (1991). *Stochastic programming in water management: A case study and a comparison of solution techniques*. *European Journal of Operational Research* 52, 28-44.



TABLE 6. Penalty function problems

		$\phi_N^S$				Penalty term			
$S$	$N$	min	max	mean	st.dev	min	max	mean	st.dev
100	0	2725.06	6353.81	4013.20	641.07	2725.06	6353.81	4013.20	641.07
100	0.1	-162909	-7245.51	-45319.08	32662.98	93461	9611898	3355484	3375082
100	1	24011.20	45692.02	33403.52	4311.27	2622.61	15554.88	7731.06	2530.95
100	10	30739.36	67854.82	42830.41	7489.58	0.00	309.34	3.09	30.93
100	100	30739.36	67854.82	42902.79	7484.36	0.00	0.00	0.00	0.00
100	1000	30739.36	67854.82	42903.93	7474.20	0.00	0.00	0.00	0.00
250	0	2868.20	11230.04	4165.97	941.94	2868.20	11230.04	4165.97	941.94
250	0.1	-88791.11	-3047.66	-33455.23	16216.02	181656	9781201	5354172	3108784
250	1	29745.59	83386.57	37382.48	6017.16	5787.60	20394.96	9673.10	2115.72
250	10	37848.38	118742.60	52156.49	9360.82	162.04	12197.59	2570.37	2255.62
250	100	39630.90	122252.72	53493.47	9862.21	0.00	0.00	0.00	0.00
250	1000	39630.90	122252.72	53458.34	9898.87	0.00	0.00	0.00	0.00
500	0	3448.54	6252.90	4202.39	596.15	3448.54	6252.90	4202.39	596.15
500	0.1	-58968.10	-8316.52	-23030.03	8951.17	233508	9846163	5698009	3318424
500	1	32345.07	71348.70	43537.39	8424.45	6682.76	27887.84	11865.88	3520.77
500	10	45481.55	110479.11	63886.92	13472.75	886.09	12248.70	5109.98	2719.34
500	100	46085.97	126638.34	68995.38	15851.31	0.00	0.00	0.00	0.00
500	1000	46085.97	126638.34	68914.67	15748.83	0.00	0.00	0.00	0.00
750	0	3337.36	6177.69	4121.26	451.58	3337.36	6177.69	4121.26	451.58
750	0.1	-38745.72	-7073.09	-20443.37	6677.07	266041	9885991	6204434	3085053
750	1	33415.20	94959.23	44922.49	9914.34	7304.04	43597.69	13191.21	5100.68
750	10	47249.47	150732.32	68251.45	17167.97	1995.04	18457.88	7328.08	3405.62
750	100	49562.73	157103.91	75669.31	19379.62	0.00	0.00	0.00	0.00
750	1000	49562.73	157103.91	75541.31	19234.11	0.00	0.00	0.00	0.00
1000	0	3567.85	5396.22	4124.17	387.42	3567.85	5396.22	4124.17	387.42
1000	0.1	-32111.10	-10628.79	-18340.84	4784.46	277825	9800651	6495290	3049268
1000	1	34061.30	98653.13	51840.01	12169.11	8886.75	42469.68	16647.91	7013.17
1000	10	51622.86	162568.42	82550.78	23493.47	2795.02	29325.36	9591.71	5108.62
1000	100	59121.39	182075.76	94331.08	27977.78	0.00	0.00	0.00	0.00
1000	1000	59121.39	182561.86	94357.45	28209.17	0.00	0.00	0.00	0.00

- [9] Y.M. Ermoliev, T.Y. Ermolieva, G.J. Macdonald, V.I. Norkin (2000). *Stochastic optimization of insurance portfolios for managing exposure to catastrophic risks*. Annals of Operations Research **99**, pp. 207–225.
- [10] J. Luedtke, S. Ahmed (2008). *A sample approximation approach for optimization with probabilistic constraints*. SIAM Journal on Optimization, vol.19, pp.674-699.
- [11] J. Nocedal, S.J. Wright (2000). *Numerical optimization*. Second Edition, Springer, New York.
- [12] B. Pagnoncelli, S. Ahmed, A. Shapiro (2008). *Computational study of a chance constrained portfolio selection problem*. Optimization Online.
- [13] B. Pagnoncelli, S. Ahmed, A. Shapiro (2009). *Sample Average Approximation Method for Chance Constrained Programming: Theory and Applications*. Journal Optimization Theory and Applications, vol. 142, pp. 399-416.
- [14] A. Prékopa (1973). *Contributions to the theory of stochastic programming*. Mathematical programming **4**, 202–221.
- [15] A. Prékopa (1990). *Dual method for a one-stage stochastic programming problem with random rhs obeying a discrete probability distribution*. Z. Oper. Res. 34, 441461.
- [16] A. Prékopa (1995). *Stochastic Programming*. Kluwer, Dordrecht and Akadémiai Kiadó, Budapest.
- [17] A. Prékopa (2003). *Probabilistic Programming*. In Stochastic Programming (A. Ruszczyński, A. Shapiro eds.), Handbook in Operations Research and Management Science Vol. 10, Elsevier, Amsterdam, 483-554.

- [18] R.T. Rockafellar, S. Uryasev (2002). *Conditional Value-at-Risk for General Loss Distributions*. Journal of Banking and Finance, 26, 1443-1471.
- [19] R.T. Rockafellar, R. Wets (2004). *Variational analysis*. 2nd edition, Springer-Verlag, Berlin.
- [20] A. Shapiro (2003). *Monte Carlo Sampling Methods*. In Stochastic Programming (A. Ruszczyński, A. Shapiro eds.), Handbook in Operations Research and Management Science Vol. 10, Elsevier, Amsterdam, 483-554.
- [21] S.W. Wallace, W.T. Ziemba (2005). *Applications of stochastic programming*. MPS-SIAM Book Series on Optimization 5.

Martin Branda  
Department of Probability and Mathematical Statistics  
Faculty of Mathematics and Physics  
Charles University in Prague  
Czech Republic  
branda@karlin.mff.cuni.cz