

6-1-1993

Nutrient Content of Runoff Water From Rice Fields

C. E. Wilson Jr.

Southeast Research and Extension Center

P. A. Moore Jr.

USDA Agricultural Research Service


R. J. Norman

Rice Research and Extension Center

B. R. Wells

University of Arkansas, Fayetteville

Follow this and additional works at: <http://scholarworks.uark.edu/awrctr>

 Part of the [Fresh Water Studies Commons](#), [Hydrology Commons](#), [Soil Science Commons](#), and the [Water Resource Management Commons](#)

Recommended Citation

Wilson, C. E. Jr.; Moore, P. A. Jr.; Norman, R. J.; and Wells, B. R.. 1993. Nutrient Content of Runoff Water From Rice Fields. Arkansas Water Resources Center, Fayetteville, AR. PUB 164. 29

This Technical Report is brought to you for free and open access by the Arkansas Water Resources Center at ScholarWorks@UARK. It has been accepted for inclusion in Technical Reports by an authorized administrator of ScholarWorks@UARK. For more information, please contact scholar@uark.edu, ccmiddle@uark.edu.

NUTRIENT CONTENT OF RUNOFF WATER FROM RICE FIELDS

C.E. Wilson, Jr.
Southeast Research and Extension Center

P.A. Moore, Jr.
USDA Agricultural Research Service

R.J. Norman
Rice Research and Extension Center

and

B.R. Wells
Department of Agronomy

Research Project Technical Completion Report

Project G-1549-05

The research upon which this report is based was financed in part by the United States Department of the Interior as authorized by the Water Research and Development Act of 1984 (P.L. 98-242).

Arkansas Water Resources Center
University of Arkansas
113 Ozark Hall
Fayetteville, AR 72701

Publication No. 164

July 1, 1992 - June 30, 1993

Contents of this publication do not necessarily reflect the views and policies of the U.S. Department of the Interior, nor does mention of trade names or commercial products constitute their endorsement or recommendation of or use by the U.S. Government.

The University of Arkansas, in compliance with federal and state laws and regulations governing affirmative action and non-discrimination, does not discriminate in the recruitment, admission, and employment of students, faculty, and staff in the operation of any of its educational programs and activities as defined by law. Accordingly, nothing in this publication should be viewed as directly or indirectly expressing any limitation, specification, or discrimination as to race, religion, color, or national origin; or to handicap, age, sex, or status as a disabled Vietnam-era veteran; except as provided by law. Inquiries concerning this policy may be directed to the Affirmative Action Officer.

TABLE OF CONTENTS

Abstract. 3

List of Figures 5

Acknowledgements. 6

Introduction. 7

A. Purpose and Objectives. 7

B. Related Research and Activities 7

Methods and Procedures. 9

Principle Findings and Significance 10

A. Inorganic N 10

B. Soluble P 13

C. Metals. 15

D. Comparison of Water Sources 17

Conclusions 18

Literature Cited. 19

ABSTRACT

NUTRIENT CONTENT OF RUNOFF WATER FROM RICE FIELDS

Current perception is that nutrient runoff from croplands is a significant contributor to poor water quality in some areas. While extensive research has been conducted to survey and ameliorate this problem for several upland crops, little work has been done to evaluate the problem with flooded rice (*Oryza sativa*, L.) soils. Since rice production utilizes a major portion of the total irrigation water usage for certain areas, it is important to understand the contribution of rice production to non-point source N and P in surface water. Several production fields were selected to evaluate the concentrations of nutrients in the floodwater at selected distances across the field, including inlet and exit. The fields were evaluated in either 1990, 1991, or 1992 and were managed by the individual rice producer. Water samples were collected from several locations within each field weekly following establishment of the permanent flood and analyzed for inorganic N ($\text{NH}_4\text{-N}$, $\text{NO}_3\text{-N}$, and $\text{NO}_2\text{-N}$) and soluble P. The N concentrations in the floodwater normally peaked following N fertilizer application but rapidly declined and remained below 1 mg N L^{-1} . Water management resulted in some variation among locations with respect to the timing and magnitude of these peaks. The P concentrations were usually highest near the well and declined to less than 0.05 mg P L^{-1} as the water moved across the field. This was attributed to plant uptake, uptake by algae, and sediment deposition. The data indicates that rice fields have the potential to be utilized as a

filtration system to reduce the nutrient load of irrigation water similar to constructed wetlands. Use of catfish pondwater, in comparison to well water, resulted in only slightly higher total N and total P levels with higher amounts of the nutrients in the organic form. Although the P levels were high enough to potentially contribute to eutrophication of surface water, the water exiting the field was lower than at the entry point irrespective of the source. Also, the total P (organic + inorganic) concentration was less than 0.05 mg P L^{-1} .

LIST OF FIGURES

- Fig. 1. Seasonal changes in $\text{NH}_4\text{-N}$ concentration in rice irrigation water.
Page 22
- Fig. 2. Seasonal changes in $\text{NO}_3\text{-N}$ concentration in rice irrigation water.
Page 23
- Fig. 3. Changes in $\text{NH}_4\text{-N}$ concentration in rice irrigation water with respect to relative distance from the well.
Page 24
- Fig. 4. Changes in $\text{NO}_3\text{-N}$ concentration in rice irrigation water with respect to relative distance from the well.
Page 25
- Fig. 5. Changes in soluble reactive phosphorus ($\text{PO}_4\text{-P}$) concentration in rice irrigation water with respect to distance across the field.
Page 26
- Fig. 6. Seasonal changes in soluble reactive phosphorus ($\text{PO}_4\text{-P}$) concentration in rice irrigation water.
Page 27
- Fig. 7. Concentrations of $\text{NH}_4\text{-N}$, $\text{NO}_3\text{-N}$, and total N in rice irrigation water from a well (A) or catfish pond (B) with respect relative to distance from the source.
Page 28
- Fig. 8. Concentrations of soluble reactive phosphorus ($\text{PO}_4\text{-P}$) and total P in rice irrigation water from a well (A) or catfish pond (B) with respect to relative distance from the source.
Page 29

ACKNOWLEDGEMENTS

The authors would like to extend thanks to those who assisted with collection and analysis of the many samples for this study including S. Castile, D. Brooks, S. Fischer, and A. Weaver. We would also like to extend our thanks to the producers (Fischer Farms, Summerford Farms, M&J Farms, Bradshaw Farms, and Reinhart Farms) for allowing us to conduct this research on their farms.

INTRODUCTION

A. Purpose and Objectives

Recently the concern about environmental pollution resulting from application of commercial fertilizers and pesticides in agricultural situations has increased. Because of this concern, it becomes increasingly important to know how much contamination of surface water resulting from these applications is actually present. Because information concerning N and P runoff from rice fields is limited, this study was initiated to evaluate the seasonal variation in nutrient contents in the rice floodwater with distance across the field.

B. Related Research and Activities

Nitrogen and P in runoff from agricultural lands have been investigated extensively, particularly for upland crop situations (Stanford et al., 1970; Water Resources Committee, 1970; Lin, 1972; Loehr, 1974; Chichester and Richardson, 1992). Some of the major topics of recent research include N and P runoff as influenced by tillage practices, irrigation, and various cropping situations (e.g., Sharpley et al., 1992; Lowrance, 1992; Chichester and Richardson, 1992). Most of the research has focused on upland crops such as corn (*Zea mays*, L.), wheat (*Triticum aestivum*, L.) and soybean (*Glycine max*, (L.) Merr.). However, research evaluating the runoff from rice production is limited.

The use of riparian forests for filtering nutrient runoff from crop production fields has demonstrated the ability to reduce nutrients entering the surrounding watersheds (Cooper et al., 1986; Jordon et al.,

1993; Lowrance et al., 1984a,b; Peterjohn and Correll, 1984, 1986; Jacobs and Gilliam, 1985; Haycock and Pinay, 1993). The riparian forests take up N, P, and other nutrients, thus reducing the loads in the runoff water. Similarly, wetland ecosystems act as a filter for nutrients contained in water moving through the system, particularly N and P (Boyt et al., 1977; Lakshman, 1979; Reddy and DeBusk, 1985; Tilton and Kadlec, 1979). The mechanisms for filtration include plant uptake, incorporation into the soil, and gaseous losses resulting from the reduced soil conditions associated with wetlands. Gaseous losses of N include nitrification - denitrification reactions (Patrick and Reddy, 1976; Reddy et al., 1980; Reddy and Patrick, 1986) and NH_3 volatilization (Mikkelsen and De Datta, 1979; Brandon and Wells, 1986; Wells and Turner, 1984). It is believed that rice fields can act as a filtration system similar to the constructed wetlands because of the vegetation and flooded soils associated with rice production.

Certain metals contained in irrigation well water, particularly Ca and Mg, are deposited due to the formation of carbonate precipitates when exposed to CO_2 (Gilmour et al., 1978). Due to this precipitation, the concentration in the floodwater usually decreases with increasing distance from the well. The result is the accumulation of calcium and magnesium carbonates in the soil which leads to increased soil pH. Subsequently, the potential for Zn deficiency exists during rice production. Data concerning the movement of other metals across the field is limited.

The majority of the irrigation water utilized in the rice-producing region of the Southern U.S. is used for the rice crop. Consequently, nutrient runoff from rice fields is a major concern in this region.

MATERIALS AND PROCEDURES

Seven production rice fields were selected to monitor the N and P concentrations of the floodwater during the 1990, 1991, and 1992 growing seasons (one field in 1990, three fields in 1991, and three fields in 1992). The fields were located in Ashley, Drew, Lincoln, and St. Francis counties in Arkansas. Each field was managed by the producer, including cultivar selection. The rice was seeded on each field and grown as an upland crop for 4-6 wk. At the four- to five-leaf growth stage, 60% of the recommended fertilizer N was applied as urea and a permanent flood was established and maintained until maturity. Fertilizer N was applied to each field as urea at rates appropriate for the particular cultivar. The N was applied in three applications with 60% of the N applied just prior to flooding, 20% applied at 1.3-cm internode elongation (IE), and 20% 10 d following IE. The fields are designated in the discussion according to the producer's name and year.

Water samples from each field were taken weekly for the entire season at seven locations across the field from entry point to exit including tailwater. Samples were collected daily for the first wk following N fertilizer applications. Water samples were collected with a 60-ml syringe and filtered on location through a 0.45-um membrane filter. The electrical conductivity (EC), pH, and temperature of the floodwater was measured *in situ* at the time of sampling with a portable EC meter and a portable pH meter. The metal samples were acidified with one drop of concentrated HNO₃ for preservation until analysis was performed. The water samples were analyzed for soluble P (PO₄⁻³) and inorganic N (NH₄⁺-N, NO₃⁻-N,

and NO_2^- -N) with a Technicon AutoAnalyzer II (Technicon Industrial Systems, 1973a,b, 1976a). Metals (Ca, Mg, K, Na, Fe, Mn, Al, As, Cu, Co, Cd, Cr, Zn, and Pb) were determined by inductively coupled atomic plasma spectroscopy (Soltanpour et al., 1982).

An additional study was conducted during 1991 and 1992 to compare the nutrient content of runoff water from different irrigation water sources. Two fields were either irrigated with well water or with water pumped from a pond used for raising catfish. Samples were collected from the field as described above. In addition to the analyses described for the other fields, unfiltered water samples were collected to analyze for total dissolved P and total dissolved N. Total N and P were determined with a Technicon AutoAnalyzer II (Technicon Industrial Systems, 1976b).

PRINCIPLE FINDINGS AND SIGNIFICANCE

A. Inorganic N

The major peaks in the seasonal N concentrations in the floodwater were following N fertilizer applications (Fig. 1 and 2, respectively). As would be expected this N peak was usually NH_4 -N (Fig. 1). The NO_3 -N remained low throughout the season (less than 1 mg N L^{-1}) except for two locations (Fig. 2). In 1991, the M&J farm site and the Reinhart farm site in 1992 had an initially high NO_3 -N due to the time between N fertilization and application of the flood. The fields required 3 d to be completely flooded. Consequently, the lower end of the field resulted in greater nitrification and, thus, NO_3 -N accumulation ($>2.5 \text{ mg N L}^{-1}$). The NO_3 -N peak declined to less than 0.5 mg N L^{-1} for both of these locations within 10 d

following establishment of the flood. Most of the peaks observed during mid-season had fallen to below 1 mg $\text{NH}_4\text{-N L}^{-1}$ within 3 d. This is consistent with previous research indicating that mid-season N fertilizer applications are utilized by the plant, incorporated into the soil biomass, or lost within 3 d after application (Wilson et al., 1989).

The $\text{NH}_4\text{-N}$ concentration varied with flow distance and location (Fig. 3). The $\text{NH}_4\text{-N}$ concentration at the Summerford location remained below 0.5 mg L^{-1} during 1991 although the $\text{NH}_4\text{-N}$ concentration exiting the field was greater than the $\text{NH}_4\text{-N}$ concentration entering the field at the other two locations in 1991 and the Pine Tree location in 1990. The Bradshaw and Reinhart locations remained relatively constant across the field during 1992. The Fischer location increased across the first 50% of the field from 0.2 to 1.8 mg N L^{-1} but decreased to 1.1 mg $\text{NH}_4\text{-N L}^{-1}$ at the outlet. Much of the variation between locations can be attributed to differences in water management by the individual producers. The producers ability to flood and/or drain the field differed primarily due to differences in soil texture, field size, and well capacity.

The $\text{NO}_3\text{-N}$ concentration in the floodwater with respect to distance from the inflow also varied among locations (Fig. 4). At most of the locations, the $\text{NO}_3\text{-N}$ concentration remained constant at a level below 0.5 mg $\text{NO}_3\text{-N L}^{-1}$. However, at Reinhart in 1992 and M&J in 1991, the nitrate levels changed with distance. At these two locations, $\text{NO}_3\text{-N}$ increased to levels of 1.5 mg $\text{NO}_3\text{-N L}^{-1}$ or more. This change in $\text{NO}_3\text{-N}$ concentration can be attributed to wetting and drying cycles associated with the inability to apply the flood to the entire field rapidly or to maintain the flood permanently once it was established.

The N concentrations reacted much as expected. The concentrations were high immediately after N application but rapidly declined to levels below that which is considered to pose a threat to surrounding watersheds. For most of the season, the N concentrations remained below 1 mg N L^{-1} as previously reported for rice fields (Moore et al., 1981).

It is important to note the variability in the N concentration in the floodwater among the locations. These differences are associated with water management which is crucial for maximum efficiency of the fertilizer N (Wells et al., 1988). If the early N application is made into the floodwater, the amount of $\text{NH}_4\text{-N}$ in the floodwater increases, NH_3 volatilization losses increase (Fillery et al., 1986), and, with an increase in $\text{NH}_4\text{-N}$ or $\text{NO}_3\text{-N}$ in the floodwater, the potential for increased pollution exists (Moore et al., 1993). The optimum N management techniques are to apply the early N onto dry soil just prior to flooding and then establish a permanent flood as rapidly as possible (Brandon and Wells, 1986; Moore et al., 1993; Wells and Turner, 1984). This technique utilizes the floodwater to incorporate the N into the soil (Fillery et al., 1984; Wells and Turner, 1984) which reduces the buildup of N in the floodwater. This method of applying the early N fertilizer has also shown to increase yields and reduce NH_3 volatilization losses (Wells et al., 1988). By utilizing good water management, $\text{NO}_3\text{-N}$ formation is reduced, less N accumulates in the floodwater, and less potential for N in runoff water exists (Fig. 2 and 4). However, when poor water management is used (i.e. M&J farms - 1991, Reinhart farms - 1992) more nitrification occurs which results in greater denitrification losses and potentially greater $\text{NO}_3\text{-N}$ released into the surrounding surface water.

B. Soluble P

The soluble reactive phosphorus (SRP) decreased with increased distance across the field in all fields except for Bradshaw-1992 (Fig. 5). The SRP increased initially at this site but decreased as the water moved across the field (Fig. 5a). The SRP in all of the fields was less than 0.05 mg P L^{-1} at the end of the field, the level established as the minimum level upon which eutrophication of surface water is possible (Sharpley et al., 1992; Sharpley and Smith, 1993). The Summerford site contained the highest initial SRP level at 0.5 mg P L^{-1} . However, this level dropped to approximately 0.15 mg P L^{-1} and remained relatively constant through the field. Although the level at this site was above the limits recognized as stimulating eutrophication of surface waters (10 ug L^{-1}) (Sharpley and Smith, 1993), the irrigation water contained less SRP at the outlet than the water entering the field from the well. Consequently, the water quality improved as a result of moving across the field.

For most locations, the SRP concentration in the well water was above that which is considered to cause eutrophication of surface water (Fig. 5). However, as the water progressed across the field, the SRP concentration declined. For the Summerford location, the SRP declined from 0.5 mg L^{-1} to approximately 0.09 mg L^{-1} within half the distance of the field (Fig. 5a). The SRP measured at Reinhart-1992 declined from 0.15 mg L^{-1} entering the field to as little as 0.02 mg L^{-1} within 50% of the flow distance (Fig. 5b). Two conclusions can be obtained from these data. First, the inherent level of soluble P in irrigation entering rice fields is variable (0.01 to 0.5 mg L^{-1}). Secondly, rice fields can potentially act as a filter for P similar to the constructed wetland ecosystems (Boyt

et al., 1977; Lakshman, 1979; Reddy and DeBusk, 1985).

The seasonal concentration of soluble P in the floodwater was variable (Fig. 6). The peaks and valleys are a reflection of when the field was irrigated, particularly for Summerford farms in 1991. The well at this particular field was inherently high in soluble P as denoted by the P concentration with distance across the field (Fig. 5b). When water was being applied to the field, the soluble P concentration was relatively high. After the field was flooded and the flood stabilized, the soluble P concentration declined. When water was added to the field because of evaporation, transpiration, runoff, and plant utilization, the amount of P in the water increased.

The P concentrations were typically higher at the well than other locations across the field. As the water flowed across the field, the concentration of P generally decreased to less than 0.05 mg P L^{-1} . The pH of the incoming water was normally close to 7 (data not shown), due to a high CO_2 partial pressure. As the water moves across the field, the CO_2 degasses resulting in CaCO_3 precipitation and PO_4^{3-} co-precipitation or calcium phosphate precipitation (Gilmour et al., 1978). Consequently, the P in the floodwater was reduced. Also, P uptake and utilization by algae, rice plants, and any other vegetation could also have contributed to the reduction in P concentrations as the water moved across the field (Sharpley et al., 1992; Sharpley and Smith, 1993). Although the soluble P concentrations were generally in excess of the limits recognized as stimulating accelerated eutrophication of surface waters (0.01 mg P L^{-1}), they were significantly reduced from the concentration upon entering the field.

C. Metals

As expected, the Ca concentrations decreased or remained the same as the floodwater moved across the field (Table 1). The Ca concentrations ranged from 11 to 78 mg L⁻¹ entering the field and ranged from 11 to 54 mg L⁻¹ exiting the field. The summary presented in Table 1 also indicates the Mg was relatively low but did not change appreciably between the inlet and exit points.

As expected the Fe and Mn concentrations were relatively high at some locations (Table 1). Although most locations resulted in a decrease in the concentration as the water moved across the field, some locations actually increased. The Fe concentrations ranged from 10 to 2404 ug L⁻¹ entering the field and ranged from 90 to 410 ug L⁻¹. The Mn ranged from 3 to 195 ug L⁻¹ entering the field and ranged from 8 to 1745 ug L⁻¹ exiting the field.

Due to the changes associated with soil reduction following flooding, Fe and Mn generally become more soluble. Consequently, it is feasible that an increase in the Fe and Mn concentrations would result. If water management for the rice crop is not optimum, it is also possible that the reduction necessary to solubilize an appreciable amount of Fe and Mn may not occur. Subsequently, Fe and Mn may be deposited in the fields as the water moves across the field because of the oxidized soil conditions. At the Summerford site, the well water contained a large amount of Fe (2404 ug L⁻¹) relative to the other locations. Poor water management leads to reduced N efficiency due to wetting and drying cycles. The wetting and drying cycles at this site resulted in, in addition to the poor N efficiency (Fig. 1), a significant amount of Fe precipitation due

to the oxidized soil conditions.

Table 1. Comparison of inlet and outlet metal concentrations in the floodwater of four production rice fields.

Metal	Bradshaw		Reinhart		Fischer		Summerford	
	In	Out	In	Out	In	Out	In	Out
	mg L ⁻¹							
Ca	41	36	11	11	78	54	45	24
K	1	2	1	2	5	8	2	5
Mg	8	7	2	2	17	16	10	11
Na	13	12	11	10	29	46	28	28
S	3	3	1	1	*	*	*	*
	ug L ⁻¹							
Al	0	63	62	255	0	0	0	73
As	0	0	137	108	0	0	0	0
B	26	29	11	11	94	63	69	76
Cd	40	3	0	0	0	0	0	0
Co	9	6	0	0	0	0	0	0
Cr	0	0	0	0	0	0	0	0
Cu	0	0	0	0	0	0	0	5
Fe	150	90	716	324	10	410	2404	229
Mn	183	31	71	8	3	1745	195	106
Ni	0	0	0	0	0	0	0	0
Pb	*	*	*	*	0	0	0	0
Zn	168	128	289	330	84	187	118	101
pH	6.75	7.69	6.34	8.09	7.15	6.96	6.65	7.43

* = not determined

D. Comparison of Water Sources

Water from ponds is used to raise catfish is used in some areas of the rice producing region of the Southern U.S. In fact, rice is also grown in rotation with catfish on some farms. When the nutrient load of water from catfish ponds was compared to irrigation from well water, the total N concentration was very similar between the two sources (Fig. 7). The $\text{NO}_3\text{-N}$ concentration was very small for both sources. The $\text{NH}_4\text{-N}$ concentration was higher in the pondwater field than in the well water field. Thus, more of the N in the well water field consisted of organic N. However, both sites had less than 3 mg N L^{-1} of total N exiting the field in the runoff with less than 0.5 mg N L^{-1} difference between the two sites.

The total P content declined from as much as 0.1 mg P L^{-1} to slightly more than 0.04 mg P L^{-1} as the water moved across the field from the catfish pond (Fig. 8). However, the well water exiting the field contained less than 0.04 mg P L^{-1} . More of the P in the catfish pond was in the organic form (total - inorganic) than in the well water. However, as the water moved across the field, the concentration of the water exiting the field was very similar for the two locations with about 50 % of the total P in the $\text{PO}_4\text{-P}$ form.

This data confirms the data from the other locations concerning the decline in P and N concentration as rice irrigation water flows across the field. The total P and organic P in all of the fields was less than 0.05 mg P L^{-1} at the end of the field but is higher than the critical limits (0.01 and 0.02 mg P L^{-1} of soluble and total P, respectively) established as contributing to accelerated eutrophication of surface water (Sharpley

et al., 1992; Sharpley and Smith, 1993). The water exiting the field irrigated by pondwater contained slightly higher concentrations of organic P than the field irrigated by well water, indicating that the contribution to surface water eutrophication is more likely used catfish pondwater for irrigation water. However, the amount of P in the water declined significantly by flowing across the rice field. Had the water been released directly from the catfish ponds, the level would have exceeded four times the critical limit for total P. Consequently, the idea was confirmed that rice fields can act similarly to constructed wetlands (Lakshman, 1979) and filter nutrients before entering surface water.

CONCLUSIONS

Since rice production utilizes a major portion of the irrigation water in the Southern Rice Belt, it is necessary to understand the potential for pollution resulting from field runoff. The results of this study emphasize the importance of water management with respect to N concentrations in the floodwater. The $\text{NO}_3\text{-N}$ levels remained low when optimum water management was utilized. The soluble P concentrations were generally lower in the water exiting the field than in the water entering the field from the well. Although the P levels were sufficient at some locations to potentially contribute to eutrophication of surrounding surface waters, the amount of P was generally higher entering the field than exiting the field. Removal of Fe and Mn apparently depends upon water management and the oxidation state of the soil.

LITERATURE CITED

- Boyt, F.L., S.E. Bayley, and J. Zoltek. 1977. Removal of nutrients from treatment municipal wastewater by wetland vegetation. *J. Water Pollut. Control Fed.* 48:789-799.
- Brandon, D.M., and B.R. Wells. 1986. Improving nitrogen fertilizer in mechanized rice culture. P. 161-170. *In* S.K. De Datta and W.H. Patrick, Jr. (ed.) *Nitrogen economy of flooded rice soils*. Martinus Nijhoff Publishing Co., Dordrecht, The Netherlands.
- Chichester, F.W., and C.W. Richardson. 1992. Sediment and nutrient loss from clay soils as affected by tillage. *J. Envir. Qual.* 21:587-590.
- Cooper, J.R., J.W. Gilliam, and T.C. Jacobs. 1986. Riparian areas as a control of nonpoint pollutants. p. 166-192. *In* D.L. Correll (ed.) *Watershed research perspectives*. Smithsonian Institution Press, Washington, D.C.
- Fillery, I.R.P., P.A. Roger, and S.K. De Datta. 1986. Ammonia volatilization from nitrogen sources applied to rice fields: II. Floodwater properties and submerged photosynthetic biomass. *Soil Sci. Soc. Am. J.* 50:86-91.
- Fillery, I.R.P., J.R. Simpson, and S.K. De Datta. 1984. Influence of field environment and fertilizer management on ammonia loss from flooded rice. *Soil Sci. Soc. Am. J.* 48:914-920.
- Gilmour, J.R., K.S. Shirk, J.A. Ferguson, and C.L. Griffis. 1978. A kinetic study of the CaCO_3 precipitation reaction. *Agric. Water Manage.* 1:253-262.
- Haycock, N.E., and G. Pinay. 1993. Groundwater nitrate dynamics in grass and poplar vegetated riparian buffer strips during the winter. *J. Environ. Qual.* 22:273-278.
- Jacobs, T.C., and J.W. Gilliam. 1985. Riparian losses of nitrate from agricultural drainage waters. *J. Environ. Qual.* 14:472-478.
- Jordon, T.E., D.L. Correll, and D.E. Weller. 1993. Nutrient interception by a riparian forest receiving inputs from adjacent cropland. *J. Environ. Qual.* 22:467-473.
- Lakshman, B. 1979. An ecosystem approach to the treatment of wastewaters. *J. Environ. Qual.* 8:353-361.
- Lin, S. 1972. Nonpoint rural sources of water pollution. Dep. Registration and Educ. Circ. III. p. 1-36. State Water Survey, Urbana, Ill.

- Loehr, R.C. 1974. Characteristics and comparative magnitude of nonpoint sources. *J. Water Pollut. Control Fed.* 45:1849-1872.
- Lowrance, R.R. 1992. Nitrogen outputs from a field-size agricultural watershed. *J. Environ. Qual.* 21:602-607.
- Lowrance, R.R., R.L. Todd, and L.E. Asmussen. 1984a. Nutrient cycling in an agricultural watershed: I. Phreatic movement. *J. Environ. Qual.* 13:22-27.
- Lowrance, R.R., R.L. Todd, J. Fail, Jr., O. Hendrickson, Jr., R. Leonard, and L. Asmussen. 1984b. Riparian forests as nutrient filters in agricultural watersheds. *BioScience* 34:374-377.
- Mikkelsen, D.S., and S.K. De Datta. 1979. Ammonia volatilization from wetland rice soils. p. 135-156. *In* Nitrogen in rice. IRRI, Los Banos, Laguana, Philippines.
- Moore, P.A., Jr., J.T. Gilmour, and B.R. Wells. 1981. Seasonal patterns of growth and soil nitrogen uptake by rice. *Soil Sci. Soc. Am. J.* 45:875-879.
- Moore, P.A., Jr., K.K. Baugh, R.J. Norman, B.R. Wells, and R.S. Helms. 1993. Nutrient runoff from rice fields. p. 142-149. *In* B.R. Wells (ed.) Arkansas Rice Research Studies 1992. Ark. Agri. Exp. Stn. Res. Ser. 431.
- Patrick, W.H., Jr., and K.R. Reddy. 1976. Fate of fertilizer nitrogen in a flooded rice soil. *Soil Sci. Soc. Am. J.* 40:678-681.
- Peterjohn, W.T., and D.L. Correll. 1984. Nutrient dynamics in an agricultural watershed: Observations on the role of a riparian forest. *Ecology* 65:1466-1475.
- Peterjohn, W.T., and D.L. Correll. 1986. The effect of riparian forest on the volume and chemical composition of baseflow in an agricultural watershed. p. 244-262. *In* D.L. Correll (ed.) Watershed research perspectives. Smithsonian Institution Press, Washington, D.C.
- Reddy, K.R., and W.F. DeBusk. 1985. Nutrient removal potential of selected aquatic macrophytes. *J. Environ. Qual.* 14:459-462.
- Reddy, K.R., and W.H. Patrick, Jr. 1986. Denitrification losses in flooded rice fields. p. 99-116. *In* S.K. De Datta and W.H. Patrick, Jr. (ed.) Nitrogen economy of flooded rice soils. Martinus Nijhoff Publishing Co., Dordrecht, The Netherlands.
- Reddy, K.R., W.H. Patrick, Jr., and R.E. Phillips. 1980. Evaluation of selected processes controlling nitrogen loss in a flooded soil. *Soil Sci. Soc. Am. J.* 44:1241-1246.

- Sharpley, A.N., and S.J. Smith. 1993. Prediction of bioavailable phosphorus loss in agricultural runoff. *J. Environ. Qual.* 22:32-37.
- Sharpley, A.N., S.J. Smith, O.R. Jones, W.A. Berg, and G.A. Coleman. 1992. The transport of bioavailable phosphorus in agricultural runoff. *J. Environ. Qual.* 21:30-35.
- Soltanpour, P.N., J.B. Jones, Jr., and S.M. Workman. 1982. Optical emission spectrometry. *In* A.L. Page et al. (ed.) *Methods of soil analysis. Part 2.* 2nd ed. Agronomy 9:29-67.
- Stanford, G., C.B. England, and A.W. Taylor. 1970. Fertilizer use and water quality. ARS publ. 41-168. USDA, Washington, D.C.
- Technicon Industrial Systems. 1973a. Ammonia in water and waste water. Industrial method no. 98-70w. Technicon Instruments Corp., Tarrytown, NY.
- Technicon Industrial Systems. 1973b. Nitrate and nitrite in water and waste water. Industrial method no. 100-70w. Technicon Instruments Corp., Tarrytown, NY.
- Technicon Industrial Systems. 1976a. Ortho-phosphate in water and waste water. Industrial method no. 94-70w/b. Technicon Instruments Corp., Tarrytown, NY.
- Technicon Industrial Systems. 1976b. Individual/simultaneous determination of nitrogen and/or phosphorus in BD acid digests. Industrial method no. 344-74a. Technicon Instruments Corp., Tarrytown, NY.
- Tilton, D.L., and R.H. Kadlec. 1979. The utilization of a freshwater wetland for nutrient removal from secondarily treated wastewater effluent. *J. Environ. Qual.* 8:328-334.
- Water Resources Committee. 1970. Soil science in relation to water resources development: IV. Responsibility of soil science in water quality improvement. *Soil Sci. Soc. Am. Proc.* 34:542-548.
- Wells, B.R., R.J. Norman, and R.S. Helms. 1988. Rice grain and dry matter response to nitrogen and water management. *Ark. Farm Res.* 37(2):13.
- Wells, B.R., and F.T. Turner. 1984. Nitrogen use in flooded rice soils. p. 349-362. *In* R.D. Hauck (ed.) *Nitrogen in crop production.* ASA, CSSA, and SSSA, Madison, WI.
- Wilson, Jr., C.E., R.J. Norman, and B.R. Wells. 1989. Seasonal uptake patterns of fertilizer nitrogen applied in split applications to rice. *Soil Sci. Soc. Am. J.* 53:1884-1887.




Fig. 1. Seasonal changes in $\text{NH}_4\text{-N}$ concentration in rice irrigation water during 1991 (A) and 1992 (B).




Fig. 2. Seasonal changes in NO₃-N concentration in rice irrigation water during 1991 (A) and 1992 (B).





Fig. 3. Changes in $\text{NH}_4\text{-N}$ concentration in rice irrigation water with respect to relative distance from the well during 1991 (A) and 1992 (B).





Fig. 4. Changes in $\text{NO}_3\text{-N}$ concentration in rice irrigation water with respect to relative distance from the well during 1991 (A) and 1992 (B).




Fig. 5. Changes in soluble reactive phosphorus ($PO_4\text{-P}$) concentration in rice irrigation water with respect to distance across the field during 1991 (A) and 1992 (B).




Fig. 6. Seasonal changes in soluble reactive phosphorus ($PO_4\text{-P}$) concentration in rice irrigation water during 1991 (A) and 1992 (B).




Fig. 7. Concentrations of NH₄-N, NO₃-N, and total N in rice irrigation water from a well (A) or catfish pond (B) with respect to relative distance from the source.




Fig. 8. Concentrations of soluble reactive phosphorus (PO₄-P) and total P in rice irrigation water from a well (A) or catfish pond (B) with respect to relative distance from the source.