



Strathprints Institutional Repository

Langer, Matthias and Woracek, Harald (2014) Stability of the derivative of a canonical product. Complex Analysis and Operator Theory, 8 (6). pp. 1183-1224. ISSN 1661-8254 , <http://dx.doi.org/10.1007/s11785-013-0315-5>

This version is available at <http://strathprints.strath.ac.uk/44380/>

Strathprints is designed to allow users to access the research output of the University of Strathclyde. Unless otherwise explicitly stated on the manuscript, Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Please check the manuscript for details of any other licences that may have been applied. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<http://strathprints.strath.ac.uk/>) and the content of this paper for research or private study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to Strathprints administrator: strathprints@strath.ac.uk

Stability of the derivative of a canonical product

MATTHIAS LANGER, HARALD WORACEK

Abstract

With each sequence $\alpha = (\alpha_n)_{n \in \mathbb{N}}$ of pairwise distinct and non-zero points which are such that the canonical product

$$P_\alpha(z) := \lim_{r \rightarrow \infty} \prod_{|\alpha_n| \leq r} \left(1 - \frac{z}{\alpha_n}\right)$$

converges, the sequence

$$\alpha' := (P'_\alpha(\alpha_n))_{n \in \mathbb{N}}$$

is associated. We give conditions on the difference $\beta - \alpha$ of two sequences which ensure that β' and α' are comparable in the sense that

$$\exists c, C > 0 : c|\alpha'_n| \leq |\beta'_n| \leq C|\alpha'_n|, \quad n \in \mathbb{N}.$$

The values α'_n play an important role in various contexts. As a selection of applications we present: an inverse spectral problem, a class of entire functions and a continuation problem.

AMS MSC 2010: Primary 30D20; Secondary 30D15, 42A82, 34B20

Keywords: Canonical product, perturbation of zeros, inverse spectral problem, Kreĭn class, positive definite function

1 Introduction

Our aim in the present paper is to show a stability result for the derivative of a canonical product evaluated at its zeros when the zeros of the product are perturbed. To illustrate this task, let us consider a toy example.

1.1 Example. Let f be the entire function $f(z) := \frac{\sin(\pi z)}{\pi z}$. The sequence of zeros of f is $+1, -1, +2, -2, \dots$, which we denote by $\alpha = (\alpha_n)_{n \in \mathbb{N}}$. Then f is given by the canonical product

$$f(z) = \lim_{r \rightarrow \infty} \prod_{|\alpha_n| \leq r} \left(1 - \frac{z}{\alpha_n}\right). \quad (1.1)$$

Let $\beta = (\beta_n)_{n \in \mathbb{N}}$ be a small perturbation of α , say

$$\beta_n - \alpha_n = O\left(\frac{1}{n^\varepsilon}\right), \quad n \rightarrow \infty, \quad (1.2)$$

for some $\varepsilon > 0$, and consider the corresponding canonical product (we tacitly assume that the numbers β_n are pairwise distinct and non-zero):

$$g(z) := \lim_{r \rightarrow \infty} \prod_{|\beta_n| \leq r} \left(1 - \frac{z}{\beta_n}\right). \quad (1.3)$$

Then g is an entire function of finite exponential type. The function $zg(z)$ is of sine type¹, and hence the sequence $(|\beta_n g'(\beta_n)|)_{n \in \mathbb{N}}$ is bounded from above and away from zero. The same is of course true for $(|\alpha_n g'(\alpha_n)|)_{n \in \mathbb{N}}$. Since $\lim_{n \rightarrow \infty} \frac{\beta_n}{\alpha_n} = 1$, we may say that

$$|g'(\beta_n)| \asymp |f'(\alpha_n)|; \quad (1.4)$$

here and in the following we write $x_n \lesssim y_n$ if there exists a constant $c > 0$ such that $x_n \leq cy_n$ for all $n \in \mathbb{N}$, and we write $x_n \asymp y_n$ if $x_n \lesssim y_n$ and $y_n \lesssim x_n$. Note that removing or adding a zero changes the asymptotics of $f'(\alpha_n)$; so in some sense, the behaviour of $f'(\alpha_n)$ depends sensitively on α_n . //

Now consider two arbitrary sequences α and β , only assuming that the products (1.1) and (1.3) converge. *What conditions on the perturbation $\beta - \alpha$ ensure that (1.4) holds?*

In our main result, Theorem 3.3, we give conditions which guarantee that (1.4) holds. They limit the size of the perturbation in two respects: (1) relative to the location of α_n and (2) relative to the regularity of the distribution of α measured by the separation

$$s_\alpha(n) := \min\{|\alpha_k - \alpha_n| : k \neq n\}$$

and by some sort of local density (see r_α in Definition 2.6 below). The rough picture is (as one would expect): if the sequence α is sparse and well-separated, large perturbations are allowed.

In Theorem 3.3 we do not assume any regularity of the sequence α ². For sequences that are regularly distributed the conditions of Theorem 3.3 can be weakened significantly. We make this precise in the two supplementary results Theorems 3.18 and 3.19; see also Corollary 3.20³.

The conditions in our theorems appear to be more or less sharp, cf. Remark 3.5. In order to construct examples to explicitly show this, one would have to deal with very irregularly distributed sequences, cf. Remark 3.15. To do explicit computations in such cases seems to be out of reach.

Our motivation to consider the question for stability of $|f'(\alpha_n)|$ is that these quantities frequently appear in complex analysis and spectral theory. They have significance in a multitude of classical problems, e.g. weighted approximation problems like the Bernstein problem, power or trigonometric moment problems, extension theory of symmetric operators, spectral theory of canonical systems, etc. In such situations often the convergence of series of the form

$$\sum_{n \in \mathbb{N}} \frac{c_n}{|f'(\alpha_n)|^\rho}$$

¹For definition and theory of sine-type functions we refer to [L2, Lecture 22].

²Besides some weak condition in the extremal case that the product (1.1) has positive exponential type, cf. Remark 3.4.

³What we have observed in the above example is reobtained. In fact, Corollary 3.20 shows that one may even allow larger perturbations than in (1.2) (although then the function $zg(z)$ may not be of sine type anymore). See Example 3.22 for more details.

with some $c_n > 0$ and $\rho > 0$ plays a role. Hence we can invoke our present results and deduce stability under small perturbations. To illustrate this vague principle, we discuss in the present paper three applications representing typical and descriptive instances:

- spectral functions of regular strings (Theorem 4.2);
- the Kreĭn class of entire functions (Theorem 4.4);
- continuations of a positive definite function on an interval (Theorem 4.8).

All those are classical topics. Only in the last example, we make a little excursion to the indefinite (Pontryagin space) world⁴.

Another interesting (and more serious) application of Theorem 3.3 is found in the theory of the Hamburger power moment problem. There, stability of N-extremality and (in conjunction with Pontryagin space methods), stability of the index of determinacy of a measure can be studied. These results will be presented as part of the forthcoming work [LW2], [LW3].

The proof of Theorem 3.3 and its supplements is carried out by accurately estimating products and, for the case of regularly distributed sequences, using some facts about growth functions. We employ the standard procedure to split a canonical product into several parts and estimate each of them. However, it is not just “business as usual”: the splitting of the product has to be done depending on a parameter $n \in \mathbb{N}$ in a somewhat tricky way, and estimates are required to be uniform with respect to n , cf. Remark 3.17.

The structure of the present paper is straightforward. First, in Section 2, we provide some preliminary facts about growth functions. In Section 3 we formulate and prove the main Theorem 3.3 and the supplements dealing with regularly distributed sequences. Finally, in Section 4 we present the aforementioned applications.

Acknowledgement. We thank Anton Baranov for discussion on the subject, and for communicating a stability result for well-separated sequences. The challenge to cover his result motivated us to push our method further until it reached the present form.

2 Preliminaries about growth functions

In complex analysis, in particular in the theory of entire functions, the notion of growth plays a central role.

2.1 Definition. A function $\lambda : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is called a *growth function* if it satisfies the following axioms:

(gf1) the limit $\rho_\lambda := \lim_{r \rightarrow \infty} \frac{\log \lambda(r)}{\log r}$ exists and is finite and non-negative;

(gf2) for all sufficiently large values of r , the function λ is differentiable and

$$\lim_{r \rightarrow \infty} \left(r \frac{\lambda'(r)}{\lambda(r)} \Big/ \frac{\log \lambda(r)}{\log r} \right) = 1;$$

moreover, $\lim_{r \rightarrow \infty} \lambda(r) = \infty$.

⁴Within Pontryagin space theory a variety of applications arises. In order not to overload the presentation, we decided not to touch upon these topics further.

//

The conditions (gf1) and (gf2)⁵ ensure that we have Valiron's theory of proximate orders available, cf. [L1, Section I.12] or [LG, Section I.6]. Note that the logarithm of a growth function is often called proximate order.

Typical examples of growth functions are functions of the form

$$\lambda(r) = r^a \cdot (\log_{(m_1)} r)^{b_1} \cdot \dots \cdot (\log_{(m_n)} r)^{b_n} \quad (2.1)$$

for large enough r ; here $a \geq 0$, $m_i \in \mathbb{N}$, $m_1 < \dots < m_n$, $b_1, \dots, b_n \in \mathbb{R}$, with $b_1 > 0$ if $a = 0$, and $\log_{(n)}$ is defined by

$$\log_{(1)} r := \log r, \quad \log_{(k+1)} r := \log(\log_{(k)} r), \quad k \in \mathbb{N},$$

for large enough r . For the function in (2.1) we have $\rho_\lambda = a$.

As for the classical notion of order, also for general growth functions the growth of an entire function is related to the density of its zeros; see, e.g. [Ru, Theorems 13.5.2–4]. Comparison of the growth of an entire function with functions of the form (2.1) goes back as far as to some work of E. Lindelöf in the early 20th century. However, in the present context we use growth functions only to handle the distribution of sequences.

2.2 Remark. Let λ be a growth function. Then the following statements are true.

(i) $\lim_{r \rightarrow \infty} \frac{\lambda(Cr)}{\lambda(r)} = C^{\rho_\lambda}$ uniformly in C on compact subsets of $(0, \infty)$.

(ii) For sufficiently large values of r the function λ is strictly increasing.

(iii) Let $\sigma > 0$; then, for sufficiently large r , the function $\frac{\lambda(r)}{r^\sigma}$ is increasing if $\sigma < \rho_\lambda$ and decreasing if $\sigma > \rho_\lambda$.

A proof of (i) can be found in [L1, Lemma 5 in I.12] and [LG, Theorem 1.18, Proposition 1.19]. Item (ii) is a direct consequence of (gf2). Finally, (iii) follows from the relation

$$\left[\frac{\lambda(r)}{r^\sigma} \right]' = \frac{\lambda(r)}{r^{\sigma+1}} \left(-\sigma + \frac{r\lambda'(r)}{\lambda(r)} \right)$$

and the fact that $\lim_{r \rightarrow \infty} \frac{r\lambda'(r)}{\lambda(r)} = \rho_\lambda$ by (gf2). //

Property (i) in Remark 2.2 says precisely that a growth function is a regularly varying function as defined, e.g. in [S, Definition 1.1]. However, we need the more special concept of a growth function since we use the properties (ii) and (iii) and the fact that $\lambda(r) \rightarrow \infty$, $r \rightarrow \infty$; regularly varying functions do not have the latter properties in general.

The whole importance of a growth function lies in its behaviour at $+\infty$. It is thus no loss of generality to assume additionally that

(gf3) the function λ is differentiable, strictly increasing and bounded away from 0.

⁵Instead of (gf2) often the condition $\lim_{r \rightarrow \infty} \frac{r\lambda'(r)}{\lambda(r)} = \rho_\lambda$ is required. If $\rho_\lambda > 0$, then, clearly, this is equivalent to (gf2). If $\rho_\lambda = 0$, (gf2) is stronger.

From now on we always include this property in the notion of a growth function.

2.3 Remark. In Subsection 2.2 we define sequences using the inverse of a growth function. Let us therefore state that a function $\mu : (a, \infty) \rightarrow (0, \infty)$ is the inverse of a growth function (satisfying (gf1)–(gf3)) if and only if

- $a > 0$; μ is surjective and differentiable with positive derivative;
- $\rho_\lambda := \lim_{s \rightarrow \infty} \frac{\log s}{\log \mu(s)}$ exists, is finite and non-negative;
- $\lim_{s \rightarrow \infty} \left(\frac{\mu(s)}{s \mu'(s)} \Big/ \frac{\log s}{\log \mu(s)} \right) = 1$.

//

2.1 Upper and lower densities

For a sequence ξ of real numbers and a growth function λ , several densities are defined.

2.4 Definition. For a sequence $\xi = (\xi_n)_{n \in \mathbb{N}}$ of real numbers, set

$$\begin{aligned} n_\xi^+(r) &:= \#\{n \in \mathbb{N} : 0 < \xi_n \leq r\}, \\ n_\xi^-(r) &:= \#\{n \in \mathbb{N} : -r \leq \xi_n < 0\}, \\ n_\xi(r) &:= \#\{n \in \mathbb{N} : |\xi_n| \leq r\}. \end{aligned}$$

The *upper* and *lower right* λ -densities of ξ are defined as

$$\Delta_\lambda^+(\xi) := \limsup_{r \rightarrow \infty} \frac{n_\xi^+(r)}{\lambda(r)}, \quad \delta_\lambda^+(\xi) := \liminf_{r \rightarrow \infty} \frac{n_\xi^+(r)}{\lambda(r)}.$$

Similarly, the *upper* and *lower left* λ -densities are

$$\Delta_\lambda^-(\xi) := \limsup_{r \rightarrow \infty} \frac{n_\xi^-(r)}{\lambda(r)}, \quad \delta_\lambda^-(\xi) := \liminf_{r \rightarrow \infty} \frac{n_\xi^-(r)}{\lambda(r)},$$

and the *upper* and *lower* λ -densities are

$$\Delta_\lambda(\xi) := \limsup_{r \rightarrow \infty} \frac{n_\xi(r)}{\lambda(r)}, \quad \delta_\lambda(\xi) := \liminf_{r \rightarrow \infty} \frac{n_\xi(r)}{\lambda(r)}.$$

//

The following facts are elementary and are proved in the same way as [Bo, Lemma 1.5.1], which is nothing but the case when $\lambda(r) = r$. We skip the details.

2.5 Lemma. *Let λ be a growth function and let ξ be a sequence of real numbers. Denote by ξ^+ and ξ^- the (finite or infinite) subsequences of ξ consisting of the positive or negative, respectively, elements of ξ arranged according to increasing modulus and indexed with $n = 1, 2, \dots$. Then⁶*

$$\begin{aligned} \delta_\lambda^+(\xi) &= \liminf_{n \rightarrow \infty} \frac{n}{\lambda(\xi_n^+)}, & \Delta_\lambda^+(\xi) &= \limsup_{n \rightarrow \infty} \frac{n}{\lambda(\xi_n^+)}, \\ \delta_\lambda^-(\xi) &= \liminf_{n \rightarrow \infty} \frac{n}{\lambda(|\xi_n^-|)}, & \Delta_\lambda^-(\xi) &= \limsup_{n \rightarrow \infty} \frac{n}{\lambda(|\xi_n^-|)}. \end{aligned}$$

⁶We tacitly understand the limit of a finite sequence as 0.

For a sequence ξ we introduce two measures for the regularity of its distribution, the below defined values $s_\xi(n)$ and $r_\xi(\rho, n)$. The first is just the separation of the sequence, the second measures whether large lumps of points appear in ξ .

2.6 Definition. Let $\xi = (\xi_n)_{n \in \mathbb{N}}$ be a sequence of real non-zero numbers and let $\rho > 1$. Then we define

$$s_\xi(n) := \inf \{ |\xi_k - \xi_n| : k \in \mathbb{N}, \xi_k \neq \xi_n \}, \quad n \in \mathbb{N},$$

$$r_\xi(\rho, n) := \# \left\{ k \in \mathbb{N} : \frac{\xi_k}{\xi_n} \in \left(\frac{1}{\rho}, \rho \right) \right\}, \quad n \in \mathbb{N}.$$

//

For each infinite sequence ξ of real numbers that has no finite accumulation point, there exists a growth function λ with $0 < \Delta_\lambda(\xi) < \infty$, cf. [L1, Theorem 16 in I.12]. However, it need not be possible to choose λ such that also $\delta_\lambda(\xi) > 0$. This is related to the possible existence of large clusters of points in ξ . A quantitative statement is the following lemma.

2.7 Lemma. Let $\xi = (\xi_n)_{n \in \mathbb{N}}$ be an increasing sequence of positive real numbers that has no finite accumulation point. Let λ be a growth function and assume that ξ has finite upper and positive lower λ -densities. Then, for each $\rho > 1$,

$$r_\xi(\rho, n) = O(n) \quad \text{and} \quad \sum_{\substack{k \in \mathbb{N} \\ \frac{\xi_k}{\xi_n} \in (\frac{1}{\rho}, \rho)}} \frac{1}{k} = O(1), \quad n \rightarrow \infty. \quad (2.2)$$

Proof. We set

$$d_1 := \inf_{r \geq \xi_1} \frac{n_\xi(r)}{\lambda(r)}, \quad d_2 := \sup_{r \geq \xi_1} \frac{n_\xi(r)}{\lambda(r)}, \quad c_\rho := \sup_{r \geq \xi_1} \frac{\lambda(\rho r)}{\lambda(r)}. \quad (2.3)$$

Our assumption implies that $d_1 > 0$ and $d_2 < \infty$, and Remark 2.2 (i) and (gf3) imply that $1 \leq c_\rho < \infty$. Thus we can estimate

$$r_\xi(\rho, n) \leq n_\xi(\rho \xi_n) \leq d_2 \lambda(\rho \xi_n) \leq d_2 c_\rho \lambda(\xi_n) \leq \frac{d_2 c_\rho}{d_1} n_\xi(\xi_n) = \frac{d_2 c_\rho}{d_1} \cdot n,$$

which shows the first relation in (2.2). Next, set

$$k_+(n) := \max \{ k \in \mathbb{N} : \xi_k \leq \rho \xi_n \}, \quad k_-(n) := \min \{ k \in \mathbb{N} : \frac{1}{\rho} \xi_n < \xi_k \}. \quad (2.4)$$

Then

$$k_+(n) = n_\xi(\rho \xi_n), \quad k_-(n) = n_\xi\left(\frac{1}{\rho} \xi_n\right) + 1.$$

Moreover, let $n_0 \in \mathbb{N}$ be such that $\frac{1}{\rho} \xi_{n_0} \geq \xi_1$. Then $k_-(n) \geq 2$ if $n \geq n_0$. For such n we estimate

$$\frac{k_+(n)}{k_-(n) - 1} = \frac{n_\xi(\rho \xi_n)}{n_\xi\left(\frac{\xi_n}{\rho}\right)} \leq \frac{d_2 \lambda(\rho \xi_n)}{d_1 \lambda\left(\frac{\xi_n}{\rho}\right)} \leq \frac{d_2 c_\rho \lambda(\xi_n)}{d_1 \frac{1}{c_\rho} \lambda(\xi_n)} = \frac{d_2 c_\rho^2}{d_1}.$$

Clearly, $\lim_{n \rightarrow \infty} k_+(n) = \lim_{n \rightarrow \infty} k_-(n) = \infty$, and hence (here γ denotes the Euler–Mascheroni constant)

$$\begin{aligned} \sum_{k \in \mathbb{N}: \frac{1}{\rho} \xi_n < \xi_k \leq \rho \xi_n} \frac{1}{k} &= \sum_{k=k_-(n)}^{k_+(n)} \frac{1}{k} \\ &= [\log k_+(n) + \gamma + o(1)] - [\log(k_-(n) - 1) + \gamma + o(1)] \\ &= \log \frac{k_+(n)}{k_-(n) - 1} + o(1) = O(1) \end{aligned}$$

as $n \rightarrow \infty$. □

2.8 Definition. We call a sequence ξ of *convergence class with respect to a growth function* λ if

$$\sum_{n \in \mathbb{N}} \frac{1}{\lambda(|\xi_n|)} < \infty.$$

//

For example, the sequence $\xi_n := n^{\frac{1}{\rho}}$ where $\rho > 0$ is of convergence class with respect to the growth function λ that satisfies $\lambda(r) := r^\rho \log r (\log \log r)^2$ for large r , but it is not of convergence class with respect to $\lambda(r) := r^\rho$.

The next statement is an analogue of the classical case when $\lambda(r) = r^\rho$ and is proved in the same way, cf. [Bo, Proof of Lemma 2.5.5]. We again skip the details.

2.9 Lemma. *Assume that ξ is of convergence class with respect to the growth function λ . Then*

$$n_\xi(r) = O(\lambda(r)), \quad r \rightarrow \infty.$$

The converse of this fact is not true (as already seen from the above mentioned example).

2.2 The standard sequence $\mathring{\lambda}$

With a growth function we associate a sequence that has most regular behaviour with respect to λ .

2.10 Definition. Let λ be a growth function. Then we define the *standard sequence* $\mathring{\lambda} = (\mathring{\lambda}_n)_{n \in \mathbb{N}}$ associated with λ by

$$\mathring{\lambda}_n := \lambda^{-1}(n), \quad n \in \mathbb{N}.$$

//

First, we collect some simple properties of this sequence. Recall that the *convergence exponent* of a sequence ξ of non-zero numbers is defined as

$$\inf \left\{ \rho > 0 : \sum_{n=1}^{\infty} \frac{1}{|\xi_n|^\rho} < \infty \right\}.$$

2.11 Lemma. *Let λ be a growth function, and let $\overset{\circ}{\lambda}$ be the standard sequence associated with λ . Moreover, let ρ_λ be as in (gf1). Then the following statements hold.*

- (i) *The sequence $\overset{\circ}{\lambda}$ is strictly increasing.*
- (ii) *We have $n_{\overset{\circ}{\lambda}}(r) = \lfloor \lambda(r) \rfloor$; here $\lfloor x \rfloor$ denotes the largest integer less than or equal to x .*
- (iii) *We have $\Delta_\lambda(\overset{\circ}{\lambda}) = \delta_\lambda(\overset{\circ}{\lambda}) = 1$.*
- (iv) *The convergence exponent of $\overset{\circ}{\lambda}$ is equal to ρ_λ .*

Proof. Item (i) is clear from our additional axiom (gf3). For item (ii), let $n \in \mathbb{N}$. Since $\overset{\circ}{\lambda}_n = \lambda^{-1}(n)$, we have $\overset{\circ}{\lambda}_n \leq r$ if and only if $n \leq \lambda(r)$. This shows that $n_{\overset{\circ}{\lambda}}(r) = \lfloor \lambda(r) \rfloor$. Item (iii) is obvious from (ii), and item (iv) follows since the convergence exponent of $\overset{\circ}{\lambda}$ can be computed as $\rho_1 = \limsup_{r \rightarrow \infty} \frac{\log n_{\overset{\circ}{\lambda}}(r)}{\log r}$, see, e.g. [Bo, Theorem 2.5.8]. \square

For standard sequences we can control $s_{\overset{\circ}{\lambda}}$ and $r_{\overset{\circ}{\lambda}}$. These estimates are used in the proof of Theorem 3.19.

2.12 Lemma. *Let λ be a growth function, and let $\rho > 1$. Then*

$$\frac{1}{s_{\overset{\circ}{\lambda}}(n)} = O\left(\frac{n}{\overset{\circ}{\lambda}_n}\right), \quad \sum_{\substack{k \in \mathbb{N}: k \neq n \\ \frac{\overset{\circ}{\lambda}_k}{\overset{\circ}{\lambda}_n} \in (\frac{1}{\rho}, \rho)}} \frac{1}{|\overset{\circ}{\lambda}_k - \overset{\circ}{\lambda}_n|} = O\left(\frac{n \log n}{\overset{\circ}{\lambda}_n}\right), \quad n \rightarrow \infty. \quad (2.5)$$

Proof. Let $k, n \in \mathbb{N}$. We use the mean value theorem to obtain a point $\theta_{k,n}$ between k and n with

$$\overset{\circ}{\lambda}_k - \overset{\circ}{\lambda}_n = \lambda^{-1}(k) - \lambda^{-1}(n) = \frac{k - n}{\lambda'(\lambda^{-1}(\theta_{k,n}))}. \quad (2.6)$$

Due to (gf2) we have $\lambda'(r) \leq C \frac{\lambda(r)}{r}$ with some $C > 0$, and hence

$$\lambda'(\lambda^{-1}(\theta_{k,n})) \leq C \frac{\theta_{k,n}}{\lambda^{-1}(\theta_{k,n})}. \quad (2.7)$$

For $k = n + 1$ this yields

$$\begin{aligned} \frac{1}{|\overset{\circ}{\lambda}_{n+1} - \overset{\circ}{\lambda}_n|} &= \lambda'(\lambda^{-1}(\theta_{n+1,n})) \leq C \frac{\theta_{n+1,n}}{\lambda^{-1}(\theta_{n+1,n})} \\ &\leq C \frac{n+1}{\lambda^{-1}(n)} = C \frac{n+1}{\overset{\circ}{\lambda}_n}. \end{aligned} \quad (2.8)$$

If $\overset{\circ}{\lambda}_{n-1} \geq \frac{1}{\rho} \overset{\circ}{\lambda}_n$, then (2.6) and (2.7) with $k = n - 1$ yield

$$\frac{1}{|\overset{\circ}{\lambda}_n - \overset{\circ}{\lambda}_{n-1}|} \leq C \frac{n}{\overset{\circ}{\lambda}_{n-1}} \leq C \rho \frac{n}{\overset{\circ}{\lambda}_n}. \quad (2.9)$$

If $\overset{\circ}{\lambda}_{n-1} < \frac{1}{\rho} \overset{\circ}{\lambda}_n$, then

$$\frac{1}{|\overset{\circ}{\lambda}_n - \overset{\circ}{\lambda}_{n-1}|} = \frac{1}{(1 - \frac{\overset{\circ}{\lambda}_{n-1}}{\overset{\circ}{\lambda}_n}) \overset{\circ}{\lambda}_n} \leq \frac{1}{(1 - \frac{1}{\rho}) \overset{\circ}{\lambda}_n}.$$

This together with (2.8) and (2.9) shows the first relation in (2.5).

Due to Lemma 2.11 (iii) we can apply Lemma 2.7 and obtain $r_{\overset{\circ}{\lambda}}(\rho, n) = O(n)$. Assume that $\frac{\overset{\circ}{\lambda}_k}{\overset{\circ}{\lambda}_n} \in (\frac{1}{\rho}, \rho)$. With c_ρ as in (2.3) we have

$$\frac{k}{n} = \frac{\lambda(\overset{\circ}{\lambda}_k)}{\lambda(\overset{\circ}{\lambda}_n)} \leq \frac{\lambda(\rho \overset{\circ}{\lambda}_n)}{\lambda(\overset{\circ}{\lambda}_n)} \leq c_\rho,$$

which implies that $\theta_{k,n} \leq c_\rho n$. Moreover, $\lambda^{-1}(\theta_{k,n})$ is some point between $\overset{\circ}{\lambda}_k$ and $\overset{\circ}{\lambda}_n$, and hence $\lambda^{-1}(\theta_{k,n}) \geq \frac{1}{\rho} \overset{\circ}{\lambda}_n$. Hence

$$\begin{aligned} \sum_{\substack{k \in \mathbb{N}: k \neq n \\ \frac{\overset{\circ}{\lambda}_k}{\overset{\circ}{\lambda}_n} \in (\frac{1}{\rho}, \rho)}} \frac{1}{|\overset{\circ}{\lambda}_k - \overset{\circ}{\lambda}_n|} &= \sum_{\substack{k \in \mathbb{N}: k \neq n \\ \frac{\overset{\circ}{\lambda}_k}{\overset{\circ}{\lambda}_n} \in (\frac{1}{\rho}, \rho)}} \frac{\lambda'(\lambda^{-1}(\theta_{k,n}))}{|k - n|} \leq C \sum_{\substack{k \in \mathbb{N}: k \neq n \\ \frac{\overset{\circ}{\lambda}_k}{\overset{\circ}{\lambda}_n} \in (\frac{1}{\rho}, \rho)}} \frac{1}{|k - n|} \cdot \frac{\theta_{k,n}}{\lambda^{-1}(\theta_{k,n})} \\ &\leq C \frac{c_\rho n}{\frac{1}{\rho} \overset{\circ}{\lambda}_n} \sum_{\substack{k \in \mathbb{N}: k \neq n \\ \frac{\overset{\circ}{\lambda}_k}{\overset{\circ}{\lambda}_n} \in (\frac{1}{\rho}, \rho)}} \frac{1}{|k - n|} \leq C \frac{\rho c_\rho n}{\overset{\circ}{\lambda}_n} \cdot 2 \sum_{l=1}^{r_{\overset{\circ}{\lambda}}(\rho, n)} \frac{1}{l} \lesssim \frac{n \log n}{\overset{\circ}{\lambda}_n} \end{aligned}$$

as $n \rightarrow \infty$. □

Next, let us investigate growth functions Λ such that $\overset{\circ}{\lambda}$ is of convergence class with respect to Λ . Since

$$\sum_{n \in \mathbb{N}} \frac{1}{\lambda(\overset{\circ}{\lambda}_n)} = \sum_{n \in \mathbb{N}} \frac{1}{n} = \infty,$$

one has to choose a slightly larger growth function in order to achieve convergence class. For example, the choices $\Lambda(r) = \lambda(r)(\log \lambda(r))^2$ and $\Lambda(r) = \lambda(r) \log \lambda(r)(\log \log \lambda(r))^2$, for large r , will always do the job.

The next statement can be seen as a refinement of Lemma 2.9. Thinking of the example below Definition 2.8, this fact is no surprise. This estimate is used in Corollary 3.20 below.

2.13 Lemma. *Let λ and Λ be growth functions such that the standard sequence $\overset{\circ}{\lambda}$ is of convergence class with respect to Λ . Assume that $\frac{\lambda}{\Lambda}$ is, for sufficiently large values of r , non-increasing.*

If $\rho_\lambda > 0$ or the function $\frac{\log \lambda(r)}{\log r}$ is non-increasing for large r , then

$$\lambda(r) \log \lambda(r) = O(\Lambda(r)), \quad r \rightarrow \infty. \quad (2.10)$$

Proof. Let $n \in \mathbb{N}$. Then

$$\begin{aligned} \int_{\overset{\circ}{\lambda}_n}^{\overset{\circ}{\lambda}_{n+1}} \frac{1}{\Lambda(r)} \lambda'(r) dr &\leq \frac{1}{\Lambda(\overset{\circ}{\lambda}_n)} \int_{\overset{\circ}{\lambda}_n}^{\overset{\circ}{\lambda}_{n+1}} \lambda'(r) dr \\ &= \frac{1}{\Lambda(\overset{\circ}{\lambda}_n)} (\lambda(\overset{\circ}{\lambda}_{n+1}) - \lambda(\overset{\circ}{\lambda}_n)) = \frac{1}{\Lambda(\overset{\circ}{\lambda}_n)}, \end{aligned}$$

which implies that $\int_1^\infty \frac{1}{\Lambda(r)} \lambda'(r) dr < \infty$.

First, assume that $\rho_\lambda > 0$. Then we find $r_0 > 1$ such that $\frac{\lambda(r)}{\Lambda(r)}$ is non-increasing for $r \geq r_0$ and, by (gf2),

$$\lambda'(r) \geq \frac{\rho_\lambda}{2} \frac{\lambda(r)}{r}, \quad r \geq r_0.$$

Hence, $\int_{r_0}^\infty \frac{1}{\Lambda(r)} \frac{\lambda(r)}{r} dr < \infty$, and integrating by parts gives ($R \geq r_0$)

$$\begin{aligned} \infty &> \int_{r_0}^\infty \frac{1}{\Lambda(r)} \frac{\lambda(r)}{r} dr \geq \int_{r_0}^R \frac{\lambda(r)}{\Lambda(r)} \cdot \frac{1}{r} dr \\ &= \frac{\lambda(R)}{\Lambda(R)} \log R - \frac{\lambda(r_0)}{\Lambda(r_0)} \log r_0 - \int_{r_0}^R \underbrace{\left(\frac{\lambda(r)}{\Lambda(r)} \right)'}_{\leq 0} \log r dr. \end{aligned}$$

We see that $\frac{\lambda(R)}{\Lambda(R)} \log R$ remains bounded when R tends to ∞ , and hence (2.10) holds since λ satisfies (gf1).

Assume now that $\frac{\log \lambda(r)}{\log r}$ is non-increasing (for large r). Then we can choose $r_0 > 1$ such that $\frac{\lambda(r)}{\Lambda(r)}$ and $\frac{\log \lambda(r)}{\log r}$ are both non-increasing for $r \geq r_0$ and that

$$\lambda'(r) \geq \frac{1}{2} \frac{\lambda(r) \log \lambda(r)}{r \log r}, \quad r \geq r_0.$$

Again we integrate by parts and obtain

$$\begin{aligned} \infty &> 2 \int_{r_0}^\infty \frac{1}{\Lambda(r)} \lambda'(r) dr \geq \int_{r_0}^R \frac{\lambda(r) \log \lambda(r)}{\Lambda(r) \log r} \cdot \frac{1}{r} dr \\ &= \frac{\lambda(R) \log \lambda(R)}{\Lambda(R)} - \frac{\lambda(r_0) \log \lambda(r_0)}{\Lambda(r_0)} - \int_{r_0}^R \underbrace{\left(\frac{\lambda(r) \log \lambda(r)}{\Lambda(r) \log r} \right)'}_{\leq 0} \cdot \log r dr, \end{aligned}$$

which, also in this case, shows that $\frac{\lambda(R) \log \lambda(R)}{\Lambda(R)}$ is bounded. \square

2.14 Remark. The hypothesis of this Lemma 2.13 is satisfied in ‘most’ cases. (1) In order to construct a growth function Λ which makes λ of convergence class, it is natural to multiply λ with some growing factor, in which case $\frac{\lambda(r)}{\Lambda(r)}$ is non-increasing. (2) Assume that $\rho_\lambda = 0$. Then $\frac{\log \lambda(r)}{\log r}$ is a positive function (for large r), which tends to zero. Thinking of λ as a regularly behaving function, requiring monotonicity appears to be not too restrictive.

If λ is given by its inverse function μ , then one can characterize the assumptions in Lemma 2.13 in terms of μ . Namely, $\frac{\lambda(r)}{\Lambda(r)}$ is non-increasing if and only if $\frac{\Lambda(\mu(s))}{s}$ is non-decreasing; $\frac{\log \lambda(r)}{\log r}$ is non-increasing if and only if $\frac{\log \mu(s)}{\log s}$ is non-decreasing. Note also that if $\mu(s) = O(s^\sigma)$, $s \rightarrow \infty$, for some $\sigma > 0$, then $\rho_\lambda \geq \frac{1}{\sigma} > 0$. //

3 Stability theorems

Let us introduce the precise setup.

3.1 Definition. We denote by \mathbb{S} the set of all sequences $\xi = (\xi_n)_{n=1}^\infty$ of real numbers that satisfy the following conditions.

- (S1) The sequence ξ consists of pairwise distinct non-zero points and has no finite accumulation point.
- (S2) Denote by ξ^+ and ξ^- the (finite or infinite) subsequences of ξ consisting of all positive or negative, respectively, elements of ξ arranged according to increasing modulus⁷. Then

$$\lim_{n \rightarrow \infty} \frac{n}{\xi_n^+} = \lim_{n \rightarrow \infty} \frac{n}{|\xi_n^-|} \in [0, \infty),$$

where we tacitly understand the limit of a finite sequence as being equal to 0.

- (S3) The limit

$$\lim_{r \rightarrow \infty} \sum_{|\xi_n| \leq r} \frac{1}{\xi_n}$$

exists in \mathbb{R} .

//

Note that we assume no particular ordering of ξ ; only ξ^+ and ξ^- are ordered with increasing modulus. We also mention that (S2) is equivalent to

$$\delta_\lambda^+(\xi) = \delta_\lambda^-(\xi) = \Delta_\lambda^+(\xi) = \Delta_\lambda^-(\xi) \in [0, \infty)$$

for $\lambda(r) = r$.

With each sequence $\xi \in \mathbb{S}$ we associate an entire function P_ξ , namely the canonical product connected with ξ .

3.2 Definition. Let $\xi \in \mathbb{S}$. Then we set

$$P_\xi(z) := \lim_{r \rightarrow \infty} \prod_{|\xi_n| \leq r} \left(1 - \frac{z}{\xi_n}\right), \quad z \in \mathbb{C}.$$

//

Because of (S1)–(S3), this limit exists locally uniformly on \mathbb{C} and represents an entire function of finite exponential type whose zeros are all simple and located exactly at the points ξ_n , $n \in \mathbb{N}$.

The following statement is the main result of this paper.

3.3 Theorem. *Let $\alpha \in \mathbb{S}$ and let Λ be a growth function such that, for sufficiently large r , the function $\frac{\Lambda(r)}{r}$ is either non-increasing or non-decreasing. Moreover, assume that α is of convergence class with respect to Λ , i.e.*

$$\sum_{n \in \mathbb{N}} \frac{1}{\Lambda(|\alpha_n|)} < \infty. \quad (3.1)$$

⁷Both sequences are supposed to have finite or infinite index sets of the form $1, 2, \dots$

Further, let $\beta \in \mathbb{S}$ be a small perturbation of α in the sense that the difference $\gamma := \beta - \alpha$ satisfies the following conditions:

$$(A) \quad |\gamma_n| = O\left(\frac{|\alpha_n|}{\Lambda(|\alpha_n|)}\right), \quad n \rightarrow \infty; \quad (3.2)$$

$$(B) \quad \left(\frac{\gamma_n}{s_\alpha(n)}\right)_{n \in \mathbb{N}} \in \ell^1, \quad (3.3)$$

$$\exists \rho > 1 : \frac{\gamma_n}{s_\alpha(n)} = O\left(\frac{1}{r_\alpha(\rho, n)}\right), \quad n \rightarrow \infty, \quad (3.4)$$

where s_α and r_α are defined as in Definition 2.6. Let P_α and P_β be the canonical products associated with α and β , respectively. Then

$$|P'_\alpha(\alpha_n)| \asymp |P'_\beta(\beta_n)|, \quad n \rightarrow \infty. \quad (3.5)$$

3.4 Remark.

- (i) If $\rho_\Lambda < 1$, then $\frac{\Lambda(r)}{r}$ is decreasing for large r by Remark 2.2 (iii). In this case, Condition (A) allows γ to grow (although the conditions in (B) still have to be satisfied, which may prohibit this). If $\frac{\Lambda(r)}{r}$ is non-decreasing, then γ must be a bounded sequence.
- (ii) Let us comment on the role of the condition (S2) and (S3) which are required from the start. What we want to do (and what we are able to do) is to estimate the derivative of functions which are of finite exponential type and represented as a product P_ξ , and not of canonical products which include exponential factors. In order to ensure that P_ξ converges and is of finite exponential type, of course some requirements on ξ are necessary, namely at least that the genus of the sequence ξ does not exceed 1 and that (S3) holds. Condition (S2) on existence and equality of angular densities is of course not necessary purely for convergence. It is related to regular growth: in the applications which motivated us to study this stability question at all, sequences are zero sequences of functions of Cartwright class and thus satisfy (S2).

We should point out that, as soon as we deal with (zero sequences of) functions of minimal exponential type, both conditions (S2) and (S3) are automatic (by Lindelöf's theorem).

//

3.5 Remark. The following intuitive picture can be regarded as commonly accepted (believed):

At places in the vicinity of which the sequence α is well separated the associated sequence $\alpha' = (P'_\alpha(\alpha_n))_{n \in \mathbb{N}}$ behaves regularly and can be controlled. On the other hand, points of α being close to each other give rise to peaks in α' , and lumps of points being close to each other produce peaks which even may spread out over neighbouring points.

Hence, it is to be expected that a perturbation of α which does not influence the behaviour of α' should be asymptotically smaller than the separation s_α of α . The perturbation must certainly be limited by the separation of α

because otherwise, we could remove or add zeros, which definitely changes the asymptotic behaviour of α' .

The conditions (A) and (B) are quantitative instantiations of this idea. For (B) this is obvious: (3.3) rules out perturbations which produce close points, and (3.4) rules out that lumps of points appear in the vicinity of α_n . To understand (A), it is advisable to consider an example of a very regularly distributed sequence where all involved quantities can be computed explicitly. Take, e.g. the sequence $\alpha_n := n^\sigma$ with some $\sigma > 1$. Then $s_\alpha(n) \asymp n^{\sigma-1}$, and we may choose for Λ , e.g. $\Lambda(r) := r^{\frac{1}{\sigma}} \log r (\log \log r)^2$. Condition (3.2) becomes

$$\frac{|\gamma_n|}{s_\alpha(n)} = O\left(\frac{1}{\log n (\log \log n)^2}\right).$$

We see that (A) requires the perturbation to be only slightly smaller than the separation, but, contrasting (3.3), that $\frac{|\gamma_n|}{s_\alpha(n)}$ tends to zero in a controllably regular way. We discuss more examples in Subsection 3.3. //

We split the proof of Theorem 3.3 in two subsections.

3.1 The basic estimates

First note that

$$P'_\xi(\xi_n) = -\frac{1}{\xi_n} \lim_{r \rightarrow \infty} \prod_{\substack{|\xi_k| \leq r \\ k \neq n}} \left(1 - \frac{\xi_n}{\xi_k}\right), \quad n \in \mathbb{N}. \quad (3.6)$$

In this subsection we provide two general estimates for quotients of such products; see Propositions 3.6 and 3.9 below. The first one deals with factors where ξ_k is small.

3.6 Proposition. *Let the following data be given:*

- a sequence $\xi = (\xi_n)_{n \in \mathbb{N}}$ of non-zero real numbers which has no finite accumulation point;
- a growth function Λ such that ξ is of convergence class with respect to Λ ;
- a sequence $\nu = (\nu_n)_{n \in \mathbb{N}}$ of real numbers with $\nu_n \neq -\xi_n$, $n \in \mathbb{N}$.

Denote by r_n the unique positive numbers with $r_n \Lambda(r_n) = |\xi_n|$, let $c > 0$ and set

$$J_n(c, \Lambda) := \{k \in \mathbb{N} : |\xi_k| \leq cr_n\}.$$

Moreover, set $\eta_n := \xi_n + \nu_n$, $n \in \mathbb{N}$.

If the sequence ν is subject to the condition

$$|\nu_n| = O\left(\frac{|\xi_n|}{\Lambda(|\xi_n|)}\right), \quad n \rightarrow \infty, \quad (3.7)$$

then

$$0 < \liminf_{n \rightarrow \infty} \left| \prod_{k \in J_n(c, \Lambda)} \frac{1 - \frac{\eta_n}{\eta_k}}{1 - \frac{\xi_n}{\xi_k}} \right| \leq \limsup_{n \rightarrow \infty} \left| \prod_{k \in J_n(c, \Lambda)} \frac{1 - \frac{\eta_n}{\eta_k}}{1 - \frac{\xi_n}{\xi_k}} \right| < \infty. \quad (3.8)$$

It is shown in the proof below that $n \notin J_n(c, \Lambda)$ if n is large enough, so that the products are well defined for such n .

We frequently use the standard procedure to estimate products by taking logarithms. The following remark is of course trivial; however, since our estimates have to be uniform with respect to several parameters, it is better to be precise (we return to this note in Remark 3.17).

3.7 Remark. Let I be a finite subset of \mathbb{N} , and let x_n , $n \in I$, be real numbers with $|x_n| \leq \frac{1}{2}$. Then

$$\left| \log \left| \prod_{n \in I} (1 + x_n) \right| \right| \leq \log 4 \cdot \sum_{n \in I} |x_n|. \quad (3.9)$$

This is obvious from the fact that

$$\log |1 + x| \begin{cases} \leq \log(1 + |x|) \leq |x|, & x \in \mathbb{R}, \\ \geq \log(1 - |x|) \geq -(\log 4)|x|, & |x| \leq \frac{1}{2}. \end{cases}$$

Of course, the inequality in (3.9) then leads to the estimate

$$e^{-\log 4 \sum_{n \in I} |x_n|} \leq \left| \prod_{n \in I} (1 + x_n) \right| \leq e^{\log 4 \sum_{n \in I} |x_n|}. \quad (3.10)$$

//

Proof of Proposition 3.6. First, we collect some simple facts.

- (i) The numbers r_n are indeed well defined since the function $r\Lambda(r)$ is a bijection from $(0, \infty)$ onto itself. Moreover, $\lim_{n \rightarrow \infty} r_n = \infty$.
- (ii) We have $\frac{r_n}{|\xi_n|} = \frac{1}{\Lambda(r_n)} \rightarrow 0$. In particular, for n sufficiently large, $r_n \leq |\xi_n|$. For such values of n ,

$$\frac{r_n}{|\xi_n|} = \frac{1}{\Lambda(r_n)} \geq \frac{1}{\Lambda(|\xi_n|)},$$

and hence $r_n \geq \frac{|\xi_n|}{\Lambda(|\xi_n|)}$.

- (iii) We have $\frac{\nu_n}{|\xi_n|} = O\left(\frac{1}{\Lambda(|\xi_n|)}\right)$, and hence

$$\lim_{n \rightarrow \infty} \frac{\nu_n}{|\xi_n|} = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{|\eta_n|}{|\xi_n|} = 1. \quad (3.11)$$

- (iv) Let $c > 0$. Then, by (ii), there exists an $n_0 \in \mathbb{N}$ such that $r_n \leq \frac{\xi_n}{4c}$, $n \geq n_0$, and hence

$$|\xi_k| \leq \frac{1}{4} |\xi_n|, \quad k \in J_n(c, \Lambda), \quad n \geq n_0.$$

Using (3.11) we conclude that there exists an $n_1 \in \mathbb{N}$ such that

$$|\eta_k| \leq \frac{1}{2} |\eta_n|, \quad k \in J_n(c, \Lambda), \quad n \geq n_1.$$

(v) For $n \geq n_0$ we have $k \notin J_n(c, \Lambda)$ whenever $|\xi_k| = |\xi_n|$. In particular, $n \notin J_n(c, \Lambda)$. For such indices n , the products in (3.8) are well defined.

(vi) For each finite subset $M \subseteq \mathbb{N}$, there exists an $n_2 \in \mathbb{N}$ such that

$$M \subseteq J_n(c, \Lambda), \quad n \geq n_2.$$

(vii) By Lemma 2.9 we have

$$\#J_n(c, \Lambda) = n_\xi(cr_n) = O(\Lambda(r_n)), \quad n \rightarrow \infty.$$

Now we come to the actual proof of Proposition 3.6. Choose $N \in \mathbb{N}$ such that

$$\begin{aligned} \frac{|\nu_n|}{|\xi_n|} &\leq \left[4 \sup_{l \in \mathbb{N}} \frac{|\xi_l|}{|\eta_l|} \right]^{-1}, \quad \frac{|\nu_n|}{|\eta_n|} \leq \frac{1}{4}, \quad n \geq N, \\ \frac{|\xi_k|}{|\xi_n|} &\leq \frac{1}{2}, \quad \frac{|\eta_k|}{|\eta_n|} \leq \frac{1}{2}, \quad n \geq N, \quad k \in J_n(c, \Lambda), \end{aligned}$$

and choose $N_1 \in \mathbb{N}$, $N_1 \geq N$, such that

$$\{k \in \mathbb{N} : k < N\} \subseteq J_n(c, \Lambda), \quad n \geq N_1.$$

We rewrite (using the shorthand $J_n := J_n(c, \Lambda)$)

$$\prod_{k \in J_n} \frac{1 - \frac{\eta_n}{\eta_k}}{1 - \frac{\xi_n}{\xi_k}} = \frac{\prod_{k \in J_n} \left(\frac{\eta_k}{\eta_n} - 1 \right)}{\prod_{k \in J_n} \left(\frac{\xi_k}{\xi_n} - 1 \right)} \cdot \prod_{\substack{k \in J_n \\ k \geq N}} \frac{\xi_k \eta_n}{\xi_n \eta_k} \cdot \prod_{k < N} \frac{\xi_k \eta_n}{\xi_n \eta_k}, \quad n \geq N_1, \quad (3.12)$$

and consider each factor separately.

For the product in the denominator of the first factor, we estimate

$$\sum_{k \in J_n} \left| \frac{\xi_k}{\xi_n} \right| \leq \frac{\max_{k \in J_n} |\xi_k|}{|\xi_n|} \cdot (\#J_n) \lesssim \frac{cr_n}{|\xi_n|} \cdot \Lambda(r_n) = c.$$

For the product in the numerator, note that

$$\left| \frac{\eta_k}{\eta_n} \right| \leq \sup_{k \in \mathbb{N}} \left| \frac{\eta_k}{\xi_k} \right| \cdot \sup_{n \in \mathbb{N}} \left| \frac{\xi_n}{\eta_n} \right| \cdot \left| \frac{\xi_k}{\xi_n} \right|.$$

Hence, the same estimate applies.

Next, we write

$$\frac{\xi_k \eta_n}{\xi_n \eta_k} = 1 + \frac{\xi_k(\eta_n - \xi_n) - (\eta_k - \xi_k)\xi_n}{\xi_n \eta_k} = 1 + \frac{\xi_k}{\eta_k} \frac{\nu_n}{\xi_n} - \frac{\nu_k}{\eta_k},$$

and, using (ii), (iii) and (vii), we can estimate

$$\sum_{k \in J_n} \left| \frac{\xi_k}{\eta_k} \frac{\nu_n}{\xi_n} \right| \leq \frac{|\nu_n|}{|\xi_n|} \cdot (\#J_n) \sup_{k \in \mathbb{N}} \left| \frac{\xi_k}{\eta_k} \right| \lesssim \frac{|\nu_n|}{|\xi_n|} \Lambda(|r_n|) \lesssim \frac{|\nu_n|}{|\xi_n|} \Lambda(|\xi_n|) = O(1),$$

$$\sum_{k \in J_n} \left| \frac{\nu_k}{\eta_k} \right| \leq \sup_{k \in \mathbb{N}} \left| \frac{\xi_k}{\eta_k} \right| \cdot \sum_{k \in J_n} \left| \frac{\nu_k}{\xi_k} \right| \lesssim \sum_{k \in J_n} \frac{1}{\Lambda(|\xi_k|)} \leq \sum_{k \in \mathbb{N}} \frac{1}{\Lambda(|\xi_k|)} < \infty.$$

By our choice of N , we have, for $n \geq N$, $k \in J_n$, $k \geq N$, that

$$\left| \frac{\xi_k \nu_n}{\eta_k \xi_n} - \frac{\nu_k}{\eta_k} \right| \leq \frac{|\nu_n|}{|\xi_n|} \sup_{l \in \mathbb{N}} \frac{|\xi_l|}{|\eta_l|} + \sup_{l \geq N} \frac{|\nu_l|}{|\eta_l|} \leq \frac{1}{2}$$

and hence Remark 3.7 can be applied to the second factor in (3.12). The same is true for the products in the first factor. Hence the estimate (3.10) implies that the products in the first three factors are bounded from above and away from zero uniformly in n .

It remains to notice that

$$\lim_{n \rightarrow \infty} \prod_{k < N} \frac{\xi_k \eta_n}{\xi_n \eta_k} = \prod_{k < N} \frac{\xi_k}{\eta_k} \neq 0.$$

□

Proposition 3.9 below contains a key estimate. In its proof we use the following fact.

3.8 Lemma. *Let Λ be a growth function, $\rho > 1$ and $R > 0$ such that $\frac{\Lambda(r)}{r}$ is non-decreasing and $\frac{\Lambda(r)}{r^2}$ is non-increasing on $[R, \infty)$. Then*

$$\left| \frac{b}{a(a-b)} \right| \leq \frac{\rho}{\rho-1} \cdot \frac{\Lambda(|b|)}{|b|\Lambda(|a|)}, \quad a, b \in \mathbb{R}, \quad |a|, |b| \geq R, \quad \frac{a}{b} \notin \left(\frac{1}{\rho}, \rho \right).$$

Proof. We first consider the case when $a, b > 0$. If $\frac{a}{b} \leq \frac{1}{\rho}$, then $|a-b| = (1 - \frac{a}{b})b \geq (1 - \frac{1}{\rho})b$, and hence

$$\left| \frac{b}{a(a-b)} \right| \leq \frac{1}{1 - \frac{1}{\rho}} \cdot \frac{1}{a} = \frac{\rho}{\rho-1} \cdot \left(\frac{\Lambda(a)}{a} / \frac{\Lambda(b)}{b} \right) \cdot \frac{\Lambda(b)}{b\Lambda(a)}.$$

Since $R \leq a \leq \frac{1}{\rho}b \leq b$, it follows from the monotonicity of $\frac{\Lambda(r)}{r}$ that the second factor is bounded by 1.

If $\frac{a}{b} \geq \rho$, then $|a-b| = (1 - \frac{b}{a})a \geq (1 - \frac{1}{\rho})a$, and hence

$$\left| \frac{b}{a(a-b)} \right| \leq \frac{1}{1 - \frac{1}{\rho}} \frac{b}{a^2} = \frac{\rho}{\rho-1} \cdot \left(\frac{\Lambda(a)}{a^2} / \frac{\Lambda(b)}{b^2} \right) \cdot \frac{\Lambda(b)}{b\Lambda(a)}.$$

Since $R \leq b \leq \frac{1}{\rho}a \leq a$, the second factor is bounded by 1, where we used the monotonicity of $\frac{\Lambda(r)}{r^2}$. Putting this together we obtain the required estimate.

Next assume that $a, b < 0$. The already proved case, applied to $|a|, |b|$, yields

$$\left| \frac{b}{a(a-b)} \right| = \left| \frac{|b|}{|a|(|a|-|b|)} \right| \leq \frac{\rho}{\rho-1} \cdot \frac{\Lambda(|b|)}{|b|\Lambda(|a|)}.$$

It remains to consider the case when a and b have different signs. Then, using the same estimates as above, we obtain

$$\left| \frac{b}{a(a-b)} \right| = \frac{|b|}{|a|(|a|+|b|)} \leq \begin{cases} \frac{1}{|a|} & \text{if } |b| \geq |a| \\ \frac{|b|}{|a|^2} & \text{if } |b| < |a| \end{cases} \leq \frac{\Lambda(|b|)}{|b|\Lambda(|a|)},$$

which finishes the proof. □

3.9 Proposition. *Let the following data be given:*

- a finite subset I of \mathbb{N} ;
- a sequence $\xi = (\xi_k)_{k \in I}$ of pairwise distinct and non-zero real numbers;
- a growth function Λ and $R_0 > 0$ such that $\frac{\Lambda(r)}{r}$ is either non-increasing or non-decreasing on $[R_0, \infty)$; in the latter case assume, in addition, that $\frac{\Lambda(r)}{r^2}$ is decreasing on $[R_0, \infty)$;
- a number $\rho > 1$;
- a sequence $\nu = (\nu_k)_{k \in I}$ of real numbers with $\nu_k \neq -\xi_k$, $k \in I$;
- an element $n \in I$.

Set

$$\delta := \frac{1}{4(2\frac{\rho^2}{\rho-1} + 1)}, \quad (3.13)$$

$$\eta_k := \xi_k + \nu_k, \quad k \in I,$$

and

$$S_1(n) := \sum_{k \in I} \frac{|\nu_k|}{|\xi_k|}, \quad S_2(n) := \sum_{\substack{k \in I, k \neq n \\ \frac{\xi_k}{\xi_n} \in (\frac{1}{\rho}, \rho)}} \frac{|\nu_k|}{|\xi_k - \xi_n|},$$

$$S_3(n) := \frac{|\nu_n|}{|\xi_n|} \Lambda(|\xi_n|) \sum_{k \in I} \frac{1}{\Lambda(|\xi_k|)}, \quad S_4(n) := |\nu_n| \sum_{\substack{k \in I, k \neq n \\ \frac{\xi_k}{\xi_n} \in (\frac{1}{\rho}, \rho)}} \frac{1}{|\xi_k - \xi_n|},$$

$$T(n) := \frac{|\nu_n|}{|\xi_n|} n_\xi(|\xi_n|),$$

$$\Theta(n) := \frac{\log 4}{1 - \delta} \left[\frac{\rho}{\rho - 1} S_1(n) + \rho S_2(n) + \frac{\rho}{\rho - 1} T(n) + \frac{\rho^2}{\rho - 1} S_3(n) + S_4(n) \right],$$

$$\tilde{\Theta}(n) := \frac{\log 4}{1 - 2\delta} \left[\frac{\rho}{\rho - 1} S_1(n) + \rho S_2(n) + \frac{\rho}{\rho - 1} S_3(n) + \rho S_4(n) \right].$$

Assume that

$$\forall k \in I : \quad \frac{|\nu_n|}{|\xi_k|} \leq \delta, \quad \frac{|\nu_k|}{|\xi_k|} \leq \delta, \quad \frac{|\nu_k|}{s_\xi(k)} \leq \delta, \quad |\xi_k| \geq \rho R_0, \quad (3.14)$$

where $s_\xi(n)$ is defined as in Definition 2.6.

(i) If $\frac{\Lambda(r)}{r}$ is non-increasing on $[R_0, \infty)$, then

$$\left| \log \left| \prod_{\substack{k \in I \\ k \neq n}} \frac{1 - \frac{\eta_n}{\xi_n}}{1 - \frac{\eta_k}{\xi_k}} \right| \right| \leq \Theta(n). \quad (3.15)$$

(ii) If $\frac{\Lambda(r)}{r}$ is non-decreasing on $[R_0, \infty)$, then

$$\left| \log \left(\left| \prod_{\substack{k \in I \\ k \neq n}} \frac{1 - \frac{\eta_n}{\eta_k}}{1 - \frac{\xi_n}{\xi_k}} \right| \middle/ \left| \prod_{\substack{k \in I \\ k \neq n}} \left(1 - \frac{\nu_n}{\eta_k} \right) \right| \right) \right| \leq \tilde{\Theta}(n). \quad (3.16)$$

Proof.

Step 1: rewriting products. For $k \neq n$ we can rewrite

$$\begin{aligned} \frac{1 - \frac{\eta_n}{\eta_k}}{1 - \frac{\xi_n}{\xi_k}} &= \frac{\xi_k(\eta_k - \eta_n)}{\eta_k(\xi_k - \xi_n)} = 1 + \frac{\xi_k(\eta_k - \eta_n) - \eta_k(\xi_k - \xi_n)}{\eta_k(\xi_k - \xi_n)} \\ &= 1 + \frac{\xi_n \eta_k - \xi_k \eta_n}{\eta_k(\xi_k - \xi_n)} = 1 + \frac{\xi_n(\eta_k - \xi_k) - \xi_k(\eta_n - \xi_n)}{\eta_k(\xi_k - \xi_n)} \\ &= 1 + \frac{\xi_n \nu_k - \xi_k \nu_n}{\eta_k(\xi_k - \xi_n)}. \end{aligned} \quad (3.17)$$

The latter expression can also be written as

$$\begin{aligned} 1 - \frac{\nu_n}{\eta_k} + \frac{\xi_n \nu_k - \xi_k \nu_n + \nu_n(\xi_k - \xi_n)}{\eta_k(\xi_k - \xi_n)} \\ &= 1 - \frac{\nu_n}{\eta_k} + \frac{\xi_n(\nu_k - \nu_n)}{\eta_k(\xi_k - \xi_n)} \\ &= \left(1 - \frac{\nu_n}{\eta_k} \right) \left[1 - \left(1 - \frac{\nu_n}{\eta_k} \right)^{-1} \frac{\xi_n(\nu_k - \nu_n)}{\eta_k(\xi_k - \xi_n)} \right]. \end{aligned} \quad (3.18)$$

Using (3.17) and (3.18), respectively, we obtain

$$\prod_{\substack{k \in I \\ k \neq n}} \frac{1 - \frac{\eta_n}{\eta_k}}{1 - \frac{\xi_n}{\xi_k}} = \prod_{\substack{k \in I \\ k \neq n}} \left[1 + \frac{\xi_n \nu_k - \xi_k \nu_n}{\eta_k(\xi_k - \xi_n)} \right], \quad (3.19)$$

$$\frac{\prod_{\substack{k \in I \\ k \neq n}} \frac{1 - \frac{\eta_n}{\eta_k}}{1 - \frac{\xi_n}{\xi_k}}}{\prod_{\substack{k \in I \\ k \neq n}} \left(1 - \frac{\nu_n}{\eta_k} \right)} = \prod_{\substack{k \in I \\ k \neq n}} \left[1 - \left(1 - \frac{\nu_n}{\eta_k} \right)^{-1} \frac{\xi_n(\nu_k - \nu_n)}{\eta_k(\xi_k - \xi_n)} \right]. \quad (3.20)$$

Step 2: Remark 3.7 is applicable. Our assumptions imply that, for each $k \in I$,

$$\begin{aligned} \left| \frac{\eta_k}{\xi_k} \right| &= \left| 1 + \frac{\nu_k}{\xi_k} \right| \geq 1 - \left| \frac{\nu_k}{\xi_k} \right| \geq 1 - \delta, \\ \left| 1 - \frac{\nu_n}{\eta_k} \right| &\geq 1 - \left| \frac{\xi_k}{\eta_k} \right| \cdot \left| \frac{\nu_n}{\xi_k} \right| \geq 1 - \frac{1}{1 - \delta} \delta = \frac{1 - 2\delta}{1 - \delta}. \end{aligned}$$

Moreover, Lemma 3.8 applied with the growth function $\Lambda(r) = r$ gives

$$\left| \frac{\xi_n}{\xi_k(\xi_k - \xi_n)} \right| \leq \begin{cases} \frac{\rho}{\rho - 1} \frac{1}{|\xi_k|} & \text{if } \frac{\xi_n}{\xi_k} \notin \left(\frac{1}{\rho}, \rho \right), \\ \frac{\rho}{|\xi_k - \xi_n|} & \text{if } \frac{\xi_n}{\xi_k} \in \left(\frac{1}{\rho}, \rho \right), \end{cases} \quad (3.21)$$

and hence

$$\left| \frac{\xi_n \nu_k}{\xi_k (\xi_k - \xi_n)} \right| \leq \begin{cases} \frac{\rho}{\rho-1} \frac{|\nu_k|}{|\xi_k|} \leq \delta \frac{\rho}{\rho-1} & \text{if } \frac{\xi_n}{\xi_k} \notin \left(\frac{1}{\rho}, \rho \right), \\ \rho \frac{|\nu_k|}{s_\xi(k)} \leq \delta \rho & \text{if } \frac{\xi_n}{\xi_k} \in \left(\frac{1}{\rho}, \rho \right), \end{cases}$$

$$\left| \frac{\xi_n \nu_n}{\xi_k (\xi_k - \xi_n)} \right| \leq \begin{cases} \frac{\rho}{\rho-1} \frac{|\nu_n|}{|\xi_k|} \leq \delta \frac{\rho}{\rho-1} & \text{if } \frac{\xi_n}{\xi_k} \notin \left(\frac{1}{\rho}, \rho \right), \\ \rho \frac{|\nu_n|}{s_\xi(n)} \leq \delta \rho & \text{if } \frac{\xi_n}{\xi_k} \in \left(\frac{1}{\rho}, \rho \right). \end{cases}$$

Both expressions are bounded by $\delta \frac{\rho^2}{\rho-1}$ in each case. The definition of δ yields $\delta \leq \frac{1}{4}$ and $\frac{\rho^2}{\rho-1} \leq \frac{1}{8\delta}$, and thus we obtain

$$\begin{aligned} \left| \frac{\xi_n \nu_k - \xi_k \nu_n}{\eta_k (\xi_k - \xi_n)} \right| &= \left| \frac{\xi_k}{\eta_k} \left(\frac{\xi_n \nu_k}{\xi_k (\xi_k - \xi_n)} - \frac{\nu_n}{\xi_k - \xi_n} \right) \right| \\ &\leq \frac{1}{1-\delta} \left(\left| \frac{\xi_n \nu_k}{\xi_k (\xi_k - \xi_n)} \right| + \frac{|\nu_n|}{s_\xi(n)} \right) \\ &\leq \frac{1}{1-\delta} \left(\delta \frac{\rho^2}{\rho-1} + \delta \right) \leq \frac{1}{1-\delta} \left(\frac{1}{8} + \delta \right) \leq \frac{1}{1-\frac{1}{4}} \left(\frac{1}{8} + \frac{1}{4} \right) = \frac{1}{2} \end{aligned}$$

and

$$\begin{aligned} \left| \left(1 - \frac{\nu_n}{\eta_k} \right)^{-1} \frac{\xi_n (\nu_k - \nu_n)}{\eta_k (\xi_k - \xi_n)} \right| &= \left| 1 - \frac{\nu_n}{\eta_k} \right|^{-1} \cdot \left| \frac{\xi_k}{\eta_k} \right| \cdot \left| \frac{\xi_n \nu_k}{\xi_k (\xi_k - \xi_n)} - \frac{\xi_n \nu_n}{\xi_k (\xi_k - \xi_n)} \right| \\ &\leq \frac{1-\delta}{1-2\delta} \cdot \frac{1}{1-\delta} \cdot 2\delta \frac{\rho^2}{\rho-1} \leq \frac{1}{1-2 \cdot \frac{1}{4}} \cdot \frac{2}{8} = \frac{1}{2}. \end{aligned}$$

Therefore we can apply Remark 3.7 to the right-hand sides of (3.19) and (3.20) and obtain the estimates

$$\begin{aligned} \left| \log \left| \prod_{\substack{k \in I \\ k \neq n}} \frac{1 - \frac{\eta_n}{\eta_k}}{1 - \frac{\xi_n}{\xi_k}} \right| \right| &= \left| \log \left| \prod_{\substack{k \in I \\ k \neq n}} \left[1 + \frac{\xi_n \nu_k - \xi_k \nu_n}{\eta_k (\xi_k - \xi_n)} \right] \right| \right| \\ &\leq \log 4 \sum_{\substack{k \in I \\ k \neq n}} \left| \frac{\xi_n \nu_k - \xi_k \nu_n}{\eta_k (\xi_k - \xi_n)} \right| = \log 4 \sum_{\substack{k \in I \\ k \neq n}} \left| \frac{\xi_k}{\eta_k} \right| \cdot \left| \frac{\xi_n \nu_k}{\xi_k (\xi_k - \xi_n)} - \frac{\nu_n}{\xi_k - \xi_n} \right| \\ &\leq \frac{\log 4}{1-\delta} \left(\sum_{\substack{k \in I \\ k \neq n}} \left| \frac{\xi_n \nu_k}{\xi_k (\xi_k - \xi_n)} \right| + \sum_{\substack{k \in I \\ k \neq n}} \left| \frac{\nu_n}{\xi_k - \xi_n} \right| \right) \end{aligned} \quad (3.22)$$

and

$$\begin{aligned}
& \left| \log \left(\left| \prod_{\substack{k \in I \\ k \neq n}} \frac{1 - \frac{\eta_n}{\eta_k}}{1 - \frac{\xi_n}{\xi_k}} \right| / \left| \prod_{\substack{k \in I \\ k \neq n}} \left(1 - \frac{\nu_n}{\eta_k} \right) \right| \right) \right| \\
&= \left| \log \left| \prod_{\substack{k \in I \\ k \neq n}} \left[1 - \left(1 - \frac{\nu_n}{\eta_k} \right)^{-1} \frac{\xi_k}{\eta_k} \cdot \frac{\xi_n(\nu_k - \nu_n)}{\xi_k(\xi_k - \xi_n)} \right] \right| \right| \\
&\leq \log 4 \sum_{\substack{k \in I \\ k \neq n}} \left| 1 - \frac{\nu_n}{\eta_k} \right|^{-1} \cdot \left| \frac{\xi_k}{\eta_k} \right| \cdot \left| \frac{\xi_n \nu_k - \xi_n \nu_n}{\xi_k(\xi_k - \xi_n)} \right| \\
&\leq \frac{\log 4}{1 - 2\delta} \left(\sum_{\substack{k \in I \\ k \neq n}} \left| \frac{\xi_n \nu_k}{\xi_k(\xi_k - \xi_n)} \right| + \sum_{\substack{k \in I \\ k \neq n}} \left| \frac{\xi_n \nu_n}{\xi_k(\xi_k - \xi_n)} \right| \right). \tag{3.23}
\end{aligned}$$

Step 3: estimating sums. It is relatively straightforward to estimate the expression that appears as the first sum in both (3.22) and (3.23). To this end we split the summation into two parts: the first inequality in (3.21) gives

$$\sum_{\substack{k \in I \\ \frac{\xi_k}{\xi_n} \notin (\frac{1}{\rho}, \rho)}} \left| \frac{\xi_n \nu_k}{\xi_k(\xi_k - \xi_n)} \right| \leq \frac{\rho}{\rho - 1} \sum_{\substack{k \in I \\ \frac{\xi_k}{\xi_n} \notin (\frac{1}{\rho}, \rho)}} \frac{|\nu_k|}{|\xi_k|} \leq \frac{\rho}{\rho - 1} S_1(n)$$

and, by the definition of $S_2(n)$, we have

$$\sum_{\substack{k \in I, k \neq n \\ \frac{\xi_k}{\xi_n} \in (\frac{1}{\rho}, \rho)}} \left| \frac{\xi_n \nu_k}{\xi_k(\xi_k - \xi_n)} \right| \leq \rho S_2(n).$$

Next, let us estimate the second sum in (3.22) under the assumption that $\frac{\Lambda(r)}{r}$ is non-increasing. If $\frac{\xi_k}{\xi_n} \in [-\frac{1}{\rho}, \frac{1}{\rho}]$, then $|\xi_k - \xi_n| = |\frac{\xi_k}{\xi_n} - 1| \cdot |\xi_n| \geq (1 - \frac{1}{\rho})|\xi_n|$. Hence

$$\sum_{\substack{k \in I \\ \frac{\xi_k}{\xi_n} \in [-\frac{1}{\rho}, \frac{1}{\rho}]}} \left| \frac{\nu_n}{\xi_k - \xi_n} \right| \leq \frac{\rho}{\rho - 1} \frac{|\nu_n|}{|\xi_n|} n_{\xi} \left(\frac{1}{\rho} |\xi_n| \right) \leq \frac{\rho}{\rho - 1} T(n).$$

If $\frac{\xi_k}{\xi_n} < -\frac{1}{\rho}$, then $|\xi_k - \xi_n| \geq |\xi_k|$ and $|\xi_k| \geq \frac{1}{\rho} |\xi_n| \geq R_0$. Hence, using that $\frac{\Lambda(r)}{r}$ is non-increasing on $[R_0, \infty)$ we obtain

$$\begin{aligned}
\sum_{\substack{k \in I \\ \frac{\xi_k}{\xi_n} < -\frac{1}{\rho}}} \left| \frac{\nu_n}{\xi_k - \xi_n} \right| &\leq |\nu_n| \sum_{\substack{k \in I \\ \frac{\xi_k}{\xi_n} < -\frac{1}{\rho}}} \frac{1}{|\xi_k|} = |\nu_n| \sum_{\substack{k \in I \\ \frac{\xi_k}{\xi_n} < -\frac{1}{\rho}}} \frac{\Lambda(|\xi_k|)}{|\xi_k|} \cdot \frac{1}{\Lambda(|\xi_k|)} \\
&\leq |\nu_n| \frac{\Lambda(\frac{1}{\rho} |\xi_n|)}{\frac{1}{\rho} |\xi_n|} \cdot \sum_{\substack{k \in I \\ \frac{\xi_k}{\xi_n} < -\frac{1}{\rho}}} \frac{1}{\Lambda(|\xi_k|)} \leq \rho S_3(n).
\end{aligned}$$

If $\frac{\xi_k}{\xi_n} \geq \rho$, then $|\xi_k - \xi_n| \geq (1 - \frac{1}{\rho})|\xi_k|$ and $|\xi_k| \geq \rho|\xi_n| \geq |\xi_n|$. We can again use the monotonicity of $\frac{\Lambda(r)}{r}$ for the estimate

$$\begin{aligned} \sum_{\substack{k \in I \\ \frac{\xi_k}{\xi_n} \geq \rho}} \left| \frac{\nu_n}{\xi_k - \xi_n} \right| &\leq \frac{\rho}{\rho - 1} |\nu_n| \sum_{\substack{k \in I \\ \frac{\xi_k}{\xi_n} \geq \rho}} \frac{1}{|\xi_k|} = \frac{\rho}{\rho - 1} |\nu_n| \sum_{\substack{k \in I \\ \frac{\xi_k}{\xi_n} \geq \rho}} \frac{\Lambda(|\xi_k|)}{|\xi_k|} \cdot \frac{1}{\Lambda(|\xi_k|)} \\ &\leq \frac{\rho}{\rho - 1} |\nu_n| \frac{\Lambda(|\xi_n|)}{|\xi_n|} \cdot \sum_{\substack{k \in I \\ \frac{\xi_k}{\xi_n} \geq \rho}} \frac{1}{\Lambda(|\xi_k|)} \leq \frac{\rho}{\rho - 1} S_3(n). \end{aligned}$$

By definition

$$\sum_{\substack{k \in I, k \neq n \\ \frac{\xi_k}{\xi_n} \in (\frac{1}{\rho}, \rho)}} \left| \frac{\nu_n}{\xi_k - \xi_n} \right| = S_4(n)$$

and hence

$$\sum_{\substack{k \in I \\ k \neq n}} \left| \frac{\nu_n}{\xi_k - \xi_n} \right| \leq \frac{\rho}{\rho - 1} T(n) + \left(\rho + \frac{\rho}{\rho - 1} \right) S_3(n) + S_4(n)$$

if $\frac{\Lambda(r)}{r}$ is non-increasing on $[R_0, \infty)$.

Finally, we have to estimate the second sum in (3.23) under the assumption that $\frac{\Lambda(r)}{r}$ is non-decreasing. Since $\frac{\Lambda(r)}{r^2}$ is decreasing on $[R_0, \infty)$ in this case by assumption, we can apply Lemma 3.8 with the growth function Λ , which yields

$$\sum_{\substack{k \in I \\ \frac{\xi_k}{\xi_n} \notin (\frac{1}{\rho}, \rho)}} \left| \frac{\xi_n \nu_n}{\xi_k (\xi_k - \xi_n)} \right| \leq \frac{\rho}{\rho - 1} |\nu_n| \sum_{\substack{k \in I \\ \frac{\xi_k}{\xi_n} \notin (\frac{1}{\rho}, \rho)}} \frac{\Lambda(|\xi_n|)}{|\xi_n| \Lambda(|\xi_k|)} \leq \frac{\rho}{\rho - 1} S_3(n).$$

Furthermore,

$$\sum_{\substack{k \in I, k \neq n \\ \frac{\xi_k}{\xi_n} \in (\frac{1}{\rho}, \rho)}} \left| \frac{\xi_n \nu_n}{\xi_k (\xi_k - \xi_n)} \right| \leq \rho |\nu_n| \sum_{\substack{k \in I, k \neq n \\ \frac{\xi_k}{\xi_n} \in (\frac{1}{\rho}, \rho)}} \frac{1}{|\xi_k - \xi_n|} = \rho S_4(n).$$

Putting these formulae together we obtain the required estimates (3.15) and (3.16). \square

3.2 Finishing the proof of Theorem 3.3

Throughout this subsection, let α, β, γ and Λ be as in Theorem 3.3.

3.10 Remark. In the next subsection (Section 3.3) we show that, for regularly distributed sequences, Hypothesis (B) in Theorem 3.3 can be weakened. Therefore it is important to keep track at which places (B) is used. We mark those places with \clubsuit . //

3.11 Remark. For the proof we may assume, without loss of generality, that $\rho_\Lambda < 2$. In fact, since, by Condition (S2), α must grow at least linearly, we can always use $\Lambda(r) = r(\log r)^2$, for which we have $\rho_\Lambda = 1$. Clearly, Condition

(A) becomes less restrictive if Λ is chosen smaller. In particular, if Λ_1, Λ_2 are two growth functions with $\rho_{\Lambda_1} < \rho_{\Lambda_2}$, then it follows from Remark 2.2 (iii) that $\frac{\Lambda_2(r)}{\Lambda_1(r)}$ is increasing for large r .

Moreover, note that if $\rho_\Lambda < 2$, then $\frac{\Lambda(r)}{r^2}$ is decreasing for large r again by Remark 2.2 (iii). //

The strategy is to cut off the sequences α and β at a very large radius r , split the product into two parts, apply the estimates for finite sequences from the previous subsection and show that these are uniform in r . Then we pass to the whole sequences again.

In order to be able to apply Proposition 3.9, we must make sure that the assumption (3.14) is satisfied. Let r_n and $J_n(c, \Lambda)$ be defined as in Proposition 3.6 using the sequence α instead of ξ , i.e.

$$r_n \Lambda(r_n) = |\alpha_n|, \quad J_n(c, \Lambda) = \{k \in \mathbb{N} : |\alpha_k| \leq cr_n\}.$$

3.12 Lemma. *Let Λ be a growth function and $R_0 > 0$ such that $\frac{\Lambda(r)}{r}$ is non-increasing or non-decreasing on $[R_0, \infty)$ and that $\frac{\Lambda(r)}{r^2}$ is decreasing on $[R_0, \infty)$. Moreover, let $\delta > 0$ and $\rho > 1$ be given.*

Then there exist $c > 0$, $r_0 \geq 1$ and $N_0 \in \mathbb{N}$ such that, for each $n \geq N_0$ and $r \geq |\alpha_n|$, the following data satisfy the hypotheses of Proposition 3.9 (i.e. it satisfies (3.14) and $n \in I$):

- $I := \{k \in \mathbb{N} : r_0 \leq |\alpha_k| \leq r\} \setminus J_n(c, \Lambda)$;
- $\xi := (\alpha_k)_{k \in I}$;
- Λ and R_0 ;
- ρ ;
- $\nu := (\gamma_k)_{k \in I}$;
- n .

Proof. Assumption (A) implies that $\frac{|\gamma_k|}{|\alpha_k|} = O\left(\frac{1}{\Lambda(|\alpha_k|)}\right)$, $k \rightarrow \infty$. In particular, $\lim_{k \rightarrow \infty} \frac{|\gamma_k|}{|\alpha_k|} = 0$. Moreover, by (3.3), we have

$$\lim_{k \rightarrow \infty} \frac{|\gamma_k|}{s_\alpha(k)} \clubsuit 0.$$

Hence we can choose $r_0 \geq \rho R_0$ such that

$$\frac{|\gamma_k|}{|\alpha_k|} \leq \delta \quad \text{and} \quad \frac{|\gamma_k|}{s_\alpha(k)} \leq \delta \quad \text{for all } k \text{ with } |\alpha_k| \geq r_0, \quad (3.24)$$

which shows that the second, third and fourth condition in (3.14) are satisfied.

Now set $c_1 := \sup_{k \in \mathbb{N}} \frac{|\gamma_k|}{|\alpha_k|} \Lambda(|\alpha_k|)$, define

$$c := \frac{c_1}{\delta},$$

and choose $N_0 \in \mathbb{N}$ such that

$$\forall n \geq N_0 : \quad \{k \in \mathbb{N} : |\alpha_k| < r_0\} \subseteq J_n(c, \Lambda) \quad \text{and} \quad n \notin J_n(c, \Lambda), \quad (3.25)$$

which is possible by items (vi) and (v) in the proof of Proposition 3.6. This shows, in particular, that $n \in I$.

Now let $k \in I$. Then $|\alpha_k| > cr_n$ since $k \notin J_n(c, \Lambda)$. Remembering that $r_n \geq \frac{|\alpha_n|}{\Lambda(|\alpha_n|)}$ by item (ii) in the proof of Proposition 3.6, we obtain that

$$|\gamma_n| \leq c_1 \frac{|\alpha_n|}{\Lambda(|\alpha_n|)} \leq c_1 r_n < \frac{c_1}{c} |\alpha_k| = \delta |\alpha_k|, \quad (3.26)$$

which proves the first condition in (3.14). \square

In order to proceed with the proof of Theorem 3.3, choose $\rho > 1$ such that the condition in (3.4) holds. \clubsuit

Define δ as in (3.13). Then $\delta > 0$, and we can use ρ and δ in Lemma 3.12. Note that the assumption on Λ in this lemma is satisfied because of Remark 3.11. If we use the data from Lemma 3.12 in Proposition 3.9, then we obtain the bounds in (3.15) or (3.16), depending on the monotonicity of $\frac{\Lambda(r)}{r}$. Of course, these bounds depend on r because the set I in Lemma 3.12 depends on r ; let us write $I(n; r)$ instead of I in the following in order to make this explicit. Moreover, we write

$$\Theta(n; r), T(n; r) \text{ etc.}$$

for the expressions in Proposition 3.9.

3.13 Lemma. *The following relations hold:*

$$\vartheta := \sup_{n \geq N_0} \sup_{\substack{r \in \mathbb{R} \\ r \geq |\alpha_n|}} \Theta(n; r) < \infty, \quad \tilde{\vartheta} := \sup_{n \geq N_0} \sup_{\substack{r \in \mathbb{R} \\ r \geq |\alpha_n|}} \tilde{\Theta}(n; r) < \infty.$$

Proof. Set $c_1 := \sup_{k \in \mathbb{N}} \frac{|\gamma_k|}{|\alpha_k|} \Lambda(|\alpha_k|)$, which is finite by Hypothesis (A). Then we have the following r -independent bounds (o and O notation is for $n \rightarrow \infty$):

$$\begin{aligned} S_1(n; r) &= \sum_{k \in I(n; r)} \left| \frac{\gamma_k}{\alpha_k} \right| \leq c_1 \sum_{k \in I(n; r)} \frac{1}{\Lambda(|\alpha_k|)} \leq c_1 \sum_{\substack{k \in \mathbb{N} \\ |\alpha_k| > cr_n}} \frac{1}{\Lambda(|\alpha_k|)} = o(1), \\ S_2(n; r) &= \sum_{\substack{k \in I(n; r), k \neq n \\ \frac{\alpha_k}{\alpha_n} \in (\frac{1}{\rho}, \rho)}} \frac{|\gamma_k|}{|\alpha_k - \alpha_n|} \leq \sum_{\substack{k \in I(n; r), k \neq n \\ \frac{\alpha_k}{\alpha_n} \in (\frac{1}{\rho}, \rho)}} \frac{|\gamma_k|}{s_\alpha(k)} \leq \sum_{\substack{k \in \mathbb{N} \\ |\alpha_k| > cr_n}} \frac{|\gamma_k|}{s_\alpha(k)} \stackrel{\clubsuit}{=} o(1), \\ S_3(n; r) &= \frac{|\gamma_n|}{|\alpha_n|} \Lambda(|\alpha_n|) \sum_{k \in I(n; r)} \frac{1}{\Lambda(|\alpha_k|)} \leq c_1 \sum_{\substack{k \in \mathbb{N} \\ |\alpha_k| > cr_n}} \frac{1}{\Lambda(|\alpha_k|)} = o(1), \\ S_4(n; r) &= |\gamma_n| \sum_{\substack{k \in I(n; r), k \neq n \\ \frac{\alpha_k}{\alpha_n} \in (\frac{1}{\rho}, \rho)}} \frac{1}{|\alpha_k - \alpha_n|} \leq |\gamma_n| \cdot r_\alpha(\rho, n) \frac{1}{s_\alpha(n)} \stackrel{\clubsuit}{=} O(1), \\ T(n; r) &= \frac{|\gamma_n|}{|\alpha_n|} n_\alpha(|\alpha_n|) \leq c_1 \frac{n_\alpha(|\alpha_n|)}{\Lambda(|\alpha_n|)} = O(1); \end{aligned}$$

the last estimate follows from Lemma 2.9. \square


3.14 Remark. If we slightly strengthen (3.4) to $\frac{|\gamma_n|}{s_\alpha(n)} = o(\frac{1}{r_\alpha(\rho, n)})$ and assume, in addition, that $n_\alpha(r) = o(\Lambda(r))$, then we may assert in the above lemma that

$$\lim_{n \rightarrow \infty} \sup_{\substack{r \in \mathbb{R} \\ r \geq |\alpha_n|}} \Theta(n; r) = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \sup_{\substack{r \in \mathbb{R} \\ r \geq |\alpha_n|}} \tilde{\Theta}(n; r) = 0,$$


respectively. //

3.15 Remark. The estimates used for $S_2(n; r)$ and $S_4(n; r)$ may seem rather careless. However, for wildly behaving sequences α , they can be sharp. Such sequences are for example constructed by mixing lumps of points of the following two kinds leaving sufficiently large empty intervals in between them:

- Type 1:



- Type 2:



//

Having available the uniform estimate from Lemma 3.13, we can now complete the proof of Theorem 3.3. We only have to take care of the following slight subtlety.

3.16 Remark. Let $l, m \in \mathbb{N}$ be such that $\alpha_l < \alpha_m$ are two consecutive members of the point set $\{\alpha_k : k \in \mathbb{N}\}$, and assume that $|\alpha_l|, |\alpha_m| \geq r_0$. By our choice of δ in (3.13), it follows from the second inequality in (3.24) that $\frac{|\gamma_k|}{s_\alpha(k)} < \frac{1}{2}$ for $k = l$ and $k = m$, and hence

$$\beta_l < \beta_m, \quad \alpha_l < \beta_m, \quad \beta_l < \alpha_m,$$

i.e. the pair $\{\alpha_l, \beta_l\}$ is separated from the pair $\{\alpha_m, \beta_m\}$. This implies that, for each sufficiently large r (where ‘ Δ ’ denotes the symmetric difference),

$$\#\left(\{k \in \mathbb{N} : |\alpha_k| \leq r\} \Delta \{k \in \mathbb{N} : |\beta_k| \leq r\}\right) \leq 2.$$

//

Let $r > r_0$ be given and assume that $\{k \in \mathbb{N} : r_0 \leq |\alpha_k| \leq r\} \neq \emptyset$. Then we define k_+ as the index with

$$\alpha_{k_+} = \max_{\alpha_k \leq r} \alpha_k.$$

If the set $\{k \in \mathbb{N} : \alpha_k > r\}$ is non-empty, we define k'_+ as the index with

$$\alpha_{k'_+} = \min_{\alpha_k > r} \alpha_k$$

and say that ' k'_+ exists'. If $\{k \in \mathbb{N} : \alpha_k > r\} = \emptyset$, we say that k'_+ does not exist. With this notation we set

$$Q_r^+(z) := \begin{cases} \left(1 - \frac{z}{\beta_{k_+}}\right)^{-1}, & \beta_{k_+} > r, \\ 1 - \frac{z}{\beta_{k'_+}}, & k'_+ \text{ exists and } \beta_{k'_+} \leq r, \\ 1, & \text{otherwise.} \end{cases}$$

Indices k_-, k'_- and a function $Q_r^-(z)$ are defined in the similar way, taking care of the left endpoint of the interval $[-r, r]$.

Proof of Theorem 3.3: the case when $\frac{\Lambda(r)}{r}$ is non-increasing. Let $n \geq N_0$ and $r \geq |\alpha_n|$. Then, using (3.25), we can write

$$\{k \in \mathbb{N} : |\alpha_k| \leq r\} = J_n(c, \Lambda) \cup \left[\{k \in \mathbb{N} : r_0 \leq |\alpha_k| \leq r\} \setminus J_n(c, \Lambda) \right],$$

and this is a disjoint union. In accordance with the paragraph before Lemma 3.13, we denote the second set in this union by $I(n; r)$. Hence

$$\begin{aligned} & \frac{-\frac{1}{\beta_n} \prod_{\substack{|\beta_k| \leq r \\ k \neq n}} \left(1 - \frac{\beta_n}{\beta_k}\right)}{-\frac{1}{\alpha_n} \prod_{\substack{|\alpha_k| \leq r \\ k \neq n}} \left(1 - \frac{\alpha_n}{\alpha_k}\right)} = \prod_{k \in J_n(c, \Lambda)} \frac{1 - \frac{\beta_n}{\beta_k}}{1 - \frac{\alpha_n}{\alpha_k}} \times \frac{\prod_{\substack{k \in I(n; r) \\ k \neq n}} \left(1 - \frac{\beta_n}{\beta_k}\right)}{\prod_{\substack{k \in I(n; r) \\ k \neq n}} \left(1 - \frac{\alpha_n}{\alpha_k}\right)} \times \\ & \quad \times \frac{\alpha_n}{\beta_n} \times Q_r^+(\beta_n) Q_r^-(\beta_n). \end{aligned}$$

If $r \rightarrow \infty$, then the left-hand side tends to $\frac{P'_\beta(\beta_n)}{P'_\alpha(\alpha_n)}$, cf. (3.6). The first and the third factors on the right-hand side do not depend on r . By Proposition 3.9, the second factor remains bounded from above and away from zero with the bounds $\exp \vartheta_n$ and $\exp(-\vartheta_n)$, respectively, where

$$\vartheta_n := \sup_{\substack{r \in \mathbb{R} \\ r \geq |\alpha_n|}} \Theta(n; r),$$

which is finite and uniformly bounded in n by Lemma 3.13. The last factor tends to 1. We conclude that

$$\left| \prod_{k \in J_n(c, \Lambda)} \frac{1 - \frac{\beta_n}{\beta_k}}{1 - \frac{\alpha_n}{\alpha_k}} \right| e^{-\vartheta_n} \frac{|\alpha_n|}{|\beta_n|} \leq \left| \frac{P'_\beta(\beta_n)}{P'_\alpha(\alpha_n)} \right| \leq \left| \prod_{k \in J_n(c, \Lambda)} \frac{1 - \frac{\beta_n}{\beta_k}}{1 - \frac{\alpha_n}{\alpha_k}} \right| e^{\vartheta_n} \frac{|\alpha_n|}{|\beta_n|}.$$

Now let n tend to ∞ . The last factors on the left- and the right-hand sides tend to 1. The respective second factors remain bounded, namely on the right-hand side by e^ϑ from above and on the left-hand side by $e^{-\vartheta}$ from below; for the definition of ϑ see Lemma 3.13. By Proposition 3.6, the first factor remains bounded from above and away from zero. \square

3.17 Remark. Maybe it is good to pause and review the proof (thereby also explain the necessity to be precise about constants in O-estimates).

For each fixed n we cut the product into two pieces. In order to have a bound of the partial product corresponding to large indices which is uniform in n , we use some uniform estimate of the product by the corresponding sum (we decided to use the ‘log 4’-estimate in Remark 3.7). Application of this uniform estimate is, however, only legitimate if we cut at a point that is so large that (3.14) holds.

The earliest legitimate cutting point depends on n . Since the perturbation γ need not be bounded, it will in general increase to infinity with n . And now Proposition 3.6 and (3.26) come into play, which say that we can control beginning sections of the (n -dependent) product $\frac{P'_\beta(\beta_n)}{P'_\alpha(\alpha_n)}$ which are getting longer sufficiently fast. \parallel

Proof of Theorem 3.3: the case when $\frac{\Lambda(r)}{r}$ is non-decreasing. Using the same argument as in the previous part of the proof with (3.16) instead of (3.15) and $\tilde{\vartheta}$ instead of ϑ we obtain that the quotient

$$\frac{|P'_\beta(\beta_n)|}{|P'_\alpha(\alpha_n)|} \bigg/ \left| \prod_{k \notin J_n(c, \Lambda)} \left(1 - \frac{\gamma_n}{\beta_k}\right) \right|$$

is bounded from above and away from zero uniformly in n .

The limit $\lim_{n \rightarrow \infty} \prod_{k \notin J_n(c, \Lambda)} \left(1 - \frac{z}{\beta_k}\right)$ exists locally uniformly on \mathbb{C} and equals 1. Since $\frac{\Lambda(r)}{r}$ is non-decreasing, Condition (A) implies that γ is bounded. Thus

$$\lim_{n \rightarrow \infty} \prod_{k \notin J_n(c, \Lambda)} \left(1 - \frac{\gamma_n}{\beta_k}\right) = 1,$$

which proves the assertion also in this case. \square

3.3 Supplements: regularly distributed sequences

For regularly distributed sequences α we have more control over s_α and r_α . In addition, we can give better estimates for $S_2(n)$ and $S_4(n)$ from Proposition 3.9 since irregular behaviour as in Remark 3.15 is ruled out.

The conditions required in Theorem 3.3 do not depend on the arrangement of α as long as β is arranged in the same way. In the theorems below, however, we need to refer to a specific arrangement of α . For this we need one more notation. Let $\alpha \in \mathbb{S}$ and consider the (finite or infinite) subsequence α^+ of α consisting of all positive elements of α arranged increasingly as in Definition 3.1. For each n such that the term α_n^+ exists, let $\chi^+(n)$ be the unique index with $\alpha_n^+ = \alpha_{\chi^+(n)}$. If α^+ is an infinite sequence, then χ^+ is a bijective map from \mathbb{N} onto $\{n \in \mathbb{N} : \alpha_n > 0\}$. Let $\nu^+ : \{n \in \mathbb{N} : \alpha_n > 0\} \rightarrow \mathbb{N}$ be its inverse. Note

that $\alpha_n = \alpha_{\nu^+(n)}^+$ if $\alpha_n > 0$. Similar notation is used for the subsequence of negative terms of α .

First, we assume only a mild regularity property of α .

3.18 Theorem. *Let α and Λ be given as in Theorem 3.3. Further, let $\beta \in \mathbb{S}$ be a small perturbation of α in the sense that $\gamma := \beta - \alpha$ satisfies (A) from Theorem 3.3 and the following conditions.*

(C1) *If α contains infinitely many positive terms, then there exists a growth function λ^+ such that α^+ has finite upper and positive lower λ^+ densities.*

If α contains infinitely many negative terms, then there exists a growth function λ^- such that α^- has finite upper and positive lower λ^- densities.

$$(C2) \quad \frac{|\gamma_{\chi^+(n)}|}{s_{\alpha^+}(n)} = O\left(\frac{1}{n}\right), \quad \frac{|\gamma_{\chi^-(n)}|}{s_{\alpha^-}(n)} = O\left(\frac{1}{n}\right), \quad n \rightarrow \infty. \quad (3.27)$$

Then (3.5) holds.

The condition (3.27) is of course only a minor weakening of (3.3), but the condition (3.4) can be dropped (in fact, it holds automatically).

Proof of Theorem 3.18. We have to check all places where Condition (B) was used (which were marked with ‘♣’).

(1) *Proof of Lemma 3.12:* Clearly, (3.27) implies that $\lim_{k \rightarrow \infty} \frac{|\gamma_k|}{s_\alpha(k)} = 0$.

(2) *Choice of ρ :* Choose $\rho > 1$ arbitrarily.

(3) *Proof of Lemma 3.13:* We have to provide suitable bounds for $S_2(n; r)$ and $S_4(n; r)$. We consider the case when n ranges over those indices with $\alpha_n > 0$; the set of indices n with $\alpha_n < 0$ is treated in the same way.

If the sequence α contains only finitely many positive elements, it is clear that $S_2(n; r)$ and $S_4(n; r)$ remain bounded. Hence, assume that α^+ is an infinite sequence. Let $n \in \mathbb{N}$ such that $\alpha_n > 0$. Then, clearly,

$$\left\{ \alpha_k : \frac{\alpha_k}{\alpha_n} \in \left(\frac{1}{\rho}, \rho\right) \right\} = \left\{ \alpha_l^+ : \frac{\alpha_l^+}{\alpha_{\nu^+(n)}^+} \in \left(\frac{1}{\rho}, \rho\right) \right\},$$

and this shows that $r_\alpha(\rho, n) = r_{\alpha^+}(\rho, \nu^+(n))$. Unless $\nu^+(n) = 1$, we also have $s_\alpha(n) = s_{\alpha^+}(\nu^+(n))$.

Using (3.27) and Lemma 2.7 we obtain (with $c := \sup_{n \in \mathbb{N}} \frac{|\gamma_{\chi^+(n)}|}{s_{\alpha^+}(n)} n$)

$$S_2(n; r) \leq \sum_{\substack{k \in \mathbb{N} \\ \frac{\alpha_k}{\alpha_n} \in (\frac{1}{\rho}, \rho)}} \frac{|\gamma_k|}{s_\alpha(k)} = \sum_{\substack{l \in \mathbb{N} \\ \frac{\alpha_l^+}{\alpha_{\nu^+(n)}^+} \in (\frac{1}{\rho}, \rho)}} \frac{|\gamma_{\chi^+(l)}|}{s_{\alpha^+}(l)} \leq c \sum_{\substack{l \in \mathbb{N} \\ \frac{\alpha_l^+}{\alpha_{\nu^+(n)}^+} \in (\frac{1}{\rho}, \rho)}} \frac{1}{l} = O(1),$$

$$S_4(n; r) \leq \frac{|\gamma_n| \cdot r_\alpha(\rho, n)}{s_\alpha(n)} = \frac{|\gamma_{\chi^+(\nu^+(n))}|}{s_{\alpha^+}(\nu^+(n))} \cdot r_{\alpha^+}(\rho, \nu^+(n)) \lesssim \frac{1}{\nu^+(n)} \cdot \nu^+(n) = 1$$

as $n \rightarrow \infty$ uniformly in r . \square

Second, we assume that α behaves quite regularly. For the definition of the standard sequence λ corresponding to a growth function λ see Section 2.2.

3.19 Theorem. *Let α and Λ be given as in Theorem 3.3. Further, let $\beta \in \mathbb{S}$ be a small perturbation of α in the sense that $\gamma := \beta - \alpha$ satisfies (A) from Theorem 3.3 and the following conditions.*

(D1) *If α contains infinitely many positive terms, then there exists a growth function λ^+ such that $\alpha^+ = \lambda^+$.*

If α contains infinitely many negative terms, then there exists a growth function λ^- such that $\alpha^- = -\lambda^-$.

$$(D2) \quad |\gamma_{\chi^+(n)}| = O\left(\frac{\alpha_n^+}{n \log n}\right), \quad |\gamma_{\chi^-(n)}| = O\left(\frac{|\alpha_n^-|}{n \log n}\right), \quad n \rightarrow \infty.$$

Then (3.5) holds.

Proof. Also here we just have to check the places with \clubsuit . Again, we restrict the explicit proof to the sequence α^+ and to the case when this sequence is infinite.

(1) *Proof of Lemma 3.12:* Using the first formula in Lemma 2.12 and (D2) we obtain

$$\frac{\gamma_n^+}{s_{\alpha^+}(n)} \lesssim \frac{\alpha_n^+}{n \log n} \cdot \frac{n}{\alpha_n^+} = \frac{1}{\log n} = o(1), \quad n \rightarrow \infty.$$

(2) *Choice of ρ :* Choose $\rho > 1$ arbitrarily.

(3) *Proof of Lemma 3.13:* For $n \in \{l \in \mathbb{N} : \alpha_l > 0\}$ and $r > 0$, we can use the second formula in Lemma 2.12 to estimate the following sum:

$$\begin{aligned} \sum_{\substack{k \in I(n;r), k \neq n \\ \frac{\alpha_k}{\alpha_n} \in (\frac{1}{\rho}, \rho)}} \frac{1}{|\alpha_k - \alpha_n|} &\leq \sum_{\substack{k \in \mathbb{N}, k \neq n \\ \frac{\alpha_k}{\alpha_n} \in (\frac{1}{\rho}, \rho)}} \frac{1}{|\alpha_k - \alpha_n|} = \sum_{\substack{l \in \mathbb{N}, l \neq \nu^+(n) \\ \frac{\alpha_l^+}{\alpha_{\nu^+(n)}^+} \in (\frac{1}{\rho}, \rho)}} \frac{1}{|\alpha_l^+ - \alpha_{\nu^+(n)}^+|} \\ &\leq C \frac{\nu^+(n) \log(\nu^+(n))}{\alpha_{\nu^+(n)}^+} \end{aligned}$$

with some $C > 0$ independent of n . This together with (D2) yields

$$S_4(n; r) = |\gamma_{\chi^+(\nu^+(n))}| \sum_{\substack{k \in I(n;r), k \neq n \\ \frac{\alpha_k}{\alpha_n} \in (\frac{1}{\rho}, \rho)}} \frac{1}{|\alpha_k - \alpha_n|} = O(1)$$

for $n \rightarrow \infty$ uniformly in r . For $S_2(n; r)$ we can estimate

$$\begin{aligned} S_2(n; r) &= \sum_{\substack{k \in I(n;r), k \neq n \\ \frac{\alpha_k}{\alpha_n} \in (\frac{1}{\rho}, \rho)}} \left| \frac{\gamma_k}{\alpha_k - \alpha_n} \right| \leq \max_{\substack{k \in I(n;r), k \neq n \\ \frac{\alpha_k}{\alpha_n} \in (\frac{1}{\rho}, \rho)}} |\gamma_k| \cdot \sum_{\substack{k \in I(n;r), k \neq n \\ \frac{\alpha_k}{\alpha_n} \in (\frac{1}{\rho}, \rho)}} \frac{1}{|\alpha_k - \alpha_n|} \\ &\leq \max_{\substack{l \in \mathbb{N}, l \neq \nu^+(n) \\ \frac{\alpha_l^+}{\alpha_{\nu^+(n)}^+} \in (\frac{1}{\rho}, \rho)}} |\gamma_{\chi^+(l)}| \cdot C \frac{\nu^+(n) \log(\nu^+(n))}{\alpha_{\nu^+(n)}^+}. \end{aligned} \quad (3.28)$$

In order to estimate the maximum, let $l \in \mathbb{N}$ such that $\frac{\alpha_l^+}{\alpha_{\nu^+(n)}^+} \in (\frac{1}{\rho}, \rho)$. Then $\alpha_l^+ > \frac{1}{\rho} \alpha_{\nu^+(n)}^+$ and hence, by Lemma 2.11 (ii) and Remark 2.2 (i),

$$l > n_{\alpha^+} \left(\frac{1}{\rho} \alpha_{\nu^+(n)}^+ \right) \geq \left\lceil \lambda^+ \left(\frac{1}{\rho} \alpha_{\nu^+(n)}^+ \right) \right\rceil \gtrsim \lambda^+ (\alpha_{\nu^+(n)}^+) = \nu^+(n),$$

which gives (with some $C' > 0$)

$$|\gamma_{\lambda^+(l)}| \leq C' \frac{\alpha_l^+}{l \log l} \leq C' \frac{\rho \alpha_{\nu^+(n)}^+}{l \log l} \lesssim \frac{\alpha_{\nu^+(n)}^+}{\nu^+(n) \log(\nu^+(n))}.$$

This together with (3.28) implies that $S_2(n; r)$ is bounded uniformly in n and r . \square

Under slightly stronger assumptions on Λ , λ^+ and λ^- one can even drop Condition (D2) as the following corollary shows.

3.20 Corollary. *Let α and Λ be given as in Theorem 3.3. Further, let $\beta \in \mathbb{S}$ be a small perturbation of α in the sense that $\gamma := \beta - \alpha$ satisfies (A) from Theorem 3.3 and the following condition.*

(E) *If α contains infinitely many positive terms, then there exists a growth function λ^+ such that $\alpha^+ = \lambda^+$, that $\frac{\lambda^+(r)}{\Lambda(r)}$ is non-increasing for large r and that either $\rho_{\lambda^+} > 0$ or $\frac{\log \lambda^+(r)}{\log r}$ is non-increasing for large r .*

If α contains infinitely many negative terms, then there exists a growth function λ^- such that $\alpha^- = -\lambda^-$, that $\frac{\lambda^-(r)}{\Lambda(r)}$ is non-increasing for large r and that either $\rho_{\lambda^-} > 0$ or $\frac{\log \lambda^-(r)}{\log r}$ is non-increasing for large r .

Then (3.5) holds.

3.21 Remark. As already mentioned in Remark 2.14 the assumptions in (E) are satisfied in most cases if α^+ and α^- are standard sequences, cf. that remark for more details.

In particular, consider the situation that either α^- is the empty sequence (i.e. $\alpha_n > 0$ for all $n \in \mathbb{N}$) or that $\alpha^- = -\alpha^+$ (i.e. the set $\{\alpha_n : n \in \mathbb{N}\}$ is symmetric). Moreover, assume that α^+ is a standard sequence: $\alpha^+ = \lambda^+ =: \lambda$ and that either $\rho_\lambda > 0$ or $\frac{\log \lambda(r)}{\log r}$ is non-increasing for large r . In this case we can choose $\Lambda := \lambda(r) \cdot \log \lambda(r) \cdot (\log \log \lambda(r))^2$ for large r . Then Condition (E) is satisfied and α is of convergence class with respect to Λ since

$$\sum_{n \in \mathbb{N}} \frac{1}{\Lambda(\lambda_n)} = \sum_{n \in \mathbb{N}} \frac{1}{n \log n \cdot (\log \log n)^2} < \infty.$$

Hence, if γ satisfies Condition (A) from Theorem 3.3, i.e.

$$|\gamma_{\lambda^\pm(n)}| = O\left(\frac{\alpha_n^+}{n \log n \cdot (\log \log n)^2}\right), \quad n \rightarrow \infty,$$

then (3.5) holds. $\quad //$

3.22 Example. In order to illustrate the power of this supplement, let us reconsider the example already discussed in Remark 3.5. That is

$$\alpha_n := n^\sigma \quad \text{with } \sigma > 1, \quad \Lambda(r) := r^{\frac{1}{\sigma}} \log r \cdot (\log \log r)^2, \quad \text{for large } r.$$

Then Corollary 3.20 (see also Remark 3.21) states that

$$\frac{|\gamma_n|}{n^{\sigma-1}} = O\left(\frac{1}{\log n \cdot (\log \log n)^2}\right)$$

is already enough to have (3.5). Theorem 3.18 requires

$$\frac{|\gamma_n|}{n^{\sigma-1}} = O\left(\frac{1}{n}\right),$$

and Theorem 3.3 even requires

$$\sum_{n=1}^{\infty} \frac{|\gamma_n|}{n^{\sigma-1}} < \infty.$$

This comparison also reflects the fact that the regularity assumption in Theorem 3.18 is very weak, whereas the one in Theorem 3.19 is quite strong.

One can also consider a linearly growing sequence α_n . In this case the sequence must be symmetric because of Condition (S2) in Definition 3.1. Consider α_n such that $\alpha_n^+ = n$, $\alpha_n^- = -n$, $n \in \mathbb{N}$. Then Corollary 3.20 (see also Remark 3.21) implies that

$$|\gamma_{\chi^+(n)}| = O\left(\frac{1}{\log n \cdot (\log \log n)^2}\right), \quad |\gamma_{\chi^-(n)}| = O\left(\frac{1}{\log n \cdot (\log \log n)^2}\right)$$

is sufficient for (3.5) to hold.

We note that one can apply the theorems also to very sparse sequences like exponentially growing ones. //

3.4 Symmetry of conditions

We close this section with a general note on the nature of the conditions appearing in our results.

In Theorems 3.3, 3.18 and 3.19 we consider β as a perturbation of α by γ , and the conditions (A), (B) etc. relate the perturbation γ to the sequence α . Strictly speaking we must therefore say that *the ordered pair (α, β) satisfies the given conditions*.

Actually, Theorems 3.3 and 3.18 are symmetric in α and β as the following proposition shows.

3.23 Proposition. *Let $\alpha, \beta \in \mathbb{S}$ and assume that the hypotheses of one of Theorems 3.3 and 3.18 are fulfilled for the pair (α, β) . Then also the pair (β, α) satisfies the corresponding hypotheses.*

Proof. It follows from (3.2) that $\lim_{n \rightarrow \infty} \frac{\beta_n}{\alpha_n} = 1$. Hence $\Lambda(|\alpha_n|) \asymp \Lambda(|\beta_n|)$ by Remark 2.2 (i), and we see that (3.1) holds for β . Moreover, also (3.2) holds with β and $-\gamma$ in place of α and γ . For sufficiently large n , the numbers α_n

and β_n have the same sign. Remembering Lemma 2.5 we thus obtain that (C1) also holds for β (with the same growth functions λ^+ and λ^-).

For the conditions in (B) and (C2) it is not so obvious that the roles of α and β can be exchanged. First we investigate the relation between s_α and s_β . Let us show that $s_\alpha(n) \lesssim s_\beta(n)$. Assume that α and γ satisfy (B) or (C2). Let us consider positive β_n and assume that there are infinitely many positive β_k and hence also infinitely many positive α_k . Choose $n_0 \in \mathbb{N}$ such that

$$\frac{|\gamma_k|}{s_\alpha(k)} \leq \frac{1}{4}, \quad k \geq n_0,$$

and let n_1 be such that $\beta_{n_1} \geq \beta_k$ for all $k < n_0$ and $\beta_{n_1} > 0$. Moreover, choose $n_2 > n_1$ such that $|\beta_k| > \beta_{n_1}$ for all $k \geq n_2$. Now let $n \geq n_2$ be such that $\beta_n > 0$. Then, for $k \geq n_0$,

$$\begin{aligned} |\beta_k - \beta_n| &= |\alpha_k - \alpha_n + \gamma_k - \gamma_n| \geq |\alpha_k - \alpha_n| - |\gamma_k| - |\gamma_n| \\ &\geq |\alpha_k - \alpha_n| - \frac{1}{4}s_\alpha(k) - \frac{1}{4}s_\alpha(n) \\ &\geq |\alpha_k - \alpha_n| - \frac{1}{4}|\alpha_n - \alpha_k| - \frac{1}{4}|\alpha_k - \alpha_n| = \frac{1}{2}|\alpha_k - \alpha_n|, \end{aligned}$$

and hence

$$s_\beta(n) = \min\{|\beta_k - \beta_n| : k \geq n_0, k \neq n\} \geq \frac{1}{2}s_\alpha(n).$$

The proof for negative β_n is similar and therefore $s_\alpha(n) \lesssim s_\beta(n)$, $n \rightarrow \infty$. This relation also implies that $\lim_{n \rightarrow \infty} \frac{|\gamma_n|}{s_\beta(n)} = 0$. With this property one can prove in the same way as above that $s_\alpha(n) \geq \frac{1}{2}s_\beta(n)$ for all sufficiently large indices n . Together, thus $s_\beta(n) \asymp s_\alpha(n)$.

Now it is clear that the condition (3.3) also holds for the sequence β and the perturbation $-\gamma$. Again using that, for all sufficiently large n , the points α_n and β_n have the same sign, we can conclude that (3.27) holds with β and $-\gamma$ in place of α and γ .

Next, we turn to the relation between $r_\alpha(\rho, n)$ and $r_\beta(\rho, n)$. Let $\rho' \in (1, \rho)$, and choose $n_0 \in \mathbb{N}$ such that

$$\left(\frac{\rho'}{\rho}\right)^{\frac{1}{2}} \leq \frac{\beta_k}{\alpha_k} \leq \left(\frac{\rho}{\rho'}\right)^{\frac{1}{2}}, \quad k \geq n_0.$$

Moreover, let $n_1 \geq n_0$ be such that $\frac{1}{\rho'}|\beta_n| \geq \max_{l \leq n_0} |\beta_l|$ for $n \geq n_1$. Now fix $n \geq n_1$. If $k \in \mathbb{N}$ is such that $\frac{\beta_k}{\beta_n} \in (\frac{1}{\rho'}, \rho')$, then $k \geq n_0$ and therefore

$$\frac{\alpha_k}{\alpha_n} = \frac{\alpha_k}{\beta_k} \cdot \frac{\beta_n}{\alpha_n} \cdot \frac{\beta_k}{\beta_n} \begin{cases} \leq \left(\frac{\rho'}{\rho}\right)^{-\frac{1}{2}} \left(\frac{\rho}{\rho'}\right)^{\frac{1}{2}} \rho' = \rho, \\ \geq \left(\frac{\rho}{\rho'}\right)^{-\frac{1}{2}} \left(\frac{\rho'}{\rho}\right)^{\frac{1}{2}} \frac{1}{\rho'} = \frac{1}{\rho}. \end{cases}$$

Hence

$$\left\{k \in \mathbb{N} : \frac{\beta_k}{\beta_n} \in \left(\frac{1}{\rho'}, \rho'\right)\right\} \subseteq \left\{k \in \mathbb{N} : \frac{\alpha_k}{\alpha_n} \in \left(\frac{1}{\rho}, \rho\right)\right\},$$

and this gives $r_\beta(\rho', n) \leq r_\alpha(\rho, n)$. Together with the already proved fact about separations, it follows that (3.4) holds for β and $-\gamma$. \square

Theorem 3.19 is not symmetric in α and β , but this is only a matter of formulation. In fact, we can restate Theorem 3.19 as follows.

3.24 Theorem. *Let α , β and Λ be given as in Theorem 3.3. Assume that the following condition is satisfied.*

(F) *The sequence α contains infinitely many positive terms if and only if β does. In this case there exists a growth function λ^+ such that*

$$\begin{aligned} |\alpha_n^+ - \lambda_n^+| &= O\left(\frac{\lambda_n^+}{\Lambda(\lambda_n^+)}\right), & |\beta_n^+ - \lambda_n^+| &= O\left(\frac{\lambda_n^+}{\Lambda(\lambda_n^+)}\right), & n &\rightarrow \infty, \\ |\alpha_n^+ - \lambda_n^+| &= O\left(\frac{\lambda_n^+}{n \log n}\right), & |\beta_n^+ - \lambda_n^+| &= O\left(\frac{\lambda_n^+}{n \log n}\right), & n &\rightarrow \infty. \end{aligned}$$

If α contains infinitely many negative terms, the analogous statement holds.

Then (3.5) holds.

We decided to present the ‘asymmetric formulation’ of Theorem 3.19 as the principle formulation in order to emphasize that strong regularity is assumed.

4 A selection of applications

4.1 The inverse spectral problem for a string

A string is a pair $S[L, m]$ that consists of a number L , $0 < L \leq \infty$, and a non-negative and non-decreasing function m defined on $[0, L)$. The string $S[L, m]$ gives rise to an operator model, namely the Kreĭn–Feller differential operator $-D_m D_x$ acting in the space $L^2(dm)$. The eigenvalue equation for one of its self-adjoint realizations can be written in integral form as

$$\begin{cases} f(x) - f(0) + z \int_{[0, x]} (x-y)f(y) dm(y) = 0, & x \in (0, L), \\ f'(0-) = 0, \end{cases}$$

where $z \in \mathbb{C}$ is the eigenvalue parameter; if the limit circle case prevails at L , i.e. if $\int_{[0, L)} x^2 dm(x) < \infty$, then a boundary condition is also needed at L . The operator $-D_m D_x$ arises when Fourier’s method is applied to the partial differential equation

$$\frac{\partial}{\partial m(s)} \left(\frac{\partial v(s, t)}{\partial s} \right) - \frac{\partial^2}{\partial t^2} v(s, t) = 0$$

describing the vibrations of an inhomogeneous string with mass distribution m and a free left endpoint.

The spectrum of the Kreĭn–Feller operator $-D_m D_x$ is fully described by one analytic function: the principle Titchmarsh–Weyl coefficient q_S associated with the string $S = S[L, m]$; see, e.g. [KaKr]. This function belongs to the Stieltjes class \mathcal{S} , i.e. q_S is analytic in the region $\mathbb{C} \setminus [0, \infty)$, has non-negative imaginary

part throughout the upper half-plane, and takes non-negative values along the negative real half-line. A Fourier transform can be constructed which maps $-D_m D_x$ to the multiplication operator by the independent variable in $L^2(\mu_S)$, where μ_S is the measure in the representation of q_S as a Cauchy integral:

$$q_S(z) = a + \int_{\mathbb{R}} \frac{d\mu_S(t)}{t-z}. \quad (4.1)$$

The cornerstones of the spectral theory of strings were established by M. G. Kreĭn in the early 1950s, see [Kr3] or [KaKr, Theorem 11.2]⁸; a presentation from a slightly different viewpoint can be found in [DK]. Most notably, an inverse spectral theorem was proved. For this we have to normalize L and m , i.e. we assume that $m(0) = 0$, that m is left continuous and that $m(x) < \sup_{t \in [0, L]} m(t)$ for all $x \in [0, L)$. Then the inverse spectral theorem reads as follows.

For each function $q \in \mathcal{S}$, there exists a unique string $S[L, m]$ such that q is the principle Titchmarsh–Weyl coefficient of $S[L, m]$.

The number a in (4.1) gives the length of a massless initial section of the string, i.e. $a = \sup\{x : m(x) = 0\}$. In the following we assume that $a = 0$, i.e. that $m(x) > 0$ for $x > 0$.

A string $S[L, m]$ is called *regular* if

$$L < \infty \quad \text{and} \quad m(L) := \lim_{x \nearrow L} m(x) < \infty,$$

and *singular* otherwise⁹. Thinking of direct and inverse spectral relations, the problem arises to describe the totality \mathcal{S}_{reg} of all Stieltjes class functions that are principle Titchmarsh–Weyl coefficients of regular strings. The solution of this problem is known. It follows from [Kr2], see also [KaKr, 11.11^o]¹⁰.

4.1 Theorem ([Kr2]). *Let μ be a positive measure with $\text{supp } \mu \subseteq (0, \infty)$ and $\int_{\mathbb{R}} \frac{d\mu(t)}{1+t} < \infty$. Consider the Cauchy transform $q_\mu(z) := \int_{\mathbb{R}} \frac{d\mu(t)}{t-z}$ of μ , and let $S[L, m]$ be the string whose principle Titchmarsh–Weyl coefficient is equal to q_μ . Then $S[L, m]$ is regular if and only if*

(i) *the measure μ is discrete, say $\mu = \sum_{n=1}^N \sigma_n \delta_{\alpha_n}$ with $N \in \mathbb{N} \cup \{\infty\}$, $\sigma_n > 0$ and $0 < \alpha_1 < \alpha_2 < \dots$;*

(ii) *the limit $\lim_{n \rightarrow \infty} \frac{n}{\sqrt{\alpha_n}}$ exists and is finite (we tacitly understand this limit as 0 if N is finite);*

(iii) *we have*

$$\sum_{n=1}^N \frac{1}{\alpha_n^2 (P'_\alpha(\alpha_n))^2 \sigma_n} < \infty.$$

It is clear that a sequence α satisfying (i)–(iii) belongs to \mathbb{S} introduced in Definition 3.1. Applying Theorem 3.3 (or Theorems 3.18, 3.19) we immediately obtain a stability result for the class \mathcal{S}_{reg} , which says that sufficiently small shifts of poles do not lead out of the class \mathcal{S}_{reg} . The precise formulation reads as follows.

⁸For the reason of physical interpretation, in [KaKr] the principle Titchmarsh–Weyl coefficient is called ‘coefficient of dynamic compliance’.

⁹In [DK] the maybe more descriptive terminology ‘short’ and ‘long’ is used.

¹⁰For simplicity we restrict ourselves to the case that $0 \notin \text{supp } \mu$.

4.2 Theorem. Let μ be a discrete measure, $\mu = \sum_{n=1}^{\infty} \sigma_n \delta_{\alpha_n}$, with

$$0 < \alpha_1 < \alpha_2 < \dots, \quad \lim_{n \rightarrow \infty} \frac{n}{\sqrt{\alpha_n}} \in [0, \infty), \quad (4.2)$$

$$\sigma_n > 0, \quad \sum_{n=1}^{\infty} \frac{\sigma_n}{\alpha_n} < \infty,$$

and set $q_\mu(z) := \int_{\mathbb{R}} \frac{d\mu(t)}{t-z}$.

Let $\gamma = (\gamma_n)_{n \in \mathbb{N}}$ be a sequence of real numbers such that the points $\beta_n := \alpha_n + \gamma_n$ are all positive and pairwise distinct, let τ_n , $n \in \mathbb{N}$, be positive real numbers, and set

$$\nu = \sum_{n=1}^{\infty} \tau_n \delta_{\beta_n}, \quad q_\nu(z) := \int_{\mathbb{R}} \frac{d\nu(t)}{t-z}.$$

If the hypotheses of one of Theorems 3.3, 3.18, 3.19 are fulfilled and $\tau_n \asymp \sigma_n$, then

$$q_\mu \in \mathcal{S}_{\text{reg}} \iff q_\nu \in \mathcal{S}_{\text{reg}}.$$

4.2 The Kreĭn class of entire functions

Let f be an entire function with $f(0) = 1$, and denote its sequence of zeros by $\alpha = (\alpha_n)_{n \in \mathbb{N}}$, which are assumed to be simple. Then f is said to belong to the *Kreĭn class* \mathbb{K} if

$$(\mathbf{Kr1}) \quad \sum_{n \in \mathbb{N}} \left| \operatorname{Im} \frac{1}{\alpha_n} \right| < \infty;$$

(**Kr2**) there exists a number $l \in \mathbb{N}$ such that

$$\sum_{n \in \mathbb{N}} \frac{1}{|\alpha_n|^l |f'(\alpha_n)|} < \infty, \quad (4.3)$$

and, on the domain $\mathbb{C} \setminus \{\alpha_n : n \in \mathbb{N}\}$, the function $\frac{1}{f}$ is represented as

$$\frac{1}{f(z)} = p(z) + \sum_{n \in \mathbb{N}} \frac{1}{f'(\alpha_n)} \left(\frac{1}{z - \alpha_n} + \frac{1}{\alpha_n} + \dots + \frac{z^{l-2}}{\alpha_n^{l-1}} \right) \quad (4.4)$$

with some polynomial p .

This class was first studied by M. G. Kreĭn in the 1940s. It appears, e.g. in the context of extension theory of symmetric operators in a Hilbert space, cf. [Kr1]; for properties of functions in \mathbb{K} see, e.g. [L1, Section V.6]. M. G. Kreĭn proved that every function from the class \mathbb{K} is of exponential type and has completely regular growth; see, e.g. [L1, Theorem V.13].

Often functions that have only real zeros are of particular interest. We therefore introduce the following subclasses.

4.3 Definition. Let $l \in \mathbb{N}$. We say that a function f belongs to the class \mathbb{K}_l if

- (i) f is entire, takes real values along the real axis, satisfies $f(0) = 1$ and has only real and simple zeros which are all non-zero;

(ii) the condition (4.3) and the representation (4.4) hold for the number l and with some polynomial p of degree at most $l - 2$; here we understand that p , as well as the regularising terms in the sum, are not present when $l = 1$.

//

Note that $\mathbb{K}_l \subseteq \mathbb{K}_{l+1} \subseteq \mathbb{K}$, $l \in \mathbb{N}$, and $\bigcup_{l \in \mathbb{N}} \mathbb{K}_l$ is the set of all real $f \in \mathbb{K}$ with only real zeros.

It follows from [L1, Theorems V.11 and V.13] that every function $f \in \bigcup_{l \in \mathbb{N}} \mathbb{K}_l$ is a canonical product and its zeros build a sequence belonging to the class \mathbb{S} , i.e. $f = P_\alpha$ with $\alpha \in \mathbb{S}$. As an application of our main theorems we prove the following perturbation result for the classes \mathbb{K}_l .

4.4 Theorem. *Let $\alpha, \beta \in \mathbb{S}$, and let $l \in \mathbb{N}$. If the hypotheses of one of Theorems 3.3, 3.18, 3.19 are fulfilled, then*

$$P_\alpha \in \mathbb{K}_l \iff P_\beta \in \mathbb{K}_l.$$

In order to establish the theorem, we use the following fact. Since an explicit reference is not known to us, we provide its proof.

4.5 Lemma. *Let $l \in \mathbb{N}$, let $\alpha \in \mathbb{S}$ and assume that*

$$\sum_{n \in \mathbb{N}} \frac{1}{|\alpha_n|^l |P'_\alpha(\alpha_n)|} < \infty. \quad (4.5)$$

Then $P_\alpha \in \mathbb{K}_l$.

Proof. It is clear that P_α satisfies (i). By [LW1, Lemma 5.5], convergence of the series (4.5) implies that P_α is of bounded type in the upper and lower half-planes \mathbb{C}^+ and \mathbb{C}^- ; for the definition of functions of bounded type see, e.g. [dB, Section 8].

Consider the function

$$\begin{aligned} g(z) &:= \sum_{k \in \mathbb{N}} \frac{1}{P'_\alpha(\alpha_k)} \left(\frac{1}{z - \alpha_k} + \frac{1}{\alpha_k} + \dots + \frac{z^{l-2}}{\alpha_k^{l-1}} \right) \\ &= z^{l-1} \sum_{k \in \mathbb{N}} \frac{1}{P'_\alpha(\alpha_k)(z - \alpha_k)\alpha_k^{l-1}}. \end{aligned}$$

Due to (4.5), this series converges absolutely and locally uniformly on the set $\mathbb{C} \setminus \{\alpha_n : n \in \mathbb{N}\}$, and thus represents an analytic function on this domain. At the points α_n it has simple poles with residua $\frac{1}{P'_\alpha(\alpha_n)}$. The function g can be written as

$$g(z) = z^{l-1} \int_{\mathbb{R}} \frac{d\mu}{z - t},$$

where μ is the discrete complex measure having point masses at the points α_k with masses $\frac{1}{P'_\alpha(\alpha_k)\alpha_k^{l-1}}$. Hence g is of bounded type in both half-planes \mathbb{C}^+ and \mathbb{C}^- .

We conclude that the difference

$$h(z) := \frac{1}{P_\alpha(z)} - g(z)$$

is an entire function which is of bounded type in \mathbb{C}^+ and \mathbb{C}^- . By Kreĭn's theorem [RR, Theorems 6.17, 6.18], h is of finite exponential type equal to the maximum of the mean types in \mathbb{C}^+ and \mathbb{C}^- .

Since $y \mapsto |1 - \frac{iy}{\alpha_n}|$ is increasing on $(0, \infty)$ and decreasing on $(-\infty, 0)$ and $\lim_{y \rightarrow \pm\infty} |1 - \frac{iy}{\alpha_n}| = \infty$ for each $n \in \mathbb{N}$, we have $\lim_{y \rightarrow \pm\infty} \frac{1}{P_\alpha(iy)} = 0$. Using dominated convergence, we see that $\lim_{y \rightarrow \pm\infty} \frac{1}{y^{l-1}} g(iy) = 0$, and together thus

$$\lim_{y \rightarrow \pm\infty} \frac{1}{y^{l-1}} h(iy) = 0.$$

We conclude that h is of minimal exponential type and, applying the Phragmén–Lindelöf principle [Bo, Theorem 1.4.3], that h is in fact a polynomial of degree at most $l - 2$ (here we understand that h vanishes identically if $l = 1$). Thus $\frac{1}{f}$ is represented as required in (4.4), and we have shown that $f \in \mathbb{K}_l$. \square

Proof of Theorem 4.4. Assume that $P_\alpha \in \mathbb{K}_l$. Then the function P_α satisfies (4.5). Since $\lim_{n \rightarrow \infty} \frac{\alpha_n}{\beta_n} = 1$ and the quotient $|\frac{P'_\alpha(\alpha_n)}{P'_\beta(\beta_n)}|$ is bounded from above and away from zero, also the function P_β satisfies (4.5). Hence, we may apply Lemma 4.5 and conclude that $P_\beta \in \mathbb{K}_l$.

For the converse, regard α as a perturbation of β , cf. Proposition 3.23 and Theorem 3.24. \square

4.3 The continuation problem for a positive definite function

A continuous function $f : \mathbb{R} \rightarrow \mathbb{C}$ is called *positive definite* if

(PD1) $f(-t) = \overline{f(t)}$, $t \in \mathbb{R}$;

(PD2) the kernel $f(t-s)$ is positive semi-definite; this means that, for each choice of $n \in \mathbb{N}$ and $t_1, \dots, t_n \in \mathbb{R}$, the quadratic form

$$Q(\xi_1, \dots, \xi_n) = \sum_{i,j=1}^n f(t_i - t_j) \xi_i \overline{\xi_j} \quad (4.6)$$

is positive semi-definite.

By Bochner's theorem the set of all positive definite functions coincides with the set of all Fourier transforms of finite positive measures on the real line, i.e. a function f is positive definite if and only if it can be represented as

$$f(t) = \int_{\mathbb{R}} e^{-itx} d\mu(x), \quad t \in \mathbb{R},$$

with some finite positive measure μ .

Let f be a positive definite function, let $a > 0$, and consider the restriction

$$\varphi := f|_{[-2a, 2a]}.$$

Then it may happen that there exist also other positive definite functions whose restriction to the interval $[-2a, 2a]$ equals φ . In fact, it is a classical result that either (I) or (II) holds:

(I) f is the only positive definite function with $f|_{[-2a, 2a]} = \varphi$.

(II) There exist infinitely many positive definite functions \tilde{f} that satisfy $\tilde{f}|_{[-2a, 2a]} = \varphi$.

A proof of this alternative and a parameterization of the set of all functions \tilde{f} in case (II) can be given. One possible approach proceeds via operator theoretic methods; see, e.g. [GG, §3.2].

Assume that case (I) prevails for f and a . When the requirement that the extension is positive definite is slightly weakened, it may happen that φ can still be extended in infinitely many ways to the whole real line. To make this precise, we say that a continuous function $f : \mathbb{R} \rightarrow \mathbb{C}$ is Hermitian indefinite with negative index $\kappa \in \mathbb{N}$ if it satisfies (PD1) and

(PD2 $_{\kappa}$) the kernel $f(t-s)$ has κ negative squares; this means that, for each choice of $n \in \mathbb{N}$ and $t_1, \dots, t_n \in \mathbb{R}$, the quadratic form (4.6) has at most κ negative squares, and, for some choice of n, t_1, \dots, t_n , this upper bound is attained.

4.6 Definition. Let f be a positive definite function, let $a > 0$, and assume that case (I) prevails for f and a . Then we set

$$\Delta(f, a) := \inf \{ \kappa \in \mathbb{N} : \exists \tilde{f} \text{ with (PD1), (PD2}_{\kappa}) \text{ s.t. } \tilde{f}|_{[-2a, 2a]} = \varphi \}.$$

Here the infimum of the empty set is understood as being equal to ∞ . //

4.7 Example. The function $f(t)$ that equals $1 - |t|$ on $[-2, 2]$ and is continued periodically to \mathbb{R} is positive definite as can be seen from its Fourier series. For $a = 1$, case (I) prevails and $\Delta(f, 1) = 1$; see [LLS]. An extension of $f|_{[-2, 2]}$ with one negative square is, e.g. the function $\tilde{f}(t) = 1 - |t|, t \in \mathbb{R}$. //

Consider f and a such that case (I) prevails, and let μ be the inverse Fourier transform of f . A characterization of ' $\Delta(f, a) < \infty$ ' in terms of μ can be given; in fact the actual value of $\Delta(f, a)$ can be computed, cf. [W, Proposition 6.11].

4.8 Theorem ([W]). *Let f be a positive definite function, let $a > 0$, and assume that case (I) prevails for f and a . Moreover, let μ be the inverse Fourier transform of f , i.e. μ is the positive finite measure with $f(t) = \int_{\mathbb{R}} e^{-itx} d\mu(x)$. Then $\Delta(f, a) < \infty$ if and only if*

(i) *the measure μ is discrete, say $\mu = \sum_{n=1}^N \sigma_n \delta_{\alpha_n}$ with $N \in \mathbb{N} \cup \{\infty\}$ and $\sigma_n > 0$;*

(ii) *the sequence $\alpha = (\alpha_n)_{n \in \mathbb{N}}$ belongs to \mathbb{S} ;*

(iii) *there exists a number $k \in \mathbb{N}$ such that*

$$\sum_{n=1}^N \frac{1}{\alpha_n^{2k} (P'_{\alpha}(\alpha_n))^2 \sigma_n} < \infty. \quad (4.7)$$

If $\Delta(f, a) < \infty$ and k_0 denotes the smallest natural number such that (4.7) holds, then

$$\Delta(f, a) = k_0 - 1.$$

We invoke the present stability results.

4.9 Theorem. *Let f be a positive definite function, let $a > 0$, and let μ be the inverse Fourier transform of f . Assume that case (I) prevails for f and a , that μ is discrete, say $\mu = \sum_{n=1}^{\infty} \sigma_n \delta_{\alpha_n}$ with $\sigma_n > 0$, and that $\alpha \in \mathbb{S}$. Let $\beta \in \mathbb{S}$ and $\tau_n > 0$, and consider*

$$\nu := \sum_{n \in \mathbb{N}} \tau_n \delta_{\beta_n}, \quad g(t) := \int_{\mathbb{R}} e^{-itx} d\nu(x).$$

If the hypotheses of one of Theorems 3.3, 3.18, 3.19 are fulfilled, $\tau_n \asymp \sigma_n$ and case (I) prevails for g and a , then

$$\Delta(g, a) = \Delta(f, a).$$

4.10 Remark. It is an open problem whether the hypotheses of Theorem 4.9 already imply that case (I) prevails for g and a . //

References

- [Bo] R. BOAS: *Entire Functions*, Academic Press, New York, 1954.
- [dB] L. DE BRANGES: *Hilbert Spaces of Entire Functions*, Prentice-Hall, London, 1968.
- [DK] H. DYM, H. MCKEAN: *Gaussian Processes, Function Theory, and the Inverse Spectral Problem*, Academic Press, New York, 1976.
- [GG] M.L. GORBACHUK, V.I. GORBACHUK: *M.G. Kreĭn's Lectures on Entire Operators*, Oper. Theory Adv. Appl. 97, Birkhäuser Verlag, Basel, 1997.
- [KaKr] I.S. KAC, M.G. KREĬN: *On the spectral functions of a string*, in F.V. Atkinson, *Discrete and Continuous Boundary Problems (Russian translation)*, Mir, Moscow, 1968, 648–737 (Addition II). English translation: Amer. Math. Soc. Transl. (2) 103 (1974), 19–102.
- [Kr1] M.G. KREĬN: *The fundamental propositions of the theory of representations of Hermitian operators with deficiency index (m, m) . (Russian)*, Ukrain. Mat. Zhurnal 1 (1949), no. 2, 3–66.
- [Kr2] M.G. KREĬN: *Determination of the density of a nonhomogeneous symmetric cord by its frequency spectrum. (Russian)*, Doklady Akad. Nauk SSSR (N.S.) 76 (1951), 345–348.
- [Kr3] M.G. KREĬN: *On a generalization of investigations of Stieltjes. (Russian)*, Doklady Akad. Nauk SSSR (N.S.) 87 (1952), 881–884.
- [LLS] H. LANGER, M. LANGER, Z. SASVÁRI: *Continuations of Hermitian indefinite functions and corresponding canonical systems: an example*, Methods Funct. Anal. Topology 10 (2004), 39–53.
- [LW1] M. LANGER, H. WORACEK: *A characterization of intermediate Weyl coefficients*, Monatsh. Math. 135 (2002), 137–155.
- [LW2] M. LANGER, H. WORACEK: *Stability of N -extremal measures*, submitted, Preprint: ASC Report 05/2013, Vienna University of Technology, 2013, <http://www.asc.tuwien.ac.at/preprint/2013/asc05x2013.pdf>.
- [LW3] M. LANGER, H. WORACEK: *A Pontryagin space approach to the index of determinacy of a measure*, manuscript in preparation.

- [LG] P. LELONG, L. GRUMAN: *Entire Functions of Several Variables*, Springer-Verlag, Berlin, 1986.
- [L1] B.J. LEVIN: *Distribution of Zeros of Entire Functions (Russian)*, Gosudarstv. Izdat. Tehn.-Teor. Lit., Moscow, 1956. English translation: *Translations of Mathematical Monographs*, vol. 5, Amer. Math. Soc., Providence, R.I., 1964.
- [L2] B.YA. LEVIN: *Lectures on Entire Functions*, Translated from the Russian manuscript by V. Tkachenko. *Translations of Mathematical Monographs*, vol. 150. Amer. Math. Soc., Providence, RI, 1996.
- [RR] M. ROSENBLUM, J. ROVNYAK: *Topics in Hardy Classes and Univalent Functions*, Birkhäuser Verlag, Basel, 1994.
- [Ru] L.A. RUBEL: *Entire and Meromorphic Functions*, Springer-Verlag, New York, 1996.
- [S] E. SENETA: *Regularly Varying Functions*, *Lecture Notes in Mathematics*, vol. 508, Springer, Berlin–Heidelberg–New York, 1976.
- [W] H. WORACEK: *Existence of zero-free functions N -associated to a de Branges Pontryagin space*, *Monatsh. Math.* 162 (2011), 453–506.

M. Langer
 Department of Mathematics and Statistics
 University of Strathclyde
 26 Richmond Street
 Glasgow G1 1XH
 UNITED KINGDOM
 email: m.langer@strath.ac.uk

H. Woracek
 Institut für Analysis und Scientific Computing
 Technische Universität Wien
 Wiedner Hauptstraße. 8–10/101
 1040 Wien
 AUSTRIA
 email: harald.woracek@tuwien.ac.at