

Sedimentary membrane lipids recycled by deep-sea benthic archaea

Yoshinori Takano^{1*}, Yoshito Chikaraishi¹, Nanako O. Ogawa¹, Hidetaka Nomaki¹,

Yuki Morono², Fumio Inagaki², Hiroshi Kitazato¹, Kai-Uwe Hinrichs³

5 **& Naohiko Ohkouchi^{1,4,5}**

¹Institute of Biogeosciences, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2-15 Natsushima, Yokosuka 237-0061, Japan.

10 *²Kochi Institute for Core Sample Research, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Monobe B200, Nangoku, Kochi 783-8502, Japan.*

³Department of Geosciences and MARUM Center for Marine Environmental Sciences, University of Bremen, D-28359, Bremen, Germany

⁴Department of Environmental Science and Technology, Tokyo Institute of Technology, 4259 Nagatsuda, Midori-ku, Yokohama, Kanagawa 226-8502, Japan

15 *⁵Department of Earth and Planetary Sciences, University of Tokyo, 7-3-1 Hongo, Tokyo 113-0033, Japan*

E-mail: takano@jamstec.go.jp; nohkouchi@jamstec.go.jp

Fax: +81-46-867-9775

Since the work of Woese *et al.*¹, Archaea are recognized as one of three domains of life and cosmopolitan microbes in marine environments²⁻⁴. In the deep ocean (>200m), cells of the archaeal phylum *Crenarchaeota* are dominant while Bacteria
25 mainly occupy the sea surface⁵. Recent molecular evidence based on lipids and DNA/RNA suggests that uncultured heterotrophic archaea relying on relic organic carbon dominate biomass in marine sediment⁶. Here we report constraints on the ecology of these benthic archaea, derived from *in situ* ¹³C-tracer experiments over 405 days on the deep-sea floor. The addition of ¹³C-glucose as substrate resulted in
30 pronounced ¹³C incorporation into the ether-linked glycerol unit (>2100‰) of archaeal tetraether membrane lipids while the isopranyl unit remained nearly unlabeled. The intramolecular ¹³C distribution and changes in the phylogenetic composition of the archaeal community during the experiment suggest that benthic archaea reutilized exogenous lipids from relic cells and detritus to build their
35 endogenous membrane lipids. Our findings imply that at least some benthic archaea build their membranes by recycling sedimentary organic compounds in order to minimize the *de novo* lipid synthesis for growth and maintenance.

Deep-sea sediments harbor a novel and vast biosphere with yet unconstrained
40 importance in the global carbon cycle⁶⁻¹⁴. Carbon isotopic signatures of archaeal polar lipids from sediments dominated by benthic archaeal communities indicate utilization of sedimentary organic carbon^{6,15}. On-shore laboratory studies aimed at understanding microbial processes in deep-sea sediments remain problematic because most indigenous

microbes appear viable but non-culturable¹⁶. To further constrain the substrates utilized
45 by benthic archaea, we conducted an *in situ* ¹³C-tracer experiment to provide direct
evidence of their metabolism *via* an investigation of membrane lipid biomarkers.

The experiment was performed on the seafloor at the central part of Sagami Bay,
Japan (35°00.8'N, 139°21.6'E; 1453 m depth)¹⁷ for 0-405 days (Supplementary Figure 1).
Sedimentary processes relevant to carbon burial and benthic activity have been
50 previously documented¹⁸. In our experiment, an aqueous solution of 16.5 mg of
¹³C-labeled glucose (¹³C₆H₁₂O₆) was injected into the headspace of the chamber using
dual 5-mL syringes attached to the top (Figure 1). The incubation cores and a reference
core were recovered at 0, 9 and 405 days after deployment (hereafter, the culture cores
are referred to as G-*i*, with *i* indicating the length of incubation in days).

55 Already after 9 days, ¹³C-tracer incorporation was unambiguously detected in
caldarchaeol ($\delta^{13}\text{C} = +46\text{‰}$) and crenarchaeol (+22‰) from the surface sediment (0–1
cm depth), reflecting the active metabolism and growth of the benthic archaeal
community (Figure 2). ¹³C uptake increased after 405 days. Depth profiles of carbon
isotopic compositions of caldarchaeol and crenarchaeol are positively correlated ($R^2 <$
60 0.83 ; $p\text{-value} > 0.07$) with those of organic matter in the sediment. $\delta^{13}\text{C}_{\text{caldarchaeol}}$ ($< -21\text{‰}$)
and $\delta^{13}\text{C}_{\text{crenarchaeol}}$ ($< -22\text{‰}$) in the experimental blank are consistent with values
previously observed in benthic¹⁵ and planktonic¹⁹ archaeal communities (Figure 2).
¹³C-tracer uptake process is either due to heterotrophic assimilation of glucose or
autotrophic assimilation of ¹³CO₂ after glucose has been oxidized by undefined microbial
65 community members (cf. DIC, $\delta^{13}\text{C}_{\text{DIC}} > +4200\text{‰}$ in G-9 experiment). In the latter case,

the signal could result from assimilation of $^{13}\text{CO}_2$ by marine group I (MGI) crenarchaeota. 16S rRNA genes affiliated with the MGI group dominate the corresponding clone library in the G-9 experiment (Figure 3). However, given previous demonstration of rapid ^{13}C -DIC uptake into the biphytanyl moieties of caldarchaeol and crenarchaeol by closely
70 related planktonic MGI crenarchaeota⁴, this autotrophic scenario appears less likely (see Supplementary Information).

The carbon isotopic compositions of the acyclic, bicyclic and tricyclic archaeal biphytanes (hereafter referred to as BP[0], BP[2] and BP[3], respectively, see Methods) released from caldarchaeol and crenarchaeol were around -22‰ in experiment G-405,
75 whereas the bulk GDGT core lipids caldarchaeol ($+69\text{‰}$) and crenarchaeol ($+132\text{‰}$) were significantly ^{13}C -enriched (Figure 2, Supplementary Tables 1 & 2). This disparity indicates that the ^{13}C tracer is unevenly distributed in the core lipid molecule, *i.e.*, ^{13}C -tracer is concentrated in the lipid's 2,3-*sn*-glycerol backbone. We can estimate the carbon isotopic composition of 2,3-*sn*-glycerol ($\delta^{13}\text{C}_{2,3\text{-}sn\text{-glycerol}}$) by mass balance
80 (Supplementary Information).

$$86 \times \delta^{13}\text{C}_{\text{caldarchaeol}} = 80 \times \delta^{13}\text{C}_{\text{BP[0]}} + 6 \times \delta^{13}\text{C}_{2,3\text{-}sn\text{-glycerol}} \quad \text{Eq. [1]}$$

$$86 \times \delta^{13}\text{C}_{\text{crenarchaeol}} = 40 \times \delta^{13}\text{C}_{\text{BP[2]}} + 40 \times \delta^{13}\text{C}_{\text{BP[3]}} + 6 \times \delta^{13}\text{C}_{2,3\text{-}sn\text{-glycerol}} \quad \text{Eq. [2]}$$

85 Accordingly the 2,3-*sn*-glycerol moieties in caldarchaeol and crenarchaeol in the 0–1 cm interval are enriched in ^{13}C by 900‰ and 500‰, respectively, after 9 days, and by 1300‰ and 2200‰ after 405 days (Figure 2). For further confirmation of our observation,

the validity of the estimates was confirmed by direct measurement of the carbon isotopic compositions of 2,3-*sn*-glycerol ([Supplementary Information](#)). Then, the carbon isotopic composition of glycerol moieties derived from the purified caldarchaeol and crenarchaeol in G-405 samples were up to +1900‰ and +2780‰, respectively. These values are consistent with ¹³C-mass balance estimations. The large heterogeneity suggests a predominance of the benthic archaeal metabolic pathway toward 2,3-*sn*-glycerol rather than isoprenoid lipid biosynthesis. The variable differences of $\delta^{13}\text{C}_{2,3\text{-}sn\text{-glycerol}}$ values of caldarchaeol and crenarchaeol are consistent with changes in community composition and associated changes of the sources of these two compounds during the course of the experiment. Structural and biosynthetic studies of archaeal lipids²⁰ suggest the presence of a pathway leading to GDGTs *via* formation of 2,3-*sn*-glycerol and isoprenoid moieties ([Supplementary Information](#)). When synthesizing *sn*-glycerate, Archaea utilize either the Embden-Meyerhof (EM), the modified EM or the Entner-Doudoroff (ED) pathway²¹. In contrast, they produce isoprenoids through either the mevalonate (MVA) or the modified MVA pathway from acetyl-CoA²².

16S rRNA and quantitative PCR (qPCR) analysis indicated a community shift in the composition of the archaeal community and its abundance (10^5 - 10^7 copies g-sed⁻¹) during the course of the experiment, whereas the relative abundances (%) of tetraether archaeal lipids were almost constant during 405 days ([Figure 3](#)). The initial community consisted predominantly of *Crenarchaeota* including MGI, the Soil Crenarchaeota Group (SCG), and the Miscellaneous Crenarchaeotal Group (MCG). After 9 days of incubation, the relative proportion of individual archaeal groups, and especially MGI and MCG, changed

110 substantially. After 405 days, the relative proportions of clones of these two groups resemble again those at initial G-0. Benthic MGI were previously identified in organic-rich (*e.g.*, Peru Margin, ODP Leg 201) and also organic-poor (*e.g.*, Equatorial Pacific, ODP Leg 201) marine sediment¹³. Furthermore, genes of the Methanosarcinales accounted for approximately 20% of the archaeal assemblages after 405 days.

115 Given the properties of the outer membrane of prokaryotic cells²³ in combination with the extremely energy-limited sedimentary archaeal communities¹¹, the following scenario is consistent with the intramolecular isotopic distribution. Benthic archaea could synthesize caldarchaeol and crenarchaeol via recycling of exogenous precursor compounds, such as biphytane diols resulting from decomposition of decayed archaeal
120 cells, while only 2,3-*sn*-glycerol is synthesized *de novo* process as signaled by exclusive ¹³C-label uptake in this molecular moiety. Price and Sowers²⁴ estimated that the metabolic energy required for growth of microbial cell is 6 orders of magnitude larger than that needed for survival. Therefore, the recycling of fossil molecules could be a crucial strategy for coping with conditions of energy starvation in sedimentary
125 environments and aid in minimizing energy expenditures for growth and maintenance^{12,15}. Transport of these exogenous building blocks may take place *via* permeable channels in the outer membrane²³. Passage of solutes across the outer membrane channel is generally permeable to hydrophobic molecules with molecular weight in excess of 1000 Da²³. Specifically, transport initiated by adsorption of hydrophobic functional groups (*e.g.*,
130 hydroxyl groups) onto porin proteins are proposed for long-chain fatty acids and other organic molecules across the outer membrane channel ([Supplementary Information](#)).

Relic archaeal tetraether core lipids are among the most abundant molecular types in marine sediments¹⁹; these compounds could potentially serve as substrate for supplying the building-blocks, although this would require biochemical scission of ether-bonds²⁵, a process associated with an appreciable investment of energy (<360 kJ mol⁻¹). This experiment strongly suggests that deep-sea benthic archaea utilize yet unknown mechanisms to minimize energy expenditures.

Methods

140 Bulk and compound-specific isotope analysis

We extracted archaeal core GDGTs (glycerol dialkyl glycerol tetraethers) from dried sediments using an improved method²⁶ and purified them by preparative high-performance liquid chromatography combined with atmospheric-pressure chemical ionization mass spectrometry (HPLC/APCI-MS) without ether-bond cleavage treatment (Supplementary Figure 2). To verify an analytical bias during HPLC, we carefully evaluated potential co-elution by fraction collection both before caldarchaeol peak and after crenarchaeol peak on the chromatogram (Supplementary Figure 3). This CSIA method was performed with an elemental analyzer coupled to an isotope ratio mass spectrometer (EA/IRMS). Bulk isotope analysis ($\delta^{13}\text{C}_{\text{TOC}}$) and total organic carbon (TOC) in the sediment were determined by conventional methods (see Supplementary Information).

To analyze the biphytanes of caldarchaeol and crenarchaeol, we performed an ether cleavage treatment²⁷, followed by reduction²⁸. We determined the carbon isotopic

compositions of resulting biphytane and glycerol derivatives using an online gas
155 chromatograph/combustion/isotope ratio mass spectrometer (GC/C/IRMS), enabling the
(i) calculation of 2,3-*sn*-glycerol ($\delta^{13}\text{C}_{2,3\text{-sn-glycerol}}$) using the mass balance equation
between caldarchaeol, crenarchaeol and their isoprenoid moieties, (ii) direct measurement
of $\delta^{13}\text{C}_{2,3\text{-sn-glycerol}}$. The validity of this method was verified using another marine
sediment sample prior to application ([Supplementary Information](#)).

160

2. Phylogenetic analysis of 16S rRNA and qPCR for benthic archaeal community

Bulk genomic DNA was extracted from 0.2–0.5 g of sediments using the ISOIL for
Beads Beating Kit (Nippon Gene, Tokyo, Japan). Extracted DNA was further purified
using the Montage PCR Kit (Millipore, Billerica, MA), and archaeal 16S rRNA gene
165 fragments were amplified with Arch21F-1492R primers^{2,29}. The PCR conditions were as
follows: denaturation at 95°C for 60 sec, annealing at 50°C for 60 sec, and extension at
72°C for 120 sec, for 25–30 cycles. A total of 528 clones (more than 160 clones per
sample) were selected from the libraries, and the inserted 16S rRNA gene sequences were
determined using a 3130xl Genetic Analyser (Life Technologies, Carlsbad, CA). For
170 analysing phylogenetic affiliation, the sequences were aligned and grouped using ARB
software (AB561312-AB561824; 513 entries)³⁰. To estimate copy numbers of archaeal
16S rRNA genes, quantitative PCR (qPCR) was performed with a SYBR[®] Premix
DimerEraser[®] (Takara Bio Inc.) by StepOnePlus[™] real-time PCR system⁶ according to the
manufacturer's instructions (Applied Biosystems). Three replicate reactions per sample
175 were performed with the primer set of ARC806F and ARC958R and the thermal condition

Takano *et al.*, *Nature Geoscience*, **3**, 858-861 (2010).

of 95°C for 30 sec for initial denaturation, up to 50 cycles of 95°C for 5 sec, 55°C for 30 sec, 72°C for 30 sec. The standard curve for archaeal 16S rRNA genes were obtained from genome DNA of *Pyrococcus horikoshii* ($R^2= 0.995$).

180 **References**

- 1 Woese, C. R., Kandler, O. & Wheelis, M. L. Towards a natural system of organisms - proposal for the domains Archaea, Bacteria, and Eucarya. *Proc. Natl Acad. Sci. USA.* **87**, 4576-4579 (1990).
- 2 DeLong, E. Archaea in coastal marine environments. *Proc. Natl Acad. Sci. USA.* **89**,
185 5685-5689 (1992).
- 3 Ouverney, C. C. & Fuhrman, J. A. Marine planktonic Archaea take up amino acids. *Appl. Environ. Microbiol.* **66**, 4829-4833 (2000).
- 4 Wuchter, C., Schouten, S., Boschker, H. T. S. & Sinninghe Damste, J. S. Bicarbonate uptake by marine Crenarchaeota. *FEMS Microbiol. Lett.* **219**, 203-207 (2003).
- 190 5 Karner, M. B., DeLong, E. F. & Karl D. M. Archaeal dominance in the mesopelagic zone of the Pacific Ocean. *Nature* **409**, 507-510 (2001).
- 6 Lipp, J. S., Morono, Y., Inagaki, F. & Hinrichs, K.-U. Significant contribution of Archaea to extant biomass in marine subsurface sediments. *Nature* **454**, 991-994 (2008).
- 195 7 Parkes, R. J., Cragg, B. A. & Wellsbury, P. Recent studies on bacterial populations and processes in seafloor sediments: A review. *Hydrogeol. J.* **8**, 11-28 (2000).

Takano *et al.*, *Nature Geoscience*, **3**, 858-861 (2010).

- 8 Rothschild, L. J. & Mancinelli, R. L. Life in extreme environments. *Nature* **409**, 1092-1101 (2001).
- 9 D'Hondt, S. *et al.* Distributions of microbial activities in deep subseafloor sediments. *Science* **306**, 2216-2221 (2004).
- 200
- 10 Schippers, A. *et al.* Prokaryotic cells of the deep sub-seafloor biosphere identified as living bacteria. *Nature* **433**, 861-864 (2005).
- 11 Jorgensen, B. & Boetius, A. Feast and famine-microbial life in the deep-sea bed. *Nature Rev. Microbiol.* **5**, 770-781 (2007).
- 205
- 12 Valentine, D. L. Adaptations to energy stress dictate the ecology and evolution of the Archaea. *Nature Rev. Microbiol.* **5**, 316-323 (2007).
- 13 Inagaki, F. *et al.* Biogeographical distribution and diversity of microbes in methane hydrate-bearing deep marine sediments on the Pacific Ocean Margin. *Proc. Natl Acad. Sci. USA.* **103**, 2815-2820 (2006).
- 210
- 14 Roussel, E. G. *et al.* Extending the sub-sea-floor biosphere. *Science* **320**, 1046-1046 (2008).
- 15 Biddle, J. F. *et al.* Heterotrophic archaea dominate sedimentary subsurface ecosystems off Peru. *Proc. Natl Acad. Sci. USA.* **103**, 3846-3851 (2006).
- 16 Teske, A. & Sorensen, K. B. Uncultured archaea in deep marine subsurface sediments: have we caught them all? *ISME J.* **2**, 3-18 (2008).
- 215
- 17 Nomaki, H., Heinz, P., Nakatsuka, T., Shimanaga, M. & Kitazato, H. Species-specific ingestion of organic carbon by deep-sea benthic foraminifera and meiobenthos: In situ tracer experiments. *Limnol. Oceanogr.* **50**, 134-146 (2005).

- 18 Kitazato, H. "The project Sagami" - dynamic sedimentary processes of both organic and
220 inorganic materials at continental margins with active tectonic forcing. *Prog. Oceanogr.* **57**, 1-2 (2003).
- 19 Shah, S. R., Mollenhauer, G., Ohkouchi, N., Eglinton, T. I. & Pearson, A. Origins of
archaeal tetraether lipids in sediments: insights from radiocarbon analysis. *Geochim. Cosmochim. Acta* **72**, 4577-4594 (2008).
- 225 20 Koga, Y. & Morii, H. Biosynthesis of ether-type polar lipids in archaea and
evolutionary considerations. *Microbiol. Mol. Biol. Rev.* **71**, 97-120 (2007).
- 21 Verhees, C. H. *et al.* The unique features of glycolytic pathways in archaea. *Biochem. J.*
375, 231-246 (2003).
- 22 Grochowski, L., Xu, H. & White, R. *Methanocaldococcus jannaschii* uses a modified
230 mevalonate pathway for biosynthesis of isopentenyl diphosphate. *J. Bacteriol.* **188**,
3192-3198 (2006).
- 23 Lodish *et al.*, The outer membrane and cell wall. *In* Molecular Cell Biology, W. H.
Freeman & Co Ltd (1995).
- 24 Price, P. B. & Sowers, T. Temperature dependence of metabolic rates for microbial
235 growth, maintenance, and survival. *Proc. Natl Acad. Sci. USA.* **101**, 4631-4636
(2004).
- 25 White, G., Russell, N. & Tidswell, E. Bacterial scission of ether bonds. *Microbiol. Rev.*
60, 216-232 (1996).
- 26 Hopmans, E. C., Schouten, S., Pancost, R. D., van der Meer, M. T. J. & Sinninghe
240 Damste, J. S. Analysis of intact tetraether lipids in archaeal cell material and

sediments by high performance liquid chromatography/atmospheric pressure
chemical ionization mass spectrometry. *Rap. Comm. Mass Spectr.* **14**, 585-589
(2000).

27 Schouten S., Hoefs M. J. L., Koopmans M. P., Bosch H. J. & Sinninghe Damste, J. S.
245 Structural characterization, occurrence and fate of archaeal ether-bound acyclic and
cyclic biphytanes and corresponding diols in sediments. *Org. Geochem.* **29**,
1305-1319 (1998).

28 Elvert, M., Suess, E., Greinert, J. & Whiticar, M. J. Archaea mediating anaerobic
methane oxidation in deep-sea sediments at cold seeps of the eastern Aleutian
250 subduction zone. *Org. Geochem.* **31**, 1175-1187 (2000).

29 Lane, D. J. *et al.* Rapid-determination of 16S Ribosomal-RNA sequences for
phylogenetic analyses. *Proc. Natl Acad. Sci. USA.* **82**, 6955-6959 (1985).

30 Ludwig, W. *et al.* ARB: a software environment for sequence data. *Nucl. Acids. Res.* **32**,
1363-1371 (2004).

255

Acknowledgements

The authors express their sincere thanks to three anonymous reviewers for critical and
invaluable reviewing comments. The authors thank T. Terada (Marine work Japan), the
shipboard scientists for their experimental and logistic support during cruises
260 NT06-04,-05,-22 and NT08-02 by R/V *Natsushima* and for providing bathymetric data.
This research was supported in part by a grant from the Japan Society for the Promotion

Takano *et al.*, *Nature Geoscience*, **3**, 858-861 (2010).

of Science, Grant-in-Aid for Creative Scientific Research (19GS0211 & 22684030), and the internship program between Japan and Germany (FY2009).

265 **Author contributions**

Y.T. performed present lipids analysis and wrote the paper; Y.C. & N.O.O. supported on carbon isotope standard reagents and IRMS analysis; H.N. & H.K. ¹³C-substrate set up, *in-situ* deployment of the chamber, core processing during NT06-04, -05, -22 and NT08-02 cruise; Y.M. and F.I. phylogenic molecular analysis of 16S rRNA and qPCR;

270 K-U.H. contributed technical aspects and was involved in study design; Y.T. and N.O. contributed to this study and all authors discussed the results and commented on the manuscript.

Additional information

275 Supplementary Information accompanies this paper on

<http://www.nature.com/ngeo/journal/v3/n12/extref/ngeo983-s1.pdf>

Reprints and permissions information is available online at

<http://npg.nature.com/reprintsandpermissions>.

Correspondence and requests for materials should be addressed to Y.T.

280

Figure captions

285 **Figure 1. Distribution of benthic archaeal lipids and their carbon isotopic compositions during *in situ* ¹³C-tracer experiment during 405 days. a, b, Total organic carbon (TOC, wt%) and carbon isotopic composition ($\delta^{13}\text{C}_{\text{TOC}}$, ‰ vs. PDB). c, Carbon isotopic compositions of caldarchaeol and crenarchaeol ($\delta^{13}\text{C}_{\text{caldarchaeol}}$ and $\delta^{13}\text{C}_{\text{crenarchaeol}}$) for 0 day (G-0), 9 days (G-9), and 405 days (G-405) experiments.**

290 The experiments using I-K incubation chamber¹⁷ were performed using the ROV *Hyper-Dolphin* and the R/V *Natsushima*. Bottom-water properties at the station are stable throughout the year, with temperatures of $2.3\pm 0.1^\circ\text{C}$ and salinity of 34.5 ± 0.2 ‰ (see also Supplementary Figure 1).

Figure 2. Results of carbon isotopic compositions of $\delta^{13}\text{C}_{\text{caldarchaeol}}$, $\delta^{13}\text{C}_{\text{crenarchaeol}}$, $\delta^{13}\text{C}_{\text{biphytane}}$ and $\delta^{13}\text{C}_{2,3\text{-sn-glycerol}}$ *in situ* ¹³C-tracer experiments during 405 days. The sample analyzed for the time-course profiles was 0–1 cm depth below the seafloor each. Here, $\delta^{13}\text{C}_{2,3\text{-sn-glycerol}}$ of caldarchaeol and crenarchaeol (see molecular structure) were determined as follows: $\delta^{13}\text{C}_{2,3\text{-sn-glycerol}} = (86 \times \delta^{13}\text{C}_{\text{caldarchaeol}} - 80 \times \delta^{13}\text{C}_{\text{BP}[0]}) / 6$, where $\delta^{13}\text{C}_{\text{caldarchaeol}}$ and $\delta^{13}\text{C}_{\text{BP}[0]}$ were

300 independently determined by GC/C/IRMS. Furthermore, $\delta^{13}\text{C}_{2,3\text{-sn-glycerol}} = (86 \times \delta^{13}\text{C}_{\text{crenarchaeol}} - 40 \times \delta^{13}\text{C}_{\text{BP}[2]} - 40 \times \delta^{13}\text{C}_{\text{BP}[3]}) / 6$, where $\delta^{13}\text{C}_{\text{crenarchaeol}}$, $\delta^{13}\text{C}_{\text{BP}[2]}$ and $\delta^{13}\text{C}_{\text{BP}[3]}$ were determined by CSIA. $\Delta\delta^{13}\text{C}_{\text{glycerol-BP}}$ stands for the difference in $\delta^{13}\text{C}$ values between 2,3-*sn*-glycerol and isoprenoid moieties.

Figure 3. Long-term monitoring for abundances of archaeal lipids and phylogenic analysis of the benthic archaeal community. a, Relative abundances (%, $n=3$) of

305

GDGTs during 405 days. **b**, Natural assemblage of benthic archaea in the sampling site (mid-depth 1.5 cm), *in situ* experiment for 9 days and 405 days. Abbreviations; ANME2-d, Anaerobic methanotroph 2-d; MBG-D, Marine Benthic Group D; MEG/TMEG, Miscellaneous Euryarchaeotal Group/Terrestrial Miscellaneous Euryarchaeotal Group; GSAG, Deep-Sea Archaeal Group; SAGMEG, South African Goldmine Euryarchaeotal Groups; MCG, Miscellaneous Crenarchaeotic Group. Species richness, community diversity, and abundance of the benthic archaea were also monitored during ^{13}C -incubation.

315

Supplementary Information:

<http://www.nature.com/ngeo/journal/v3/n12/extref/ngeo983-s1.pdf>




Figure 1





Figure 2

Figure 3



Supplementary Information:

<http://www.nature.com/ngeo/journal/v3/n12/extref/ngeo983-s1.pdf>