

2017

An efficient platform for astrocyte differentiation from human induced pluripotent stem cells

Julia TCW

Mount Sinai School of Medicine

Minghui Wang

Mount Sinai School of Medicine

Anna A. Pimenova

Mount Sinai School of Medicine

Kathryn R. Bowles

Mount Sinai School of Medicine

Brigham J. Hartley

Mount Sinai School of Medicine

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Recommended Citation

TCW, Julia; Wang, Minghui; Pimenova, Anna A.; Bowles, Kathryn R.; Hartley, Brigham J.; Lacin, Emre; Machlovi, Saima I.; Abdelaal, Rawan; Karch, Celeste M.; Phatnani, Hemali; Slesinger, Paul A.; Zhang, Bin; Goate, Alison M.; and Brennand, Kristen J., "An efficient platform for astrocyte differentiation from human induced pluripotent stem cells." *Stem Cell Reports*.9,2. 600-614. (2017). https://digitalcommons.wustl.edu/open_access_pubs/6066

Authors

Julia TCW, Minghui Wang, Anna A. Pimenova, Kathryn R. Bowles, Brigham J. Hartley, Emre Lacin, Saima I. Machlovi, Rawan Abdelaal, Celeste M. Karch, Hemali Phatnani, Paul A. Slesinger, Bin Zhang, Alison M. Goate, and Kristen J. Brennand

An Efficient Platform for Astrocyte Differentiation from Human Induced Pluripotent Stem Cells

Julia TCW,^{1,2} Minghui Wang,^{4,7} Anna A. Pimenova,^{1,2,7} Kathryn R. Bowles,^{1,2,7} Brigham J. Hartley,^{3,7} Emre Lacin,^{1,7} Saima I. Machlovi,^{1,2} Rawan Abdelaal,⁵ Celeste M. Karch,⁶ Hemali Phatnani,⁵ Paul A. Slesinger,¹ Bin Zhang,⁴ Alison M. Goate,^{1,2,4,*} and Kristen J. Brennand^{1,3,*}

¹Department of Neuroscience & Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, 1425 Madison Avenue, New York, NY 10029, USA

²Ronald M. Loeb Center for Alzheimer's disease, Icahn School of Medicine at Mount Sinai, 1425 Madison Avenue, New York, NY 10029, USA

³Department of Psychiatry, Icahn School of Medicine at Mount Sinai, 1425 Madison Avenue, New York, NY 10029, USA

⁴Department of Genetics and Genomic Sciences, Icahn Institute of Genomics and Multiscale Biology, Icahn School of Medicine at Mount Sinai, 1470 Madison Avenue, New York, NY 10029, USA

⁵New York Genome Center, 101 Avenue of the Americas, New York, NY 10013, USA

⁶Department of Psychiatry, Washington University in St. Louis, St. Louis, MO 63130, USA

⁷These authors contributed equally

*Correspondence: alison.goate@mssm.edu (A.M.G.), kristen.brennand@mssm.edu (K.J.B.)

<http://dx.doi.org/10.1016/j.stemcr.2017.06.018>

SUMMARY

Growing evidence implicates the importance of glia, particularly astrocytes, in neurological and psychiatric diseases. Here, we describe a rapid and robust method for the differentiation of highly pure populations of replicative astrocytes from human induced pluripotent stem cells (hiPSCs), via a neural progenitor cell (NPC) intermediate. We evaluated this protocol across 42 NPC lines (derived from 30 individuals). Transcriptomic analysis demonstrated that hiPSC-astrocytes from four individuals are highly similar to primary human fetal astrocytes and characteristic of a non-reactive state. hiPSC-astrocytes respond to inflammatory stimulants, display phagocytic capacity, and enhance microglial phagocytosis. hiPSC-astrocytes also possess spontaneous calcium transient activity. Our protocol is a reproducible, straightforward (single medium), and rapid (<30 days) method to generate populations of hiPSC-astrocytes that can be used for neuron-astrocyte and microglia-astrocyte co-cultures for the study of neuropsychiatric disorders.

INTRODUCTION

Astrocytes are the most abundant cell type in the CNS, rivaling the diversity of neurons in cellular morphologies, gene expression profiles, developmental origins, physiological properties, functions, and responses to injury and disease (Zhang and Barres, 2010). Within the human brain, astrocytes have a variety of essential functions including glutamate biology, axonal guidance, trophic support, inflammatory response and wound healing, formation of the blood-brain barrier, and neuronal synapse formation, and plasticity (Barres, 2008; Eroglu and Barres, 2010; Freeman and Rowitch, 2013). Although the full contribution of astrocytes to neurological disease remains unresolved, astrocyte cell-autonomous deficits have been implicated in a variety of neurological disorders (Seifert et al., 2006; Tong et al., 2014). The most significant genetic risk factor for Alzheimer's disease (AD), apolipoprotein E4 (APOE4), is predominantly synthesized and secreted by astrocytes (Xu et al., 2006). Furthermore, astrocytes derived from human induced pluripotent stem cell (hiPSC)- or mouse-based models of amyotrophic lateral sclerosis (Di Giorgio et al., 2008; Marchetto et al., 2008; Papadeas et al., 2011), Rett syndrome (Ballas et al., 2009), and Huntington disease (Bradford et al., 2009) damage neurons in co-culture or after transplantation.

Evolutionarily, the astrocyte-to-neuron ratio increases from low vertebrates to rodents and to primates (Sherwood

et al., 2006). Human cortical astrocytes are larger, structurally more complex and diverse, and propagate calcium waves 4-fold faster than their rodent counterparts (Oberheim et al., 2009). Transplantation of human glia into mice enhances activity-dependent plasticity and learning (Bi et al., 2013). Given the unique biology of human astrocytes, it is critical that improved human-specific cell-based systems be established to enable the study of human astrocyte function in health and disease.

Because of their ability to model all of the (known and unknown) genetic risk factors underlying neuropsychiatric disease, hiPSCs are routinely used as a source of various types of neurons and astrocytes for study (Mertens et al., 2016). Current hiPSC-based methods for the differentiation of astrocytes typically rely on either a neural progenitor cell (NPC) (Haidet-Phillips et al., 2014; Krencik et al., 2011; McGivern et al., 2013; Serio et al., 2013; Shaltouki et al., 2013) or oligodendrocyte progenitor cell (Jiang et al., 2013) intermediate. While it has been widely demonstrated that hiPSCs can be differentiated to functional astrocytes for cell-based models of neuropsychiatric disorders *in vitro* (Haidet-Phillips et al., 2014; McGivern et al., 2013; Serio et al., 2013; Shaltouki et al., 2013) or engraftment *in vivo* (Chen et al., 2015; Haidet-Phillips et al., 2014; Jiang et al., 2013; Krencik et al., 2011), existing methods are slow (up to 6 months) (Jiang et al., 2013; Krencik et al., 2011; Shaltouki et al., 2013) and/or require sorting to reduce

**Table 1. Screening Conditions for Astrocyte Differentiation**

ID	Reference	Basal Medium	Supplement 1	Supplement 2	Growth Factors			
1	Brennand and Gage (2011)	DMEM/F12	N2, B27	GlutaMAX + sodium bicarbonate + sodium pyruvate	BDNF	GDNF	cAMP	AA
2	ScienCell 1801	ND (AM)	AGS				2% FBS	
3	Thermo Fisher A1261301	DMEM	N2	GlutaMAX + D-glucose + sodium pyruvate			10% FBS	
4	Lonza CC-3186	ND	–	L-glutamine	EGF		3% FBS	insulin + AA
5	C.M.K., unpublished data	neural basal glutamine	2 × N2		FGF2 + CNTF			
6	modified from C.M.K., unpublished data	neural basal glutamine	N2, B27	GlutaMAX, NEAA, bME	FGF2 + CNTF			
7	Krencik et al. (2011)	DMEM/F12	N2, B27	GlutaMAX, NEAA, bME	FGF2 + CNTF			
8	Jiang et al. (2013)	DMEM/F12	N2, B27	GlutaMAX, NEAA, bME	FGF2 + CNTF	BMP4		
9	reduction of % FBS from no. 10	DMEM/F12	N2, B27	GlutaMAX, NEAA, bME	FGF2 + CNTF	BMP4	3% FBS	
10	Roybon et al. (2013)	DMEM/F12	N2, B27	GlutaMAX, NEAA, bME	FGF2 + CNTF	BMP4	10% FBS	
11	Shaltouki et al. (2013)	DMEM/F12	N2, B27	GlutaMAX, NEAA, bME	FGF2 + CNTF	BMP4	3% FBS	neuregulin
12	adding insulin + AA to no. 9	DMEM/F12	N2, B27	GlutaMAX, NEAA, bME	FGF2 + CNTF	BMP4	3% FBS	insulin + AA

The 11 screened conditions of astrocyte differentiation from hiPSCs, including 3 commercially available astrocyte media (nos. 2–4) and 8 conditions modified from published protocols (nos. 5–12), with a forebrain neuron differentiation protocol (no. 1). NEAA, non-essential amino acid; bME, 2-mercaptoethanol; AA, ascorbic acid; ND, non-disclosure; AM, astrocyte medium; AGS, astrocyte growth supplement.


heterogeneity ([Chaboub and Deneen, 2013](#); [Yuan et al., 2011](#)). Here, we screened a number of published protocols, along with commercially available media for primary human astrocyte culture, identifying a robust and straightforward differentiation protocol for generating astrocytes from hiPSCs. By co-culture with microglia, we compared the function of primary human fetal astrocytes and hiPSC-astrocytes in assays for neuroinflammatory response, phagocytosis, and spontaneous calcium activity, concluding that hiPSC-astrocytes are highly similar to their primary counterparts. Altogether, our rapid differentiation protocol, co-culture strategy, and scalable phenotypic assays will serve as a robust platform for queries of healthy and diseased human astrocytes.

RESULTS

30-Day Exposure of hiPSC-Derived NPCs to Commercial Astrocyte Media Is Sufficient to Robustly Generate hiPSC-Astrocytes

We first screened 11 different media conditions on forebrain-patterned NPCs ([Brennand et al., 2015](#); [Brennand](#)

[and Gage, 2011](#)) derived from hiPSCs ([Table 1](#)). The screening conditions, based on recently published hiPSC-astrocyte differentiation protocols ([Chen et al., 2015](#); [Haidet-Phillips et al., 2014](#); [Jiang et al., 2013](#); [Krencik et al., 2011](#); [McGivern et al., 2013](#); [Serio et al., 2013](#); [Shaltouki et al., 2013](#)), included different combinations of fibroblast growth factor 2 (FGF2) ([Haidet-Phillips et al., 2014](#)), ciliary neurotrophic factor (CNTF), ([Krencik et al., 2011](#); [Shaltouki et al., 2013](#)), bone morphogenetic protein 4 (BMP4) ([Han et al., 2013](#); [Jiang et al., 2013](#); [Shaltouki et al., 2013](#)), fibroblast bovine serum (FBS) ([Han et al., 2013](#); [Shaltouki et al., 2013](#)), neuregulin ([Pinkas-Kramarski et al., 1994](#); [Shaltouki et al., 2013](#)), insulin ([Heni et al., 2011](#)), and ascorbic acid (AA) ([Palm et al., 2015](#)), as well as three commercial astrocyte media (ScienCell, Gibco, and Lonza) for the culture of primary human fetal astrocytes ([Table 1](#)). Screening criteria included immunoreactivity for two classical markers of astrocyte identity, S100 β and glial fibrillary acidic protein (GFAP) ([Ludwin et al., 1976](#)), astrocyte morphology, survival, replicative ability, and cell line variability ([Table S1](#); [Figure S1A](#)). When tested on NPCs, most conditions resulted in limited cell proliferation and expression of neuronal markers ([Table S1](#)); however,



(legend on next page)



two commercial media, ScienCell and Lonza, yielded S100 β - and GFAP-positive astrocyte-like cells (Figures S1B–S1D). These results were confirmed across four representative NPC lines by both flow cytometry and immunocytochemistry by 30 days (Figures 1A and S1E–S1G). Culture of NPCs in both media, when combined with low initial seeding density (nearly single cells: 15,000 cells/cm²) and minimal serum exposure (1%–2%), resulted in astrocyte morphology within 10 days (Figure S1H); star-shaped astrocyte morphologies were evident within 30 days (Figure S1I). Although ScienCell and Lonza astrocyte media showed equivalent efficiencies (Figures S1B–S1D), ScienCell medium was selected owing to its lower cost and relative simplicity.

We next tested the efficacy of this protocol across 42 NPC lines from 30 individuals (including healthy controls as well as cases with schizophrenia and frontotemporal dementia; 16 male and 14 female) generated as part of three unique hiPSC cohorts (reprogrammed and differentiated through different protocols in independent laboratories) (Table S2). All 42 NPC lines ultimately yielded replicative cells with an astrocyte-like morphology (although not necessarily on the first attempt), with an average composition of 90% S100 β ⁺ and 82% GFAP⁺ cells by flow cytometry, relative to the appropriate secondary antibody control (Figures 1A and 1B; hiPSCs served as a negative control, lacking S100 β - and GFAP-positive cells), and confirmed by immunocytochemistry (Figures 1A, 1C, S1J, and S1K). As with primary astrocytes (Figures 1G and S1O), there is substantial variability in mean fluorescent intensity between S100 β - and GFAP-positive hiPSC-astrocyte populations (Figure 1A).

Within 30 days, hiPSC-astrocytes were immunopositive for the astrocyte markers ALDH1L1 (Cahoy et al., 2008)

and Vimentin (VIM) (Schnitzer et al., 1981), as well as the glutamate transporters GLAST (EAAT1 and GLUT1) (Rothstein et al., 1994) (Figures 1C, S1L, and S1M). Compared with NPCs or excitatory neurons (hNGN2-induced neurons) (Brennand et al., 2015), hiPSC-astrocytes expressed GFAP (Ludwin et al., 1976), S100 β (Ludwin et al., 1976), VIM (Schnitzer et al., 1981), AQU4 (Hubbard et al., 2015), ACSBG1 (Chaboub and Deneen, 2013), and APOE (Boyles et al., 1985) by qPCR (Figure 1D), although expression levels between individual hiPSC-astrocytes and primary astrocyte lines varied substantially. In addition, hiPSC-astrocytes expressed low levels of the neuronal markers *TUJ1*, *MAP2AB*, *RELN*, and *CACNA1C* (Figure 1E).

Across a larger panel of neural lineage markers (Table S4) in 23 hiPSC-astrocyte lines and their isogenic NPC lines, principal component analysis (PCA) revealed that the NPCs grouped together, while the astrocytes (both hiPSC-derived and primary human fetal astrocytes) were more dispersed (Figures 1F and S1N). Because it was apparent in our hiPSC-astrocytes as well as the primary human fetal astrocytes (obtained from different donors and brain regions), this variability in lineage marker expression may reflect inter-individual variability and/or differences in regional patterning (Figures 1G and S1O).

hiPSC-Astrocytes and Primary Human Fetal Astrocytes Share Similar Transcriptional Profiles

To query the extent to which global gene expression in hiPSC-astrocytes resembles primary human fetal brain astrocytes, we performed RNA sequencing (RNA-seq) transcriptomic analyses on four control hiPSC-astrocytes and primary human fetal astrocytes from two brain regions (cerebral cortex and midbrain), together with isogenic

Figure 1. Rapid Differentiation of hiPSC-Derived NPCs to Astrocyte-like Identity

(A) Representative flow-cytometry analysis of S100 β (top) and GFAP (bottom) for four 30-day hiPSC-astrocyte differentiations. Arrows indicate the cells positive for each marker protein. Appropriate secondary-only control is shown in black.

(B) Graphs of flow-cytometry analysis across 35 hiPSC-astrocyte differentiations from 26 NPC lines from three independent hiPSC cohorts. S100 β (left) and GFAP (right) immunostaining is shown, with primary human fetal astrocytes (positive control) and hiPSCs (negative control).

(C) Representative immunofluorescence images of hiPSC-astrocytes stained for astrocyte markers, glutamate transporters GLAST (green), VIMENTIN (green), ALDH1L1 (red), and APOE (green). Scale bars, 100 μ m.


(D) mRNA levels of astrocyte markers: *S100 β* , *GFAP*, *VIM*, *AQU4*, *ACSBG1*, and *APOE* in hiPSC-astrocytes (n = 3 from four different lines) and primary human fetal astrocytes (pAstrocytes; n = 3 from cerebral cortex astrocytes). Primer sequences are listed in Table S3. n, the number of independent experiments.

(E) mRNA levels of neuronal markers: *TUJ1*, *MAP2AB*, *RELN*, and *CACNA1C* in hiPSC-astrocytes (n = 3 from four different lines) and pAstrocytes (n = 3 from cerebral cortex astrocytes), relative to hNGN2-induced neurons (n = 3 from two lines). Primer sequences are listed in Table S3.

(F) Principal component analysis of lineage-specific marker expression in 23 pairs of hiPSC-astrocytes and isogenic source NPCs, together with four pAstrocyte lines isolated from fetal cerebral cortex, midbrain, cerebellum, and whole brain.

(G) Flow-cytometry analysis of S100 β (top) and GFAP (bottom) for pAstrocytes from the cerebral cortex, midbrain, cerebellum, and whole brain. Appropriate secondary-only control is shown in black. CC, cerebral cortex; Mid, midbrain; Cereb, cerebellum; Mixed, whole brain; hiPSC-Astros, hiPSC-astrocytes; pAstros, primary human fetal astrocytes.

Data are presented as mean \pm SD using two-tailed homoscedastic Student's t test. **p < 0.01, ***p < 0.001.



(legend on next page)



hiPSC-derived NPCs and neurons (Figures 2 and S2), comparing them with *in vivo* human and rodent astrocyte transcriptomic datasets (Figures 2, 3, S2, and S3). hiPSC-astrocytes showed transcriptional profiles most similar to those of fetal brain astrocytes. Using PCA and hierarchical clustering, all four hiPSC-astrocytes clustered together with the primary human fetal astrocytes and distinct from the NPC and neuron clusters (Figures 2A, 2B, and S2A).

There were nearly 6-fold fewer differentially expressed genes (DEGs) between hiPSC-astrocytes and primary human fetal astrocytes (900 genes) than between hiPSC-astrocytes and isogenic hiPSC-derived neurons (10,000 genes) or NPCs (5,500 genes) (Figures 2C and S2B). hiPSC-astrocytes are highly similar to primary human fetal astrocytes ($r = 0.945$); the majority of both expressed genes (counts per million [CPM] > 1) and enriched genes (CPM > 5) were shared between hiPSC-astrocytes and primary human fetal astrocytes (87.0% and 90.3%, respectively) (Figure 2D). Functional enrichment analyses (using MSigDB) demonstrated that signals regulating neuronal maturation, such as synapse or ion channel formation, were downregulated in hiPSC-astrocytes and primary human fetal astrocytes, whereas signals promoting extracellular cell adhesion and interaction were upregulated (Figure 2E). When we specifically considered just the top 100 most varying genes distinguishing hiPSC-astrocytes from hiPSC-derived NPCs and neurons, functional enrichment analysis identified a group of 19 genes related to reactivity, cytokine, interferon, T cell receptor (TCR), and antigen-processing signaling that were enriched in astrocytes (G2) (Figures 3A and S2C; Table 2).

To assess the extent to which hiPSC-astrocytes are related to human brain astrocytes *in vivo*, we compared expression profiles of hiPSC-derived astrocytes, NPCs,

and neurons from four healthy controls, as well as primary human fetal astrocytes, with the Allen BrainSpan Atlas of the Developing Human Brain (<http://www.brainspan.org>) (Miller et al., 2014) using a Spearman rank correlation analysis. Heatmaps generated through a Wilcoxon's rank-sum test revealed that gene expression in hiPSC-astrocytes and primary human fetal astrocytes was highly correlated with gene expression in human fetal cortical brain tissue (8–16 weeks post conception) (Figure 2F). Unsupervised hierarchical cell-type-specific cluster analysis demonstrated that hiPSC-astrocytes closely cluster with astrocytes purified by immunopanning from human brain tissue, particularly hippocampal astrocytes, rather than oligodendrocytes, endothelial cells, myeloid cells, or neurons (Zhang et al., 2016) (Figure 3A); pathway analysis further revealed that many astrocyte-enriched genes were related to immune signaling (Figure 2G).

Expression Profile of hiPSC-Astrocytes Resembles Quiescent Astrocytes

To determine whether the gene expression pattern of our control hiPSC-astrocytes more resembles quiescent or reactive astrocytes, we compared them with an RNA-seq dataset of “proinflammatory” A1-type and “immunoregulatory” A2-type murine astrocytes, as well as their saline- and sham-treated controls (Zamanian et al., 2012). (Because these are *in vivo* cell types, whereby A1-type astrocytes are induced by a bacterial lipopolysaccharide [LPS] infection and A2-type astrocytes by middle cerebral artery occlusion [MCAO], no comparable dataset exists for human astrocytes.) hiPSC-astrocyte gene expression best clusters with the control conditions (Figures 3B and S3A), suggesting that hiPSC-astrocytes may be closer to the quiescent state than to reactive astrocytes.

Figure 2. Transcriptional Profile of hiPSC-Astrocytes and Primary Human Fetal Astrocytes

RNA-seq analysis of hiPSC-derived NPCs ($n = 8$), neurons ($n = 6$), and astrocytes ($n = 4$) together with pAstrocytes from human fetal cerebral cortex and midbrain.

(A and B) Principal component (P.Co) analysis (A) and clustering diagram (B) of hiPSC-derived NPCs, neurons, and astrocytes, together with pAstrocytes.


(C) Volcano plot comparison of hiPSC-astrocytes to pAstrocytes (top), as well as to hiPSC-derived NPCs (middle) and neurons (bottom). Average \log_2 (fold change) versus $-\log_{10}$ (FDR) is shown for all genes. Genes upregulated and downregulated by 2-fold change and FDR < 0.05 are labeled by red dots. The number of genes differentially expressed between different cell types is indicated by the red color density and quantified in Figure S2B.

(D) Scatterplot (left) comparing gene expression in hiPSC-astrocytes and pAstrocytes. r represents the Spearman correlation coefficient. Venn diagram (right) of overlapping gene expression (CPM > 5) between hiPSC-astrocytes and pAstrocytes.

(E) Functional pathway enrichment analysis of differentially expressed genes between hiPSC-astrocytes and pAstrocytes (left), hiPSC-astrocytes and NPCs (middle), and pAstrocytes and NPCs (right); hiPSC-astrocytes and pAstrocytes express increased extracellular cell communication signals, but decreased neuronal signals relative to NPCs.

(F) Heatmap produced by Wilcoxon's rank-sum comparisons of hiPSC-derived NPCs, neurons, and astrocytes, as well as pAstrocytes, relative to the Allen BrainSpan Atlas.

(G) Fold enrichment from functional pathway analysis of astrocyte-enriched genes, group G2 sorted from top 100 most variable genes from Figure S3A and Table 2.



(legend on next page)



hiPSC-Astrocytes Secrete Cytokines in Response to Inflammatory Stimuli

A key effector of the astrocyte neuroinflammatory response in neurodegenerative diseases is interleukin 6 (IL-6) (Zhao and Schwartz, 1998). Consistent with effects observed in primary human fetal astrocytes (Figure 3C), 24-hr treatment with polyinosinic-polycytidylic acid (poly(I:C)) (50 ng/mL or 100 ng/mL), LPS (10 μ g/mL or 50 μ g/mL), and β -amyloid 42 (A β 42) (5 μ M or 10 μ M) led to dose-dependent increases in IL-6 secretion in control hiPSC-astrocytes (Figure 3D), but not the source hiPSC-derived NPCs (Figure 3C).

To more fully characterize the release of inflammatory mediators from control hiPSC-astrocytes following 24-hr treatment with 5 μ M A β 42, a main component of the amyloid plaques in AD, we measured proinflammatory (IL-1 β , IL-4, IL-6, IL-8, IL-10, and tumor necrosis factor α) and anti-inflammatory cytokines (IL-1 α , IL-2, IL-12, IL-17 α , interferon- γ , and granulocyte macrophage colony-stimulating factor) by Multi-Analyte ELISArray, confirming that A β 42 treatment primarily increased IL-6 secretion (Garwood et al., 2011), as well as a second proinflammatory cytokine, IL-8 (Figures S3B and S3C). We also measured 36 cytokines, chemokines, and acute-phase proteins using the Proteome Profiler Human Cytokine Array (Table S5) in baseline conditions and following 24-hr treatment with 5 μ M A β 42 (Figures 3D and S3C–S3E); A β 42 increased IL-6 release in both hiPSC-astrocytes and primary human fetal astrocytes (Figure S3D). Together, our findings indicate that hiPSC-astrocytes are capable of secreting proinflammatory cytokines in response to neuroinflammatory cues.

hiPSC-Astrocytes Display Phagocytic Capacity and Promote Phagocytic Function of Microglia

Astrocytes can phagocytose and degrade β -amyloid in AD (Chung et al., 2013; Wyss-Coray et al., 2003). We used flow cytometry to examine the ability of control hiPSC-astrocytes to phagocytose pHrodo red conjugated myelin or zymosan bioparticles (pHrodo red dye fluoresces red in

the acidic environment of phagosomes; zymosan is an agonist of Toll-like receptor 2 [TLR2] [Marinelli et al., 2015; Richard et al., 2008]). hiPSC-astrocytes, primary human fetal astrocytes, and BV2 microglia showed a similar capacity to internalize myelin purified from brain homogenates, while zymosan bioparticle uptake was much greater in microglia (consistent with higher levels of TLR2 reported in microglia) (Jana et al., 2008) (Figures 4A, 4B, and S4A–S4C; Movies S1–S3). The specificity of bioparticle uptake was confirmed by treating cells with cytochalasin D, an inhibitor of β -actin polymerization that reduces phagocytosis (Figures 4A and 4B).

Although microglia are the major phagocytic cells in the brain, astrocytes mediate microglial response to inflammatory stimuli (Krencik et al., 2011; Lee and Landreth, 2010; Skripuletz et al., 2013). We repeated the pHrodo red zymosan assay using microglia-astrocyte co-cultures; both hiPSC-astrocytes and primary human fetal astrocytes enhanced phagocytic capacity of BV2 mouse microglial cells (Figures 4C and S4D). We discriminated microglia and astrocyte phagocytic activity by labeling CD11b- and GFAP-positive cells, respectively (Figure S4E). After 3 hr of labeling, only the CD11b⁺ BV2 microglial cells phagocytosed the pHrodo red zymosan bioparticles (Figure S4E). Finally, we treated BV2 microglial cells with astrocyte conditioned medium (ACM) for 24 hr prior to analyzing their phagocytic capacity; pretreatment of BV2 microglial cells with ACM from either hiPSC-astrocytes or primary human fetal astrocytes significantly increased microglial phagocytic capacity (Figures 4D and S4F). Taken together, these findings show that both hiPSC-astrocytes and primary human fetal astrocytes secrete factors that increase the capacity of microglia to phagocytose zymosan bioparticles.

hiPSC-Astrocytes Display Spontaneous Calcium Transient Activity

We next evaluated whether control hiPSC-astrocytes exhibit spontaneous calcium transients, as has been

Figure 3. Characterization of the Neuroinflammatory Status and Reactivity of hiPSC-Astrocytes

(A) Cluster analysis of hiPSC-derived NPCs, neurons, and astrocytes, as well as pAstrocytes, combined with a human adult brain dataset (Zhang et al., 2016). hiPSC-Astros, hiPSC-astrocytes; pAstros, primary human fetal astrocytes; FDR, false discovery rate; CPM, counts per million.

(B) Cluster diagram of hiPSC-astrocytes and pAstrocytes compared with the astrocyte reactivity dataset (Zamarian et al., 2012), which was sorted by reactivity genes enriched in the A1 (LPS-treatment), A2 (MCAO ischemia), and pan-reactive phenotypes, or related controls (saline and sham, respectively).

(C) IL-6 secretion from pAstrocytes (cerebral cortex, midbrain, cerebellum and whole brain [mixed regions]) and negative controls (fibroblasts and NPCs), after 24-hr treatment with 50 ng/mL or 100 ng/mL poly(I:C), 10 μ g/mL or 50 μ g/mL LPS, or 5 μ M or 10 μ M A β 42 and its relative vehicle controls (saline for poly(I:C) and LPS, and Tris-HCl [pH 8] for A β (1–42)), as measured by ELISA.

(D) IL-6 secretion following 24-hr treatment with poly(I:C), LPS, and A β 42 across hiPSC-astrocyte differentiations from nine NPC lines from three independent hiPSC cohorts. hiPSC-Astros, hiPSC-astrocytes; pAstros, primary human fetal astrocytes.

Data are presented as mean \pm SD using one-way ANOVA with Tukey multiple comparison test. n.s., not significant; * p < 0.05, *** p < 0.001.



Table 2. Annotation of the Top 100 Most Variable Genes

MSigDB	Group	Overlap Size	Group Size	MSigDB Size	FE	p Value	FDR	Overlap Genes
Antigen processing and presentation	G2	7	19	57	151.3	1.71×10^{-14}	8.49×10^{-11}	CD74; HLA-DQA1; HLA-DQB1; HLA-DRA; HLA-DRB1; HLA-DRB5
Cell adhesion molecules (CAMs)	G2	6	19	115	64.3	3.17×10^{-10}	3.61×10^{-7}	HLA-DQA1; HLA-DQB1; HLA-DRA; HLA-DRB1; HLA-DRB5
Reactome MHC class II antigen presentation	G2	5	19	86	71.6	6.64×10^{-9}	6.94×10^{-6}	CD74; HLA-DQA1; HLA-DQB1; HLA-DRB1; HLA-DRB5
Reactome translocation of ZAP70 to immunological synapse	G2	3	19	7	528	1.58×10^{-8}	1.53×10^{-5}	HLA-DQA1; HLA-DRB1; HLA-DRB5
Reactome phosphorylation of CD3 and TCR ζ chains	G2	3	19	10	369.6	5.42×10^{-8}	4.85×10^{-5}	HLA-DQA1; HLA-DRB1; HLA-DRB5
Reactome PD1 signaling	G2	3	19	11	336	7.45×10^{-8}	6.22×10^{-5}	HLA-DQA1; HLA-DRB1; HLA-DRB5
Reactome generation of second messenger molecules	G2	3	19	18	205.3	3.67×10^{-7}	0.000287	HLA-DQA1; HLA-DRB1; HLA-DRB5
Reactome downstream TCR signaling	G2	3	19	27	136.9	1.31×10^{-6}	0.000965	HLA-DQA1; HLA-DRB1; HLA-DRB5
Neuron differentiation	G1	4	28	73	45.8	1.68×10^{-6}	0.001205	CNTN4; LMX1B; OTX2; SLIT1
Generation of neurons	G1	4	28	79	42.3	2.31×10^{-6}	0.001574	CNTN4; LMX1B; OTX2; SLIT1
Axon guidance	G1	3	28	22	114	2.32×10^{-6}	0.001574	CNTN4; OTX2; SLIT1
Neurogenesis	G1	4	28	89	37.6	3.73×10^{-6}	0.002458	CNTN4; LMX1B; OTX2; SLIT1
Nervous system development	G1	6	28	366	13.7	3.95×10^{-6}	0.002538	CNTN4; LMX1B; LY6H; OTX2; SHOX2; SLIT1
Reactome TCR signaling	G2	3	19	41	90.1	4.74×10^{-6}	0.002876	HLA-DQA1; HLA-DRB1; HLA-DRB5
Nervous system development	S1	5	17	366	18.8	4.82×10^{-6}	0.002876	CNTN4; LMX1B; LY6H; SHOX2; SLIT1
Biocarta TCRA pathway	G2	2	19	5	492.8	6.23×10^{-6}	0.003549	HLA-DRA; HLA-DRB1
Reactome co-stimulation by the CD28 family	G2	3	19	52	71.1	9.77×10^{-6}	0.005322	HLA-DQA1; HLA-DRB1; HLA-DRB5
Reactome interferon- γ signaling	G2	3	19	57	64.8	1.29×10^{-5}	0.006553	HLA-DQA1; HLA-DRB1; HLA-DRB5
Biocarta IL-5 pathway	G2	2	19	7	352	1.31×10^{-5}	0.006553	HLA-DRA; HLA-DRB1
Axonogenesis	G1	3	28	42	59.7	1.71×10^{-5}	0.007939	CNTN4; OTX2; SLIT1
Biocarta asbcell pathway	G2	2	19	8	308	1.74×10^{-5}	0.007939	HLA-DRA; HLA-DRB1
Neuron differentiation	S1	3	17	73	56.6	1.92×10^{-5}	0.00858	CNTN4; LMX1B; SLIT1

MSigDB (<http://software.broadinstitute.org/gsea/msigdb>) is a collection of gene annotations, including gene ontology, and functional pathways. Overlap size: the number of genes shared between a query gene set (e.g., genes in group 1, G1) and a gene set in MSigDB (e.g., antigen processing and presentation). Group size: the number of genes in a query gene set. MSigDB size: the number of genes in MSigDB gene set. FE, fold enrichment; FDR, false discovery rate.

described for astrocytes (Volterra et al., 2014). We used the calcium indicator Fluo-4AM to monitor calcium signaling under basal conditions and in response to a pulse of extracellular glutamate (3 μ M) (Zhang et al., 2016). A single pulse of glutamate produced a slow calcium response in

both hiPSC-astrocytes and primary human fetal astrocytes (Figure 5). In addition, some cells exhibited spontaneous calcium spikes, suggesting the presence of a network of connected astrocytes (Scemes and Giaume, 2006) (Figures 5A–5D). To quantify these responses, we measured the




Figure 4. Impact of hiPSC-Astrocytes on the Phagocytic Capacity of BV2 Microglial Cells

(A) Phagocytic indices of BV2 cells, hiPSC-astrocytes, and pAstrocytes incubated with 20 μ g of pHrodo-labeled myelin for 3 hr and analyzed by flow cytometry.

(B) Amnis images representative of pHrodo red myelin after engulfment in hiPSC-astrocytes, pAstrocytes, and BV2 cells.

(C) Phagocytic indices of BV2 microglia co-cultured with hiPSC-astrocytes and pAstrocytes that were treated with 30 μ g of pHrodo-labeled zymosan for 3 hr and analyzed by flow cytometry, $F(3, 33) = 79.33$.

(D) Phagocytic indices of BV2 microglia treated with astrocyte conditioned medium for 20 hr and then incubated with 30 μ g of pHrodo-labeled zymosan for 3 hr for analysis by flow cytometry, $F(3, 13) = 30.80$. Data are representative of three independent experiments from 6–8 different control hiPSC-astrocytes and 2–4 different

pAstrocytes and are shown as mean \pm SEM. Similar significance was obtained in two other independent experiments. hiPSC-Astros, hiPSC-astrocytes; pAstros, primary human fetal astrocytes. Treatment with 2 μ M cytochalasin D (Cyt) was used as a negative control for phagocytosis inhibition.

frequency of spontaneous activity, the number of spontaneous spikes per time, and the amplitude of the calcium spike (Figures 5E–5G). There was no statistical difference in the frequency and number of spikes between hiPSC-astrocytes and primary astrocytes. Interestingly, the amplitude of the spontaneous calcium spike was significantly higher in hiPSC-astrocytes compared with primary astrocytes. Taken together, the excitability of hiPSC-astrocytes was largely indistinguishable from that of primary human astrocytes.

DISCUSSION

We screened 11 methods to differentiate hiPSC-derived NPCs into astrocytes, identifying a medium with low FBS (1%–2%) that produces populations of hiPSC-astrocytes within 30 days. Although this is a commercial medium sold for the culture of primary astrocytes, it has not previously been demonstrated to support differentiation of hiPSC to astrocytes. Our method is fast and robust: unlike previous reports, it does not require prolonged culture (~6 months) or a serial sorting process (Krencik et al., 2011; Shaltouki et al., 2013; Yuan et al., 2011). This protocol was evaluated across 42 NPC lines from 30 individuals (16 males and 14 females) generated from three independent hiPSC cohorts (Table S2). We caution that the quality of

the starting NPC population is a critical predictor of success and note that for a few particularly intransient NPC lines, starting from very low-passage stocks proved critical to the ultimate successful differentiation of hiPSC-astrocytes. By both fluorescence-activated cell sorting and qPCR, GFAP seems to be a more variable marker of astrocyte fate (Figures 1A and 1B) and we recommend instead using S100 β when evaluating hiPSC-astrocyte populations; moreover, more than any single gene, using panels of markers and/or global transcriptomic analysis more fully reveals the extent of astrocyte patterning of each line. We also suggest that hiPSC-astrocyte response to inflammatory stimuli might represent a simple and easy platform by which to confirm the functionality of hiPSC-astrocytes; astrocytes undergo stimulation-dependent cytokine secretion, while NPCs do not secrete cytokines.

To our knowledge, our method is distinct from others recently reported (Chen et al., 2015; Haidet-Phillips et al., 2014; Jiang et al., 2013; Krencik et al., 2011; McGivern et al., 2013; Serio et al., 2013; Shaltouki et al., 2013), its major advantage being the ease by which hiPSC-astrocytes can be concurrently differentiated from many NPCs from multiple individuals, owing to the simplicity of the protocol, and furthermore that astrocytes can be expanded for months and cryopreserved, serving as a source of cells to support an array of experiments. Therefore, because our protocols for NPC and astrocyte culture are robust and




Figure 5. Spontaneous and Glutamate-Responsive Calcium Transients in hiPSC-Astrocytes and Primary Astrocytes

(A and C) Representative Fluo-4-stained hiPSC-astrocytes and pAstrocytes. Note similarity in shape of cells.

(B and D) Plots of fluorescence (RFUs) versus time for 25 regions of interest from hiPSC-astrocytes and pAstrocytes.

(E) Average number of spontaneously active cells per field ($318 \mu\text{m}^2$) for hiPSC-astrocytes ($n = 22$ fields from four different lines) and pAstrocytes ($n = 12$ fields from three different preparations of cerebral cortex astrocytes). n, the number of fields.

(F) Average number of calcium spikes per 348-s trace for hiPSC-astrocytes ($n = 22$) and pAstrocytes ($n = 12$).

(G) Average amplitude of calcium spike in each 348-s trace for hiPSC-astrocytes ($n = 22$) and pAstrocytes ($n = 12$). Peak excludes the amplitude of the glutamate-induced spike. Each point represents the average multiple regions of interest per $318 \mu\text{m}^2$. Line shows mean \pm SEM. hiPSC-Astrocytes, hiPSC-astrocytes, pAstrocytes, primary human fetal astrocytes. Using a two-tailed Student's t test, n.s., not significant; ** $p < 0.05$.



easily scaled, this methodology is both preferable for large case-control studies from dozens of idiopathic patients and highly amenable to automated culture and future high-throughput drug screens using patient-derived astrocytes (Xu et al., 2016).

We have established a platform for querying astrocyte-specific contributions to disease predisposition in hiPSC-based models. hiPSC-astrocytes closely resemble primary human fetal astrocytes, purified adult brain astrocytes (Zhang et al., 2016), and brain tissue homogenate (Miller et al., 2014) (Figures 2 and 3A). Although the transcriptional profile of our hiPSC-astrocytes most closely resembles a quiescent state (Figure 3B), hiPSC-astrocytes secrete various cytokines and chemokines in response to neuroinflammatory stimuli (Figures 3C and 3D). hiPSC-astrocytes displayed phagocytic capacity (Figure 4A) and enhanced the phagocytic function of microglia in a co-culture assay (Figure 4C). They also showed spontaneous calcium transient activity and responses to glutamate stimulus (Figure 5). Moreover, because nearly pure populations of excitatory neurons and astrocytes from the same individual, hiPSC clone, and even NPC batch can now be compared, our protocol may reduce experimental variation in studies of the cell-autonomous and non-cell-autonomous factors underlying disease predisposition.

While many studies of neurodegeneration and synaptic dysfunction use either primary mouse or human fetal glia for human neuron co-culture (Johnson et al., 2007; Kiskinis et al., 2014; Verheyen et al., 2015), these methods are limited by species-specific differences in astrocyte functions and limited access to human fetal samples. Our hiPSC-astrocytes overcome these drawbacks, making possible hiPSC-based astrocyte-neuron (Jiang et al., 2013) and astrocyte-microglia (Muffat et al., 2016; Pandya et al., 2017) co-culture platforms for uncovering disease-related mechanisms *in vitro*, which should help to reveal how the crosstalk between these three neural cell types contributes to neurological and psychiatric disease.

EXPERIMENTAL PROCEDURES

Human subject work on these de-identified control hiPSCs was approved by the Institutional Review Board at Icahn School of Medicine at Mount Sinai. Control mouse brain tissue was obtained from the Center for Comparative Medicine and Surgery at Icahn School of Medicine at Mount Sinai.

NPCs and Astrocyte Cell Culture

Forebrain NPCs were maintained at high density, grown on Matrigel (BD Bioscience) in NPC medium (DMEM/F12, 1× N2, 1× B27-RA [Invitrogen], 1 mg/mL laminin [Invitrogen], and 20 ng/mL FGF2 [Invitrogen]), and split approximately 1:3 to 1:4 every week with Accutase (Millipore). NPCs could be expanded

up to 14 passages. Forebrain NPCs were differentiated to astrocytes by seeding dissociated single cells at 15,000 cells/cm² density on Matrigel-coated plates in astrocyte medium (ScienCell: 1801, astrocyte medium [1801-b], 2% fetal bovine serum [0010], astrocyte growth supplement [1852] and 10 U/mL penicillin/streptomycin solution [0503]). Initial NPC quality, seeding density, and single-cell dissociation are critical, particularly during the first 30 days of differentiation, in order to efficiently generate hiPSC-astrocytes. See detailed protocol for astrocyte differentiation from hiPSC-derived NPCs in the [Supplemental Information](#).

Immunocytochemistry

Cultures were washed with PBS and fixed using 4% paraformaldehyde (no. 15714, Electron Microscopy Sciences) for 20 min, then washed three times with PBS 1× plus 0.01% Triton X-100 prior to preincubation with blocking solution made of PBS 1× plus 0.01% Triton X-100 and 10% donkey serum (Sigma-Aldrich), for at least 1 hr. Cultures were then incubated in primary antibody solution overnight at 4°C, followed by an incubation with secondary antibodies for 1 hr at room temperature. See [Supplemental Experimental Procedures](#) for antibody details and confocal microscopy.

Real-Time qPCR

The total RNA was extracted from cultured cells with an RNeasy mini kit (no. 74106, Qiagen) and reverse transcribed into cDNA with an iScript cDNA Synthesis kit (no. 1708890, Bio-Rad). cDNA was used as template for the qPCR using a 7900 Real-Time PCR system (Applied Biosystems) with Power SYBR Green PCR Master Mix. The primers used are listed in [Table S3](#). Gene expression was analyzed using the $\Delta\Delta C_t$ method. All results were normalized to *GAPDH*, *β -ACTIN*, and *TBP* (TATA-box binding protein) expression, and the values of uninduced fibroblasts were set to 1. Three replicates were used to determine the error bars. See the [Supplemental Information](#) for RNA expression studies for qRT-PCR and primer sequences.

Analysis of RNA-Seq Data

Pair-ended RNA-seq data was generated using the Illumina HiSeq 2500 platform for hiPSC-astrocytes (four lines), neurons (six lines), and NPCs (eight lines) ([Table S2](#)), as well as for two primary astrocyte samples. The pair-ended sequencing reads were aligned to human hg19 genome using Star Aligner (version 2.5.0b). Following read alignment, featureCounts ([Liao et al., 2014](#)) was used to quantify the gene expression at the gene level based on Ensembl gene model GRCh37.70. Gene expression data preprocessing and downstream analyses, including differential gene expression and functional enrichment, are detailed in [Supplemental Experimental Procedures](#).

IL-6 ELISA, Multi-Analyte ELISArray, and Protein Array

hiPSC-astrocytes and primary astrocytes were seeded in astrocyte medium 1 day before the experiment. Cells were treated for 24 hr with 50 ng/mL or 100 ng/mL of poly(I:C) (InvivoGen, no. ttrl-pic), 10 μ g/mL or 50 μ g/mL of LPS (Sigma-Aldrich, no.



L5886), and 5 μM or 10 μM human β -amyloid (A β 42) (California-Peptide, no. 641-15) and vehicle control solutions (saline for poly(I:C) and LPS or Tris-HCl for A β 42). Samples were analyzed with IL-6 ELISA assay (Affymetrix eBioscience, no. 88-7066), Human Inflammatory Cytokines Multi-Analyte ELISArray (Qiagen, no. MEH-004A), RT2 Profiler PCR Arrays (Qiagen), and Proteome Profiler Human Cytokine Array (R&D Systems, no. ARY005B). For a detailed protocol see [Supplemental Experimental Procedures](#).

Phagocytosis Assay

Phagocytic capacity of hiPSC-astrocytes, primary astrocytes, and BV2 cells was analyzed by incubating cells with pHrodo-labeled zymosan (Thermo Fisher, P35364) or myelin purified from mouse brains, and then measured by flow cytometry. Detailed protocols for co-culture experiments and astrocyte conditioned media treatment are provided in the [Supplemental Experimental Procedures](#).

Calcium Imaging

Cells were incubated with 2 μM Fluo-4AM (Molecular Probes) and 0.02% Pluronic F 127 detergent in Krebs HEPES buffer (KHB). Time-lapse image sequences ($\times 40$ magnification) were acquired at 0.9 Hz and analyzed by FluoroSNNAP software operated by MATLAB. For a detailed protocol see [Supplemental Experimental Procedures](#).

Statistical Analysis

For all experiments, data are represented as mean \pm SD or SEM of three to five biological replicates. Statistical significance was determined using a two-tailed homoscedastic Student's *t* test or one-way ANOVA and Tukey-Kramer post hoc test for differences of means between each group of data with parametric distribution. Significant comparisons are labeled in the figures as **p* < 0.05, ***p* < 0.01, and ****p* < 0.001.

Extended experimental procedures and any associated references are available in the online version of this paper.

ACCESSION NUMBERS

The accession number for the RNA-seq data reported in this paper is GEO: GSE97904.

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, four figures, and five tables and can be found with this article online at <http://dx.doi.org/10.1016/j.stemcr.2017.06.018>.

AUTHOR CONTRIBUTIONS

J.TCW, K.J.B., and A.M.G. designed the experiments and wrote the manuscript. J.TCW developed the astrocyte differentiation protocol. J.TCW, K.R.B., B.J.H., and R.A. (with H.P.) differentiated the hiPSC-astrocyte lines. M.W. and B.Z. conducted the RNA-seq analysis. J.TCW performed the neuroinflammatory assays. A.A.P., J.TCW, and S.I.M. performed the phagocytosis assays. J.TCW, A.A.P., and B.J.H. executed flow-cytometry analysis on astrocytes and NPCs. J.TCW and K.R.B. performed real-time qPCR and immu-

nofluorescence staining. E.L. and P.A.S. conducted and analyzed the calcium imaging analysis. C.M.K. shared prepublication information about astrocyte differentiation conditions.

ACKNOWLEDGMENTS

BV2 cell line was kindly provided by Marc Diamond (UT Southwestern Medical Center). We are grateful for the continuous support of the Flow Cytometry Core at the Icahn School of Medicine at Mount Sinai Hospital. This work was supported in part by the JPB Foundation (A.M.G.), the Rainwater Foundation (A.M.G.), NIMH R01MH101454 (K.J.B.), NIA U01P50AG005138-30-1 (Alzheimer's Disease Research Center: Pilot 30-1) (K.J.B.), NIA/NIH U01AG046170 (K.J.B., M.W., and B.Z.), a component of the AMP-AD Target Discovery and Preclinical Validation Project, NIMH R01MH11499 (P.A.S.), NIAAA R01AA018734 (P.A.S.), NIDA R01DA037170 (P.A.S.), K01AG046374 (C.M.K.), and the New York Stem Cell Foundation (K.J.B.) and Project ALS (H.P.). A.M.G. serves on the Scientific Advisory Board for Denali Therapeutics.

Received: April 30, 2017

Revised: June 28, 2017

Accepted: June 29, 2017

Published: July 27, 2017

REFERENCES

- Ballas, N., Liou, D.T., Grunseich, C., and Mandel, G. (2009). Non-cell autonomous influence of MeCP2-deficient glia on neuronal dendritic morphology. *Nat. Neurosci.* *12*, 311–317.
- Barres, B.A. (2008). The mystery and magic of glia: a perspective on their roles in health and disease. *Neuron* *60*, 430–440.
- Bi, F., Huang, C., Tong, J., Qiu, G., Huang, B., Wu, Q., Li, F., Xu, Z., Bowser, R., Xia, X.G., et al. (2013). Reactive astrocytes secrete lcn2 to promote neuron death. *Proc. Natl. Acad. Sci. USA* *110*, 4069–4074.
- Boyles, J.K., Pitas, R.E., Wilson, E., Mahley, R.W., and Taylor, J.M. (1985). Apolipoprotein E associated with astrocytic glia of the central nervous system and with nonmyelinating glia of the peripheral nervous system. *J. Clin. Invest.* *76*, 1501–1513.
- Bradford, J., Shin, J.Y., Roberts, M., Wang, C.E., Li, X.J., and Li, S. (2009). Expression of mutant huntingtin in mouse brain astrocytes causes age-dependent neurological symptoms. *Proc. Natl. Acad. Sci. USA* *106*, 22480–22485.
- Brennand, K.J., and Gage, F.H. (2011). Concise review: the promise of human induced pluripotent stem cell-based studies of schizophrenia. *Stem Cells* *29*, 1915–1922.
- Brennand, K., Savas, J.N., Kim, Y., Tran, N., Simone, A., Hashimoto-Torii, K., Beaumont, K.G., Kim, H.J., Topol, A., Ladrán, I., et al. (2015). Phenotypic differences in hiPSC NPCs derived from patients with schizophrenia. *Mol. Psychiatry* *20*, 361–368.
- Cahoy, J.D., Emery, B., Kaushal, A., Foo, L.C., Zamanian, J.L., Christopherson, K.S., Xing, Y., Lubischer, J.L., Krieg, P.A., Krumpal, S.A., et al. (2008). A transcriptome database for astrocytes, neurons, and oligodendrocytes: a new resource for understanding brain development and function. *J. Neurosci.* *28*, 264–278.



- Chaboub, L.S., and Deneen, B. (2013). Astrocyte form and function in the developing central nervous system. *Semin. Pediatr. Neurol.* *20*, 230–235.
- Chen, H., Qian, K., Chen, W., Hu, B., Blackbourn, L.W., Du, Z., Ma, L., Liu, H., Knobel, K.M., Ayala, M., et al. (2015). Human-derived neural progenitors functionally replace astrocytes in adult mice. *J. Clin. Invest.* *125*, 1033–1042.
- Chung, W.S., Clarke, L.E., Wang, G.X., Stafford, B.K., Sher, A., Chakraborty, C., Joung, J., Foo, L.C., Thompson, A., Chen, C., et al. (2013). Astrocytes mediate synapse elimination through MEGF10 and MERTK pathways. *Nature* *504*, 394–400.
- Di Giorgio, F.P., Boulting, G.L., Bobrowicz, S., and Eggan, K.C. (2008). Human embryonic stem cell-derived motor neurons are sensitive to the toxic effect of glial cells carrying an ALS-causing mutation. *Cell Stem Cell* *3*, 637–648.
- Eroglu, C., and Barres, B.A. (2010). Regulation of synaptic connectivity by glia. *Nature* *468*, 223–231.
- Freeman, M.R., and Rowitch, D.H. (2013). Evolving concepts of gliogenesis: a look way back and ahead to the next 25 years. *Neuron* *80*, 613–623.
- Garwood, C.J., Pooler, A.M., Atherton, J., Hanger, D.P., and Noble, W. (2011). Astrocytes are important mediators of Abeta-induced neurotoxicity and tau phosphorylation in primary culture. *Cell Death Dis.* *2*, e167.
- Haidet-Phillips, A.M., Roybon, L., Gross, S.K., Tuteja, A., Donnelly, C.J., Richard, J.P., Ko, M., Sherman, A., Eggan, K., Henderson, C.E., et al. (2014). Gene profiling of human induced pluripotent stem cell-derived astrocyte progenitors following spinal cord engraftment. *Stem Cells Transl. Med.* *3*, 575–585.
- Han, X., Chen, M., Wang, F., Windrem, M., Wang, S., Shanz, S., Xu, Q., Oberheim, N.A., Bekar, L., Betstadt, S., et al. (2013). Forebrain engraftment by human glial progenitor cells enhances synaptic plasticity and learning in adult mice. *Cell Stem Cell* *12*, 342–353.
- Heni, M., Hennige, A.M., Peter, A., Siegel-Axel, D., Ordelheide, A.M., Krebs, N., Machicao, F., Fritsche, A., Haring, H.U., and Staiger, H. (2011). Insulin promotes glycogen storage and cell proliferation in primary human astrocytes. *PLoS One* *6*, e21594.
- Hubbard, J.A., Hsu, M.S., Seldin, M.M., and Binder, D.K. (2015). Expression of the astrocyte water channel aquaporin-4 in the mouse brain. *ASNNeuro* *7*. <http://dx.doi.org/10.1177/1759091415605486>.
- Jana, M., Palencia, C.A., and Pahan, K. (2008). Fibrillar amyloid-beta peptides activate microglia via TLR2: implications for Alzheimer's disease. *J. Immunol.* *181*, 7254–7262.
- Jiang, P., Chen, C., Wang, R., Chechneva, O.V., Chung, S.H., Rao, M.S., Pleasure, D.E., Liu, Y., Zhang, Q., and Deng, W. (2013). hESC-derived Olig2+ progenitors generate a subtype of astroglia with protective effects against ischaemic brain injury. *Nat. Commun.* *4*, 2196.
- Johnson, M.A., Weick, J.P., Pearce, R.A., and Zhang, S.C. (2007). Functional neural development from human embryonic stem cells: accelerated synaptic activity via astrocyte coculture. *J. Neurosci.* *27*, 3069–3077.
- Kiskinis, E., Sandoe, J., Williams, L.A., Boulting, G.L., Moccia, R., Wainger, B.J., Han, S., Peng, T., Thams, S., Mikkilineni, S., et al. (2014). Pathways disrupted in human ALS motor neurons identified through genetic correction of mutant SOD1. *Cell Stem Cell* *14*, 781–795.
- Krencik, R., Weick, J.P., Liu, Y., Zhang, Z.J., and Zhang, S.C. (2011). Specification of transplantable astroglial subtypes from human pluripotent stem cells. *Nat. Biotechnol.* *29*, 528–534.
- Lee, C.Y., and Landreth, G.E. (2010). The role of microglia in amyloid clearance from the AD brain. *J. Neural Transm. (Vienna)* *117*, 949–960.
- Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* *30*, 923–930.
- Ludwin, S.K., Kosek, J.C., and Eng, L.F. (1976). The topographical distribution of S-100 and GFA proteins in the adult rat brain: an immunohistochemical study using horseradish peroxidase-labelled antibodies. *J. Comp. Neurol.* *165*, 197–207.
- Marchetto, M.C., Muotri, A.R., Mu, Y., Smith, A.M., Cezar, G.G., and Gage, F.H. (2008). Non-cell-autonomous effect of human SOD1 G37R astrocytes on motor neurons derived from human embryonic stem cells. *Cell Stem Cell* *3*, 649–657.
- Marinelli, C., Di Liddo, R., Facci, L., Bertalot, T., Conconi, M.T., Zusso, M., Skaper, S.D., and Giusti, P. (2015). Ligand engagement of Toll-like receptors regulates their expression in cortical microglia and astrocytes. *J. Neuroinflammation* *12*, 244.
- McGivern, J.V., Patitucci, T.N., Nord, J.A., Barabas, M.E., Stucky, C.L., and Ebert, A.D. (2013). Spinal muscular atrophy astrocytes exhibit abnormal calcium regulation and reduced growth factor production. *Glia* *61*, 1418–1428.
- Mertens, J., Marchetto, M.C., Bardy, C., and Gage, F.H. (2016). Evaluating cell reprogramming, differentiation and conversion technologies in neuroscience. *Nat. Rev. Neurosci.* *17*, 424–437.
- Miller, J.A., Ding, S.L., Sunkin, S.M., Smith, K.A., Ng, L., Szafer, A., Ebbert, A., Riley, Z.L., Royall, J.J., Aiona, K., et al. (2014). Transcriptional landscape of the prenatal human brain. *Nature* *508*, 199–206.
- Muffat, J., Li, Y., Yuan, B., Mitalipova, M., Omer, A., Corcoran, S., Bakiasi, G., Tsai, L.H., Aubourg, P., Ransohoff, R.M., et al. (2016). Efficient derivation of microglia-like cells from human pluripotent stem cells. *Nat. Med.* *22*, 1358–1367.
- Oberheim, N.A., Takano, T., Han, X., He, W., Lin, J.H., Wang, F., Xu, Q., Wyatt, J.D., Pilcher, W., Ojemann, J.G., et al. (2009). Uniquely hominid features of adult human astrocytes. *J. Neurosci.* *29*, 3276–3287.
- Palm, T., Bolognin, S., Meiser, J., Nickels, S., Trager, C., Meilenbrock, R.L., Brockhaus, J., Schreitmuller, M., Missler, M., and Schwamborn, J.C. (2015). Rapid and robust generation of long-term self-renewing human neural stem cells with the ability to generate mature astroglia. *Sci. Rep.* *5*, 16321.
- Pandya, H., Shen, M.J., Ichikawa, D.M., Sedlock, A.B., Choi, Y., Johnson, K.R., Kim, G., Brown, M.A., Elkhouloun, A.G., Maric, D., et al. (2017). Differentiation of human and murine induced pluripotent stem cells to microglia-like cells. *Nat. Neurosci.* *20*, 753–759.
- Papadeas, S.T., Kraig, S.E., O'Banion, C., Lepore, A.C., and Maragakis, N.J. (2011). Astrocytes carrying the superoxide dismutase 1 (SOD1G93A) mutation induce wild-type motor neuron degeneration in vivo. *Proc. Natl. Acad. Sci. USA* *108*, 17803–17808.



- Pinkas-Kramarski, R., Eilam, R., Spiegler, O., Lavi, S., Liu, N., Chang, D., Wen, D., Schwartz, M., and Yarden, Y. (1994). Brain neurons and glial cells express Neu differentiation factor/heregulin: a survival factor for astrocytes. *Proc. Natl. Acad. Sci. USA* *91*, 9387–9391.
- Richard, K.L., Filali, M., Prefontaine, P., and Rivest, S. (2008). Toll-like receptor 2 acts as a natural innate immune receptor to clear amyloid beta 1-42 and delay the cognitive decline in a mouse model of Alzheimer's disease. *J. Neurosci.* *28*, 5784–5793.
- Rothstein, J.D., Martin, L., Levey, A.I., Dykes-Hoberg, M., Jin, L., Wu, D., Nash, N., and Kuncl, R.W. (1994). Localization of neuronal and glial glutamate transporters. *Neuron* *13*, 713–725.
- Roybon, L., Lamas, N.J., Garcia-Diaz, A., Yang, E.J., Sattler, R., Jackson-Lewis, V., Kim, Y.A., Kachel, C.A., Rothstein, J.D., Przedborski, S., et al. (2013). Human stem cell-derived spinal cord astrocytes with defined mature or reactive phenotypes. *Cell Rep.* *4*, 1035–1048.
- Scemes, E., and Giaume, C. (2006). Astrocyte calcium waves: what they are and what they do. *Glia* *54*, 716–725.
- Schnitzer, J., Franke, W.W., and Schachner, M. (1981). Immunocytochemical demonstration of vimentin in astrocytes and ependymal cells of developing and adult mouse nervous system. *J. Cell Biol.* *90*, 435–447.
- Seifert, G., Schilling, K., and Steinhauser, C. (2006). Astrocyte dysfunction in neurological disorders: a molecular perspective. *Nat. Rev. Neurosci.* *7*, 194–206.
- Serio, A., Bilican, B., Barmada, S.J., Ando, D.M., Zhao, C., Siller, R., Burr, K., Haghi, G., Story, D., Nishimura, A.L., et al. (2013). Astrocyte pathology and the absence of non-cell autonomy in an induced pluripotent stem cell model of TDP-43 proteinopathy. *Proc. Natl. Acad. Sci. USA* *110*, 4697–4702.
- Shaltouki, A., Peng, J., Liu, Q., Rao, M.S., and Zeng, X. (2013). Efficient generation of astrocytes from human pluripotent stem cells in defined conditions. *Stem Cells* *31*, 941–952.
- Sherwood, C.C., Stimpson, C.D., Raghanti, M.A., Wildman, D.E., Uddin, M., Grossman, L.I., Goodman, M., Redmond, J.C., Bonar, C.J., Erwin, J.M., et al. (2006). Evolution of increased glia-neuron ratios in the human frontal cortex. *Proc. Natl. Acad. Sci. USA* *103*, 13606–13611.
- Skipuletz, T., Hackstette, D., Bauer, K., Gudi, V., Pul, R., Voss, E., Berger, K., Kipp, M., Baumgartner, W., and Stangel, M. (2013). Astrocytes regulate myelin clearance through recruitment of microglia during cuprizone-induced demyelination. *Brain* *136*, 147–167.
- Tong, X., Ao, Y., Faas, G.C., Nwaobi, S.E., Xu, J., Hausteiner, M.D., Anderson, M.A., Mody, I., Olsen, M.L., Sofroniew, M.V., et al. (2014). Astrocyte Kir4.1 ion channel deficits contribute to neuronal dysfunction in Huntington's disease model mice. *Nat. Neurosci.* *17*, 694–703.
- Verheyen, A., Diels, A., Dijkmans, J., Oyelami, T., Meneghello, G., Mertens, L., Versweyveld, S., Borgers, M., Buist, A., Peeters, P., et al. (2015). Using human iPSC-derived neurons to model TAU aggregation. *PLoS One* *10*, e0146127.
- Volterra, A., Liaudet, N., and Savtchouk, I. (2014). Astrocyte Ca(2)(+) signalling: an unexpected complexity. *Nat. Rev. Neurosci.* *15*, 327–335.
- Wyss-Coray, T., Loike, J.D., Brionne, T.C., Lu, E., Anankov, R., Yan, F., Silverstein, S.C., and Husemann, J. (2003). Adult mouse astrocytes degrade amyloid-beta in vitro and in situ. *Nat. Med.* *9*, 453–457.
- Xu, Q., Bernardo, A., Walker, D., Kanegawa, T., Mahley, R.W., and Huang, Y. (2006). Profile and regulation of apolipoprotein E (ApoE) expression in the CNS in mice with targeting of green fluorescent protein gene to the ApoE locus. *J. Neurosci.* *26*, 4985–4994.
- Xu, M., Lee, E.M., Wen, Z., Cheng, Y., Huang, W.K., Qian, X., Tcw, J., Kouznetsova, J., Ogden, S.C., Hammack, C., et al. (2016). Identification of small-molecule inhibitors of Zika virus infection and induced neural cell death via a drug repurposing screen. *Nat. Med.* *22*, 1101–1107.
- Yuan, S.H., Martin, J., Elia, J., Flippin, J., Paramban, R.I., Hefferan, M.P., Vidal, J.G., Mu, Y., Killian, R.L., Israel, M.A., et al. (2011). Cell-surface marker signatures for the isolation of neural stem cells, glia and neurons derived from human pluripotent stem cells. *PLoS One* *6*, e17540.
- Zamanian, J.L., Xu, L., Foo, L.C., Nouri, N., Zhou, L., Giffard, R.G., and Barres, B.A. (2012). Genomic analysis of reactive astrogliosis. *J. Neurosci.* *32*, 6391–6410.
- Zhang, Y., and Barres, B.A. (2010). Astrocyte heterogeneity: an underappreciated topic in neurobiology. *Curr. Opin. Neurobiol.* *20*, 588–594.
- Zhang, Y., Sloan, S.A., Clarke, L.E., Caneda, C., Plaza, C.A., Blumenthal, P.D., Vogel, H., Steinberg, G.K., Edwards, M.S., Li, G., et al. (2016). Purification and characterization of progenitor and mature human astrocytes reveals transcriptional and functional differences with mouse. *Neuron* *89*, 37–53.
- Zhao, B., and Schwartz, J.P. (1998). Involvement of cytokines in normal CNS development and neurological diseases: recent progress and perspectives. *J. Neurosci. Res.* *52*, 7–16.