

Original Article

Crosstalk between MLO-Y4 osteocytes and C2C12 muscle cells is mediated by the Wnt/ β -catenin pathway[†]

Jian Huang, MS, PhD^{1#}, Sandra Romero-Suarez, PhD¹, Nuria Lara, PhD², Chenglin Mo, MS, PhD^{1#}, Simon Kaja, PhD^{3,4}, Leticia Brotto, MD^{1#}, Sarah L. Dallas, PhD², Mark L. Johnson, PhD², Katharina Jähn, PhD^{2##}, Lynda F. Bonewald, PhD^{2,5} and Marco Brotto, BSN, MS, PhD^{1#,†}

¹ Muscle Biology Research Group-MUBIG, School of Nursing & Health Studies, University of Missouri- Kansas City, 2464 Charlotte Street, Kansas City, MO 64108, USA;

² Department of Oral and Craniofacial Sciences, School of Dentistry, University of Missouri-Kansas City, MO, 64108, USA;

³ Department of Ophthalmology, Vision Research Center, School of Medicine, University of Missouri-Kansas City, 2411 Holmes St., Kansas City, MO, 64108, USA;

⁴ Current address: Department of Ophthalmology, Stritch School of Medicine, Loyola University Chicago, 2160 S First Ave., Maywood, IL, 60153, USA;

⁵ Current address: Indiana Center for Musculoskeletal Health, School of Medicine, Indiana University, Barnhill Drive, Indianapolis, In 46202, USA;

Current address: College of Nursing and Health Innovation, University of Texas at Arlington, 411 S. Nedderman Dr, Arlington, TX 76019, USA;

Current address: Emmy Noether Research Group for Bone Material Quality. Department of Osteology and Biomechanics (IOBM), University Medical Center Hamburg-Eppendorf (UKE) Lottestr. 55a, 22529 Hamburg, Germany;

† Correspondence to Dr. Marco Brotto, College of Nursing and Health Innovation, University of Texas at Arlington, 411 S. Nedderman Dr, Arlington, TX 76019, telephone: 817-272-5018, Email: Marco.brotto@uta.edu

There is one Supplemental Figure has been included with the submission.

† This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/jbm4.10015]

Additional Supporting Information may be found in the online version of this article.

Initial Date Submitted March 8, 2017; Date Revision Submitted July 13, 2017; Date Final Disposition Set July 29, 2017

JBMR Plus

**This article is protected by copyright. All rights reserved
DOI 10.1002/jbm4.10015**

Abstract

We examined the effects of osteocyte secreted factors on myogenesis and muscle function. MLO-Y4 osteocyte-like cell conditioned media (CM) (10%) increased *ex vivo* soleus muscle contractile force by ~25%. MLO-Y4 and primary osteocyte CM (1-10%) stimulated myogenic differentiation of C2C12 myoblasts, but 10% osteoblast CMs did not enhance C2C12 cell differentiation. Since WNT3a and WNT1 are secreted by osteocytes, and the expression level of *Wnt3a* is increased in MLO-Y4 cells by fluid flow shear stress, both were compared, showing WNT3a more potent than WNT1 in inducing myogenesis. Treatment of C2C12 myoblasts with WNT3a at concentrations as low as 0.5ng/mL mirrored the effects of both primary osteocyte and MLO-Y4 CM by inducing nuclear translocation of β -catenin with myogenic differentiation, suggesting that Wnts might be potential factors secreted by osteocytes that signal to muscle cells. Knocking down *Wnt3a* in MLO-Y4 osteocytes inhibited the effect of CM on C2C12 myogenic differentiation. Sclerostin (100ng/mL) inhibited both the effects of MLO-Y4 CM and WNT3a on C2C12 cell differentiation. RT-PCR array results supported the activation of the Wnt/ β -catenin pathway by MLO-Y4 CM and WNT3a. These results were confirmed by qPCR showing up-regulation of myogenic markers and two Wnt/ β -catenin downstream genes, *Numb* and *Flh1*. We postulated that MLO-Y4 CM/WNT3a could modulate intracellular calcium homeostasis as the trigger mechanism for the enhanced myogenesis and contractile force. MLO-Y4 CM and WNT3a increased caffeine-induced Ca^{2+} release from the sarcoplasmic reticulum (SR) of C2C12 myotubes and the expression of genes directly associated with intracellular Ca^{2+} signaling and homeostasis. Together, these data show that *in vitro* and *ex vivo*, osteocytes can stimulate myogenesis and enhance muscle contractile function and suggest that Wnts could be mediators of bone to muscle signaling, likely via modulation of intracellular Ca^{2+} signaling and the Wnt/ β -Catenin pathway. This article is protected by copyright. All rights reserved

Key words: Wnt/ β -Catenin/LRPs; bone-muscle interactions; molecular pathways; osteocytes; general muscle physiology.

Introduction

Muscle and bone are connected both anatomically and functionally and interact as a functional unit ⁽¹⁾. Musculoskeletal diseases are the most common cause of chronic disabilities worldwide. Osteoporosis/osteopenia (loss of bone tissue and function) and sarcopenia (progressive muscle loss with a greater and disproportional loss of muscle force/strength) are inevitable consequences of the aging process and prevalent in the frail elderly and have become a growing public health concern ⁽²⁾.

The emerging field of bone-muscle crosstalk is helping to clarify the role of bones and muscles as endocrine organs because they produce and secrete “hormone-like factors” that can mutually influence each other and other tissues ^(3,4). Recently, evidence has emerged that osteocytes secrete factors that have profound effects on C2C12 muscle cells in culture, and vice versa ^(3,5), suggesting that muscle and bone communicate at a molecular level via biochemical factors that are reciprocally important for optimal function ⁽⁶⁾.

Bones secrete a host of factors that have clear paracrine and endocrine effects, such as FGF23, Sclerostin, Osteocalcin, WNTs, Prostaglandin E₂ (PGE₂), etc. (reviewed in ⁽⁷⁾). Sclerostin is one of these factors, which has gained much attention because of recent clinical applications and trials ⁽⁸⁾. Sclerostin is one of the osteocyte-specific proteins, and is product of the *SOST* gene ⁽⁹⁾, which is expressed by mature osteocytes ^(10,11). Sclerostin is a negative regulator of Wnt/ β -catenin signaling pathway by binding to the Wnt coreceptors, low-density lipoprotein receptor-related proteins 5 and 6 (LRP5 and LRP6) ⁽¹²⁾. In the presence of Sclerostin, Wnt-receptor interaction is inhibited, and β -catenin is phosphorylated by glycogen synthase kinase 3 and targeted for ubiquitination and degradation via the proteasome pathway ⁽¹³⁾. Studies using loss of function and gain of function mouse models of *Sost* have demonstrated increased

and decreased bone mass, respectively ⁽¹⁰⁾. Our earlier studies demonstrated that many of the effects of bone cell conditioned medium in triggering acceleration of myogenesis could be partially mimicked by low nanomolar range concentrations of PGE₂ ⁽³⁾.

The Wnt / β -catenin signaling pathway is important for cell and tissue homeostasis, since secreted WNTs act through specific receptors to control and modulate cell proliferation, differentiation, apoptosis, survival, migration, and polarity (reviewed in ⁽¹⁴⁾). They play critical roles during embryonic development (including muscle and skeletal patterning) as well as in postnatal health and diseases, including cancer and degenerative disorders. The Wnt/ β -catenin signal pathway has been shown to be an important component of bone mass accrual, regulation and maintenance ⁽¹⁵⁾, and accumulating data show that the Wnt/ β -catenin signal pathway is strongly implicated in the skeletal muscle development, growth and regeneration ⁽¹⁶⁾.

To determine whether osteocytes can potentially regulate muscle function, we tested the effects of MLO-Y4 conditioned medium (MLO-Y4 CM) on muscle contractility in soleus (SOL) muscles using a murine *ex vivo* muscle contractility assay and found out that MLO-Y4 CM increased the contractile force of SOL muscles. To gain new insight into the mechanisms of bone to muscle signaling we have used MLO-Y4 osteocyte-like cells, osteoblast cells, primary osteocytes and C2C12 myoblasts as *in vitro* models. We report that MLO-Y4 cells and primary osteocytes secrete factors that potently stimulate myogenesis, accompanied by enhanced β -catenin translocation, suggesting that the effect may be mediated via Wnt/ β -catenin signaling. However, 10% osteoblast CMs did not enhance C2C12 cell differentiation. We therefore investigated the expression of Wnts in osteocytes and showed that WNT3a, which is expressed in osteocytes, mirrored the effects of osteocyte conditioned medium on myogenesis. Knocking down *Wnt3a* in MLO-Y4 osteocytes inhibited the effect of CM on C2C12 myogenic

differentiation. Sclerostin inhibited the effects of CM or WNT3a on C2C12 cell differentiation.

To determine potential mechanisms of contractile force enhancement, we examined the effects of osteocyte conditioned medium on calcium release from the sarcoplasmic reticulum (SR). Our *in vitro* and *ex vivo* data show that osteocytes secrete soluble factors that enhance myogenic differentiation, enhance both contractile force and calcium release from the SR, and provide evidence that WNT3a is a potential factor from osteocytes with the intrinsic potential to modulate these effects of bone-muscle crosstalk.

Materials and methods

Materials

DMEM high glucose media, α MEM media, penicillin-streptomycin (P/S) 10,000 U/mL each and trypsin-EDTA 1 \times solution were obtained from Mediatech Inc. (Manassas, VA, USA); calf serum (CS), fetal bovine serum (FBS), horse serum (HS) and caffeine were obtained from Thermo Fischer Scientific Inc. (Waltham, MA, USA); bovine serum albumin (BSA), diamidino-2-phenylindole (DAPI) were from Sigma-Aldrich (St Louis, MO, USA); rat tail collagen type I was purchased from BD Biosciences (San Jose, CA, USA); 16% paraformaldehyde was from Alfa Aesar (Ward Hill, MA, USA); Recombinant Mouse Wnt-3a and Recombinant Mouse SOST/Sclerostin protein was purchased from R&D Systems Inc. (Minneapolis, MN, USA); WNT1 Recombinant Human Protein was obtained from Life Technologies. (Grand Island, NY, USA); Cy-3 donkey anti-mouse were purchased from Invitrogen. (Carlsbad, CA, USA); This antibody has been previously validated ⁽⁵⁾. Mouse anti-active β -catenin antibody was from Millipore. (Billerica, MA, USA); This antibody has been previously validated ⁽⁵⁾. Lipofectamine[®] RNAiMAX Transfection Reagent was from ThermoFisher Scientific. (Waltham, MA, USA); *Wnt3a* siRNA (antisense strand: 5'-

GCAUCCGCUCUGACACUAAUACTC-3'), negative control siRNA and TYE 563 DS Transfection Control was from Integrated DNA Technologies. (Coralville, Iowa, USA); Tri reagent was obtained from Molecular Research Center, Inc. (Cincinnati, OH, USA); High capacity cDNA reverse transcription kit was from Applied Biosystems. (Foster city, CA, USA); RT² Real-Time™ SYBR green/Rox PCR master mix was from SABiosciences. (Valencia, CA, USA); Anti-human Myosin Heavy Chain-CFS, Carboxyfluorescein (CFS)-conjugated mouse monoclonal anti-human Myosin Heavy Chain antibody was from R&D Systems Inc. (Minneapolis, MN, USA); This antibody has been previously validated ⁽⁶⁾. Fura-2/AM was obtained from Life Technologies (Grand Island, NY, USA). C2C12 cells were obtained from American Type Culture Collection (ATCC) (Manassas, VA, USA). Gentamycin and 2-APB; Wingless Type MMTV Integration Site Family, Member 3A (WNT3A) ELISA Kit was purchased from MyBioSource. (San Diego, CA, USA).

Methods

No *in vivo* experiments were conducted in this study.

Animals

Five month old male C57BL/6 mice from Jackson Laboratory were used for isolation of bone cells and isolation of intact muscles for *ex vivo* contractility studies, following humane euthanasia cervical dislocation. All animal procedures were performed according to an approved IACUC protocol at the University of Missouri-Kansas City (UMKC), and conformed to relevant federal guidelines. The UMKC animal facility is operated as a specific pathogen-free, AAALAC approved facility. Animal care and husbandry at UMKC meet the requirements in the Guide for the Care and use of Laboratory Animals (8th edition), National Research Council. Animals were group housed and maintained on a 12-hour light/dark cycle with ad libitum food and water at a

constant temperature of 72°F and humidity of 45-55%. Daily health check inspections were performed by qualified veterinary staff and/or animal care technicians.

***Ex vivo* muscle contractility**

Dissection and setup: Mice were humanely sacrificed and soleus muscles were carefully removed for contractility analysis, as previously reported ⁽¹⁷⁾ and following the Protocol of the Treat-NMD Network (DMD_M.1.2.002). Dissected soleus muscles were immediately placed in a dish containing a HEPES Ringer solution (143 mM NaCl; 5 mM KCl; 1.8 mM MgCl₂; 10 mM HEPES; 2.5 mM CaCl₂; pH 7.40) with 10 mM glucose. This solution was continuously aerated with 100% O₂. Muscles were mounted vertically between two stimulating platinum electrodes and immersed in a 25 mL bathing chamber containing the Ringer solution. Via the tendons, the muscles were suspended from adjustable isometric force transducers above the chambers and secured to the base of the tissue support within the chambers. The analog output of the force transducer was digitized, stored and analyzed with PowerLab[®] Software (ADInstruments Inc., Colorado Springs, CO, USA). For each muscle the stimulatory voltage was provided by an S8800 dual pulse digital stimulator (Grass Products, West Warwick, RI, USA) (pulse duration, 1 ms; train duration, 500 ms). Optimal muscle length (L_0) was determined by tetanic stimulations of 100 Hz at an interval of 1 minute. *Equilibration:* The intact muscles were allowed a 30 minutes equilibration period during which time they were stimulated with pairs of alternating high (100 Hz) and low (20 Hz) frequency pulse-trains administered with a periodicity of 1 minute. Utilization of the proposed paradigm of stimulation helps with the study of the relative contributions of the contractile proteins (100 Hz) and the SR (20 Hz) to contractile function ⁽¹⁷⁾. After equilibration and stabilization of submaximal and maximal forces, blank medium or MLO-Y4 CM was added to the control and test muscle, respectively and force was followed for 30

minutes after the addition of the media. *Force normalization*: Muscle force was reported as relative force (mN) and force normalized to muscle physiological cross sectional area (N/cm²) as previously reported by our group⁽¹⁷⁻¹⁹⁾. Five animals and 10 muscles were used for final analyses.

Cell cultures and generation of conditioned medium (CM) from primary osteocytes (PO CM), MLO-Y4 cells (MLO-Y4 CM), 2T3 cells (2T3 CM) and MC3T3-E1 (MC3T3-E1 CM)

Cell culture: C2C12 myoblasts were cultured following our own previously published protocols^(3, 5, 6). Briefly, cells were grown at 37°C in a controlled humidified 5% CO₂ atmosphere in growth medium (GM), DMEM/high glucose +10% FBS (100U/mL P/S), and maintained at 40 - 70% cell density. Under these conditions, myoblasts proliferate, but do not differentiate into myotubes. During myoblast proliferation, medium was changed every 48 hours. To induce differentiation of myoblasts into myotubes, when cells reached 75% confluence, the culture media were switched from GM to differentiation medium (DM), DMEM/high glucose + 2.5% horse serum (HS) (100U/mL P/S), and DM was changed every 48 hours. Fully differentiated, functional, contracting myotubes were formed within 5-7days.

MLO-Y4, a murine long bone-derived osteocyte-like cell line (referred to hereafter as MLO-Y4 cells), were cultured as previously described⁽²⁰⁾. Briefly, cells were seeded onto type I collagen-coated plates and cultured in α MEM + 2.5% FBS + 2.5% CS (100U/mL P/S). Cells were maintained at ~60% confluence throughout the culture period.

2T3 osteoblast-like cells (referred to hereafter as 2T3 cells) were cultured as previously described⁽²¹⁾ with a few modifications. Briefly, cells were cultured in α MEM + 10% FBS (100U/mL P/S). Cells were maintained at ~60% confluence throughout the culture period.

MC3T3-E1 preosteoblast cells (referred to hereafter as MC3T3-E1 cells) were cultured as previously described⁽²²⁾. Briefly, cells were cultured in α MEM + 10% FBS (100U/mL P/S).

Cells were maintained at ~60% confluence throughout the culture period.

Generation of CMs from primary osteocytes (PO CM), MLO-Y4 cells (MLO-Y4 CM), 2T3 cells (2T3 CM) and MC3T3-E1 cells (MC3T3-E1 CM): The method of isolation of primary osteocytes was described by Stern et al.^(23, 24), with minor modifications. Briefly, long bones were dissected and muscles were removed taking care not to remove the periosteum. Epiphyses were cut and bone marrow was then flushed out. The bones were cut into small pieces (1-2 mm²) and incubated for 25 minutes in collagenase solution (300 units/mL) with constant shaking in an incubator at 37°C and 5% CO₂. Solution was then removed and centrifuged. Cells were then re-suspended in 10% FBS containing α MEM and plated in collagen-coated plates (fraction 1-fibroblast-enriched). Bone chips were then subsequently incubated with fresh collagenase solution to obtain fractions 2 to 4 (osteoblast-enriched). Then bone chips were digested in alternating EDTA (5 mM) and collagenase solutions to obtain fractions 5 to 9 (osteocyte-enriched). The remaining bone pieces were transferred into collagen-coated dishes and cells migrated from the bone pieces after 3–5 days of culture (fraction 10-bone particles). All of the cells obtained from fractions 5 to 9 that displayed an osteocyte-like morphology stained positive for the osteocyte marker E11/GP38. The osteocyte phenotype was further confirmed by a lack of staining for alkaline phosphatase and the absence of collagen1a1 expression. The osteocytes also expressed additional osteocyte-specific genes such as *Sost* and *Mepe*. To generate PO CM, cells of fractions 5 to 9 were cultured in 60 cm² collagen-coated plates with α MEM + 5 % FBS + 5 % CS (1,000 U/mL P/S) culture media, 3600 cells/cm² and allowed to attach and grow for 48 hours.

The medium was collected and centrifuged at $500 \times g$ for 10 minutes. Conditioned medium (CM) was aliquoted and flash frozen in liquid nitrogen then stored at -80°C .

To prepare MLO-Y4 CM, 2T3 CM and MC3T3-E1 CM, MLO-Y4, 2T3 and MC3T3-E1 cells were plated in 10cm dishes, $5000 \text{ cells}/\text{cm}^2$ and allowed to attach and grow for 48 hours. The medium was collected and centrifuged at $500 \times g$ for 10 minutes. Conditioned medium (CM) was aliquoted and flash frozen in liquid nitrogen then stored at -80°C .

Fluid flow shear stress (FFSS) treatment of MLO-Y4 and 2T3 cells

At day 1, MLO-Y4 and 2T3 cells were seeded onto glass slides coated with type I collagen at a density of $5,600 \text{ cells}/\text{cm}^2$ in alpha-MEM supplemented with either 2.5% calf serum (CS), 2.5% fetal bovine serum (FBS) and 1% P/S (for MLO-Y4 cells) or 10% FBS and 1% P/S (for 2T3 cells) for 24 hours. At day 2, shear stress was applied to the slides at $2 \text{ dynes}/\text{cm}^2$ for 2 hours in alpha-MEM supplemented with 0.1% CS, 0.1% FBS and 1% P/S. For fluid flow shear stress, 350 mL media was used to run the system (6 slides per run). For static controls, three slides were placed in a 150 mm round tissue culture dish with 175 mL of alpha-MEM media (1% P/S) containing 0.1% CS and 0.1% FBS. Media was collected after 2 hours of shear stress. Three slides of seeded cells were used for RNA isolation at the end of the 2 hours of shear stress (or static culture). The other 3 slides were incubated in regular media (alpha-MEM supplemented with 2.5% CS, 2.5% FBS and 1% P/S) for 24 hours. RNA was extracted from both static and shear stressed slides, 2 hours and 24 hours after FFSS for RT-qPCR detection of gene expression.

ELISA for WNT3a in PO CM and MLO-Y4 CM

WNT3a in PO CM and MLO-Y4 CM was detected by Wingless Type MMTV Integration Site Family, Member 3A (WNT3A) ELISA Kit according to the manufacturer's protocol. Serum-free

CM were collected from the cells that were cultured for twenty four hours, and 100 μ l CM was added into the ELISA plate wells coated with a biotin-conjugated WNT3a antibody. Next, Avidin conjugated to Horseradish peroxidase is added and incubated for 1 hour. After TMB substrate solution is added, the color change is measured spectrophotometrically at 450 nm. The concentration of WNT3a is then determined by comparing the O.D. of the samples to the standard curve. Three independent experiments with 2 replicates were performed for these experiments.

Immunolabeling of active β -catenin for nuclear translocation

These studies followed our previous protocols ⁽²⁵⁾ with minor modifications for C2C12 cells.

Briefly, C2C12 cells were seeded onto positively-charged glass slides and cultured overnight.

CMs or blank media was added to produce 10% final concentration. Cells were incubated for 30 minutes, 60 minutes and 90 minutes respectively, rinsed two times in cold PBS, fixed in 2% paraformaldehyde containing 0.2% Triton X-100 for 10 minutes and then washed three times in PBS at ambient temperature for 10 minutes each. Slides were blocked with 2.5 % BSA and 1% normal donkey serum in PBS overnight at 4 °C. Primary antibody against the active form of β -catenin was used at 1:100 dilution in blocking buffer for 4 hours at ambient temperature. Slides were washed three times with PBS and incubated for 1 hour with Cy3 -conjugated donkey anti-mouse antibody. The quantification of the intensity of β -catenin immunofluorescence was performed as described previously using a Leica TCS SP5 II confocal microscopy system on an inverted microscope platform ⁽²⁶⁾. 16 bit images were obtained and were pseudo colored in green and analyzed using ImageJ Software (NIH). The nuclear and cytoplasmic histograms from each image were used to quantify staining intensity. Three independent experiments with 5-10 cells monitored on each individual experiments.

C2C12 Cell Morphometry and Immunostaining

Cell Morphology: Phase contrast images were taken with a LEICA DMI-4000B inverted microscope equipped with a 14-BIT CoolSNAP CCD camera (Photometrics), using the LEICA LAS imaging software for calibration (Leica microsystems).

Immunostaining: Experiments were performed following our published protocols ^(5, 6, 27).

Briefly, C2C12 cells were fixed with neutral buffered formalin and permeabilized with 0.1% Triton X-100. Myosin heavy chain (MHC) was detected with Anti-Human Myosin Heavy Chain-CFS (1:50) at room temperature for 30 minutes and counter stained with DAPI. Images were taken using a 20X LEICA FLUO objective with the LEICA system described above. *Fusion index:* To quantify myogenic differentiation of C2C12 cells after treatments, the Fusion Index (FI) was calculated, where FI is defined as: (nuclei within myosin heavy chain-expressing myotubes/total number of myogenic nuclei) $\times 100$ ⁽²⁸⁾. We conducted 3-5 independent experiments, 2 replicates, and 3-5 areas per replicate were randomly selected for the measurements.

Treatment of C2C12 cells with PO CM, MLO-Y4 CM, 2T3 CM, MC3T3-E1 CM and WNT3a

C2C12 cells were plated in 6-well plates, 10×10^4 /well, and allowed to attach and grow overnight. Medium of C2C12 myoblasts were changed from GM to DM with CMs and WNT3a, respectively, 48 hours later, changed medium with fresh DM. CMs and WNT3a were removed from the medium. At day 3 of differentiation, C2C12 cells were analyzed according to “C2C12 Cell Morphometry and Immunostaining” described above.

Treatment of MLO-Y4 cells with *Wnt3a* siRNA and generation of CM from *Wnt3a* siRNA treated MLO-Y4 cells (MLO-Y4 siRNA-treated CM)

MLO-Y4 cells were plated in 6-well plates, 5×10^4 cells/well, and allowed to attach and grow overnight. Cells were transfected with Lipofectamine® RNAiMAX Transfection Reagent following the manufacturer's instructions. First, TYE 563 DS Transfection Control which is tagged with a red fluorescent probe was used to detect the transfection efficiency. To knock down *Wnt3a*, MLO-Y4 cells were treated with 10 nM *Wnt3a* siRNA+ 9 μ l Lipofectamine® RNAiMAX reagent (MLO-Y4-siRNA-treated cells), while negative control cells were treated with 10 nM negative control siRNA + 9 μ l Lipofectamine® RNAiMAX reagent. We further employed a vehicle control by treating the cells with only 9 μ l Lipofectamine® RNAiMAX reagent without siRNA. After 24 hours of transfection, total RNA was isolated from the cells with Tri reagent and *Wnt3a* expression was detected by RT-qPCR. To prepare the conditioned medium (CM) of MLO-Y4 cells transfected with *Wnt3a* siRNA, 24 hours after siRNA transfection, medium was changed with fresh medium and the cells were cultured for 48 hours, then CM was collected and centrifuged at $500 \times g$ for 10 minutes. CM was aliquoted and flash frozen in liquid nitrogen then stored at -80°C . The CMs of negative control and Lipofectamine® RNAiMAX reagent-only vehicle treated cells (vehicle control) were also collected. Three independent experiments were performed.

Treatment of C2C12 cells with MLO-Y4 siRNA-treated CM

C2C12 cells were plated in 6-well plates, 10×10^4 /well, and allowed to attach and grow overnight. Medium of C2C12 myoblasts were changed from GM to DM with 10% CMs, respectively, 48 hours later, changed medium with fresh DM. CMs were removed from the medium. At day 3 of

differentiation, C2C12 cells were analyzed according to “C2C12 Cell Morphometry and Immunostaining” described above.

Treatment of C2C12 cells with Sclerostin

C2C12 cells were plated in 6-well plates, 10×10^4 /well, and allowed to attach and grow overnight.

a) C2C12 myoblasts in DM without Sclerostin treatment as a negative control. b) C2C12 myoblasts were treated with 100ng/mL Sclerostin in DM for 6 hours. c) C2C12 myoblasts were treated in DM with 10% MLO-Y4 CM for 48 hours as one of our positive controls. d) C2C12 myoblasts were treated in DM with 10ng/mL WNT3a for 48 hours as our second positive control. e) C2C12 myoblasts were treated with 100ng/mL Sclerostin in DM for 6 hours, followed by 10% MLO-Y4 CM for 48 hours; f) C2C12 myoblasts were treated with 100ng/mL Sclerostin in DM for 6 hours, followed by 10ng/mL WNT3a for 48 hours; g) C2C12 myoblasts were treated with 100ng/mL Sclerostin in DM for 6 hours, followed by the combination of 10% MLO-Y4 CM + 10ng/mL WNT3a for 48 hours. Under any experimental where Sclerostin, CM, WNT3a was used, they were maintained during treatments for 48 hours. After 48 hours, medium was changed with fresh DM. Sclerostin, CM, and WNT3a were removed from the media. At day 3 of differentiation, C2C12 cells were analyzed according to “C2C12 Cell Morphometry and Immunostaining” described above.

RNA isolation and real time quantitative PCR (RT-qPCR)

Total RNA was isolated from the cells with Tri reagent according to the manufacturer's protocol and cDNA was synthesized using a high capacity cDNA reverse transcription kit. RT-qPCR was performed using RT² Real-TimeTM SYBR green/Rox PCR master mix. RT-qPCR primers used in this study are summarized in **Table 1**. RT-qPCR was run in a 25 μ l reaction volume on 96-well

plates using a StepOnePlus instrument (Applied Biosystems, Foster city, CA, USA). Data were analyzed using RT² Profiler™ PCR Array Data Analysis (SABiosciences); C_t values were normalized to *Gadph* (Glyceraldehyde-3-phosphate dehydrogenase) as a reference gene. Gene expression was determined as fold up or down regulation of the gene of interest compared to the controls. All reactions were done in duplicate and all experiments were repeated at least three times.

RT-PCR Gene Arrays

The Mouse Signal Transduction PathwayFinder PCR Array from SABiosciences was used to simultaneously detect gene expression changes of 10 signaling pathways (see details in **Table 2**). cDNA was synthesized using the RT² First Strand Kit (This kit contains genomic DNA elimination process) and the PCR Array was run according to the manufacturer's protocol including a threshold of 0.25 and validation of each gene tested by the identification of single peaks in melting curves. As above, data were analyzed using the RT² Profiler™ PCR Array Data Analysis Software; C_t values were normalized to six built-in reference housekeeping genes, genomic DNA control, reverse transcription control, and positive PCR control. We used this analytical software (<http://www.qiagen.com/us/shop/genes-and-pathways/data-analysis-center-overview-page/>) to set the statistical significance of up/down regulation of all tested genes at 2-fold difference. Three independent experiments were performed.

Intracellular Ca²⁺ measurements

A Photon Technology International (PTI) imaging system was used to measure intracellular calcium transients. The intracellular calcium transients produced by the stimulation of calcium release from sarcoplasmic reticulum (SR) with 20 mM caffeine were measured. Each myotube imaged was loaded with Fura-2, a ratiometric calcium dye and imaged in real time with the 14-

BIT CoolSNAP CCD camera). All calcium imaging was analyzed with PTI EasyRatioPro fluorescence imaging software. We conducted 3-4 independent experiments, resulting in 12-18 cells analyzed per group ^{(6), (27)}.

Statistical analysis

Data are presented as mean \pm SD. Comparisons between more than two groups were made using one-way ANOVA followed by Tukey's post hoc test for multiple comparisons. For comparisons of differences between two groups the *t*-test was used. A value of $p < 0.05$ was considered as being significantly different. *Statistical Analyses of Muscle Contractility*: Data points that were three standard deviations away from the mean were considered outliers and removed from the data set. Kolmogorov-Smirnov test was conducted to test for the normality assumption of the outcomes variables. All variables passed the normality test under the significance level of 0.05.

A Mixed model approach was conducted to test for the treatment effect while accounting for the repeated measures on the same individual. Pairwise contrasts were estimated and tested for significance between different treatments. Due to the multiple comparisons of the contrasts, a more strict alpha level of 0.025 was used to establish the statistical significance.

Results

MLO-Y4 CM increased the contractile force of soleus muscle

SOL muscle was treated with blank media as control and with 10% MLO-Y4 CM. The muscles were stimulated with two alternating frequencies of 20 Hz and 100 Hz. MLO-Y4 CM increased the contractile force of SOL muscles at the 20 Hz frequency by ~25% (**Fig. 1A**) compared to blank media.

MLO-Y4 CM, Primary Osteocyte (PO) CM and WNT3a enhance myogenic differentiation of C2C12 cells, while Osteoblast CMs do not enhance myogenesis

C2C12 cells were treated with MLO-Y4 CM, PO CM, and WNT3a, respectively. At day 3 of differentiation, control and treatment groups cells were stained with Anti-human Myosin Heavy Chain-CFS antibody which only stains maturing myocytes/myotubes not myoblasts and DAPI that stains only the nuclei (**Fig. 1B**). We also detected the specific fusion index (FI) of each experimental group. These data are summarized in **Fig. 1C**, showing that FI is significantly increased in the treatment groups compared with control. When C2C12 cells were treated with the 10% CM deriving from two different osteoblast cell lines, 2T3 (an established osteoblast cell line) and MC3T3-E1 (a preosteoblast cell line), respectively, there is no significant effects on the myogenic differentiation (**Fig. 2A-B**).

***Wnt3a* mRNA is expressed in MLO-Y4 cell and 2T3 osteoblast cells and the expression increased after FFSS**

Wnt1, *Wnt3a*, *Wnt4*, *Wnt5a* and *Wnt7a* are involved in myogenesis (reviewed in (16)). To detect the gene expression of *Wnt1*, *Wnt3a*, *Wnt4*, *Wnt5a* and *Wnt7a*, we performed RT-qPCR in MLO-Y4 and 2T3 cells. Our results show that *Wnt3a* is expressed in bone cells and is modulated by FFSS. These results are summarized and explained in detail in the legend of **Fig. 3A-F**.

ELISA for WNT3a in PO CM and MLO-Y4 CM

The only commercially available ELISA kit was used to detect WNT3a in MLO-Y4 CM and PO CM and the factor was detected in the 100-200 pg/mL range. While this level might be considered low, we do see effects *in vitro* with WNT3a concentrations as low as 500 pg/mL. Furthermore, there are many technical reasons that might explain these results, including the fact

that it is technically challenging to isolate and purify WNT3A

(<http://web.stanford.edu/group/nusselab/cgi-bin/wnt/purification>).

Physiological concentrations of WNT3a and low concentrations of MLO-Y4 CM stimulate myogenesis of C2C12 cells

Studies were performed with concentrations of WNT3a ranging from 0.1 to 10ng/mL as well as 1% MLO-Y4 CM. A concentration as low as 0.5ng/mL to 10ng/mL WNT3a consistently enhanced the C2C12 cell differentiation (**Fig. 4A and C**). In agreement with lower concentrations of WNT3a accelerating myogenesis, we also found that the myogenic differentiation of C2C12 cells was enhanced with as low as 1% MLO-Y4 CM. After 3 days of differentiation, a significant increase of the FI of MLO-Y4 CM-treated cells was found as compared with the control cells (**Fig. 4B and D**).

MLO-Y4 CM, PO CM, and WNT3a induced translocation of β -Catenin to the nucleus of C2C12 cells

To explore the possible roles of Wnt/ β -catenin pathway in C2C12 cells, confocal microscopy was used to study the time course of the subcellular localization of β -catenin after C2C12 cells were treated with 10% MLO-Y4 CM, which demonstrated that peak translocation occurred at 60 minutes (min) (**Fig. 5A-B**). Next, we tested the effects of C2C12 growth medium (GM), C2C12 differentiation medium (DM), MLO-Y4 CM, PO CM, WNT3a and Lithium chloride as a positive control. The results demonstrated that β -catenin translocation to the nucleus was enhanced significantly by the different treatments when compared to GM. Furthermore, MLO-Y4 CM, PO CM, WNT3a and Lithium chloride treatments were more potent than DM in inducing nuclear translocation of β -catenin (**Fig. 5C-D**).

Treatment of C2C12 cells with MLO-Y4 siRNA-treated CM

To explore the roles of WNT3a from MLO-Y4 CM on C2C12 cells, MLO-Y4 osteocytes were transfected with *Wnt3a* siRNA. To detect siRNA transfection efficiency, MLO-Y4 osteocytes were transfected with 10 nM TYE 563 DS Transfection Control siRNA. Twenty four hours after transfection, images were taken, and the percentage of fluorescent positive cells was determined. The result revealed that the siRNA transfection efficiency of MLO-Y4 osteocytes was $80.02 \pm 7.18\%$ (**Fig. 6A**). RT-qPCR was conducted to examine *Wnt3a* expression after knock down by *Wnt3a* siRNA. The results indicated that the relative *Wnt3a* mRNA expression in the siRNA-treated cells was $20.47 \pm 4.72\%$ of the negative control cells when normalized to expression of *Gapdh*, which was not altered by the negative control, vehicle control or siRNA treatments (**Fig. 6A**), therefore ~80% of *Wnt3a* expression was knocked down. MLO-Y4 cells were treated with *Wnt3a* siRNA, and the CM (MLO-Y4 siRNA-treated CM) was collected. C2C12 cells were treated with MLO-Y4 siRNA-treated CM. At day 3 of differentiation, we measured the specific fusion index (FI) of each experimental group. These data are summarized in **Fig. 6B-C**, showing that in the cells treated with MLO-Y4 CM or the MLO-Y4 CM siRNA negative control, FI significantly increases, while in the cells treated with the MLO-Y4 siRNA-treated CM, FI did not increase and it is very similar to the negative control. (There was no difference between negative control and vehicle control).

Treatment of C2C12 cells with Sclerostin inhibits the effects of MLO-Y4 CM and WNT3a on myogenic differentiation.

Sclerostin is a Wnt signaling inhibitor⁽¹²⁾. To study the effect of Sclerostin on C2C12 myoblast differentiation stimulation by MLO-Y4 CM and WNT3a, C2C12 myoblasts were pre-treated with 100ng/mL Sclerostin for 6 hours prior to the addition of MLO-Y4 CM and WNT3a. At day 3 of differentiation, we measured the specific fusion index (FI) of each experimental group. FI

data of each group is summarized in **Fig. 7A-B**, showing that compared to negative control (a), FI of treatment group (b): C2C12 myoblasts were treated with 100ng/mL Sclerostin in DM for 6 hours; treatment group (e): C2C12 myoblasts were treated with 100ng/mL Sclerostin in DM for 6 hours, followed by 10% MLO-Y4 CM for 48 hours; treatment group (f): C2C12 myoblasts were treated with 100ng/mL Sclerostin in DM for 6 hours, followed by 10ng/mL WNT3a for 48 hours was not significantly changed, and significantly reduced when compared to the positive controls (c and d), 10% MLO-Y4 CM and 10ng/mL WNT3a, respectively. The FI of treatment group (g): C2C12 myoblasts were treated with 100ng/mL Sclerostin in DM for 6 hours, followed by the combination of 10% MLO-Y4 CM + 10ng/mL WNT3a for 48 hours; increased significantly compared with negative control (DM), suggesting that only the MLO-Y4 CM + WNT3a together can overcome the inhibitory effect of Sclerostin.

Modulation of Gene Expression by MLO-Y4 CM and WNT3a

We employed the Mouse Signal Transduction PathwayFinder PCR Array to detect differences in expression of signaling molecules between control versus MLO-Y4 CM (10%) and WNT3a (10ng/mL) treated C2C12 cells at 24 hours after treatment. A major advantage of this technology is that monitors 10 different pathways simultaneously and is validated at the protein level. We found a clear common shared gene regulation change in between 10% MLO-Y4 CM (**Fig. 8A**) and 10ng/mL WNT3a (**Fig. 8B**), which was the upregulation of the Wnt pathway detected by the overexpression of *Wnt1* in the cells treated with CM and *Wnt5a* and *Axin2* in the cells treated with WNT3a.

We also employed RT-qPCR to detect the specific expression of myogenic differentiation markers (*Mhc*, *MyoD* and *MyoG*) and two Wnt/ β -catenin downstream genes directly associated with C2C12 myogenesis (*Fhl1*⁽²⁹⁾, *Numb*⁽³⁰⁾). At day 3 of myotube differentiation, we found that

both 1% and 10% MLO-Y4 CM, as well as 0.5ng and 10ng/mL WNT3a (except *Numb* in 1% MLO-Y4 CM) significantly increased the expression of these five specific genes (**Fig. 8C-D**).

MLO-Y4 CM and WNT3a enhance caffeine induced SR calcium release

Calcium homeostasis is very important in myoblast differentiation⁽³¹⁾, and is a major surrogate of skeletal muscle function^{(27), (31)}. Avila et al⁽³²⁾ have demonstrated that WNT3a modulates intracellular Ca²⁺ in hippocampal neurons. Furthermore, since our *ex vivo* muscle contractility experiments showed that the force produced by the lower frequency of stimulation was increased, to determine if SR calcium release was influenced by either MLO-Y4 CM or WNT3a, we employed an effective approach by testing the direct effects of caffeine's ability to release calcium from the SR in Fura-2 loaded myotubes. In day 5 of myotube differentiation, caffeine responses were significantly larger in myotubes treated with 10% MLO-Y4 CM and 10ng/mL WNT3a as indicated by multiple and oscillatory peaks and a prolongation of the relaxation phase of the calcium transient response induced by caffeine (**Fig. 9A**).

MLO-Y4 CM and WNT3a induced common gene overexpression profile of

Ca²⁺ signaling/hypertrophy, Ca²⁺ homeostasis, mitochondrial biogenesis and oxidative stress genes

Expression of important genes related to Ca²⁺ signaling/hypertrophy, Ca²⁺ homeostasis, mitochondrial biogenesis and oxidative stress in C2C12 cells at 60 hours of differentiation after treatments was detected by a focused custom-built RT-PCR gene array using the same technology that the PathwayFinder was used. In agreement with both the enhanced contractile force and SR calcium release by MLO-Y4 CM and WNT3a, the expression of these key regulatory pathways were significantly upregulated (**Fig. 9B, and Table 3**).

Discussion

Recent studies have shown that bone (osteoblasts and osteocytes) acts as an endocrine organ by the production and secretion of at least two circulating factors, FGF23 and osteocalcin⁽⁷⁾. Osteocytes also secreted large amounts of PGE₂, which we previously demonstrated to be a promoter of C2C12 myoblasts myogenesis⁽³⁾ and proliferation⁽³³⁾.

Our concept is that bone-muscle crosstalk enhances myogenic differentiation but may also have effects on muscle function. In this study, we showed that 10% MLOY4 CM and 10% BP CM enhanced C2C12 cell differentiation *in vitro*. In contrast, the differentiation was not significantly affected when C2C12 cells were treated with 10% 2T3 CM and 10% MC3T3-E1 CM, respectively. In support of this we found that 10% MLOY4 CM increased the force production of SOL muscles stimulated *ex vivo* at lower frequencies of stimulation, suggesting that osteocytes release soluble factors that can either promote increased calcium release from the sarcoplasmic reticulum (SR) or enhance the sensitivity of the contractile machinery to calcium. Consistent with this, we showed that the caffeine-induced calcium release from the SR was significantly increased in C2C12 myotubes treated with MLO-Y4 CM and WNT3a. Another major motivation for the current study was the observation that WNT3a can activate *MyoD* and *Myogenin*⁽³⁴⁾, and these two genes are important myogenic regulatory factors. WNT3a, that is found in the extracellular matrix of different cell types⁽³⁵⁾, is expressed in C2C12 cells and 2T3 osteoblasts⁽³⁶⁻³⁸⁾, and is detected at relatively high levels in serum of both humans (16.9 ± 2.4ng/mL) and mice (0.225-3.74ng/mL)⁽³⁹⁾, suggesting a potential role of WNT3a in bone-muscle crosstalk. In this study, we used RT-qPCR to detect *Wnt3a* expression after C2C12 myoblasts were treated with MLO-Y4 CM, and found that *Wnt3a* expression was not altered (data not shown). We detected WNT3a in ~200 pg/mL range in MLO-Y4 CM and PO CM CMs

by using the only available commercial Elisa kit, and observed stimulation of myogenesis with as little as 500 pg/mL. While this difference seems large, when considering the log scale for the effects of drugs (i.e., dose-response relationship) perhaps this difference is not as profound as it appears to be. In addition, there are many technical reasons that might explain these results, including the fact that it is technically challenging to isolate and purify WNT3a (see <http://web.stanford.edu/group/nusselab/cgi-bin/wnt/purification>). Furthermore, it is possible that the continuous and chronic exposure of cells to pico-gram levels of WNT3a might lead to a host of cell biological effects.

To determine *Wnt3a* expression in MLO-Y4 and 2T3 cells, we investigated it at different time points and under static and FFSS conditions; and found that *Wnt3a* expression was modulated by FFSS. This finding is consistent with the results of Santos and colleagues who reported that MLO-Y4 cells subjected to Pulsating Fluid Flow (PFF) showed increased expression of *Wnt3a*. Jia and colleagues found similar results in osteoblast cells, stimulated with PFF^(40, 41). These results might suggest that with increased levels of activity, bone-secreted WNT3a could increase, leading to signaling to muscle cells that modulate muscle function.

When C2C12 cells were treated with either 10% MLO-Y4 CM or PO CM, and 10ng/mL WNT3a, there was an enhancement of β -catenin translocation into the nucleus, providing evidence that the Wnt/ β -catenin pathway is activated by CM and also by WNT3a. In contrast, when C2C12 cells were treated with either 10% 2T3 CM or 10% MC3T3-E1 CM (osteoblasts CMs), we did not observe any enhancement of myogenic differentiation, suggesting that factors secreted in the osteoblast CM are either different in nature or in concentration. This concept is further supported by the data from osteocytes showing that even 1% CM can enhance myogenic differentiation (**Fig. 4B and 4D**).

To directly test the involvement of WNT3a as a mediator of the effects of CM, we treated MLO-Y4 cell with *Wnt3a* siRNA and collected the CM (MLO-Y4 siRNA-treated CM) (**Fig. 6**). When C2C12 cells were treated with 10% MLO-Y4 siRNA-treated CM, there was no significant change of C2C12 cell differentiation. This indicated that WNT3a is a key factor in MLO-Y4 CM responsible for the enhancement effect of C2C12 cell differentiation.

Next, to further explore if Wnt/ β -catenin pathway was involved in the effect of MLO-Y4 CM and WNT3a on C2C12 cell differentiation enhancement, we pre-treated C2C12 cells with Sclerostin which is a negative regulator of Wnt/ β -catenin signaling pathway. Sclerostin inhibited the capacity of 10% MLO-Y4 CM and 10ng/mL WNT3a to enhance C2C12 cell differentiation, respectively, but failed to inhibit myogenic differentiation acceleration when CM and WNT3a were combined in the same treatment. These results support that Wnt/ β -catenin signaling pathway is involved in MLO-Y4 CM and WNT3a enhancement of C2C12 cell differentiation.

The reason that the combined treatment with MLO-Y4 CM and WNT3a together enhanced C2C12 cell differentiation may be because factors in MLO-Y4 CM (such as WNT3a, WNT1, *et al*) and WNT3a together reached a certain threshold that they competed with Sclerostin to bind to Lrp5 and Lrp6, then decrease the inhibition of Sclerostin on the Wnt/ β -catenin signaling pathway. Furthermore, other factors in CM such as PGE₂ could be contributing to the combined effects of CM and WNT3a.

To search for molecular insights into these effects of CM and WNT3a, we used the Mouse Signal Transduction PathwayFinder PCR Array and found that in 10% MLO-Y4 CM treatment, the expression of two protein coding genes *Tnf* and *Wnt1* were significantly increased (**Fig. 8A**). *Tnf* has been found to exert many of its effects on skeletal muscle through NF κ B signaling⁽⁴²⁾. It has been reported that mechanical stimulation of myoblasts leads to release of

TNF, which is necessary for myogenic differentiation⁽⁴³⁾. TNF is known to be one of the key regulators of skeletal muscle cell responses to injury. Whereas physiological levels of *Tnf* are transiently upregulated during myoblast regenerative responses to injury and stimulate differentiation, sustained high levels of TNF are associated with chronic inflammatory diseases and especially with muscle pathology associated with impairment of differentiation and muscle wasting. *Wnt1* is directly involved in myogenesis⁽⁴⁴⁾. Based on this, we propose that MLO-Y4 CM myogenic effect may be predominantly due to *Tnf* and *Wnt1* activation. Since our data also show that WNT1 can enhance myogenesis (**Supplemental Fig. 1A-C**), the *Wnt1* induced by MLOY4 CM might contribute to autocrine stimulation of the Wnt/ β -catenin signaling pathway. An important question raised by our experiments with CMs and WNT3a was whether other Wnts might exert similar effects. We tested this possibility by treating C2C12 myoblasts with WNT1 ranging from 0.5 to 100ng/mL. Interestingly, we found that the lower concentrations of WNT1 did not increase differentiation, but 10ng/mL, 50ng/mL and 100ng/mL of WNT1 significantly enhanced myogenic differentiation in C2C12 cells, although even at these higher concentrations WNT1 was less potent than WNT3a.

When C2C12 cells were treated with 10ng/mL WNT3a, the expression of two genes (*Axin2*, *Wnt5a*) increased significantly (**Fig. 8B**). *Axin2* associates directly with β -catenin, GSK-3 β and APC (adenomatous polyposis coli) and is induced by active Wnt signaling and acts in a negative feedback loop⁽¹⁶⁾. Thus, upregulation of this gene indicates activation of the Wnt/ β -catenin pathway. *Wnt5a* is involved in modulation of the non-canonical Wnt pathway, which regulates calcium inside cells⁽⁴⁵⁾. Therefore, providing strong evidence that WNT3a is acting via Wnt/ β -catenin pathway and mobilizing intracellular calcium.

In order to further confirm the enhancement of myogenic differentiation by both CM and WNT3a, we monitored with RT-qPCR the expression of two muscle regulatory factors (MRFs, *MyoD* and *Myogenin*) and one marker of terminal differentiation (*Mhc*) (**Fig. 8C-D**). Our results showed upregulation of these genes. To further dissect the mechanisms underlying these effects, we also measured the expression of two Wnt/ β -catenin pathway downstream genes (*Fhl1*, *Numb*), which were both upregulated. *Fhl1* was recently shown to be a downstream gene of the Wnt signaling pathway that promotes differentiation of C2C12 cells ⁽²⁹⁾, and it has been reported that the expression of *Numb* was increased during C2C12 cell differentiation and that treatment with WNT3a increased both *Numb* gene and protein expressions ⁽³⁰⁾.

Therefore, although not exactly the same genes, CM and WNT3a seem to be activating the Wnt/ β -catenin pathway and upregulation MRFs. In addition, striking similarities of overlapping gene regulation by MLO-Y4 CM and WNT3a were revealed by our custom-built muscle specific RT-PCR gene array as summarized in **Fig. 9B**. Our muscle specific –RT PCR gene array results allowed us to demonstrate that both CM and WNT3a upregulated genes associated to pathways linked to Ca^{2+} signaling/hypertrophy, Ca^{2+} homeostasis, mitochondrial biogenesis and oxidative stress (**Table 3**). In support of our observations of enhanced force generation by SOL muscles induced by MLOY4 CM and increased myogenic differentiation and enhanced calcium released induced by both CM and WNT3a, we found overexpression of specific genes associated with SR calcium release (*IP3Rs* ⁽⁴⁶⁻⁴⁸⁾ and *Fkbp1a* ^(49, 50)), SR calcium uptake (*SERCAs* ⁽⁵¹⁾), and store-operated calcium entry/myoblast differentiation (*Stim 2*) ⁽⁵²⁾. These studies also revealed that three key Ca^{2+} signaling genes (*CamkII* ^(53, 54), *Nfatc3* ⁽⁵²⁾ and *Srf* ⁽⁵⁵⁾) critical for myotube homeostasis and cellular hypertrophy were also upregulated as well as two genes associated with mitochondrial biogenesis and REDOX regulation (*PGC1 α* ⁽⁵⁶⁾ and

SOD1 ^(57, 58)). Our results are also in agreement with previous reports that suggested the maintenance of normal intracellular Ca^{2+} concentration is vital to myocyte gene expression and differentiation ⁽³¹⁾. It is also intriguing that Avila et al ⁽³²⁾ has reported that WNT3a modulates intracellular Ca^{2+} in hippocampal neurons, in fact, in a manner similar to that we have observed in muscle cells, suggesting that this might be a common mechanism in excitatory cells. We believe that the modulation in intracellular Ca^{2+} might work as the biophysical linker between the molecular-genetic adaptations and the changes (phenotypes) induced by WNT3a in muscle cells.

A recurring question in this emerging field of bone-muscle crosstalk is how secreted factors from bone and muscle can reach each other. Experiments performed by Fritton and colleagues ⁽⁵⁹⁾ show that mouse tail vein injection of small dyes and molecules up to 70 kDa can permeate the osteocyte-lacunar-canalicular network in just a few minutes, which is bigger than the size of both WNT3a and WNT1 (~40kDa), and in fact the majority of WNTs. This demonstrates that the canalicular fluid has ready access to the circulation and suggests that factors secreted by osteocytes could enter the blood and have effects on distant target cells, including muscle. Furthermore, osteocytes are known to be the main producers of FGF23, which is readily detected in the serum, supporting an endocrine role for the osteocyte (reviewed in ⁽⁷⁾). Another potential mechanism for the transport of factors between tissues is through microvesicles or extracellular vesicles (EVs), cell-derived membrane vesicles secreted by all cell types. The release of EVs is now recognized as an important modulator of crosstalk between many cells types, and in disease states ⁽⁶⁰⁾. Recently, McBride et al ⁽⁶¹⁾ demonstrated that bone marrow mesenchymal stem cell derived exosomes transport WNT3a and enhance dermal fibroblast proliferation and angiogenesis *in vitro*.

Conclusions

Obviously, a limitation of the current study is that we are not studying bone-muscle crosstalk *in vivo*, since this was beyond the scope of the current work. Importantly, the cell models and the *ex vivo* work did support the concept of bone to muscle signaling. In conclusion, our data showed that MLO-Y4 cells and primary osteocytes produce WNT3a. Bone-secreted factors contained in CM and WNT3a enhanced C2C12 cell differentiation through mechanisms associated with stimulation of the Wnt/ β -catenin pathway and muscle regulatory factors. Osteocyte CM also increased the contractile function of *ex vivo* soleus muscles, which might be explained by the increased SR calcium release and the concurrent modulation of intracellular Ca^{2+} signaling/homeostasis. Furthermore, this modulation of Ca^{2+} signaling, might serve another purpose, by linking the effects of WNT3a through the activation of Wnt/ β -catenin pathway, leading to specific phenotypic changes. From our *in vitro* and *ex vivo* data, WNT3a could be one of the bone cell secreted factor(s) that promote differentiation of C2C12 myoblast. Our results support a concept that muscle and bone are not only communicating via mechanical interaction, but also that bone can secrete factor(s) that can act in a paracrine or endocrine fashion to support muscle differentiation and function.

Acknowledgements:

Funding: This study was directly funded by NIH-NIA P01 AG039355 (LFB, MB), the George W. and Hazel M. Jay Endowment (MB), and the UT System Science and Technology Acquisition and Retention Program (STARS) (MB). Dr. Yukiko Kitase prepared MLO-Y4 CM and MC3T3E-1 CM and Ms. Jennifer Rosser prepared 2T3 CM for this study.

Authors' roles: Study design: SLD, MLJ, LFB and MB. Study conduct: JH, SR, NL, CM, SK and LB. Data collection: JH, SR, NL, CM, SK and LB. Data analysis: JH, SR, NL, LFB and MB. Data interpretation: JH, SLD, MLJ, LFB and MB. Drafting manuscript: JH and MB. Revising manuscript: JH, NL, CM, SLD, MLJ, LFB and MB. Approving final version of manuscript: MB. MB take responsibility for the integrity of the data analysis.

Disclosures: All authors state that they have no conflicts of interest.

References

1. Kaji H. Linkage between muscle and bone: common catabolic signals resulting in osteoporosis and sarcopenia. *Curr Opin Clin Nutr Metab Care*. 2013;16(3):272-7.
2. Burge R, Dawson-Hughes B, Solomon DH, Wong JB, King A, Tosteson A. Incidence and economic burden of osteoporosis-related fractures in the United States, 2005-2025. *J Bone Miner Res*. 2007;22(3):465-75.
3. Mo C, Romero-Suarez S, Bonewald L, Johnson M, Brotto M. Prostaglandin E2: from clinical applications to its potential role in bone- muscle crosstalk and myogenic differentiation. *Recent Pat Biotechnol*. 2012;6(3):223-9.
4. Hamrick MW. A role for myokines in muscle-bone interactions. *Exerc Sport Sci Rev*. 2011;39(1):43-7.
5. Jahn K, Lara-Castillo N, Brotto L, Mo CL, Johnson ML, Brotto M, et al. Skeletal muscle secreted factors prevent glucocorticoid-induced osteocyte apoptosis through activation of beta-catenin. *Eur Cell Mater*. 2012;24:197-209; discussion -10.
6. Huang J, Hsu YH, Mo C, Abreu E, Kiel DP, Bonewald LF, et al. METTL21C Is a Potential Pleiotropic Gene for Osteoporosis and Sarcopenia Acting Through the Modulation of the NF-kappaB Signaling Pathway. *J Bone Miner Res*. 2014;29(7):1531-40.
7. Dallas SL, Prideaux M, Bonewald LF. The osteocyte: an endocrine cell ... and more. *Endocr Rev*. 2013;34(5):658-90.
8. Shah AD, Shoback D, Lewiecki EM. Sclerostin inhibition: a novel therapeutic approach in the treatment of osteoporosis. *International journal of women's health*. 2015;7:565-80.
9. Brunkow ME, Gardner JC, Van Ness J, Paepfer BW, Kovacevich BR, Proll S, et al. Bone dysplasia sclerosteosis results from loss of the SOST gene product, a novel cystine knot-containing protein. *American journal of human genetics*. 2001;68(3):577-89.
10. Winkler DG, Sutherland MK, Geoghegan JC, Yu C, Hayes T, Skonier JE, et al. Osteocyte control of bone formation via sclerostin, a novel BMP antagonist. *The EMBO journal*. 2003;22(23):6267-76.
11. Poole KE, van Bezooijen RL, Loveridge N, Hamersma H, Papapoulos SE, Lowik CW, et al. Sclerostin is a delayed secreted product of osteocytes that inhibits bone formation. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. 2005;19(13):1842-4.
12. Li X, Zhang Y, Kang H, Liu W, Liu P, Zhang J, et al. Sclerostin binds to LRP5/6 and antagonizes canonical Wnt signaling. *The Journal of biological chemistry*. 2005;280(20):19883-7.
13. Semenov M, Tamai K, He X. SOST is a ligand for LRP5/LRP6 and a Wnt signaling inhibitor. *The Journal of biological chemistry*. 2005;280(29):26770-5.
14. Clevers H, Nusse R. Wnt/beta-catenin signaling and disease. *Cell*. 2012;149(6):1192-205.
15. Kato M, Patel MS, Levasseur R, Lobov I, Chang BH, Glass DA, 2nd, et al. Cbfa1-independent decrease in osteoblast proliferation, osteopenia, and persistent embryonic eye vascularization in mice deficient in Lrp5, a Wnt coreceptor. *J Cell Biol*. 2002;157(2):303-14.
16. von Maltzahn J, Chang NC, Bentzinger CF, Rudnicki MA. Wnt signaling in myogenesis. *Trends Cell Biol*. 2012;22(11):602-9.
17. Park KH, Brotto L, Lehoang O, Brotto M, Ma J, Zhao X. Ex vivo assessment of contractility, fatigability and alternans in isolated skeletal muscles. *J Vis Exp*. 2012(69):e4198.

18. Brotto M, Brotto L, Jin JP, Nosek TM, Romani A. Temporal adaptive changes in contractility and fatigability of diaphragm muscles from streptozotocin-diabetic rats. *J Biomed Biotechnol.* 2010;2010:931903.
19. de Paula Brotto M, van Leyen SA, Brotto LS, Jin JP, Nosek CM, Nosek TM. Hypoxia/fatigue-induced degradation of troponin I and troponin C: new insights into physiologic muscle fatigue. *Pflugers Arch.* 2001;442(5):738-44.
20. Kato Y, Windle JJ, Koop BA, Mundy GR, Bonewald LF. Establishment of an osteocyte-like cell line, MLO-Y4. *J Bone Miner Res.* 1997;12(12):2014-23.
21. Yang W, Lu Y, Kalajzic I, Guo D, Harris MA, Gluhak-Heinrich J, et al. Dentin matrix protein 1 gene cis-regulation: use in osteocytes to characterize local responses to mechanical loading in vitro and in vivo. *The Journal of biological chemistry.* 2005;280(21):20680-90.
22. Bonewald LF, Harris SE, Rosser J, Dallas MR, Dallas SL, Camacho NP, et al. von Kossa staining alone is not sufficient to confirm that mineralization in vitro represents bone formation. *Calcified tissue international.* 2003;72(5):537-47.
23. Stern AR, Stern MM, Van Dyke ME, Jahn K, Prideaux M, Bonewald LF. Isolation and culture of primary osteocytes from the long bones of skeletally mature and aged mice. *Biotechniques.* 2012;52(6):361-73.
24. Stern AR, Bonewald LF. Isolation of osteocytes from mature and aged murine bone. *Methods in molecular biology (Clifton, NJ).* 2015;1226:3-10.
25. Kitase Y, Barragan L, Qing H, Kondoh S, Jiang JX, Johnson ML, et al. Mechanical induction of PGE2 in osteocytes blocks glucocorticoid-induced apoptosis through both the beta-catenin and PKA pathways. *J Bone Miner Res.* 2010;25(12):2657-68.
26. Noursadeghi M, Tsang J, Hausteiner T, Miller RF, Chain BM, Katz DR. Quantitative imaging assay for NF-kappaB nuclear translocation in primary human macrophages. *J Immunol Methods.* 2008;329(1-2):194-200.
27. Shen J, Yu WM, Brotto M, Scherman JA, Guo C, Stoddard C, et al. Deficiency of MIP/MTMR14 phosphatase induces a muscle disorder by disrupting Ca(2+) homeostasis. *Nat Cell Biol.* 2009;11(6):769-76.
28. Filigheddu N, Gnocchi VF, Coscia M, Cappelli M, Porporato PE, Taulli R, et al. Ghrelin and des-acyl ghrelin promote differentiation and fusion of C2C12 skeletal muscle cells. *Mol Biol Cell.* 2007;18(3):986-94.
29. Lee JY, Chien IC, Lin WY, Wu SM, Wei BH, Lee YE, et al. Fhl1 as a downstream target of Wnt signaling to promote myogenesis of C2C12 cells. *Mol Cell Biochem.* 2012;365(1-2):251-62.
30. Liu XH, Wu Y, Yao S, Levine AC, Kirschenbaum A, Collier L, et al. Androgens up-regulate transcription of the Notch inhibitor Numb in C2C12 myoblasts via Wnt/beta-catenin signaling to T cell factor elements in the Numb promoter. *The Journal of biological chemistry.* 2013;288(25):17990-8.
31. Porter GA, Jr., Makuck RF, Rivkees SA. Reduction in intracellular calcium levels inhibits myoblast differentiation. *The Journal of biological chemistry.* 2002;277(32):28942-7.
32. Avila ME, Sepulveda FJ, Burgos CF, Moraga-Cid G, Parodi J, Moon RT, et al. Canonical Wnt3a modulates intracellular calcium and enhances excitatory neurotransmission in hippocampal neurons. *The Journal of biological chemistry.* 2010;285(24):18939-47.
33. Mo C, Zhao R, Vallejo J, Igwe O, Bonewald L, Wetmore L, et al. Prostaglandin E2 promotes proliferation of skeletal muscle myoblasts via EP4 receptor activation. *Cell Cycle.* 2015;14(10):1507-16.

34. Ridgeway AG, Petropoulos H, Wilton S, Skerjanc IS. Wnt signaling regulates the function of MyoD and myogenin. *The Journal of biological chemistry*. 2000;275(42):32398-405.
35. Willert K, Brown JD, Danenberg E, Duncan AW, Weissman IL, Reya T, et al. Wnt proteins are lipid-modified and can act as stem cell growth factors. *Nature*. 2003;423(6938):448-52.
36. Vaes BL, Dechering KJ, van Someren EP, Hendriks JM, van de Ven CJ, Feijen A, et al. Microarray analysis reveals expression regulation of Wnt antagonists in differentiating osteoblasts. *Bone*. 2005;36(5):803-11.
37. Kalajzic I, Staal A, Yang WP, Wu Y, Johnson SE, Feyen JH, et al. Expression profile of osteoblast lineage at defined stages of differentiation. *The Journal of biological chemistry*. 2005;280(26):24618-26.
38. Tanaka S, Terada K, Nohno T. Canonical Wnt signaling is involved in switching from cell proliferation to myogenic differentiation of mouse myoblast cells. *Journal of molecular signaling*. 2011;6:12.
39. Korkosz M, Gasowski J, Leszczynski P, Pawlak-Bus K, Jeka S, Kucharska E, et al. High disease activity in ankylosing spondylitis is associated with increased serum sclerostin level and decreased wntless protein-3a signaling but is not linked with greater structural damage. *BMC Musculoskelet Disord*. 2013;14:99.
40. Santos A, Bakker AD, Zandieh-Doulabi B, Semeins CM, Klein-Nulend J. Pulsating fluid flow modulates gene expression of proteins involved in Wnt signaling pathways in osteocytes. *J Orthop Res*. 2009;27(10):1280-7.
41. Jia YY, Li F, Geng N, Gong P, Huang SJ, Meng LX, et al. Fluid flow modulates the expression of genes involved in the Wnt signaling pathway in osteoblasts in 3D culture conditions. *Int J Mol Med*. 2014;33(5):1282-8.
42. Li H, Malhotra S, Kumar A. Nuclear factor-kappa B signaling in skeletal muscle atrophy. *J Mol Med (Berl)*. 2008;86(10):1113-26.
43. Zhan M, Jin B, Chen SE, Reecy JM, Li YP. TACE release of TNF-alpha mediates mechanotransduction-induced activation of p38 MAPK and myogenesis. *Journal of cell science*. 2007;120(Pt 4):692-701.
44. Porter JD, Baker RS. Absence of oculomotor and trochlear motoneurons leads to altered extraocular muscle development in the Wnt-1 null mutant mouse. *Brain Res Dev Brain Res*. 1997;100(1):121-6.
45. Miller JR, Hocking AM, Brown JD, Moon RT. Mechanism and function of signal transduction by the Wnt/beta-catenin and Wnt/Ca²⁺ pathways. *Oncogene*. 1999;18(55):7860-72.
46. Rizzuto R, Duchen MR, Pozzan T. Flirting in little space: the ER/mitochondria Ca²⁺ liaison. *Science's STKE : signal transduction knowledge environment*. 2004;2004(215):re1.
47. Hayashi T, Su TP. Sigma-1 receptor chaperones at the ER-mitochondrion interface regulate Ca²⁺ signaling and cell survival. *Cell*. 2007;131(3):596-610.
48. Powell JA, Carrasco MA, Adams DS, Drouet B, Rios J, Muller M, et al. IP(3) receptor function and localization in myotubes: an unexplored Ca²⁺ signaling pathway in skeletal muscle. *Journal of cell science*. 2001;114(Pt 20):3673-83.
49. Gant JC, Blalock EM, Chen KC, Kadish I, Porter NM, Norris CM, et al. FK506-binding protein 1b/12.6: a key to aging-related hippocampal Ca²⁺ dysregulation? *European journal of pharmacology*. 2014;739:74-82.
50. Zalk R, Lehnart SE, Marks AR. Modulation of the ryanodine receptor and intracellular calcium. *Annual review of biochemistry*. 2007;76:367-85.

51. Periasamy M, Kalyanasundaram A. SERCA pump isoforms: their role in calcium transport and disease. *Muscle & nerve*. 2007;35(4):430-42.
52. Phuong TT, Yun YH, Kim SJ, Kang TM. Positive feedback control between STIM1 and NFATc3 is required for C2C12 myoblast differentiation. *Biochem Biophys Res Commun*. 2013;430(2):722-8.
53. Tavi P, Westerblad H. The role of in vivo Ca(2)(+) signals acting on Ca(2)(+)-calmodulin-dependent proteins for skeletal muscle plasticity. *J Physiol*. 2011;589(Pt 21):5021-31.
54. Witczak CA, Jessen N, Warro DM, Toyoda T, Fujii N, Anderson ME, et al. CaMKII regulates contraction- but not insulin-induced glucose uptake in mouse skeletal muscle. *American journal of physiology Endocrinology and metabolism*. 2010;298(6):E1150-60.
55. Guerci A, Lahoute C, Hebrard S, Collard L, Graindorge D, Favier M, et al. Srf-dependent paracrine signals produced by myofibers control satellite cell-mediated skeletal muscle hypertrophy. *Cell metabolism*. 2012;15(1):25-37.
56. Vina J, Gomez-Cabrera MC, Borrás C, Froio T, Sanchis-Gomar F, Martínez-Bello VE, et al. Mitochondrial biogenesis in exercise and in ageing. *Advanced drug delivery reviews*. 2009;61(14):1369-74.
57. Buettner GR. Superoxide dismutase in redox biology: the roles of superoxide and hydrogen peroxide. *Anti-cancer agents in medicinal chemistry*. 2011;11(4):341-6.
58. Buettner GR, Ng CF, Wang M, Rodgers VG, Schafer FQ. A new paradigm: manganese superoxide dismutase influences the production of H₂O₂ in cells and thereby their biological state. *Free radical biology & medicine*. 2006;41(8):1338-50.
59. Beno T, Yoon YJ, Cowin SC, Fritton SP. Estimation of bone permeability using accurate microstructural measurements. *J Biomech*. 2006;39(13):2378-87.
60. Rufino-Ramos D, Albuquerque PR, Carmona V, Perfeito R, Nobre RJ, de Almeida LP. Extracellular vesicles: Novel promising delivery systems for therapy of brain diseases. *Journal of controlled release : official journal of the Controlled Release Society*. 2017:[Epub ahead of print].
61. McBride JD, Menocal-Rodriguez L, Candanedo A, Guzman W, Garcia-Contreras M, Badiavas EV. Bone marrow mesenchymal stem cell-derived CD63+ exosomes transport Wnt3a exteriorly and enhance dermal fibroblast proliferation, migration and angiogenesis in vitro. *Stem cells and development*. 2017:[Epub ahead of print].

Figure Legends

Fig. 1. Osteocyte CM and WNT3a modulate the function of skeletal muscle. (A) MLO-Y4

CM increased the contractile force of *ex vivo* intact SOL muscles. Original record shows the increase in 20 Hz force produced by 10% MLO-Y4 CM, on average the force increased by 25%

(5 mice, 10 muscles, $p < 0.05$). (B) MLO-Y4 CM, Primary Osteocyte (PO) CM, and WNT3a

enhance myogenic differentiation. Representative Fluorescence images of DAPI-stained nuclei

(blue) and myosin heavy chain antibody (MHC, green) stained myocytes/myotubes of C2C12

myoblasts after 3 days in 100% DM media (control), 10% PO CM, 10% MLO-Y4 CM and

10ng/mL WNT3a. (C) Summary data for FI of control, PO CM, MLO-Y4 CM and WNT3a

treatment. The treatment of 10% PO CM, 10% MLO-Y4 CM and 10ng/mL WNT3a promoted

myogenic differentiation compared to the control (n= 3-5, compared to control, 10% PO CM, $p =$

0.001; 10% MLO-Y4 CM, $p = 0.004$; 10ng/mL WNT3a, $p < 0.001$). Calibration bar =100 μm .

Fig. 2. Osteoblast CM does not increase myogenic differentiation. (A) Representative

Fluorescence images of C2C12 myoblasts after 3 days in DM media (control), 10% 2T3 CM,

and 10% MC3T3-E1 CM. (B) Summary data for FI of control, 2T3 CM and MC3T3-E1 CM

treatment. The treatment of 10% 2T3 CM and 10% MC3T3-E1 CM had no significant effect on

myogenic differentiation compared to the control (n= 3-5, compared to control, 10% 2T3 CM, p

= 0.670; 10% MC3T3-E1 CM, $p = 1.000$). Calibration bar =100 μm .

Fig. 3. Expression of *Wnt3a* and other *Wnt* genes in MLO-Y4 cells and 2T3 cells. Summary

data of RT-qPCR results at 24 and 48 hours of culture and 2 hours and 24 hours after FFSS in

MLO-Y4 and 2T3 cells (A-F). (A) *Wnt3a*, *Wnt1* and *Wnt4* gene expression was detected in

MLO-Y4 cells cultured for 24 and 48 hours while the expression of *Wnt5a* and *Wnt7a* was

undetectable. At 48 hours, compared to 24 hours, the expression of *Wnt3a* increased significantly

by 1.995 fold, while the expression of *Wnt1* and *Wnt4* was not changed. (B) In MLO-Y4 cells after FFSS treatment, compared to control, the expression of *Wnt3a* and *Wnt1* increased significantly by 1.95 and 2 fold, respectively. (C) No significant difference in expression of *Wnt3a*, *Wnt1* and *Wnt4* was detected at 24 hours after FFSS treatment, compared to control. (D) In 2T3 cells, at 48 hours, compared to 24 hours, the expression of *Wnt3a*, *Wnt1* and *Wnt4* increased significantly by 2.01, 1.99 and 4.02 fold, respectively. (E) In 2T3 cells after FFSS treatment, compared to control, the expression of *WNT3a* and *Wnt1* increased significantly by 2.02 and 7.98 fold, respectively, and expression of *Wnt4* decreased by 50% ($p = 0.004$). (F), Compared to control, 24 after FFSS, the expression of *Wnt3a* increased by 11.9 fold and *Wnt1* and *Wnt4* decreased by 50.1% ($p = 0.001$) and 50.7% ($p = 0.001$), respectively (n=3).

Fig. 4. Lower concentrations, more physiological levels of WNT3a and 1% MLO-Y4 CM enhance C2C12 myoblast differentiation. Fluorescence images of DAPI-stained nuclei (blue) and myosin heavy chain antibody (MHC, green) stained myocytes/myotubes of C2C12 cells at differentiation day 3 after WNT3a treatment (A) and 1% MLO-Y4 CM (B). (C) Summary data for FI for treatments ranging from 0.1ng/mL to 10ng/mL WNT3a. FI increased significantly for concentration ranging from 0.5 to 10ng/mL WNT3a compared to control (n=3-5; compared to control, 0.5ng/mL WNT3a, $p = 0.028$; 1ng/mL WNT3a, $p = 0.016$; 2.5ng/mL WNT3a, $p = 0.004$; 5ng/mL WNT3a, $p = 0.001$; 10ng/mL WNT3a, $p < 0.001$). (D) Summary data for 1% MLO-Y4 CM-treated C2C12 cells at differentiation day3. FI increased significantly 1% MLO-Y4 CM compared to control (n=3-5, $p = 0.014$). Calibration bar =100 μm .

Fig. 5. MLO-Y4 CM, PO CM, and WNT3a induced nuclear translocation of β -catenin in C2C12 myoblasts. (A) Immunostaining for β -catenin of C2C12 myoblasts at 30 minutes, 60 minutes and 90 minutes after treatment with 10% MLO-Y4 CM. (B) Quantification of the

nuclear β -catenin staining showed that the maximum induction of nuclear translocation of β -catenin in C2C12 myoblasts occurred at 60 minutes (n=3, 5-10 cells per experiment, $p < 0.001$).

(C) Confocal images of immunostaining for β -catenin of C2C12 cells cultured in GM, DM, treated 60 minutes in DM with 10% PO CM, 10% MLO-Y4 CM, 10ng/mL WNT3a and Lithium chloride was used as positive control. (D) Summary data show that compared to GM, DM promoted significantly higher nuclear translocation of β -catenin (n=3, 5-10 cells per experiment, $p = 0.023$) and compared to DM, PO CM, MLO-Y4 CM, WNT3a and LiCl induced significantly higher nuclear translocation of β -catenin (compared to DM, PO CM, $p < 0.001$; MLO-Y4 CM, $p = 0.003$; WNT3a, $p < 0.001$; LiCl, $p < 0.001$).

Fig. 6. *Wnt3a* knockdown in MLO-Y4 cells impairs CM capacity to enhance myogenic differentiation. MLO-Y4 cells were transfected with *Wnt3a* siRNA. (A) Twenty hours after transfection, the transfection efficiency equaled $80.02 \pm 7.18\%$, leading *Wnt3a* expression decreased to $20.47 \pm 4.72\%$ compared to negative control (n=3). (B) Fluorescence images of DAPI- and MHC stained C2C12 cells at differentiation day 3 after treatment of MLO-Y4 siRNA-treated CM. (C) Summary data for FI for treatments ranging from DM control (a), 10% MLO-Y4 siRNA-treated CM (b), 10% CM from MLO-Y4 cells treated with negative control siRNA (c) and 10% MLO-Y4 CM as positive control (d). FI significantly increased in 10% CM from MLO-Y4 cells treated with negative control siRNA and positive control compared to DM control, while no significant difference was found in 10% MLO-Y4 siRNA-treated CM treatment (n=3; compared to control, b, $p = 0.318$; c, $p < 0.001$; d, $p < 0.001$). Calibration bar =100 μm .

Fig. 7. Sclerostin inhibits the effects of MLO-Y4 CM and WNT3a on myogenic

differentiation. (A) Fluorescence images of DAPI and MHC stained C2C12 cells at differentiation day 3. (B) Summary data for FI for treatment groups: (a) DM without Sclerostin treatment as a negative control, (b) DM with 100ng/mL Sclerostin for 6 hours, (c) DM with 10% MLO-Y4 CM for 48 hours as one of our positive controls, (d) DM with 10ng/mL WNT3a for 48 hours as our second positive control, (e) DM with 100ng/mL Sclerostin for 6 hours, followed by 10% MLO-Y4 CM for 48 hours, (f) DM with 100ng/mL Sclerostin for 6 hours, followed by 10ng/mL WNT3a for 48 hours, and (g) DM with 100ng/mL Sclerostin for 6 hours, followed by the combination of 10% MLO-Y4 CM + 10ng/mL WNT3a for 48 hours. # No significant difference of FI was found for groups (b), (e) and (f), compared to negative control (a), respectively. (n=3-6; compared to negative control, b, $p = 0.925$; e, $p = 0.971$; f: $p = 1.000$). * FI increased significantly for group (g) and positive controls (c and d), compared to negative control. (n=3-6; compared to negative control, g, $p = 0.030$; c, $p = 0.007$; d, $p < 0.001$). Calibration bar = 100 μ m.

Fig. 8. RT-PCR Gene Array and RT-qPCR detection of gene alterations in C2C12 cells after MLO-Y4 CM and WNT3a treatment. Summary data from 3 independent experiments for the Mouse Signal Transduction Pathway Finder PCR Array (A) In 10% MLO-Y4 CM, the expression of *Tnf* and *Wnt1* were significantly increased ($p = 0.031$; $p = 0.022$, respectively). (B) When cells treated with 10ng/mL WNT3a, the expression of *Axin2* and *Wnt5a* increased significantly ($p < 0.001$; $p = 0.011$, respectively). Summary data of RT-qPCR detection of gene expression changes for *Mhc*, *MyoD*, *MyoG* and *Fhl1*, *Numb* at differentiation day3 after 1%, 10% MLO-Y4 CM (C) and 0.5ng/mL and 10ng/mL WNT3a treatment (D). The results demonstrated that both 1% and 10% MLO-Y4 CM, as well as 0.5ng and 10ng/mL WNT3a

(except *Numb* in 1% MLO-Y4 CM, $p = 0.084$) significantly increased the expression of these five specific genes as compared to control (n=3, 2 replicates). *Mhc*, *MyoD*, *MyoG* and *Fhl1* increased significantly after 1% MLO-Y4 CM treatment ($p = 0.002$; $p = 0.003$; $p = 0.004$; $p < 0.043$, respectively). *Mhc*, *MyoD*, *MyoG*, *Fhl1* and *Numb* increased significantly after 10% MLO-Y4 CM treatment ($p = 0.001$; $p < 0.001$; $p = 0.006$, $p = 0.002$; $p < 0.001$, respectively). After 0.5ng/mL WNT3a treatment, *Mhc*, *MyoD*, *MyoG*, *Fhl1* and *Numb* increased significantly ($p = 0.004$; $p = 0.002$; $p = 0.001$; $p = 0.001$; $p < 0.001$, respectively). After 10ng/mL WNT3a treatment, *Mhc*, *MyoD*, *MyoG*, *Fhl1* and *Numb* increased significantly ($p = 0.006$; $p = 0.001$, $p < 0.001$; $p < 0.001$, $p < 0.001$, respectively).

Fig. 9. MLO-Y4 CM and WNT3a enhance Ca^{2+} release from the SR and modulate expression of key Ca^{2+} signaling/homeostasis genes. (A) Representative Ca^{2+} transient of C2C12 myotubes loaded with Fura-2/AM exposed to 20mM caffeine (arrows). MLO-Y4 CM and WNT3a treatment increased caffeine induced Ca^{2+} release in C2C12 myotubes. In 10% MLO-Y4 CM and 10ng/mL WNT3a-treated C2C12 myotubes, multiple oscillatory Ca^{2+} peaks in response to caffeine were observed and the relaxation phase of the transient was significantly longer ($p < 0.05$). (B) Increased expression of key genes related to Ca^{2+} signaling/hypertrophy, Ca^{2+} homeostasis, mitochondrial biogenesis and oxidative stress in C2C12 cells at 60 hours of differentiation after treatments was detected by a focused muscle custom-built RT-PCR gene array. Two PCR arrays provided similar results.

Table1. RT-qPCR primers used in this study


Primer	Primer sequence (5' to 3')
GAPDHF	TGCGATGGGTGTGAACCACGAGAA
GAPDHR	GAGCCCTTCCACAATGCCAAAGTT
NumbF	AACCAGCCTTTGTCCCTACC
NumbR	GGCGACTGATGTGGATGAG
Fhl1F	TTCTGGATCCCATCCTGTGTGAG
Fhl1R	ACACGGTACCTGGTTCCAATGCC
MYHCF	CAAGTCATCGGTGTTTGTGG
MYHCR	TGTCGTA CTTGGGCGGGTTC
MyoDF	CCCCGGCGGCAGAATGGCTACG
MyoDR	GGTCTGGGTTCCTGTTCTGTGT
MyoGF	TGAGCATTGTCCAGGCCAG
MyoGR	GCTTCTCCCTCAGTGTGGCT
Wnt1F	ATGAACCTTCACAACAACGAG
Wnt1R	GGTTGCTGCCTCGGTTG
Wnt3aF	TAGATGGGTGCGACCTGTTG
Wnt3aR	GAACCCTGCTCCCGTGTTAG
Wnt4F	CGAGGAGTGCCAATACCAGT
Wnt4R	GCCACACCTGCTGAAGAGAT
Wnt5aF	GGCGAGCTGTCTACCTGTGG
Wnt5aR	GGCGAACGGGTGACCATAGT
Wnt7aF	CGACTGTGGCTGCGACAAG
Wnt7aR	CTTCATGTTCTCCTCCAGGATCTTC

Table 2. Pathways and genes in Mouse Signal Transduction PathwayFinder PCR Array

Signaling Pathways	Genes
TGFβ Pathway	<i>Atf4, Cdkn1b (p27Kip1), Emp1, Gadd45b, Herpud1, Ifrd1, Myc, Tnfrsf10</i>
WNT Pathway	<i>Axin2, Ccnd1, Ccnd2, Dab2, Fosl1 (Fra-1), Mmp7 (Matrilysin), Myc, Ppard, Wisp1</i>
NFκB Pathway	<i>Bcl2a1a (Bfl-1/A1), Bcl2l1 (Bcl-x), Birc3 (c-IAP1), Ccl5 (Rantes), Csf1 (Mcsf), Icam1, Ifng, Stat1, Tnf.</i>
JAK/STAT Pathway	
JAK1, 2 / STAT1	<i>Irf1</i>
STAT3	<i>Bcl2l1 (Bcl-x), Ccnd1, Cebpd, Lrg1, Mcl1, Socs3</i>
STAT5	<i>Bcl2l1 (Bcl-x), Ccnd1, Socs3</i>
JAK1, 3 / STAT6	<i>Fcer2a, Gata3</i>
p53 Pathway	<i>Bax, Bbc3, Btg2, Cdkn1a (p21Cip1/Waf1), Egfr, Fas (Tnfrsf6), Gadd45a, Pcna, Rb1</i>
Notch Pathway	<i>Hes1, Hes5, Hey1, Hey2, Heyl, Id1, Jag1, Lfng, Notch1</i>
Hedgehog Pathway	<i>Bcl2, Bmp2, Bmp4, Ptch1, Wnt1, Wnt2b, Wnt3a, Wnt5a, Wnt6</i>
PPAR Pathway	<i>Acsl3, Acsl4, Acsl5, Cpt2, Fabp1, Olr1, Slc27a4, Sorbs1</i>
Oxidative Stress	<i>Fth1, Gclc, Gclm, Gsr, Hmox1, Nqo1, Sqstm1, Txn1, Txnrd1</i>
Hypoxia	<i>Adm, Arnt, Car9, Epo, Hmox1, Ldha, Serpine1 (PAI-1), Slc2a1, Vegfa</i>

Table 3. Pathways and genes in focused custom-built RT-PCR dene array

Signaling Pathways	Genes
Ca²⁺ signaling/ cellular hypertrophy	<i>Nfatc3</i> ⁽⁵²⁾ , <i>CamkII</i> ^(53, 54) , <i>Srf</i> ⁽⁵⁵⁾
Ca²⁺ homeostasis	<i>IP3-R1</i> , <i>IP3-R3</i> ⁽⁴⁶⁻⁴⁸⁾ , <i>Fkbp1a</i> ^(49, 50) , <i>SERCA1a</i> , <i>SERCA2a</i> ⁽⁵¹⁾ , <i>Stim 2</i> ⁽⁵²⁾
Mitochondrial biogenesis	<i>PGC1α</i> ⁽⁵⁶⁾
Oxidative stress/Redox	<i>SOD1</i> ^(57, 58)






Fig. 2A-B




Fig. 3A-F




Fig. 4A-D




Fig. 5A-D





Fig. 6A-C






Fig. 8A-D




Fig. 9A-B