

6-28-2017

DACH1 suppresses breast cancer as a negative regulator of CD44.

Hanxiao Xu

Department of Oncology, Tongji Hospital of Tongji Medical College, Huazhong University of Science and Technology, Wuhan, 430030, China

Shengnan Yu

Department of Oncology, Tongji Hospital of Tongji Medical College, Huazhong University of Science and Technology, Wuhan, 430030, China

Xun Yuan

Department of Oncology, Tongji Hospital of Tongji Medical College, Huazhong University of Science and Technology, Wuhan, 430030, China

Jing Xiong

Department of Pathology, Tongji Hospital of Tongji Medical College, Huazhong University of Science and Technology, Wuhan, 430030, China

Dong Kuang

Department of Pathology, Tongji Hospital of Tongji Medical College, Huazhong University of Science and Technology, Wuhan, 430030, China

[Let us know how access to this document benefits you](#)

See next page for additional authors
Follow this and additional works at: <https://jdc.jefferson.edu/cbfp>

 Part of the [Oncology Commons](#)

Recommended Citation

Xu, Hanxiao; Yu, Shengnan; Yuan, Xun; Xiong, Jing; Kuang, Dong; Pestell, Richard; and Wu, Kongming, "DACH1 suppresses breast cancer as a negative regulator of CD44." (2017). *Department of Cancer Biology Faculty Papers*. Paper 116.

<https://jdc.jefferson.edu/cbfp/116>

Authors

Hanxiao Xu, Shengnan Yu, Xun Yuan, Jing Xiong, Dong Kuang, Richard Pestell, and Kongming Wu

SCIENTIFIC REPORTS



OPEN

DACH1 suppresses breast cancer as a negative regulator of CD44

Hanxiao Xu¹, Shengnan Yu¹, Xun Yuan¹, Jing Xiong², Dong Kuang², Richard G. Pestell³ & Kongming Wu¹ 

Received: 17 January 2017

Accepted: 18 May 2017

Published online: 28 June 2017

Dachshund homolog 1 (DACH1), a key cell fate determination factor, contributes to tumorigenesis, invasion, metastasis of human breast neoplasm. However, the exact molecular mechanisms for the anti-tumor roles of DACH1 in breast carcinoma are still lack of extensive understanding. Herein, we utilized immunohistochemistry (IHC) staining and public microarray data analysis showing that DACH1 was higher in normal breast, low-grade and luminal-type cancer in comparison with breast carcinoma, high-grade and basal-like tumors respectively. Additionally, both correlation analysis of public databases of human breast carcinoma and IHC analysis of mice xenograft tumors demonstrated that DACH1 inversely related to cancer stem cells (CSCs) markers, epithelial-mesenchymal transition (EMT) inducers and basal-enriched molecules, while cluster of differentiation 44 (CD44) behaved in an opposite manner. Furthermore, mice transplanted tumor model indicated that breast cancer cells Met-1 with up-regulation of DACH1 were endowed with remarkably reduced potential of tumorigenesis. Importantly, meta-analysis of 19 Gene Expression Omnibus (GEO) databases of breast cancer implicated that patients with higher *DACH1* expression had prolonged time to death, recurrence and metastasis, while *CD44* was a promising biomarker predicting worse overall survival (OS) and metastasis-free survival (MFS). Collectively, our study indicated that CD44 might be a novel target of DACH1 in breast carcinoma.

In spite of significant achievement made in early diagnosis and therapeutic strategies, breast cancer still draws great attention from the worldwide because of its high incidence rate and mortality^{1–3}. The unsatisfactory clinical outcome is mostly due to tumor recurrence, metastasis and therapy-resistance¹. Identifying novel biomarkers related to molecular subtypes, aggressive phenotypes and prognosis of breast cancer is essential for drug development, disease surveillance and precise therapy.

The retinal determination gene network (RDGN), including DACH1, EYA1 and SIX1, plays crucial roles in the development of multiple organs⁴. SIX1 and EYA1, two important RDGN members, exert favorable effects on tumor initiation and progression^{4,5}, and high expression of SIX1 and EYA1 is an adverse factor for clinical outcomes for breast cancer patients^{6–8}. On the contrary, another key RDGN member DACH1 behaved as a tumor suppressor and reduced expression of DACH1 predicts poor survival performance of breast cancer patients⁹. Several lines of evidence have demonstrated that the hypermethylation of promoter region leads to the down-regulation of DACH1, which is closely associated with proliferation, invasion and metastasis of various tumors, including breast cancer^{10–13}, lung cancer¹⁴, esophageal cancer¹⁵, renal cell carcinoma¹⁶ and hepatocellular carcinoma¹⁷. DACH1 antagonizes the transcription and translation of oncogenes and induces epithelial-mesenchymal transition (EMT) in breast cancer, resulting in the inhibition of tumor growth, invasion and migration^{9,10}. Recent studies prove that cancer stem cells (CSCs) possess potent self-renewal ability and are responsible for tumor relapse and metastasis and endogenous DACH1 participates in the negative regulation of CSCs^{18,19}.

Cluster of differentiation-44 (CD44), a ubiquitously present glycoprotein on the membrane of mammalian cells, plays essential roles in a variety of biological function such as cell division, adhesion and migration²⁰. During the past decades, the role of CD44 in cancer development has been revealed and valued. As a well-known marker of CSCs, CD44 promotes carcinogenesis, invasion, metastasis and therapy-resistance^{20–23}. It promotes proliferation and suppresses apoptosis by regulation of relative pathways, including Ras-Raf-Mek-Erk-Cyclin D1

¹Department of Oncology, Tongji Hospital of Tongji Medical College, Huazhong University of Science and Technology, Wuhan, 430030, China. ²Department of Pathology, Tongji Hospital of Tongji Medical College, Huazhong University of Science and Technology, Wuhan, 430030, China. ³Department of Cancer Biology, Sidney Kimmel Cancer Center, Thomas Jefferson University, Philadelphia, PA, 19107, USA. Correspondence and requests for materials should be addressed to K.W. (email: kmwu@tjh.tjmu.edu.cn)




Figure 1. DACH1 and CD44 were correlated with tumorigenesis and histological grade of breast carcinoma. (a) Representative images of immunohistochemistry staining of DACH1 and CD44 in noncancerous and cancerous tissues were shown. (b) Representative images of immunohistochemistry staining of DACH1 and CD44 in low-grade and high-grade breast neoplasm tissues were shown.

pathway and phosphoinositide 3-kinase (PI3K)-Akt signaling, as well as stimulates EMT, which contributes to tumor invasion and metastasis^{20, 24}.

Previous study has demonstrated that endogenous reduction of DACH1 was accompanied by down-regulation of CSC markers, such as SOX2, Nanog, KLF4¹⁸. To further evaluate the correlation between DACH1 and CSC markers and EMT inducers in breast cancer, we performed a comprehensive analysis of immunohistochemistry (IHC) staining, publicly available microarray data, RNA profiling and western blot. Our study indicated that DACH1 was inversely correlated with CD44 and CD44 might be a novel target of DACH1 in breast cancer.

Results

DACH1 and CD44 associated with tumorigenesis and histological grade of breast cancer. In order to evaluate the expression of DACH1 and CD44 in normal breast and breast malignant tissues, we carried out IHC analysis on two TMAs (BR1502-97 and BR1502-98) with normal breast and human breast cancer tissues. DACH1 was majorly found in nucleus and CD44 was mostly detected on the membrane of breast cancer cells. Representative images of IHC staining for noncancerous and cancerous tissues were shown in Fig. 1a, showing that DACH1 decreased and CD44 increased in breast neoplasm tissues in comparison with normal breast.

Additionally, we also explored the correlation between the protein abundance of DACH1 and CD44 and histological grade. Representative images of IHC staining for low-grade and high-grade cancerous tissues were shown in Fig. 1b, which indicating that DACH1 was inversely correlated with tumor grade, while CD44 was positively associated with histological grade.

Expression of DACH1 and CD44 correlated with molecular subtypes of breast cancer. We also carried out IHC staining to assess the protein abundance of DACH1 and CD44 in luminal-type and basal-like




Figure 2. The expression of DACH1 and CD44 correlated with molecular subtypes of breast carcinoma. (a) Representative images of immunohistochemistry staining of DACH1 and CD44 in luminal-type and basal-like breast cancer tissues were shown. (b) Expression analysis of public microarray dataset GSE20711 showed that the mRNA level of *DACH1* was significantly higher in luminal-type than in basal-like breast tumor. (c) Expression analysis of GSE 20711 also displayed that *CD44* mRNA expression was remarkably lower in luminal-type than in basal-like breast cancer.

breast cancer tissues. Representative images for the expression of DACH1 and CD44 in luminal and basal tissues were shown in Fig. 2a. Furthermore, expression analysis of GSE20711 including a total of 45 luminal and 22 basal-like breast tumor cases was also interrogated to evaluate the mRNA levels of *DACH1* and *CD44* in luminal and basal-like breast neoplasm tissues, which showed that *DACH1* was enriched in luminal breast carcinoma in comparison with basal-like breast cancer ($P < 0.001$) (Fig. 2b), while *CD44* exhibited an opposite tendency ($P < 0.001$) (Fig. 2c) at mRNA level. Altogether, our results indicated that luminal breast carcinoma was most likely to be *DACH1*^{high}/*CD44*^{low} type, and basal-like breast tumor tissues were majorly *DACH1*^{low}/*CD44*^{high} type.

DACH1 down-regulated some CSC and EMT markers *in vitro*, as well as blocked Met-1 tumor growth *in vivo*. Breast cancer Met-1 cells were transfected with a *DACH1* expression vector resulting in an ~4.5-fold increase in *DACH1* expression (Fig. 3a) and subsequent reduction of CSC markers *CD44*, *KLF4* and *MYC* as well as EMT markers including *FN1* and *VIM* (Fig. 3b) by mRNA analysis. Western blot also demonstrated the presence of the DACH1-tagged FLAG epitope, and the effects of DACH1 overexpression on the protein abundance of CD44, Fibronectin, Vimentin, p21 and p27, which were shown in Fig. 3c. Ectopic expression of DACH1 contributed to remarkable reduction of CD44, Fibronectin, Vimentin and significant up-regulation of p21 and p27 in Met-1 cells. Mammary tumor growth *in vivo* was assessed by subcutaneous implantation of Met-1 cells in nude mice (Fig. 3f). Met-1 cells with engineered expression of DACH1 were endowed with remarkably reduced potential of tumorigenesis in xenograft tumors. Up-regulation of DACH1 significantly reduced the volume of tumors by ~90% and slowed down tumor growth (Fig. 3d). Tumor weight was also reduced by ~90% in comparison with the control tumors (Fig. 3e). The results implicated that DACH1 suppressed the expression of some CSCs and EMT markers and serves as a potent anti-tumor factor in xenograft tumors.




Figure 3. DACH1 regulated the expression of some CSCs and EMT genes in Met-1 cells and suppressed tumor growth *in vivo*. (a) Stable expression of *DACH1* in breast cancer Met-1 cells was achieved by retrovirus infection. (b) RNA microarray and cluster analysis showed that upregulation of *DACH1* reduced the mRNA levels of *CD24*, *CD44*, *KLF4*, *MYC*, *FNI* and *VIM* in Met-1 cells. (c) Western blot indicated that upregulation of *DACH1* reduced the protein abundance of *CD44*, Fibronectin, Vimentin, p21 and p27 in Met-1 cells. (d) *DACH1* overexpression significantly reduced the volume of tumors by ~90% and slowed down the speed of tumor growth in nude mice xenograft tumors. (e) Overexpression of *DACH1* also reduced mice transplanted tumor weight by ~90%. (f) Mammary tumor growth *in vivo* was evaluated by subcutaneous implantation of *DACH1*-overexpressing Met-1 cells and the GFP controls in nude mice.

DACH1 reduced the expression of *CD44*, Fibronectin, Vimentin, Myc, Sox2, EGFR, Ki-67 *in vivo*. Immunohistochemistry analysis was conducted to assess the protein abundance of *DACH1*, *CD44*, *Myc*, *Sox2*, Fibronectin, Vimentin, EGFR, Ki-67 in nude mice xenograft tumor tissues with overexpression of *DACH1* and the GFP controls. Additionally, we also employed IHC scoring to quantize the levels of these proteins in both *DACH1*-overexpressing and the control tumors by using semi-quantitative criteria. About six 200 magnification images of each kind of protein were selected for IHC scoring by two experienced pathologists independently. Representative images of IHC staining and scoring results for *DACH1*, *CD44*, *Myc*, *Sox2*, Fibronectin, Vimentin, EGFR and Ki-67 were shown in Fig. 4a,b,c,d,e,f,g and h, respectively. Our results displayed that over-expression of *DACH1* ($P < 0.001$) remarkably reduced the expression of *CD44* ($P < 0.001$), *Myc* ($P < 0.001$), *Sox2* ($P < 0.001$), Fibronectin ($P < 0.001$), Vimentin ($P < 0.001$), EGFR ($P < 0.001$) and Ki-67 ($P < 0.001$), demonstrating that *DACH1* could potentially down-regulated the expression of some CSCs and EMT markers, basal-like factor EGFR and proliferative biomarker Ki-67 *in vivo* at protein level.




Figure 4. DACH1 reduced the expression of CD44, Myc, Sox2, Fibronectin, Vimentin, EGFR and Ki-67 *in vivo*. Both representative immunohistochemistry images and scoring results showed that overexpression of DACH1 (a) down-regulated the protein abundance of CD44 (b), Myc (c), Sox2 (d), Fibronectin (e), Vimentin (f), EGFR (g) and Ki-67 (h) in nude mice xenograft tumors.

Correlation between the expression of DACH1 and CD44 and the levels of FN1, VIM, YBX1, FOXA1, EGFR and MKI67. Previous study has implicated that DACH1 enriched in luminal A breast cancer and its expression fluctuated in direct proportion to the level of luminal-like marker FOXA1¹³. Previously experimental study demonstrated that DACH1 participated in the inhibition of Snail-induced EMT through suppressing the activity of the Y box-binding protein (YB-1)⁹. Herein, public dataset GSE20685 was interrogated to assess the association between DACH1 and CD44, FN1, VIM, FOXA1, EGFR and MKI67, as well as evaluate the correlation between CD44 and the above genes. The results showed that DACH1 mRNA expression was inversely correlated with CD44 ($R = -0.341$, $P < 0.001$) (Fig. 5a), FN1 ($R = -0.214$, $P < 0.001$) (Fig. 5b), VIM ($R = -0.229$, $P < 0.001$) (Fig. 5c), EGFR ($R = -0.390$, $P < 0.001$) (Fig. 5e) and MKI67 ($R = -0.376$, $P < 0.001$) (Fig. 5f), but




Figure 5. The expression of *DACH1* and *CD44* correlated with *VIM*, *FNI*, *YBX1*, *FOXA1*, *EGFR* and *MKI67* in breast cancer tissues. Correlation analysis of public dataset GSE 20685 showed that *DACH1* was inversely correlated with cancer stem cell marker *CD44* (a), mesenchymal markers *FNI* (b) and *VIM* (c) as well as basal-like markers *EGFR* (e) and *MKI67* (f), while positively associated with luminal marker *FOXA1* (d). *CD44* was parallel with *FNI* (g) and *VIM* (h), *YBX1* (i), *EGFR* (k) and *MKI67* (l), while negatively associated with *FOXA1* (j).

positively associated with the mRNA expression of *FOXA1* ($R = 0.608$, $P < 0.001$) (Fig. 5d). Conversely, *CD44* was found to be positively correlated with *FNI* ($R = 0.253$, $P < 0.001$) (Fig. 5g), *VIM* ($R = 0.237$, $P < 0.001$) (Fig. 5h), *YBX1* ($R = 0.446$, $P < 0.001$) (Fig. 5i), *EGFR* ($R = 0.336$, $P < 0.001$) (Fig. 5k) and *MKI67* ($R = 0.215$, $P < 0.001$) (Fig. 5l), while there was a significantly negative association between *CD44* and *FOXA1* ($R = -0.402$, $P < 0.001$) (Fig. 5j).

Breast cancer cell line data reported by Neve RM²⁵, including a total of 51 different breast cancer lines from luminal-type ($N = 25$) or basal-like ($N = 26$), were also employed to evaluate the correlation between the expression of *DACH1* and *CD44* and the levels of *FNI*, *VIM*, *YBX1*, *FOXA1*, *EGFR* and *MKI67*. The results displayed




Figure 6. The association between the expression of *DACHI* and *CD44* with *VIM*, *FNI*, *YBX1*, *FOXA1*, *EGFR* and *MKI67* in breast cancer cell lines. Correlation analysis of data with a total of 51 breast cancer cell lines showed that *DACHI* was inversely correlated with *CD44* (a), *FNI* (b), *VIM* (c), *EGFR* (e) and *MKI67* (f), while positively associated with *FOXA1* (d). In contrast, *CD44* was parallel with *FNI* (g), *VIM* (h), *YBX1* (i), *EGFR* (k) and *MKI67* (l), but negatively related to *FOXA1* (j).

that *DACHI* mRNA expression was inversely correlated with *CD44* ($R = -0.507$, $P < 0.001$) (Fig. 6a), *FNI* ($R = -0.354$, $P = 0.011$) (Fig. 6b), *VIM* ($R = -0.419$, $P = 0.002$) (Fig. 6c), *EGFR* ($R = -0.523$, $P < 0.001$) (Fig. 6e) and *MKI67* ($R = -0.336$, $P = 0.016$) (Fig. 6f) but positively associated with the mRNA level of *FOXA1* ($R = 0.539$, $P < 0.001$) (Fig. 6d), which were consistent with the results got in GSE20685. In contrast, *CD44* was parallel with *FNI* ($R = 0.406$, $P = 0.003$) (Fig. 6g), *VIM* ($R = 0.614$, $P < 0.001$) (Fig. 6h), *YBX1* ($R = 0.557$, $P < 0.001$) (Fig. 6i), *EGFR* ($R = 0.506$, $P < 0.001$) (Fig. 6k) and *MKI67* ($R = 0.391$, $P = 0.005$) (Fig. 6l) but negatively associated with *FOXA1* ($R = -0.689$, $P < 0.001$) (Fig. 6j), supporting the conclusion from the correlation analysis of human breast tumor samples (GSE20685).

First Author	GSE accession	Year	Duration (Months)	Patient Number	Detection	Platform
Desmedt C ²⁶	GSE7390	2007	163.2	198	Microarray	GPL96
Pawitan Y ²⁷	GSE1456	2005	102	159	Microarray	GPL96
Hennessy BT ²⁸	GSE10885	2009	106	89	Microarray	GPL887
Dedeurwaerder S ²⁹	GSE20711	2011	169	88	Microarray	GPL570
Kao KJ ³⁰	GSE20685	2011	156	327	Microarray	GPL570
Clarke C ³¹	GSE42568	2013	100.9	104	Microarray	GPL570
Desmedt C ³²	GSE16446	2011	60	120	Microarray	GPL570
Loi S ³³	GSE6532	2010	176.8	327	Microarray	GPL96
Bild AH ³⁴	GSE3143	2006	156	158	Microarray	GPL8300
Terunuma A ³⁵	GSE39004	2014	120	61	Microarray	GPL6244
Heikkinen T ³⁶	GSE24450	2011	120	183	Microarray	GPL6947
Hatzis C ³⁷	GSE25066	2011	120	508	Microarray	GPL96
Symmans WF ³⁸	GSE17705	2010	196	298	Microarray	GPL96
Wang Y ³⁹	GSE2034	2005	180	286	Microarray	GPL96
Tofigh A ⁴⁰	GSE58644	2014	145	321	Microarray	GPL6244
Minn AJ ⁴¹	GSE2603	2005	130	99	Microarray	GPL96
Minn AJ ⁴²	GSE5327	2007	156	58	Microarray	GPL96
Sircoulomb F ⁴³	GSE17907	2010	112	51	Microarray	GPL570
Nagalla S ⁴⁴	GSE45255	2013	127.4	139	Microarray	GPL96

Table 1. Characteristics of the included public microarray datasets in the meta-analysis.

The opposite roles of *DACH1* and *CD44* in clinical outcomes of breast cancer patients. In order to assess the prognostic value of *DACH1* and *CD44* in breast cancer, a meta-analysis enrolling a total of 19 published Gene Expression Omnibus (GEO) databases and including 3574 breast cancer patients was performed^{26–44}. The characteristics of these 19 GSE databases were displayed in Table 1. The results indicated that patients with higher mRNA expression of *DACH1* tended to enjoy longer time to death (HR: 0.72 (0.57–0.92), $I^2 = 22.9\%$, $P = 0.232$) (Fig. 7a), relapse (HR: 0.73 (0.56–0.97), $I^2 = 64.1\%$, $P = 0.003$) (Fig. 7c) and metastasis (HR: 0.73 (0.56–0.95), $I^2 = 27.2\%$, $P = 0.202$) (Fig. 7e). On the contrary, higher mRNA expression of *CD44* was directly related to worse OS (HR: 1.27 (1.03–1.57), $I^2 = 3.0\%$, $P = 0.412$) (Fig. 7b) and MFS (HR: 1.40 (1.02–1.93), $I^2 = 48.4\%$, $P = 0.050$) (Fig. 7f), but did not insignificantly contribute to worse RFS (HR: 1.19 (0.94–1.51), $I^2 = 45.7\%$, $P = 0.056$) (Fig. 7d). Our results demonstrated that *DACH1* was a promising biomarker predictive of better clinical outcomes, while *CD44* was an adverse factor for the survival performance of breast cancer patients.

Discussion

DACH1, as an important member of RDGN, is widely expressed in epithelial cells and plays critical roles in normal organ development^{45,46}. However, its absent or lower expression contributes to the initiation and progression of various tumor types, including lung adenocarcinoma¹⁴, pancreatic cancer⁴⁷, breast cancer¹⁰ and gastric cancer⁴⁸. This study provided results that expression types of *DACH1* and *CD44* were opposite in breast cancer. Overexpression of *DACH1* reduced the levels of CSC and EMT markers in breast cancer cell line and potentially inhibited the ability of tumorigenesis in xenograft model. Besides, correlation analysis exhibited that *DACH1* and *CD44* behaved completely differently in the correlations with *FN1*, *VIM*, *YBX1*, *FOXA1*, *EGFR* and *MKI67*. Importantly, *DACH1* serves as a protective factor, while *CD44* is an unfavorable element for the prognosis of breast cancer patients. Thus, we concluded that *DACH1* might exert inhibitory effects on the development of breast cancer partly by suppression of EMT inducers and CSCs markers, especially *CD44*.

Carcinogenesis may derive from the acquisition of a plethora of oncogenic mutations, sequential suppression of endogenous growth inhibitors⁴⁹ and the loss control over pivotal cellular functions⁵⁰. *DACH1* accounts for the carcinogenesis of various tumor types, including human breast cancer⁵¹. Previous studies have implicated that *DACH1* negatively regulated cellular proliferation and tumor growth by repressing cell cycle protein cyclin D1 in both breast cancer¹¹ and renal clear cell cancer¹⁶. Restoration of *DACH1* suppressed clone formation of renal clear cell cancer cells *in vitro* as well as tumor growth *in vivo* through the inhibition of cyclin D1 transcription via associating with AP-1 protein¹⁶. Furthermore, hypermethylation of *DACH1* promoter region itself led to carcinogenesis and it also exerted inhibitory effects on tumor initiation through activating transforming growth factor-beta (TGF-beta) signaling¹⁵. Besides, *DACH1* inhibited cellular growth in an NAD and p53-dependent manner by direct protein-protein association in breast cancer⁵². On the contrary, *CD44*, a family of transmembrane glycoproteins, exerted promoting roles in tumorigenesis of various cancer types²⁰. Up-regulation of the cleaved intracellular domain of *CD44* (*CD44ICD*) enriched the mammosphere formation, while the blockage of *CD44ICD* reversed the effects⁵³. Immunohistochemistry analysis on human breast cancer tissues and expression analysis of GSE42568 indicated that *CD44* remarkably increased in breast carcinoma in comparison with normal breast tissues²². *CD44* functions in carcinogenesis through binding to extracellular matrix components and messenger molecules in tumor environment⁵⁴.




Figure 7. *DACH1* and *CD44* were related to clinical outcomes of breast cancer patients. Meta-analysis of a total of 19 public databases showed that breast cancer patients with higher *DACH1* mRNA level tended to have better overall survival (a), relapse-free survival (c), metastasis-free survival (e). On the contrary, patients with higher *CD44* mRNA expression had shorter time to death (b) and metastasis (f), but did not acquire statistically significant worse relapse-free survival in comparison to patients with comparatively lower *CD44* expression (d).

Several lines of evidence indicated that *DACH1* expression correlated with tumor differentiation. Immunohistochemistry analysis of clear cell renal cancer tissues displayed that *DACH1* was inversely correlated with tumor grade¹⁶. About 60% of cells in low-grade tumors expressed *DACH1*, and less than 20% cells in grade III tumors expressed *DACH1*¹⁶. Similar phenomenon was also found in breast cancer¹³. *DACH1* was remarkably enriched in low-grade breast tumors¹³. In contrast, *CD44* positively correlated with breast tumor grade^{21,22}. Our previous study displayed a positive association between *CD44* expression and breast tumor grade at both mRNA and protein levels^{21,22}. This paper further supported this opposite tendency of *DACH1* and *CD44* in low-grade and high-grade breast carcinoma, respectively.

Majorly according to the status of ER, PR and Her2, breast cancer is grouped into four distinct subtypes including luminal, Her2-positive, basal-type and normal-like. Among these, luminal tumor patients accounted for the most part of breast carcinoma population and have relatively good clinical outcomes, while patients

with basal-like tumor endowed with malignancy features have significantly poor prognosis. Previous study has demonstrated that nuclear DACH1 is a biomarker of luminal breast cancer¹³. Breast tumors with positive ER-alpha and co-expressing PR-alpha were most likely to highly express DACH1, and nuclear DACH1 expression was positively correlated with luminal marker FOXA1 and inversely associated with basal-like markers EGFR¹³. In contrast, our previous meta-analysis showed that *CD44* mRNA was remarkably enriched in basal-like breast cancer compared with luminal-type breast tumor²¹. *CD44* was also positively correlated with basal markers *EGFR*, *KRT5* and *KRT17*, and inversely associated with luminal marker *FOXA1*²². Our results supported that luminal breast neoplasm tended to be with high DACH1 expression and low CD44 level, while basal-like tumors were most likely to be the inverse type. The correlation analysis further indicated that DACH1 was significantly inversely associated with CD44.

CSCs were composed of a subset of tumor cells with the expression of stem cell-associated markers and enhanced capacity for tumorigenesis, metastasis and therapy-resistance⁵⁵. It has been implicated that endogenous DACH1 participated in the negative regulation of CSCs¹⁸. DACH1 suppressed the expression of stem cell markers SOX2, Nanog and KLF4 through binding to the promoters of these genes¹⁸. On the contrary, CD44, as a well-known CSCs marker, positively monitored the levels of CSCs markers²⁰. Nuclear location of cleaved CD44 intracellular domain transcriptionally activated stemness factors Nanog, Sox2 and Oct4⁵³.

EMT is a complex and highly conserved process which enhances cellular invasiveness, being critical for metastasis of various solid tumors and considered to be promising therapeutic targets⁵⁶. EMT was dysregulated by a complex network during tumor development. Knock-down of DACH1 in breast cancer cells MCF-7 and T47D promoted the morphology change from epithelial phenotype to mesenchymal pattern and interfered with cell-cell contact, accompanied by down-regulation of epithelial marker E-cadherin, resulting in cell migration and invasion¹⁰. DACH1 also transcriptionally suppressed the activity of Snail, leading to the activation of E-cadherin in breast cancer cells¹⁰, but the complex of DACH1 and Snail could bind to the E-box of E-cadherin promoter then contributing to the reduction of E-cadherin¹⁰. In addition, DACH1 reduced the expression of the mesenchymal marker Snail through suppressing the activity of the Y box-binding protein, an important EMT inducer⁹. Besides, previous study has revealed that DACH1 decreased both in breast cancer cell lines and tissues accompanied by relatively high proportion of CSCs¹⁸. Inversely, CD44 not only promoted EMT, but also was upregulated by some mesenchymal markers²⁰. Previous studies have demonstrated that mesenchymal genes including *ZEB1*, *TWIST1*, *SNAI1* and *SLUG* were positively correlated with CD44 expression²⁰. The switch from CD44 variant isoforms to its standard isoform is essential to undergo EMT for normal epithelial cells⁵⁷.

DACH1 played important roles in invasion and migration of various neoplasms, such as lung adenocarcinoma¹⁴, gastric cancer⁴⁸, pancreatic cancer⁴⁷ and breast cancer¹². Several lines of evidence showed that DACH1 is absent or suppressed in poor prognosis breast cancer⁵⁸. Breast cancer patients with reduced DACH1 expression had three years shorter time to death in comparison to those with normal DACH1 levels¹¹. Immunohistochemistry analysis showed that higher nuclear level of DACH1 was predictive of longer disease-free interval, cancer relapse-free survival and distant metastasis-free survival over 5 years post diagnosis¹³. In consistency, our analysis of GSE databases showed that higher mRNA level of DACH1 was parallel with better OS, RFS and MFS. The protected effect of DACH1 in the prognosis of breast cancer patients could be explained partly by negatively interplaying CSCs and EMT^{10,59}. Besides, DACH1 inhibited breast cancer migration and invasion also via suppressing oncogene function through targeting interleukin-8¹². Altogether, these studies suggested that DACH1 could be a valuable molecular marker for prognosis, thereby detection of DACH1 level is useful for therapeutic stratification of breast cancer patients. In contrast, CD44 contributed to tumor formation and progression predictive of poor clinical outcome²⁰. CD44 functioned as an oncoprotein and knockdown of CD44 remarkably attenuated the migration and invasion of breast cancer cells MDA-MB-231 and Hs578T by modulating c-Src transcription⁶⁰. According to the analysis of 448 primary breast tumors, CD44 was parallel with enhanced distant recurrence and decreased disease-free survival⁶¹.

In this study, 19 public datasets were enrolled to assess the correlation between the mRNA levels of *DACH1* and *CD44* and the survival performance of breast cancer patients. However, there are still some limitations: 1) the overall sample size is limited. Some data were not available when the meta-analysis was conducted; 2) the platforms used to evaluate the mRNA expression of *DACH1* and *CD44* are different; 3) the results at the mRNA levels might not be consistent with those at protein abundance because there is a complex regulation network from mRNA to protein.

In conclusion, this study confirmed that DACH1 and CD44 inversely related in breast cancer, different grade tumors and different subtypes. DACH1 was negatively associated with CD44 *in vitro* and *in vivo*. DACH1 served as a good prognostic marker, and CD44 was an unfavorable element of breast cancer patients. Thus, we concluded that CD44 might be a novel target of DACH1.

Materials and Methods

Immunohistochemical staining. To evaluate the expression of DACH1 and CD44 in normal breast versus breast tumor tissues, grade 1–2 versus grade 3 tissues as well as luminal-type and basal-like human breast carcinoma tissues, two commercially available tissue microarray (TMA) slides (BR1502–97 and BR1502–98, US Biomax, Inc, Rockville, MD) containing histologically confirmed tissues were purchased for immunohistochemistry (IHC) analysis. In addition, IHC was also employed to compare the expression of CD44, Fibronectin, Vimentin, Sox2, Myc, EGFR and Ki-67 in Met-1 nude mice xenograft tumors with overexpression of DACH1 and the controls. The Specific primary antibodies against DACH1 (10914-1-AP, ProteinTech Group), CD44 (15675-1-AP, ProteinTech Group), Fibronectin (sc-8422, Santa Cruze), Vimentin (5741, Cell Signaling Technology), Sox2 (AB5603, Millipore), Myc (sc-40, Santa Cruze), EGFR (sc-03, Santa Cruze) and Ki-67 (ab15680, Abcam) were utilized for IHC with a 2-step protocol⁶².

Analysis and quantification of staining. For quantification, a total of six 200× magnifications of each kind of protein were selected for IHC scoring. Two experienced pathologists assessed the immunohistochemical score independently. Scores were calculated on intensity and proportion of positive staining tumor cells in the whole tissue stains according to the Fromowitz Standard as described above⁶³. The staining intensity was scored as 0 (no staining), 1 (weak staining, light yellow), 2 (moderate staining, yellow brown) and 3 (strong staining, brown). The proportions of stained tumor cells were classified as 1 (0–25% positive cells), 2 (26–50% positive cells), 2 (51–75% positive cells) and 3 (76–100% positive cells). The multiplication for intensity and proportion was utilized to represent the protein levels of DACH1, CD44, Myc, Sox2, Fibronectin, Vimentin, EGFR and Ki-67.

Cell culture. The breast cancer cell line Met-1, was cultured in high-glucose Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS, Life Technologies, Inc.). All cells were grown 37 °C in a humidified incubator with 5% CO₂.

Western blot. Cells were washed with cold PBS, scraped into RIPA buffer and centrifuged. The cell lysates were subjected to 10% SDS-PAGE and transferred to a polyvinylidene fluoride (PVDF) hybridization transfer membrane. The primary antibodies used were as follows: HA (Sc-7392, Santa Cruze), CD44 (15675-1-AP, ProteinTech group), Fibronectin (sc-8422, Santa Cruze), Vimentin (5741, Cell Signaling Technology), p21cip1 (sc-6246, Santa Cruze), p27kip2 (sc-1641, Santa Cruze) and β -actin (Sc-47778, Santa Cruze). Secondary staining and detection were carried out in accordance with standard protocols^{16,64}.

RNA profiling by microarray. DNA-free total RNA isolated from Met-1 cells expressing GFP or *DACH1* were used to probe Affymetrix Gene 1.0 arrays (Affymetrix, Santa Clara, CA). RNA quality was determined by gel electrophoresis. Analysis of the arrays was performed using GeneSpring. Arrays were normalized using robust multi-array analysis, and the *p* value of 0.05 was applied as a statistical criterion for differentially expressed genes.

Meta-analysis for DACH1 on published Gene Expression Omnibus (GEO) databases. We carried out a comprehensive search of relevant GEO databases for mRNA expression of *DACH1* and *CD44* through ArrayExpress and Oncomine. The datasets meeting the following criteria were included: 1) the datasets were about human breast cancer; 2) the mRNA expression of *DACH1* and *CD44* was measured in these databases; 3) clinical outcomes of patients were showed in these databases; 4) the sample capacity was more than 50. Only the latest and most complete datasets were included when several databases shared common patients. At last, a total of 19 independent human breast cancer microarray databases with the mRNA expression of *DACH1* and *CD44* and required the survival information of breast cancer patients were enrolled in this systematic analysis.

Cutoff value for *DACH1* and *CD44* was median expression. OS, RFS and MFS were evaluated by Cox proportional hazard ratio (HR) and 95% confidence interval (95% CI). HRs was employed to assess the survival outcome of breast cancer patients with high mRNA expression of *DACH1* and *CD44* and HR >1 indicated that high expression of *DACH1* and *CD44* predicted worse survival of patients. The random-effect model was employed when heterogeneity was present, and the fixed-effect model was used when homogeneity was demonstrated. Heterogeneity of publication was evaluated by means of the Chi-square-based-Q statistic and inconsistency index (I²) statistic. Statistical analysis was performed based on the guidelines of Meta-Analysis of Observational Studies. The STATA software package (version 12.0) (Stata Corp LP, College Station, TX, USA) was employed to perform the meta-analysis.

Analysis of gene expression data. GSE20711, available through GEO databases and containing 45 luminal-type and 22 basal-like breast carcinoma cases, was analyzed to evaluate the mRNA expression of *DACH1* and *CD44* in basal-like carcinoma in comparison with that in luminal-type tumors.

GSE20685, containing a total of 327 patients of breast cancer, was interrogated to assess the correlation between *DACH1* and *CD44*, *FN1*, *VIM*, *FOXA1*, *EGFR*, *MKI67* as well as the correlation between *CD44* and *FN1*, *VIM*, *YBX1*, *FOXA1*, *EGFR* and *MKI67* in breast cancer tissues.

Breast cancer cell line data of 51 breast cancer cell lines of different molecular subtypes published by Neve RM, *et al.*²⁵ were also employed to evaluate whether the correlation in breast cancer cells was consistent with that in breast tumor tissues.

Nude Mice Study. The animal protocols were approved by the ethics committee of the Tongji Hospital of Huazhong University of Science and Technology. The methods used in this section were in accordance with the relevant guidelines and regulations. 1 × 10⁵ Met-1 cells expressing GFP control or *DACH1* were implanted subcutaneously into 4–6-week-old athymic female nude mice purchased from Wuhan Hamilton Biological Polytron Technologies Inc. The tumor growth was measured twice weekly for 3 weeks by using a digital caliper. Tumor weight was measured when mice were sacrificed on day 27 after cells implantation.

Statistical analysis. Correlation analyses were performed using SPSS 20 statistical software (SPSS Inc., Chicago, IL, USA). The Student's t-test was applied to evaluate the differences in groups as appropriate and the significance level was set at 0.05. The correlation analysis was evaluated by a χ^2 -test. A two-tailed *p* value < 0.05 was considered statistically significant.

References

1. Siegel, R. L., Miller, K. D. & Jemal, A. Cancer statistics, 2016. *CA Cancer J. Clin.* **66**, 7–30 (2016).
2. Xu, H. *et al.* Recent advances of highly selective CDK4/6 inhibitors in breast cancer. *J. Hematol. Oncol.* **10**, 97 (2017).
3. Hu, X., Huang, W. & Fan, M. Emerging therapies for breast cancer. *J. Hematol. Oncol.* **10**, 98 (2017).
4. Kong, D. *et al.* The retinal determination gene network: from developmental regulator to cancer therapeutic target. *Oncotarget* **7**, 50755–50765 (2016).

5. Liu, Y. *et al.* The DACH/EYA/SIX gene network and its role in tumor initiation and progression. *Int. J. Cancer* **138**, 1067–1075 (2016).
6. Xu, H. X. *et al.* Expression profile of SIX family members correlates with clinic-pathological features and prognosis of breast cancer: A systematic review and meta-analysis. *Medicine (Baltimore)* **95**, e4085 (2016).
7. Wu, K. *et al.* EYA1 phosphatase function is essential to drive breast cancer cell proliferation through cyclin D1. *Cancer Res.* **73**, 4488–4499 (2013).
8. Liu, Q. *et al.* The expression profile and clinic significance of the SIX family in non-small cell lung cancer. *J. Hematol. Oncol.* **9**, 119 (2016).
9. Wu, K. *et al.* Cell fate factor DACH1 represses YB-1-mediated oncogenic transcription and translation. *Cancer Res.* **74**, 829–839 (2014).
10. Zhao, F. *et al.* DACH1 inhibits SNAI1-mediated epithelial-mesenchymal transition and represses breast carcinoma metastasis. *Oncogenesis* **4**, e143 (2015).
11. Wu, K. *et al.* DACH1 is a cell fate determination factor that inhibits cyclin D1 and breast tumor growth. *Mol. Cell Biol.* **26**, 7116–7129 (2006).
12. Wu, K. *et al.* Dachshund inhibits oncogene-induced breast cancer cellular migration and invasion through suppression of interleukin-8. *Proc. Natl. Acad. Sci. USA* **105**, 6924–6929 (2008).
13. Powe, D. G. *et al.* DACH1: its role as a classifier of long term good prognosis in luminal breast cancer. *PLoS One* **9**, e84428 (2014).
14. Han, N. *et al.* DACH1 inhibits lung adenocarcinoma invasion and tumor growth by repressing CXCL5 signaling. *Oncotarget* **6**, 5877–5888 (2015).
15. Wu, L. *et al.* Silencing DACH1 promotes esophageal cancer growth by inhibiting TGF-beta signaling. *PLoS One* **9**, e95509 (2014).
16. Chu, Q. *et al.* DACH1 inhibits cyclin D1 expression, cellular proliferation and tumor growth of renal cancer cells. *J. Hematol. Oncol.* **7**, 73 (2014).
17. Liu, Y. *et al.* DACH1 is a novel predictive and prognostic biomarker in hepatocellular carcinoma as a negative regulator of Wnt/beta-catenin signaling. *Oncotarget* **6**, 8621–8634 (2015).
18. Wu, K. *et al.* Cell fate determination factor Dachshund reprograms breast cancer stem cell function. *J. Biol. Chem.* **286**, 2132–2142 (2011).
19. Schulenburg, A. *et al.* Cancer stem cells in basic science and in translational oncology: can we translate into clinical application? *J. Hematol. Oncol.* **8**, 16 (2015).
20. Xu, H. *et al.* The role of CD44 in epithelial-mesenchymal transition and cancer development. *Onco. Targets Ther* **8**, 3783–3792 (2015).
21. Xu, H. *et al.* Enrichment of CD44 in basal-type breast cancer correlates with EMT, cancer stem cell gene profile, and prognosis. *Onco. Targets Ther* **9**, 431–444 (2016).
22. Xu, H. *et al.* CD44 correlates with clinicopathological characteristics and is upregulated by EGFR in breast cancer. *Int. J. Oncol.* **49**, 1343–1350 (2016).
23. Orian-Rousseau, V. CD44 Acts as a Signaling Platform Controlling Tumor Progression and Metastasis. *Front Immunol* **6**, 154 (2015).
24. Zoller, M. CD44: can a cancer-initiating cell profit from an abundantly expressed molecule? *Nat. Rev. Cancer* **11**, 254–267 (2011).
25. Neve, R. M. *et al.* A collection of breast cancer cell lines for the study of functionally distinct cancer subtypes. *Cancer Cell* **10**, 515–527 (2006).
26. Desmedt, C. *et al.* Strong time dependence of the 76-gene prognostic signature for node-negative breast cancer patients in the TRANSBIG multicenter independent validation series. *Clin. Cancer Res.* **13**, 3207–3214 (2007).
27. Pawitan, Y. *et al.* Gene expression profiling spares early breast cancer patients from adjuvant therapy: derived and validated in two population-based cohorts. *Breast Cancer Res* **7**, R953–964 (2005).
28. Hennessy, B. T. *et al.* Characterization of a naturally occurring breast cancer subset enriched in epithelial-to-mesenchymal transition and stem cell characteristics. *Cancer Res.* **69**, 4116–4124 (2009).
29. Dedeurwaerder, S. *et al.* DNA methylation profiling reveals a predominant immune component in breast cancers. *EMBO Mol. Med.* **3**, 726–741 (2011).
30. Kao, K. J., Chang, K. M., Hsu, H. C. & Huang, A. T. Correlation of microarray-based breast cancer molecular subtypes and clinical outcomes: implications for treatment optimization. *BMC Cancer* **11**, 143 (2011).
31. Clarke, C. *et al.* Correlating transcriptional networks to breast cancer survival: a large-scale coexpression analysis. *Carcinogenesis* **34**, 2300–2308 (2013).
32. Desmedt, C. *et al.* Multifactorial approach to predicting resistance to anthracyclines. *J. Clin. Oncol.* **29**, 1578–1586 (2011).
33. Loi, S. *et al.* PIK3CA mutations associated with gene signature of low mTORC1 signaling and better outcomes in estrogen receptor-positive breast cancer. *Proc. Natl. Acad. Sci. USA* **107**, 10208–10213 (2010).
34. Bild, A. H. *et al.* Oncogenic pathway signatures in human cancers as a guide to targeted therapies. *Nature* **439**, 353–357 (2006).
35. Terunuma, A. *et al.* MYC-driven accumulation of 2-hydroxyglutarate is associated with breast cancer prognosis. *J. Clin. Invest.* **124**, 398–412 (2014).
36. Heikkinen, T. *et al.* Variants on the promoter region of PTEN affect breast cancer progression and patient survival. *Breast Cancer Res* **13**, R130 (2011).
37. Hatzis, C. *et al.* A genomic predictor of response and survival following taxane-anthracycline chemotherapy for invasive breast cancer. *Jama* **305**, 1873–1881 (2011).
38. Symmans, W. F. *et al.* Genomic index of sensitivity to endocrine therapy for breast cancer. *J. Clin. Oncol.* **28**, 4111–4119 (2010).
39. Wang, Y. *et al.* Gene-expression profiles to predict distant metastasis of lymph-node-negative primary breast cancer. *Lancet* **365**, 671–679 (2005).
40. Tofigh, A. *et al.* The prognostic ease and difficulty of invasive breast carcinoma. *Cell Rep* **9**, 129–142 (2014).
41. Minn, A. J. *et al.* Genes that mediate breast cancer metastasis to lung. *Nature* **436**, 518–524 (2005).
42. Minn, A. J. *et al.* Lung metastasis genes couple breast tumor size and metastatic spread. *Proc. Natl. Acad. Sci. USA* **104**, 6740–6745 (2007).
43. Sircoulomb, F. *et al.* Genome profiling of ERBB2-amplified breast cancers. *BMC Cancer* **10**, 539 (2010).
44. Nagalla, S. *et al.* Interactions between immunity, proliferation and molecular subtype in breast cancer prognosis. *Genome Biol.* **14**, R34 (2013).
45. Davis, R. J. *et al.* Dach1 mutant mice bear no gross abnormalities in eye, limb, and brain development and exhibit postnatal lethality. *Mol. Cell Biol.* **21**, 1484–1490 (2001).
46. Heanue, T. A. *et al.* Dach1, a vertebrate homologue of Drosophila dachshund, is expressed in the developing eye and ear of both chick and mouse and is regulated independently of Pax and Eya genes. *Mech. Dev.* **111**, 75–87 (2002).
47. Bu, X. N., Qiu, C., Wang, C. & Jiang, Z. Inhibition of DACH1 activity by short hairpin RNA represses cell proliferation and tumor invasion in pancreatic cancer. *Oncol. Rep.* **36**, 745–754 (2016).
48. Yan, W. *et al.* Epigenetic silencing of DACH1 induces the invasion and metastasis of gastric cancer by activating TGF-beta signalling. *J. Cell Mol. Med.* **18**, 2499–2511 (2014).
49. Hanahan, D. & Weinberg, R. A. The hallmarks of cancer. *Cell* **100**, 57–70 (2000).
50. Chen, Y. *et al.* Identifying potential cancer driver genes by genomic data integration. *Sci. Rep* **3**, 3538 (2013).

51. Popov, V. M. *et al.* The Dachshund gene in development and hormone-responsive tumorigenesis. *Trends Endocrinol. Metab.* **21**, 41–49 (2010).
52. Chen, K. *et al.* Acetylation of the cell-fate factor dachshund determines p53 binding and signaling modules in breast cancer. *Oncotarget* **4**, 923–935 (2013).
53. Cho, Y. *et al.* Cleaved CD44 intracellular domain supports activation of stemness factors and promotes tumorigenesis of breast cancer. *Oncotarget* **6**, 8709–8721 (2015).
54. Morath, I., Hartmann, T. N. & Orian-Rousseau, V. CD44: More than a mere stem cell marker. *Int. J. Biochem. Cell Biol.* **81**, 166–173 (2016).
55. Velasco-Velazquez, M. A., Homsí, N., De La Fuente, M. & Pestell, R. G. Breast cancer stem cells. *Int. J. Biochem. Cell Biol.* **44**, 573–577 (2012).
56. Jayachandran, A., Dhungel, B. & Steel, J. C. Epithelial-to-mesenchymal plasticity of cancer stem cells: therapeutic targets in hepatocellular carcinoma. *J. Hematol. Oncol.* **9**, 74 (2016).
57. Klingbeil, P. & Isacke, C. M. The 'alternative' EMT switch. *Breast Cancer Res* **13**, 313 (2011).
58. Wu, K. *et al.* Cell fate determination factor DACH1 inhibits c-Jun-induced contact-independent growth. *Mol. Biol. Cell* **18**, 755–767 (2007).
59. Wu, K. *et al.* DACH1 inhibits transforming growth factor-beta signaling through binding Smad4. *J. Biol. Chem.* **278**, 51673–51684 (2003).
60. Nam, K., Oh, S., Lee, K. M., Yoo, S. A. & Shin, I. CD44 regulates cell proliferation, migration, and invasion via modulation of c-Src transcription in human breast cancer cells. *Cell. Signal.* **27**, 1882–1894 (2015).
61. McFarlane, S. *et al.* CD44 increases the efficiency of distant metastasis of breast cancer. *Oncotarget* **6**, 11465–11476 (2015).
62. Liu, Q. Q. *et al.* Decreased DACH1 expression in glomerulopathy is associated with disease progression and severity. *Oncotarget* (2016).
63. Fromowitz, F. B. *et al.* ras p21 expression in the progression of breast cancer. *Hum. Pathol.* **18**, 1268–1275 (1987).
64. Tian, Y. *et al.* Modification of platinum sensitivity by KEAP1/NRF2 signals in non-small cell lung cancer. *J. Hematol. Oncol.* **9**, 83 (2016).

Acknowledgements

This work was supported by the National Natural Science Foundation of China (NSFC) No. 81572608 and 81172422 (KW), 81502296 (JX), and was also supported by NIH grant R01CA132115 (RGP).

Author Contributions

H.X.X. collected and analyzed the data, did experiments and wrote the manuscript. S.N.Y., X.Y., J.X., and D.K. carried out immunohistochemical analysis. K.M.W. and R.G.P. designed the study and revised the manuscript. All authors edited and approved the final manuscript.

Additional Information

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017