#### University of Massachusetts Medical School

## eScholarship@UMMS

UMass Center for Clinical and Translational Science Research Retreat

2017 UMass Center for Clinical and Translational Science Research Retreat

May 16th, 1:45 PM

Variable Knowledge, Use and Perceptions of the IMPACT Model among Physicians during Prognostication Meetings for Criticallyill Traumatic Brain Injury Patients – Results from a Qualitative Study

Jesse Moskowitz University of Massachusetts Medical School

Et al.

# Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts\_retreat

Part of the Health Services Administration Commons, Nervous System Diseases Commons, Neurology Commons, and the Translational Medical Research Commons

Moskowitz J, Quinn T, Khan M, Shutter L, Goldberg RJ, Col N, Mazor KM, Muehlschlegel S. (2017). Variable Knowledge, Use and Perceptions of the IMPACT Model among Physicians during Prognostication Meetings for Critically-ill Traumatic Brain Injury Patients – Results from a Qualitative Study. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from https://escholarship.umassmed.edu/cts\_retreat/2017/posters/55

Creative Commons License

This work is licensed under a Creative Commons Attribution-Noncommercial-Share Alike 3.0 License. This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

### VARIABLE KNOWLEDGE, USE AND PERCEPTIONS OF THE IMPACT MODEL AMONG PHYSICIANS DURING PROGNOSTICATION MEETINGS FOR CRITICALLY-ILL TRAUMATIC BRAIN INJURY PATIENTS – RESULTS FROM A QUALITATIVE STUDY

Jesse Moskowitz, BS<sup>1</sup>;Thomas Quinn, BS<sup>1</sup>; Muhammad W. Khan, MBBS<sup>1</sup>, Lori Shutter, MD<sup>2</sup>; Robert Goldberg, PhD<sup>3</sup>; Nananda Col, MD, MPP, MPH<sup>4</sup>; Kathleen M. Mazor, EdD<sup>5,6</sup>; Susanne Muehlschlegel, MD,MPH<sup>1,7,8</sup>

<sup>1</sup>Departments of Neurology (Neurocritical Care), University of Massachusetts Medical School; <sup>2</sup>Departments of Critical Care Medicine and Neurology, University of Pittsburgh School of Medicine, Pittsburgh, PA; <sup>3</sup>Department of Quantitative Health Sciences, University of Massachusetts Medical School; <sup>4</sup>Shared Decision Making Resources, Georgetown, ME; <sup>5</sup>Meyers Primary Care Institute, <sup>6</sup>Internal Medicine, <sup>7</sup>Anesthesiology/Critical Care and <sup>8</sup>Surgery, University of Massachusetts Medical School

**Introduction:** The International Patient Decision Aid Standards, a framework for the creation of high-quality decision aids, calls for the presentation of probabilities. To inform the content of a goals-of-care decision aid in critically-ill TBI (ciTBI) patients, we examined physician's awareness, perceptions, and use of the IMPACT-model, the most widely validated ciTBI outcome model, and explored their preferences for communicating prognostic information towards families.

**Methods:** We conducted a qualitative study using semi-structured interviews in 20 attending physicians (neurocritical care, neurosurgery, trauma, palliative care) at 7 U.S. academic medical centers. We used descriptive statistics and performed qualitative content analysis of transcribed interviews to identify major themes.

**Results:** Only 12 physicians (60%) expressed awareness of the IMPACT-model; two stated that they "barely" knew the model. Seven physicians indicated using the model at least some of the time in clinical practice, although none used it exclusively to derive a patient's prognosis. Four major themes emerged: the IMPACT-model is intended for research but should not be applied to individual patients; mistrust in the IMPACT-model derivation data; the IMPACT-model is helpful in reducing prognostic variability among physicians; concern that statistical models may mislead families about a patient's prognosis.

**Conclusions**: We identified substantial variability in the awareness of, use, and attitude toward the IMPACT model among physicians. Understanding knowledge, use, and barriers to using existing models, like IMPACT, is vital for creating and implementing a meaningful shared decision-making tool to improve goals-of-care discussions.

### Contact:

Jesse Moskowitz University of Massachusetts Medical School MD Candidate ~ Class of 2018 Jesse.moskowitz@umassmed.edu