


# Surficial Geology of Black Hawk County, Iowa



## LEGEND

### CENOZOIC

#### QUATERNARY SYSTEM

##### Hudson Episode

- Qal** - Alluvium (Deforest Formation) (alluvial fan) Variable thickness of less than 1 to 5 m (3-16 ft) of very dark gray to brown, noncalcareous to calcareous, massive to stratified silty clay loam, clay loam, loam to sandy loam alluvium and colluvium in stream valleys, on hillslopes and in closed depressions. May overlie Noah Creek Formation, Wolf Creek or Alburtus Formation, or bedrock. Associated with low-level modern floodplains, closed depressions, modern drainage swales or terrace positions on the landscape. Seasonal high water table and potential for frequent flooding.
- Qalb** - Alluvium Shallow to Bedrock (Deforest Formation) (alluvial fan) Variable thickness of less than 1 to 5 m (3-16 ft) of very dark gray to brown, noncalcareous to calcareous, stratified silty clay loam, clay loam, loam to sandy loam alluvium and colluvium in stream valleys, on hillslopes and in closed depressions. May overlie Noah Creek Formation or Devonian carbonate bedrock. Bedrock surface is within 5 m (16 ft) of the land surface. Associated with low-level modern floodplains, closed depressions, modern drainage swales or terrace positions on the landscape. Seasonal high water table and potential for frequent flooding.
- Qall** - Low Terrace (Deforest Formation) (Camp Creek Mts. and Roberts Creek Mts.) Variable thickness of less than 1 to 5 m (3-16 ft) of very dark gray to brown, noncalcareous, stratified silty clay loam, clay loam, or clay loam. A associated with the modern terrace belt of the Cedar and Wapaisippian river valleys and their tributaries. Overlies the Noah Creek Formation. Occurs on low position on the floodplain, in modern channel belts. Seasonal high water table and frequent flooding potential.
- Qall-ht** - Intermediate-High Terrace (Deforest Formation) (Gander Mts.) Variable thickness of less than 1 to 5 m (3-16 ft) of very dark gray to brown, noncalcareous, silty clay loam to loam alluvium or colluvium that overlie the Noah Creek Formation. Occurs on terrace and valley margin positions. 1 to 2 m (3-7 ft) above the modern floodplain. May be associated with 2 to 3 m (7-10 ft) of well-sorted medium to fine sand derived from wind working of the alluvium. Seasonal high water table and low to moderate flooding potential.

##### Hudson and Wisconsin Episode

- Qe** - Sand Dunes and Sand Sheets (Poria Formation) (sand sheet) Generally less than 3 m (10 ft) of yellowish brown, massive, calcareous loam sandy to fine sand. It may overlie yellowish-brown sand and gravel (Noah Creek Formation) or reworked unsorted loamy sediments associated with the low an Erosion Surface and/or it may overlie yellowish-brown loam, often calcareous and fractured clay by loam from the Wolf Creek or Alburtus Formation.
- Qnw2** - Sand and Gravel (Noah Creek Formation) (Noah Creek Formation) Generally 2 to 8 m (6-26 ft) of yellowish brown to gray, poorly to well-sorted, massive to well stratified, coarse to fine (clay) quartz sand, pebbly sand and gravel with fine interlocking layers of silty clay. A thin mark of loam, overlaid loam or fine-grained siltstone may be present. This unit includes only occasional deposits derived from the adjacent map units. In places this unit is marked with the 3 m (10 ft) of well-sorted medium to fine sand derived from wind working of the loam. This unit encompasses deposits that accumulated as low-level stream valley during the Wisconsin and Hudson episodes. Seasonal high water table and some potential for flooding.

##### Wisconsin Episode

- Qpt** - Low Mounded Terrace (Poria Formation) (Poria Formation) (Poria Formation) Generally 2 to 8 m (6-26 ft) of yellowish brown to gray, massive, jointed, calcareous to noncalcareous, silty loam and interbedded fine to medium, well sorted, sand. May grade downward and to poorly to moderately well sorted, moderately to well stratified, coarse to fine (clay) quartz sand, loam, or silty loam alluvium (Late Phase High Terrace) or may overlie a Farnham Geosol developed in Phase 50a which in turn overlies a well-sorted Sangamon Geosol developed in Phase 50b (generally well-sorted, moderately well stratified, coarse to fine sand, loam, or silty loam alluvium) (Late Phase High Terrace).
- Qps1** - Loamy and Low Mounded Terrace (Poria Formation) (Poria Formation) Generally 2 to 8 m (6-26 ft) of yellowish brown to gray, massive, fractured, calcareous, fine to medium, well sorted, calcareous, silty loam and interbedded fine to medium, well sorted, sand. Sand is most abundant in the lower part of the terrace package. Overlies massive, fractured, loamy glacial till of the Wolf Creek or Alburtus Formation with or without inter-lying Farnham-Sangamon Geosol.
- Qnw** - Sand and Gravel (Noah Creek Formation) (Noah Creek Formation) 3 to 10 m (10-30 ft) more than 20 m (66 ft) of yellowish brown to gray, poorly to well-sorted, massive to well stratified, coarse to fine (clay) quartz sand, pebbly sand and gravel. In places marked with fine to medium well-sorted calcareous siltstone and shale derived from wind working of the alluvium. Fractured calcareous bedrock is less than 6 m (20 ft) below the land surface. The unit encompasses deposits that accumulated in river and stream valley during the Wisconsin and Hudson episodes of the Wolf Creek and Alburtus Formations. Deposits may be slightly thicker along the Cedar River.
- Qnw3** - Sand and Gravel Shallow to Bedrock (Noah Creek Formation) (Noah Creek Formation) 1 to 6 m (3-20 ft) of yellowish brown to gray, poorly to well-sorted, coarse to fine (clay) quartz sand, pebbly sand and gravel. In places marked with fine to medium well-sorted calcareous siltstone and shale derived from wind working of the alluvium. Fractured calcareous bedrock is less than 6 m (20 ft) below the land surface. The unit encompasses deposits that accumulated in river and stream valley during the Wisconsin and Hudson episodes of the Wolf Creek and Alburtus Formations. Deposits may be slightly thicker along the Cedar River.

##### Wisconsin Episode

- Qwa2** - Loamy and Sandy Sediment Shallow to Bedrock (Unnamed erosion surface) (Unnamed erosion surface) Generally 2 to 8 m (6-26 ft) of yellowish brown to gray, massive to well sorted, well to poorly sorted loam, sandy to silty loam surface sediment. Map unit includes some areas marked with less than 3 m (10 ft) of Poria Formation sand facies (loam sand). Eolian sand may be directly on top of bedrock in isolated areas. Overlies Farnham-Sangamon or bedrock rocks. Seasonal high water table may occur in this map unit.

##### Pre-Illinois Episode

- Qwa3** - Till (Wolf Creek or Alburtus Formation) (Wolf Creek or Alburtus Formation) Generally 3 to 7 m (10-24 ft) of very dense, massive, fractured, loamy glacial till of the Wolf Creek or Alburtus Formations with or without thin beds of sand (Poria Formation) (sand sheet) or thin loamy siltstone (Poria Formation) (sand sheet) or thin loamy siltstone (Poria Formation) (sand sheet) or thin loamy siltstone (Poria Formation) (sand sheet). This mapping unit is shown only in the cross-section and may be bounded by unnamed erosion surface collines, loam or alluvium.

##### Other Mapping Units

- Qpq** - Pits and Quarries Sand and gravel pits and rock quarries. Erosion mapped as shown county survey maps as identified on aerial imagery.
- Bedrock** - All areas of bedrock outcrop regardless of the unit are shown in red on the map, without any labels. Bedrock units are shown on the cross-section with the following description:

### PALEOZOIC

#### DEVONIAN SYSTEM

- Dgc** - Dolomite, Limestone, and Shale (Hatchery City Formation) (Middle to Upper Devonian). The total thickness of this map unit is up to 30 m (97 ft), consisting of interbedded laminated lithologies and shaly lithologies. The dolomite and limestone lithology is extensive to abundant brachiopods, corals and stromatolites. This unit occurs on some bedrock highs in the western one-third of the county.
- Dcv** - Limestone and Dolomite (Corvallis Formation) (Middle Devonian). The thickness of this map unit varies between 0 and 31 m (0-100 ft) within the county. The lower Gazer Creek Member is a fossiliferous carbonate with an abundant marine fauna and is dominated by dolomite and dolomite limestone. Being slightly argillaceous in part, with common cherty beds, the low cherty beds are characterized by crinoid stems and brachiopods and rare graptolites and tree trunks. The upper Gazer Creek Member is carbonate dominated, with laminated, brecciated, or argillaceous textures and some argillaceous marls. The coralline marine fauna is dominated by favosites corals and/or branching and colonial stromatolites. Stromatolites and brachiopods occur within and around the map area.
- Dic** - Dolomite and Limestone (Little Cedar Formation) (Middle Devonian). The thickness of this map unit ranges from 0 to 40 m (0-130 ft) within the county. The map unit is dominated by slightly argillaceous to argillaceous dolomite and dolomite limestone, usually very argillaceous and partially laminated and/or cherty. This unit is commonly fossiliferous and brachiopods are abundant in the lower portion. The upper portion (Hickory Hills) is a dense, calcareous, lithology or shaly lithology limestone or dolomite limestone, with laminated, peloidal, intertactile, and bioturbate fabrics.
- Dw** - Dolomite, Limestone, Shale, and minor Sandstone (Wapaisippian Group) (Middle Devonian). This map unit contains only the Panton Field Formation of the same name, a shaly to silty limestone with minor sandstone (Panton Field Formation) (Middle Devonian). It is dominated by laminated or brecciated, calcareous limestone and dolomite that is commonly sandy and cherty at its base. This unit occurs in some deeper bedrock valleys within the map area.

#### Silurian System

- Sl** - Limestone, Dolomite Limestone and Dolomite (Lafayette City Formation) (Upper Silurian) (Lafayette City Formation) (Upper Silurian). This unit is a limestone facies that correlates with the upper Hagenlow lower Scott Grove Formations. These rocks are unconformably covered by the Wapaisippian Group. The total thickness of the map unit is up to 43 m (140 ft). The formation is dominated by dense, fossiliferous limestone that is cherty to very cherty. Secondary lithologies include dolomite, limestone and dolomite. Minor lithologies include argillaceous to shaly clay sandstone at the top of the interval (may be basal Wapaisippian Group) and green-gray shale. This unit occurs in a bedrock valley within the southeastern part of the county.
- Shb** - Shale with Chert (Hagenlow and Blanning Formations) (Lower Silurian). The total thickness of this map unit is up to 49 m (160 ft). This unit is fossiliferous to very argillaceous and cherty to very cherty with nodular to block chert in the upper part of the Blanning Formation. Fossils include corals, brachiopods, and stromatolites. The Hagenlow rocks are generally well fossiliferous and less cherty than the underlying Blanning rocks. This unit occurs in some of the deeper portions of the bedrock valleys within the map area.


#### Ordovician System

- Om** - Shale and Dolomite (Maquoketa Formation) (Upper Ordovician). The total thickness of this map unit is up to 91 m (300 ft). This unit is comprised of interbedded gray to olive dolomite shale and olive dolomite with minor limestone, variably cherty and variably fossiliferous with brachiopods and graptolites; thin lenses to brown-gray dolomite shale layers occur in the lower 10 m (33 ft). This unit occurs in the deepest portions of the bedrock valleys within the map area.


#### Drill Holes

- Wells


## CORRELATION CHART



## GEOLOGIC CROSS-SECTION A-B



## Location Map



**SURFICIAL GEOLOGY OF  
BLACK HAWK COUNTY, IOWA**

Iowa Geological and Water Survey  
Open File Map OFM-13-4  
September 2013

prepared by  
Stephanie Tassier-Surine, Deb Quade, Robert Rowden, Robert McKay, Huaiho Liu, and James Giglierao

Iowa Geological and Water Survey, Iowa City, Iowa

**DNR**  
Iowa Department of Natural Resources, Chuck Grier, Director  
Iowa Geological and Water Survey, Robert D. Linn, State Geologist

Supported in part by the U.S. Geological Survey  
Cooperative Agreement Number G12AC02026  
National Cooperative Geologic Mapping Program (STATMAP)

**ACKNOWLEDGMENTS**

Recognized for contributions to map's production: Special thanks to Sherry Lund of BMC Aggregate, and Lee Fries of Paul Neuman Construction for allowing us access to their properties. New subsurface geologic data was mostly generated by University of Iowa students Kyle Baaden and Jaime Rice who produced descriptive logs of water wells. Michael Blank and Tom Marshall of the Iowa Geological and Water Survey (IGWS) provided additional descriptive logging of water wells. Jason Vogelgesang and Carolyn Kobel (IGWS) prepared well samples for stratigraphic logging and assisted with field work during core collection. Ray Anderson and Brian Witzke (IGWS) provided valuable background information concerning the bedrock geology, geology and stratigraphy of the area. Cody Haddock and Casey Koeltz (IGWS) provided assistance with aerial imagery and GIS mapping technical help. A special thank you to landowners who graciously allowed access to their land for drilling: John Becker, Ron Dropper, William Dropper, Lonnie Egan, Mike Egan, Wayne Friden, Richard and Dorothy Grose, Gary Hanson, Ralph Knapp, Gary Mann, Dave Myers (Hatchery Farms), John and Debra Rottmush, Craig Sharp, Mike and Christy Watson, Dave Zeeb, Mike Zuck, Len Youngblood, the Cedar Falls Gun Club, Vern Fink (Black Hawk County Conservation Board) and Scott Graham (Dunkerton Sportsman Club). Geoffry Tinker and Catherine Nichols, Black Hawk County Engineers Office, allowed access to county roads for drilling and geophysics. Roger Ricker provided assistance with locating drilling sites. Drilling was provided under contract with Cahoy Well and Pump Service of Sumner, Iowa. Special thanks to Dr. Her Mark Classen. Thank you to Joshua McNary of Aerial Services, Inc. for assistance in acquiring 2008 color photos of the Cedar River valley. The imagery was courtesy of Aerial Services, Inc. (ASIS) Cedar Falls, Iowa - AerialServicesInc.com

**Introduction to the Surficial Geology of Black Hawk County**

Black Hawk County lies within the Iowan Erosion Surface (IES) Landform Region (Prior and Koeltz, 2006) in northeast Iowa. This area has been subjected to multiple periods of Quaternary glaciations and subaerial erosion. Generally speaking, the map area consists of unsorted loamy sediments (IES materials) of variable thickness overlying Pre-Illinoian glacial sediments. The Cedar and Wapaisippian river valleys are filled with Wisconsin Episode Noah Creek Formation sand and gravel and mantled with younger Holocene terrace materials. These deposits are regionally extensive.

Previous surficial geologic mapping completed as part of the STATMAP program includes the Gilbertville (Tassier-Surine et al., 2011) and Cedar Falls (Tassier-Surine et al., 2012) quadrangles in Black Hawk County and mapping to the north in adjacent Bremer County (Tassier-Surine et al., 2007, 2009, 2010). The only other regional surficial map of the area consists of the Des Moines 4° x 6° Quadrangle at a scale of 1:1,000,000 (Halberg et al., 1991).

At least seven episodes of Pre-Illinoian glaciations occurred in this region between approximately 2.2 and 0.5 million years ago (Boettler et al., 1978a,b; Halberg, 1980, 1986). Episodic erosion during the last 500,000 years has led to the destruction of pre-existing glacial landforms associated with Pre-Illinoian glaciations. A period of intense cold occurred during the Wisconsin full glacial episode from 21,000 to 16,500 years ago (Betts, 1989). This cold episode and ensuing upland erosion led to the development of the distinctive landform recognized as the IES (Prior, 1976). A periglacial environment prevailed during this period with intensive freeze-thaw action, solifluction, strong winds, and a host of other periglacial processes (Walters, 1996). Surface soils were removed from the IES and the Pre-Illinoian till surface was significantly eroded; resulting in the development of a region-wide colluvial lag deposit referred to as a "stone line." Another common feature of this region are pits, isolated and uneroded topographic highs of loess-mantled Pre-Illinoian till with a directional orientation from northwest to southeast that exist as erosional outcrops of the once higher and older landscape. These packages of stratified loamy and sandy sediments located low in the upland landscape and adjacent to streams are remnants of solifluction lobes associated with the formation of the IES. These materials can commonly be found along tributaries of the Cedar River.

Black Hawk County is covered by various Quaternary deposits with a maximum thickness of up to 73 m (240 ft) occurring in bedrock valleys. Surficial deposits of the map area are composed of five formations: Deforest, Noah Creek, Poria, Wolf Creek, and Alburtus formations as well as unnamed erosion surface sediments. Hudson age deposits associated with fine-grained alluvial and colluvial sediments are composed of the Deforest Formation which is subdivided into the Camp Creek, Roberts Creek, Gander, and Corrigton members. The Noah Creek Formation includes coarse sand and gravel associated with outwash from the Des Moines Lake, as well as coarse to fine grained fluvial deposits associated with local stream and river valleys. Unnamed erosion surface sediments consist of reworked till and silt/clay deposits associated with periglacial activity during the Wisconsin ice advance and may be up to 8 m (26 ft) thick. Poria Formation eolian materials consist of fine sand and silt. A relatively thin (up to 3 m, 10 ft) loess mantle is present in the southwest portion of the county. Thick deposits of eolian sand are only present in the Cedar and Wapaisippian river valleys. Additional colluvial materials may be intertongued present mantling most other mapping units and are more abundant near stream valleys and on terraces. Pre-Illinoian glacial deposits in northeast Iowa consist of two formations: the younger Wolf Creek Formation and the Alburtus Formation. The Wolf Creek Formation is divided into the Winthrop, Aurora, and Hickory Hills members (Bedrock to youngest). The Alburtus Formation consists of two units: "undifferentiated" members. Pre-Illinoian till is not exposed in the map area but is mantled throughout Black Hawk County by IES materials, eolian sand, or alluvial sediments.

The Quaternary materials are underlain by Devonian and Silurian carbonate bedrock. Ekeven bedrock outcrops (five quarries, five road cuts and one excavation for a fill station) were found in the map area during the field investigation. In the mapping area, Middle Devonian rocks form the major bedrock surface, and water wells are developed in both Devonian and Silurian rocks. Ordovician Maquoketa Formation rocks occur in the deepest portions of the bedrock valleys in the northeast part of the map area and directly underlie the Silurian rocks. The stratigraphy of the regional area has been intensively studied by Iowa Geological Survey staff (e.g., Beharck, 1927; Koch, 1970) and re-studied and correlated by Witzke and Bunker (1984), Witzke and others (1988, 2010), Anderson and Bunker (1989), Groves and others (2008), etc. Other studies in the area include Anderson and Garvin (1984) and Day and others (2006). The stratigraphic nomenclature and correlation in this map follow the stratigraphic framework proposed by Witzke and others (1988). The bedrock surface of two quadrangles within Black Hawk County was recently mapped by Rowden and others (2011 and 2012).

**References**

Anderson, W.J. and Garvin, P.L. (eds), 1984. The Cedar Valley Formation (Devonian), Black Hawk and Buchanan counties: Carbonate Facies and Mineralization, *Ann. Geol. Soc. of Iowa*, Guidebook 42, 47 p.

Anderson, R.R. and Bunker, B.J. (eds), 1988. Fossil shells, piggy smells, and drainage wells: the geology of the Mason City, Iowa, area. *Geol. Soc. of Iowa, Guidebook 65*, 71 p.

Beharck, C.H., 1927. The Shelfrock Stage of the Devonian. *American Midland Naturalist*, v. 10, p. 316-370.

Betts, E.A., III, 1989. Late Quaternary history of the Iowa River Valley in the Corvallis Lake area in Picher, O.W. (ed.), *Geologic Reconnaissance of the Corvallis Lake area*. Geological Society of Iowa Guidebook 51, p. 93-100.

Boettler, J., 1978a. North American Pleistocene Stages reconsidered in light of probable Pliocene-Pleistocene continental glaciation. *Science*, v. 202, p. 303-307.

Boettler, J., 1978b. Chronology of some late Cenozoic deposits from the central United States and the ice ages. *Transactions of the Nebraska Academy of Sciences*, v. 6, p. 35-49.

Day, J., Lucraj, J., and Anderson, R. (eds), 2006. New Perspectives and Advances in the Understanding of Lower and Middle Paleozoic Carbonate Depositional Systems of the Iowa and Illinois Basins. *Guidebook for the 36th Annual Field Conference of the Great Lakes Section, Society for Sedimentary Geology (SEPM)*, and the 67th Annual Tri-State Field Conference, September 29 - October 2, 2006, 167 p.

Groves, J.R., Walters, J.C., and Day, J. (eds), 2008. Carbonate platform facies and faunas of the Middle and Upper Devonian Cedar Valley Group and Lime Creek Formation, northern Iowa. *KGS Guidebook 28*, 96 p.

Halberg, G.R., 1980. Pleistocene stratigraphy in east-central Iowa. *Iowa Geological Survey Technical Information Series*, 10, 168p.

Halberg, G.R., 1986. Pre-Wisconsinan stratigraphy of the central plains region in Iowa, Nebraska, Kansas, and Missouri: in Richmond, G.M. and Fullerton, D.S., eds., *Quaternary Glaciations in the United States of America*, Report of the International Correlation Programme Project 24, in Sirovica, V., Bowen, D.Q., and Richmond, G.M., eds., *Quaternary Science Reviews*, *Quaternary Glaciations in the Northern Hemisphere*, v. 5, p. 11-15.

Halberg, G.R., Lindeck, J.A., Melschon, D.M., Knox, J.C., Goebel, E., Hobbs, H.C., Whitfield, J.W., Ward, R.A., Boettler, J.D., and Swinchart, J.B., 1991. Quaternary geologic map of the Des Moines 4° x 6° quadrangle, United States. U.S. Geological Survey, Miscellaneous Investigations Series, Map I-420, 1:100,000 scale map sheet.

Koch, D.L., 1970. Stratigraphy of the Upper Devonian Shell Rock Formation of north-central Iowa. IGS Report of Investigations 10, the state of Iowa, 123 p.

Prior, J.C., 1976. Landforms of Iowa. Iowa City, University of Iowa Press, 154 p.

Prior, J.C. and Koeltz, C.J., 2006. The Landform Regions of Iowa. Iowa Geological Survey, digital map, available on DNR GIS Library. [http://pib.igwb.uiowa.edu/igsib\\_library/state/geologic/landform/landform\\_regions.asp](http://pib.igwb.uiowa.edu/igsib_library/state/geologic/landform/landform_regions.asp); <http://www.igwb.uiowa.edu/igsib/>

Rowden, R., McKay, R.M., Liu, H., Tassier-Surine, S., Quade, D., and Giglierao, J.D., 2011. Bedrock Geology of the Gilbertville 7.5 Quadrangle, Black Hawk County, Iowa. Iowa Geological and Water Survey Open File Map OFM-2011-03, 1:24,000 scale map sheet.

Rowden, R., McKay, R.M., Liu, H., Tassier-Surine, S., Quade, D., and Giglierao, J.D., 2012. Bedrock Geology of the Cedar Falls 7.5 Quadrangle, Black Hawk County, Iowa. Iowa Geological and Water Survey Open File Map OFM-2012-04, 1:24,000 scale map sheet.

Tassier-Surine, S., Quade, D., J. Giglierao, J.D., and Ben, E.A., III, 2007. Surficial Geology of the Bremer (Iowa) 7.5 Quadrangle. Iowa Geological Survey Open File Map OFM-07-5, 1:24,000 scale map sheet.

Tassier-Surine, S., Quade, D., Liu, H., McKay, R., and Giglierao, J., 2009. Surficial Geology of the Shell Rock 7.5 Quadrangle, Butler, Black Hawk and Bremer counties. Iowa Geological Survey Open File Map OFM 09-5, 1:24,000 scale map sheet.

Tassier-Surine, S., Quade, D., McKay, R., Liu, H., and Giglierao, J., 2010. Surficial Geology of Bremer County, Iowa. Iowa Geological Survey Open File Map OFM-10-2, 1:100,000 scale map sheet.

Tassier-Surine, S., Quade, D., Rowden, R., McKay, R.M., Liu, H., and Giglierao, J.D., 2011. Surficial Geology of the Gilbertville 7.5 Quadrangle, Black Hawk County, Iowa. Iowa Geological and Water Survey Open File Map OFM 11-7, 1:24,000 scale map sheet.

Tassier-Surine, S., Quade, D., Rowden, R., McKay, R., Liu, H., and Giglierao, J.D., 2012. Surficial Geology of the Cedar Falls 7.5 Quadrangle, Black Hawk County, Iowa. Iowa Geological and Water Survey Open File Map OFM 12-4, 1:24,000 scale map sheet.

Walters, J.C., 1996. General and Environmental Geology of the Cedar Falls/Waterloo Area. The Iowan Surface, in General and Environmental Geology of Cedar Falls/Waterloo and Surrounding Area, Northeast Iowa. Iowa Geological Survey Guidebook Series No. 22, p. 7-9.

Witzke, B.J., Anderson, R.R., and Pope, J.P., 2010. Bedrock Geologic Map of Iowa, scale: 1:500,000. Iowa Geological and Water Survey, Open File Digital Map OFM-10-1.

Witzke, B.J. and Bunker, B.J., 1984. Devonian stratigraphy of north-central Iowa. IGS Open File Report 84-2, 107-149.

Witzke, B.J., Bunker, B.J., and Rogers, F.S., 1988. Eolian through lower Farnham stratigraphy and deposition in the Iowa area, central midcontinent, U.S. in McKelvey, N.J., Embury, A.F., and Glass, D.J. (eds.) *Devolution of the World*, Canadian Soc. Of Petroleum Geologists, Memoir 14, vol. 1, p. 221-250.

Base map from Black Hawk County GIS data derived from DOT Transportation Basemap files from 2009.

Iowa Geological and Water Survey digital cartographic file: <http://www.igwb.uiowa.edu/igsib/>

Iowa Geological and Water Survey digital cartographic file: <http://www.igwb.uiowa.edu/igsib/>

Map projection and coordinate system based on Universal Transverse Mercator (UTM) Zone 15, datum NAD83.

Map and cross-sections are based on interpretations of the best available information at the time of mapping.

Map interpretations are not a substitute for detailed site specific studies.