

## ON EXTENSIONS OF NILPOTENT TORSION RINGS BY SEMISIMPLE RINGS

ISMAIL A. AMIN AND FERENC A. SZÁSZ

A class of rings in which each member is the extension of a nilpotent torsion ring by a semisimple semiartinian ring is presented.

Throughout the following  $R$  will always designate an associative ring not necessarily containing an identity element.

Problem 75 cited in the second author's monograph seeks to determine the structure of rings  $R$  satisfying the following two conditions:

- (A) if  $P$  is a prime ideal of  $R$ ,  $R/P$  is a simple ring with non-zero socle; and
- (B) the left annihilator of every homomorphic image of  $R$  is an MHR-ring (that is, a ring satisfying the minimum condition for principal right ideals).

We note that a left perfect ring is an MHR-ring with 1 (see [1] and [9]). Before giving a class of rings satisfying these two conditions, we pause to give certain definitions.

**DEFINITION 1.** An  $S$ -ring is a ring that satisfies conditions (A) and (B) simultaneously.

**DEFINITION 2.** A ring  $R$  is called a  $P$ -ring if for every homomorphic image  $B$  of  $R$  and for every right  $B$ -module  $M$ ,  $M = M_0 \oplus M_1$  where  $M_0 B = \{0\}$  and  $M_1 B = M_1$ .

---

Received 3 April 1981.

**DEFINITION 3.** A *PS*-ring is a ring which is both a *P*-ring and an *S*-ring.

**THEOREM 4.** A right Noetherian *PS*-ring *R* is the extension of a nilpotent torsion ring *B* by a semisimple ring *C*. Moreover, every non-zero right ideal of every homomorphic image of *C* contains a minimal right ideal.

*Proof.* Since *R* is right Noetherian, every ideal of *R* contains a product of finitely many prime ideals of *R* (see [1] and [9]). In particular we can find prime ideals  $P_1, \dots, P_k$  of *R* whose product is the zero ideal. There will be no loss of generality if we suppose that

$$R = P_0 \not\supseteq P_1 \not\supseteq P_1 P_2 \not\supseteq \dots \not\supseteq P_1 P_2 \dots P_k = \{0\} .$$

Having this we build the following factor rings:

$$H_j = P_1 \dots P_{j-1} / P_1 \dots P_j , \quad j = 1, 2, \dots, k+1 .$$

It is not hard to see that each  $H_j$  can be regarded as an  $R/P_j$ -module in the natural way. Our assumption shows that  $R/P_j$  is a simple MHR-ring that possesses minimal one-sided ideals. Thus, by a result of the second author [10, 11], the  $R/P_j$ -module  $H_j$  has a direct sum representation:

$$H_j = B_j \oplus C_j , \quad \text{where } B_j R = \{0\} \quad \text{and} \quad C_j R = C_j .$$

This means that the perfect  $R/P_j$ -module  $C_j$  is a perfect *R*-module too. So  $C_j$  is a completely reducible *R*-module [11]. On the other hand,

$$B_j = (0 : R/P_1 \dots P_j)_l .$$

But since the left annihilator of  $R/P_1 \dots P_j$  is an MHR-ring, the additive group of  $B_j$  is torsion [11, 12]. Moreover, building the union of the complete inverse images of the ring homomorphisms

$$\phi_j : B_j \rightarrow B_j / (B_j \cap (P_1 \dots P_j))$$

one obviously obtains a nil ideal *B* of *R*. But since *R* is right Noetherian, *B* should be nilpotent by the Levitzki theorem ([6], p. 199). Moreover, (*B*, +) is torsion as we have seen.

Finally the union  $C$  of the chain of the complete inverse images of the mappings

$$\psi : C_j \rightarrow C_j / (C_j \cap (P_1 P_2 \dots P_j))$$

is a Jacobson semisimple ring with the required properties. This completes the proof. //

The importance of Theorem 4 may be best seen if we consider rings with 1. A ring with 1 is obviously a  $P$ -ring. Moreover, a ring with 1 is an MHR-ring if and only if it is a left perfect ring. For characterizations of left perfect rings we refer to [1] and [9]. We recall that a ring  $R$  with 1 is right semiartinian if and only if every homomorphic image of right  $R$ -module  $R$  has non-zero socle. The study of such important rings can be found in [2], [3] and [9]. One can see that a right semiartinian ring is a  $PS$ -ring. On the other side, the class of  $PS$ -rings contains every right Artinian ring and every quasi-Frobenius ring. Also, a right Noetherian ring with 1 which is in the same time right seminoetherian is necessarily right Artinian. So, many of the known results concerning such rings can be drawn or reformulated in view of Theorem 4.

**THEOREM 5.** *A right Noetherian  $PS$ -ring  $R$  with 1 is right Artinian.*

*Proof.* Since  $R$  has 1, Theorem 4 asserts that  $R$  is the extension of a nilpotent torsion ring by a right semiartinian ring. This shows that  $R$  itself is right Artinian. //

**PROBLEM 6.**  $R$  is a  $PS$ -ring and  $G$  is a finite (or soluble) group. Is the group ring  $R[G]$  again a  $PS$ -ring?

**PROBLEM 7.** Give an example of a  $PS$ -ring which is not semiartinian.

## References

- [1] Carl Faith, *Algebra II: ring theory* (Die Grundlehren der mathematischen Wissenschaften, 191. Springer-Verlag, Berlin, Heidelberg, New York, 1976).

- [2] Jonathan S. Golan, *Localization of noncommutative rings* (Pure and Applied Mathematics, 30. Marcel Dekker, New York, 1975).
- [3] Jonathan S. Golan, *Decomposition and dimension in module categories* (Lecture Notes in Pure and Applied Mathematics, 33. Marcel Dekker, New York, Basel, 1977).
- [4] Oscar Goldman, "Rings and modules of quotients", *J. Algebra* 13 (1969), 10–47.
- [5] Oscar Goldman, "Elements of noncommutative arithmetic I", *J. Algebra* 35 (1975), 308–341.
- [6] Nathan Jacobson, *Structure of rings*, revised edition (American Mathematical Society Colloquium Publication, 37. American Mathematical Society, Providence, Rhode Island, 1964).
- [7] Andor Kertész, *Vorlesungen über artinsche Ringe* (Akadémiai Kiadó, Budapest, 1968).
- [8] A. Kertész, "Noethersche Ringe, die artinsch sind", *Acta Sci. Math. (Szeged)* 31 (1970), 219–221.
- [9] Bo Stenström, *Rings of quotients. An introduction to methods of ring theory* (Die Grundlehren der mathematischen Wissenschaften, 217. Springer-Verlag, Berlin, Heidelberg, New York, 1975).
- [10] F. Szász, "Über Ringe mit Minimalbedingung für Hauptideale. I", *Publ. Math. Debrecen* 7 (1960), 54–64.
- [11] F. Szász, "Über Ringe mit Minimalbedingung für Hauptideale. II", *Acta Math. Acad. Sci. Hungar.* 12 (1961), 417–439.
- [12] F. Szász, "Über Ringe mit Minimalbedingung für Hauptideale. III", *Acta Math. Acad. Sci. Hungar.* 14 (1963), 447–461.
- [13] Ferenc Szász, *Radikale der Ringe* (VEB Deutscher Verlag der Wissenschaften, Berlin, 1975).

Department of Mathematics,  
Faculty of Science,  
Cairo University,  
Giza, Egypt.

Mathematical Institute,  
Hungarian Academy of Sciences,  
Budapest,  
Hungary.