

# *Physicochemical assessment and bacteriological studies of hand-dug wells of major markets in south western, Nigeria*

**T. A. Laniyan, A. S. Olatunji,  
O. O. Bayewu, G. O. Mosuro,  
M. O. Odukoya, S. Kenjiru &  
S. T. Odunuga**

**Arabian Journal of Geosciences**


ISSN 1866-7511

Volume 9

Number 4

Arab J Geosci (2016) 9:1-8

DOI 10.1007/s12517-015-2105-z



**Your article is protected by copyright and all rights are held exclusively by Saudi Society for Geosciences. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at [link.springer.com](http://link.springer.com)".**

# Physicochemical assessment and bacteriological studies of hand-dug wells of major markets in south western, Nigeria

T. A. Laniyan<sup>1</sup> · A. S. Olatunji<sup>2</sup> · O. O. Bayewu<sup>1</sup> · G. O. Mosuro<sup>1</sup> · M. O. Odukoya<sup>3</sup> · S. Kenjiru<sup>1</sup> · S. T. Odunuga<sup>1</sup>

Received: 19 January 2015 / Accepted: 10 September 2015 / Published online: 28 March 2016  
© Saudi Society for Geosciences 2016

**Abstract** Rapid population in developing nations has imposed stress on groundwater resources, thus the need to assess physicochemical and bacteriological impact of microbes on hand-dug wells along some major markets in Ibadan South-western Nigeria. Water samples from hand-dug wells were measured sequentially, and total dissolved solute (TDS), pH, electrical conductivity (EC), salinity, and temperature were measured in situ. Water samples were analyzed at a Microbiology Laboratory. Most probable techniques used for micro-organism analysis were in three stages: presumptive test used for confirmation of *Escherichia coli*, confirmed test for total viable bacteria count (TVBC), and complete test to reconfirm the presence of coliform. Presumptive test showed high rate of *E. coli* in most of the hand-dug wells with (37.5 %). Confirmed test revealed *Staphylococcus aureus* to be 25 %, followed by *Proteus vulgaris* (14.6 %), *Bacillus species* (12.5 %), *Pseudomonas aeruginosa* (8.3 %), and *Klebsiella* spp. (2.1 %) respectively. Total viable bacteria counts are 500 to 192,000. Physicochemical results (total dissolved solute (TDS), pH, electrical conductivity (EC), salinity, temperature) when compared with WHO (2006) and SON (2007) revealed all the parameters to be within the permissible limits except pH (5.8 to 9.56), and high values of the parameters were caused by organic matter. High *E. coli* in the study area revealed influence of human and animal fecal that could

lead to urinary tract infection; *Staphylococcus aureus* that comes from sewage could lead to diarrhea, gastrointestinal disease, and multiplication in tissues. High value of pH and *E. coli* above the permissible limits showed contamination of hand-dug wells in the study area.

**Keywords** Pathogens · Bacteriological · Coliform · Disease · Epidemic

## Introduction

Rapid growth in population, which enhances increase in water demand for domestic, market, agricultural, and industrial, has imposed stress on groundwater resources, USAID (2005), by increasing contamination of the surface and groundwater in developing countries which is the most economic and valuable natural resource for various human activities especially those tapping water from shallow unconfined aquifers (Amadi 2011; Olasehinde and Amadi 2009). The amount and type of dissolved ions in groundwater are a composition of the local geology, geochemical characteristics, prevailing human activities, and application (Ezeigbo 1989). The deterioration of groundwater in terms of quality through anthropogenic interference could lead to waterborne disease. Generally, waterborne diseases are transmitted through the ingestion of water and food that are contaminated by fecal materials that carry infective dose of various microbial pathogens. Waterborne diseases remain a major cause of death and illness in various developing countries. The global spatial distribution of these diseases showed that Africa and Asia accounts for a large percentage of such that includes cholera, typhoid fever, paratyphoid, bacillary dysentery, amoebic dysentery, gastroenteritis, and infective hepatitis (Lucas and Gilles 1973; Foster and Chilton 2003).

✉ T. A. Laniyan  
ttlaniyan@yahoo.com

<sup>1</sup> Department of Earth Sciences, Olabisi Onabanjo University, Ago Iwoye, Ogun State, Nigeria

<sup>2</sup> Department of Geology, University of Ibadan, Ibadan, Nigeria

<sup>3</sup> Department of Geology, University of Lagos, Lagos, Nigeria

In drinking water, microbes pathogens are the contaminants with the greatest chance of reaching levels high enough to cause acute health effect; such microbes are *Escherichia coli*, *Staphylococcus aureus*, *Proteus vulgaris*, and *Bacillus* spp. (Jespersen 2004; Chinedu et al. 2011; EPA 2011; IDLO 2006). Pathogens are disease-causing micro-organisms that include Bacterial pathogens (*Acinetobacter*, *Aeromonas*, *Bacillus*, *Campylobacter*, *Enterobacter sakazakii*, *E. coli*, *Klebsiella*, *Legionella*, *Pseudomonas aeruginosa*, *Salmonella*, *Shigella*, *Staphylococcus aureus*, *Vibrio*, and a typical mycobacteria that can grow in water and soil (WHO 1997): Viral pathogens (*adenoviruses*, *astroviruses*, *caliciviruses*, *enteroviruses*, *hepatitis A virus*, *hepatitis E virus*, *rotaviruses*, and *orthoreoviruses* (Grabow et al. 2001): Protozoan pathogens (*Acanthamoeba*, *Balantidium coli*, *Blastocystis*, *Cryptosporidium*, *Cyclospora cayetanensis*, *Entamoeba histolytica*, *Giardia intestinalis*, *Isospora belli*, *Microsporidia*, *Naegleria fowleri*, *Toxoplasma gondii* (WHO 2003)): Helminth pathogens (dracunculiasis and fascioliasis (Cairncross et al. 2002)) which require intermediate hosts to complete their life cycles but are transmitted through drinking water by different mechanisms. Coliforms could travel through a distance of 70.7 m (232 ft) from sewage trenches intersecting groundwater, while traveling through sand and pea gravel aquifer in 35 h will cover 30.5 m (100 ft) from primary sewage-injected subsurface (Crane and Moore 1984). Presence of septic systems and well depth was found to be related to detection of coliforms in groundwater, although these relationships were not statistically significant (Francy et al. 2000). Effect of distance from pollution sources was more pronounced on fecal and total coliform counts, which decreased with increasing distance from waste dumps and defecation sites (Adekunle et al. 2007). Also, other factors such as the environment where the well was sited and the level of hygiene of the well, in terms of the use of drawers and the population of people the well is serving and its surroundings could be considered as other possible sources of contamination aside the septic tanks.

Statistics revealed that inhabitants of Ibadan Metropolis in Nigeria suffer mainly from diarrhea, gastroenteritis, malaria, measles, tuberculosis, cholera, and typhoid fever (Iyun 1994). This had led to divers' investigation on the quality of water samples drawn from private wells and springs in various southwestern regions Adekunle (2008); Shimizu et al. (1980); Snow (1894); Swerdlow (1992); Swerdlow et al. 1992. The study area could easily breed microorganisms due to diver's reasons such as dirty environment, poor town planning, dilapidated infrastructure, and indiscriminate siting of wells and boreholes (Valenzuela et al. 2009) since coliform bacteria are widely found in nature (Binnie et al. 2002). The study therefore examined various pattern of physicochemical and bacteriological impact of microbes on hand-dug wells along some major markets.

## Materials and method

### The study area

Ibadan, the study area known as the city at the edge of savannah, the third largest metropolitan area, by population, in Nigeria, after Lagos and Kano with a population of 282,585 people (139,515 male and 143,070 female) in 2006, has a relief pattern that can be described as undulating with an elevation varying between 150 and 235 m above the sea level and longitude 07° 23' 18.2" N with latitude 03° 54' 27.4" E and a dendritic type of drain located within the basement complex of south-western Nigeria, underlain by a migmatite-gneiss complex which consist of banded gneiss with associated quartzite, hornblende biotite granite gneiss, quartzite schist, mica schist, and amphibolite layers, quartzites form prominent topographic features in the study area. The area has a well-accessible road network with minor paths which provides real accessibility to the wells. The two main problems man contends with in the study area are the quantity (source and amount) and quality of water (Fig. 1). The type of wastes which constitute environmental pollution in the area is domestic refuse consisting of degradable food wastes, leaves, dead animals, corpse, and non-degradable ones such as plastics, bottles, nylon, medical, and hospital wastes, generated in households, hospitals, industries, and commercial centers (Omoleke 2004).

### Data collection


Sampling was carried out during the dry season, and a total of 22 water samples were collected in hand-dug wells in major markets around Oje, Bere, Idi-Arere, Mapo, Oja-Oba, Bode, and Molete Market respectively. The procedure of collection was divided into three:

1. Procedure done before the field sampling
2. Field sampling
3. Laboratory methods

Procedure done before the field sampling involved using clean porcelain polyethylene plastic bottles to collect the water samples before the field procedure, and bottles were autoclaved at 121 °C for 15 min and were not opened until the point of collection.

For the field sampling, total dissolved solvent (TDS), salinity, temperature, and pH were measured in situ with a pH meter instrument before the water samples were collected. Samples were collected into sterile bottles and sealed with paper tape after disinfection with spirit lamp of dispensing point with flame. The samples were then placed in a container filled with iced block to maintain the microbial population and process microbial presence.





**Fig. 1** Location map of the study area with sampling points

The laboratory method includes presumptive test, confirmation test, and complete test. The presumptive test was used to detect and enumerate fecal coliform in the underground water; nine separate bottles were randomly picked and labeled 1–9 to be a representative sample for the whole study to detect the coliform level. One hundred milliliters of double strength of lactose broth was added to bottles 1–3, 100 ml of single strength

of lactose broth was added to bottles 4–9, 10 ml of the water sample was then added to bottles 1–3, 1 ml of water sample was to bottles 4–6, and 0.1 ml of water samples were added to bottles 7–9. Durham tube was then inverted over each bottle before sterilization after which all the bottles with lactose broth the water samples were later incubated at 37 °C for 24 h. Evidence of coliform gas observed from the Durham tube with bubbles

coming at the top of the tube indicates the presence of coliform, while the absence of Durham tube coliform could be detected through the change in color of the original media from red to yellow, and the absence of gas indicates absence of coliform.

The confirmed test involves serial dilution, streaking method, and pour plate method. Serial dilution is done to reduce the microbial load present in water samples; before plating, each of the positive and negative of the presumptive test was diluted to tenfold in peptone water (0.1 % nutrient broth) before being plated on the appropriate media. In the streaking method, 36.4 g of MacConkey agar was diluted to 700 ml of water and sterilized at 121 °C for 15 min, the mixture was then poured into petri-dish and was allowed to cool and solidify before streaking, organisms were thereafter incubated inversely at 37 °C for 24 h, and results were observed after the 24th hour by identifying different organisms on each plate which includes gram-positive and gram-negative. In Pour plate method, 1 ml of water was added to sterile nutrient broth on two different bottles each with Petri-dish plate and allowed to cool at 45 °C or chick tolerance, and then, 1 mL of water was added to 9 ml of nutrient blood which was shaken thoroughly after which was poured immediately into the Petri-dish plate which was allowed to solidified and incubated inversely at 37 °C for 72 h. Total viable bacteria count (TVBC) began by dividing the plates into four segments with permanent marker for an easy counting of microorganism, counting and recording each of the four segments separately, summing up the whole count, and converting the value to the colony-forming unit (CFU/ML) by using formula number of organisms  $\times \left[\frac{1}{10}\right]^{-3}$ .

The Indole test is the final analysis of the water sample. Subculture organisms' colonies were taken from the plate with inoculating loop into the sterile peptone water after which it was incubated for 72 h, and then a drop of shelling reagent was added to peptone water, for the observation of the color to ascertain the presences of coliform or gram-staining organism. After the few minutes, there was a presence of yellow color which indicates the presence of *E. coli*.

## Results and interpretation

### Physiochemical results

Total dissolved solid (TDS), pH, temperature, biochemical oxygen demand (BOD), electrical conductivity (EC), and salinity (Table 1) of the different markets were compared with SON (2007), and WHO (2006) results revealed all parameters to be within the permissible standard with the exception of pH in some of the markets due to high rate of indiscriminate disposal of refuse and uncontrolled drain. The mean values for the parameters are 438.50±267.17, 6.59±0.940, 27.10±0.778 °C, 5.60±1.264, 525.00±218.25, and 0.04±0.039, respectively.

**Table 1** Mean ±SD concentration (mg/l) of physiochemical parameters in the various water sources in the study area

Physical parameters	Oje market	Bere market	Mapo market	Oja-oba market	Idi-arere market	Bode market	Molete market	SON limit (2007)
pH	8.20±0.917	6.59±0.141	6.86±0.940	6.76±0.530	6.90±0.336	8.00±0.183	8.29±0.212	6.5–8.5
Temp. (°C)	29.10±0.504	30.35±0.778	27.10±0.552	26.80±0.505	29.20±0.781	28.50±0.387	30.20±0.566	–
BOD (mg/l)	6.00±0.405	5.61±0.535	6.00±0.910	6.80±1.264	8.03±0.132	5.60±0.441	5.40±0.387	–
TDS (mg/l)	535.00±176.230	498.50±70.003	827.30±173.900	654.30±267.170	690.00±9.899	539.00±61.230	438.50±4.950	1500
Electrical conductivity (µS/cm)	668.30±202.130	705.00±74.450	966.60±138.684	766.60±218.250	905.00±63.640	705.00±70.470	525.00±7.070	1000
Salinity (mg/l)	0.05±0.011	0.05±0.007	0.07±0.039	0.05±0.002	0.06±0.007	0.05±0.000	0.04±0.000	–

**Table 2** Occurrence of isolated organisms from the different locations

Locations	ST	EC	BS	PA	PV	KS
1	+	+	-	-	-	-
2	-	+	-	-	+	-
3	-	+	+	+	-	-
4	+	+	-	-	-	-
5	+	+	-	-	-	-
6	+	+	-	-	+	-
7	+	-	-	-	-	-
8	+	+	-	-	+	-
9	+	-	-	-	+	-
10	+	-	-	-	-	-
11	+	+	-	+	+	-
12	+	+	-	-	-	-
13	-	+	+	-	-	-
14	-	+	+	+	-	-
15	-	+	+	-	-	-
16	-	+	-	-	+	-
17	-	+	-	-	-	-
18	+	+	-	-	-	+
19	+	-	-	-	-	-
20	-	+	+	+	-	-
21	-	+	+	-	-	-
22	-	+	-	-	+	-


Keys: *ST* *Staphylococcus aureus*, *PV* *Proteus vulgaris*, *EC* *Escherichia coli*, *KS* *Klebsiella species*, *BS* *Bacillus species*, + present, *PA* *Pseudomonas aeruginosa*, - absent

**Bacteriological results**

Microbiological quality assessment (Table 2) of hand-dug wells was conducted and was observed that none of the water samples were found to be bacteria free. The 18.2 % of the four water samples are suitable for drinking after the coliform test, while the remaining 81.8 % contains high coliform level thus importable for drinking. Total viable bacteria count was observed to also be high, while prevalence of bacteria pathogens of a total of six bacteria species (two gram positive and four



**Fig. 2** Result showing *Escherichia coli* on petri-dish




**Fig. 3** Percentage of isolated organisms in the study area

gram negative) was recovered from the water samples circulating at the study area. *E. coli* (Fig. 2) were found to be the most predominant organism with prevalence rate of 18 (37.5 %) and closely followed by the prevalence rate of 12 (25 %) *Staphylococcus aureus*, then the prevalence rate of 7 (14.6 %) of *Proteus vulgaris*, and the prevalence rate of 6 (12.5 %) of *Baccillus species* and prevalence rate of 4 (8.3 %) of *Pseudomonas aeruginosa*, respectively, while the least isolated organism in the study area was *Klebsiella species*. The overall 48 strains of bacteria isolated belonging to six species were recovered. Mapo market was found to be most contaminated, and Bode market was least contaminated with bacterial (Fig. 3).

**Discussion**

Growth and survival of micro-organisms (Thomas et al. 2012) in drinking water are of public health (CFSPH 2009), because the presence of such in water is a risk factor to developing country infections. Eighteen (81.8 %) of the examined well water were found not to be acceptable for drinking. This observation might have adduced that the examined well water is not properly managed in terms of hygiene and possibly environmental cleanliness. The fact that *E. coli* was the most prevalent in this study further stressed that the examined well water was contaminated with fecal samples (Oshoma et al. 2009). *E. coli* are well-known for causing serious gastroenteritis disorder such as watery diarrhea, hemorrhagic colitis, and/or uremic syndrome (CFSPH 2009). *Staphylococcus aureus* which was the second most prevalent in the study has been linked with various hospital acquired infection (Duquette and Nuttall 2004). Some strains of these organisms especially the meticillin-resistant strains are even more virulent due to their multidrug-resistant attributes (CFSPH 2009), and they have also been implicated as the etiology of many community



**Fig. 4** Distribution of *Bacillus species*, *Escherichia coli*, *Klebsiella species*, *Pseudomonas aeruginosa*, *Proteus vulgaris*, and *Staphylococcus aureus* found in the well locations

associated and nosocomial infection (Duquette and Nuttall 2004). Until recently, these strains were susceptible; however, resistance seems to be increasing, and multiple antibiotic resistant strains have started to emerge. *Proteus vulgaris*, one of the isolated organisms, are known as common inhabitants of the human guts and frequently implicated in urinary tract infections (Senior 1979). *Bacillus species*, known as opportunist pathogens, since the late nineteenth century and reports of cases, have appeared at the rate of about 6 years for the past decade (Logan 1988). Also, some species of this organism especially the species *subtilis* are known to produce bacitracin, a toxic antibiotic to human tissue in addition to their antagonistic effect against other pathogenic bacteria (Brooks et al. 2005). *Pseudomonas aeruginosa*, another major nosocomial pathogens recovered in the study, was found to be associated with catheter-associated urinary tract infection which cannot be overemphasized (Fluit et al. 2001). The virulence of this bacterial isolate has been linked to their cell associated factors

like alginate, lipopolysaccharide (LPS), flagellum, pilus, and non-pilus adhesins as well as with exoenzymes or secretory virulence factors like protease, elastase, phospholipase, pyocyanin, exotoxin A, exoenzyme S, hemolysins (rhamnolipids), and siderophores (Matheson et al. 2006). Another common human intestinal flora which was isolated from some of the well waters was *Klebsiella* sp. This organism is frequently associated with hospital acquired infection (Sung-sheng and Tsai 2009). According to some authors, certain underlying diseases such as malignancy, cirrhosis, biliary tract disorders, diabetes mellitus (DM), and alcoholism may impair an individual's defenses and increase the risk of *Klebsiella pneumoniae* infection (Tsay et al. 2002). These bacteria may also be implicated in bacteremia thereby causing significant morbidity and mortality in general population (Tsay et al. 2002). Metastatic infections such as pyogenic brain abscess, meningitis, and endophthalmitis are the most common and/or important characteristics of *Klebsiella* infections. Higher



counts of total viable bacteria were observed in the study area compared with coliform counts. These high counts were due to the poor level of hygiene and sanitation observed, since hand-dug wells in the environ are constantly exposed to contamination from human activities and unplanned layout of houses that led to minute distances between wells, septic tanks, and refuse dumps (Fig. 4).

## Conclusions

The type of bacteria isolated from the examined well water samples is similar to that reported by Shittu et al. (2008). This observation might have adduced that the source of such water contamination may be similar. Such source may include improper disposal of refuse, contamination of water by sewage, and surface runoff. Heterotrophic count (HPC) measures a range of bacteria that are naturally present in the environment (EPA 2002). The total bacteria counts for all the water samples were generally high exceeding the limit of  $1.0 \times 10^2$  cfu/ml which is the standard limit of heterotrophic count for drinking water (EPA 2002). The high total heterotrophic count is indicative of the presence of high organic and dissolved salts in the water. The primary source of bacterial contamination includes surface runoff, pasture, and other land area where animal waste is deposited (Shittu et al. 2008). Additional sources include seepage or discharge septic tanks, sewage treatment facilities, and nature soil/plant bacteria (EPA 2002). Thus, from the bacteriological results, the well waters are highly contaminated and are not good for human consumption, agriculture, industrial, municipal uses, among others. It is therefore recommended that wells should be properly located, human activities should be controlled to prevent water body, and diverse educational talks should be organized for the people in the environment.

## References

- Adekunle AS (2008) Impacts of industrial effluent on quality of well water within Asa dam industrial estate, Ilorin, Nigeria. *Nat Sci* 6(3):1–5
- Adekunle IM, Adetunji MT, Gbadebo AM, Banjoko OB (2007) Assessment of groundwater quality in a typical rural settlement in Southwest Nigeria. *Int J Env Res Publ Health* 4(4):307–318
- Amadi AN (2011) Assessing the effects of Aladimma dumpsite on soil and groundwater using water quality index and factor analysis. *Aust J Basic Appl Sci* 5(11):763–770
- Binnie C, Kimber M, Smethurst G (2002) Basic water treatment. Royal Society of Chemistry, Cambridge
- Brooks GF, Butel SJ, Morse SA (2005) Medical microbiology, 23rd edn. McGraw-Hill Companies Inc, Singapore
- Caimcross S, Muller R, Zagaria N (2002) *Dracunculiasis* (guinea worm disease) and the eradication initiative. *Clin Microbiol Rev* 15:223–246
- Centre for Food Security and Public Health (2009) The Centre for Food Security and Public Health (CFSPH) Enterohemorrhagic *Escherichia coli* infections, Verocytotoxin Producing *Escherichia coli* (VTEC), Shiga toxin Producing *Escherichia coli* (STEC), *Escherichia coli* 0157 : H7
- Chinedu SN, Nwinyi OC, Oluwadamisi AY, Eze VN (2011) Assessment of water quality in Canaanland, Ota, Southwest Nigeria. *Agric Biol J N Am* 2(4):577–583
- Crane SR, Moore JA (1984) Bacterial pollution of groundwater: a review. *Water Air Soil Pollut* 22:67–83
- Duquette RA, Nuttall TJ (2004) Methicillin (Oxacillin)—resistant *Staphylococcus aureus* in dog and cats: an emerging problem. *J Small Anim Pract* 45(12):591–597
- Environmental Protection Agency (2002) US Environmental Protection Agency, safe drinking water act amendment, <http://www.epa.gov/safewater/mcl.html>
- Environmental Protection Agency (2011) Basic information about pathogens and indicators in drinking water, <http://www.epa.gov/drink/contaminants/pathogens>
- Ezeigbo HI (1989) Groundwater quality problems in parts of Imo state, Nigeria. *J Min Geol* 25(2):1–9
- Fluit AC, Schmitz FJ, Verhoel J (2001) Frequency of isolation of pathogens from bloodstream, nosocomial pneumonia skin and soft tissue and urinary tract infections occurring in European patients. *Eur J Clin Microb Infect Dis* 20:188–191
- Foster SSD, Chilton PJ (2003) Groundwater: the processes and global significance of aquifer degradation. *Philos Trans R Soc B Biol Sci* 258:1957–1972
- Francy DS, Helsel DR, Nally RA (2000) Occurrence and distribution of microbiological indicators in groundwater and stream water. *Water Environ Res* 72:152–161
- Grabow WOK, Taylor MB, de Villiers JC (2001) New methods for the detection of viruses: call for review of drinking water quality guidelines. *Water Sci Technol* 43:1–8
- IDLO (2006) Water tenure reform and public access to water as a basic need, International Development Law Organization Voice of Development Jurists Series
- Iyun BF (1994) Health problems in Ibadan region. In: Filani MO, Akintola FO, Ikporukpo CO (eds) Ibadan region. Rex Charles Publishers, Ibadan, pp 256–261
- Jesperon K (2004) Public drinking water and public health: a tie that binds, On Tap Fall pp.16–22
- Logan NA (1988) *Bacillus* species of medical and veterinary importance 25: 157–165
- Lucas AC, Gilles HM (1973) A short textbook of preventive medicine for the tropics. Hodder and Stoughton, London
- Matheson NR, Potempa J, Travis J (2006) Interaction of a novel form of *Pseudomonas aeruginosa* alkaline protease (aeruginolysin) with interleukin-6 and interleukin-8. *Biol Chem* 387:911–915
- Olasehinde PI, Amadi AN (2009) Assessment of groundwater vulnerability in Owerri and its environs, southeastern Nigeria. *Niger J Technol Res* 4(1):27–40
- Omoleke II (2004) Management of environmental pollution in Ibadan, an African city: the challenges of health hazard facing government and the people. *J Hum Ecol* 15(4):265–275
- Oshoma CE, Aghimien MO, Bello ZO (2009) Growth and survival of *Escherichia coli* in Kunun Zaki, during storage. *World J Agric Sci* 5(4):447–497
- Senior BW (1979) The special affinity of particular types of *Proteus mirabilis* for the urinary tract. *J Med Microbiol* 12:1–8
- Shimizu TTM, Kigotake TM, Agatomo HN (1980) Bacteria contamination of drinking water from wells in Miyazaki, Japan. In: Bulletin for Agriculture. Miyakazi University, pp. 21–28
- Shittu OB, Olaitan JO, Amusa TS (2008) Physico-chemical and bacteriological analyses of water used for drinking and swimming purposes in Abeokuta, Nigeria. *Afr J Biomed Res* 11:285–290

- Snow ND (1894) The pathology and mode of communication of Cholera. Lond Med Gaz 9:745–753
- Standard Organization of Nigeria (2007) Nigerian standard for drinking water quality, pp. 15–16
- Sung-Sheng S, Tsai MD (2009) Characteristics of *Klebsiella pneumoniae* bacteremia in community-acquired and Nosocomial infections in diabetic patients
- Swerdlow DS (1992) Water-borne transmission of epidemic cholera in Trujillo, Peru: lessons for a continent at risk. Lancet 340:28–33
- Swerdlow DL, Woodruff BA, Brady RC (1992) A waterborne outbreak in Missouri of *Escherichia coli* 0157:H7 associated with bloody diarrhea and death. Ann Intern Med 117(10):812–819
- Thomas BT, Effedua HI, Musa OS, Afolabi O (2012) Growth and survival of gastroenteritis pathogens in dried cassava powder (Garri). N Y Sci J 5(2):9–14
- Tsay RW, Siu LK, Fung CP, Chang FY (2002) Characteristics of bacteremia between community-acquired and nosocomial *Klebsiella pneumoniae* infection: risk factor for mortality and the impact of capsular serotype as a herald for community-acquired infection. Archaeological Int Med 162:1021–1027
- USAID (2005) *USAID Health: Environmental Health, Overview*. Available at [www.usaid.gov/oue\\_work/goal\\_health/eh](http://www.usaid.gov/oue_work/goal_health/eh)
- Valenzuela M, Lagos B, Claret M, Mondaca MA, Pérez C, Parra O (2009) Fecal contamination of groundwater in a small rural dryland watershed in central Chile. Chilean J Agric Res 69(2):235–243
- World Health Organization (1997) Guidelines for drinking-water quality: surveillance and control of community supplies, 2nd ed. Vol. 3, Geneva, World Health Organization, [http://www.who.int/water\\_sanitation\\_health/dwq/guidelines4/en/](http://www.who.int/water_sanitation_health/dwq/guidelines4/en/). Accessed on 19th April 2014
- World Health Organization (2003) Emerging issues in water and infectious disease. World Health Organization, Geneva
- World Health Organization (2006) Guidelines for drinking water quality, third edition, incorporating the first and second addenda volume 1, recommendations. WHO, Geneva