



Title	PLASMA LEVEL OF FIBROBLAST GROWTH FACTOR 21 IS INDEPENDENTLY RELATED TO BLOOD PRESSURE
Author(s)	Cheung, BMY; Fong, CHY; Chen, C; Cho, AWK; Wang, Y; Law, LSC; Ong, KL; Wat, NMS; Xu, A; Lam, KSL
Citation	The 26th Meeting of the International Society of Hypertension (Hypertension Seoul 2016) in collaboration with the 12th Congress of the Asian Pacific Society of Hypertension (APSH) and the 25th Annual Scientific Meeting of the Korean Society of Hypertension (KSH), Seoul, Korea, 24-29 September 2016: Abstract Book. In Journal of Hypertension, 2016, v. 34 n. e-Supplement 1, p. e507, Abstract no. LBPS 02-06
Issued Date	2016
URL	http://hdl.handle.net/10722/247540
Rights	This is a non-final version of an article published in final form in Journal of Hypertension, 2016, v. 34 n. e-Supplement 1, p. e507, Abstract no. LBPS 02-06; This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.

Plasma level of fibroblast growth factor 21 is independently related to blood pressure

Cheung BMY,^{1,3} Fong CHY,¹ Chen C,¹ Tso AWK,¹ Wang Y,^{2,3} Law LSC,¹ Ong KL,¹ Wat NMS,¹ Xu A,^{1,2,3} Lam KSL^{1,3}

¹Department of Medicine,

²Department of Pharmacology and Pharmacy,

³State Key Laboratory of Pharmaceutical Biotechnology,
University of Hong Kong, Hong Kong

Introduction: Fibroblast growth factor 21 (FGF21) plays an important role in glucose and lipid metabolism. Elevated blood FGF21 level is associated with obesity, diabetes and atherosclerosis. We therefore investigated its relationship with blood pressure.

Methods: We measured FGF21 in the plasma of 1921 participants (891 men, 1030 women; 52±12 years) taken at the start of the Hong Kong Cardiovascular Risk Factor Prevalence Study (CRISPS) using an enzyme-linked immunosorbent assay (Antibody & Immunoassay Services, University of Hong Kong). The log of FGF21 level was analysed for relationship with systolic and diastolic blood pressure (BP) at baseline and at the 5-year follow-up.

Results: Plasma FGF21 level was 224.3±7.4 in men and 214.1±7.1 pg/ml in women. It correlated significantly ($p<0.001$) with age ($r=0.30$), waist circumference ($r=0.31$), systolic BP ($r=0.32$), diastolic BP ($r=0.22$), triglyceride ($r=0.41$), HDL-C ($r=-0.27$), fasting blood glucose ($r=0.27$) and hsCRP ($r=0.27$). In multivariate analysis, FGF21 was related to systolic and diastolic blood pressure at baseline ($\beta=0.076$, $p<0.001$ and $\beta=0.074$, $p=0.001$ respectively) and to diastolic blood pressure at follow-up ($\beta=0.06$, $p=0.025$).

Conclusions: FGF21 level in blood is related to systolic and diastolic blood pressure, independent of age, obesity, lipids and blood glucose. FGF21 is related to the components of the metabolic syndrome and may have a pathophysiological role in hypertension.

Acknowledgment: Support from the Hong Kong Research Grant Council (HKU2/CRF/12R) is gratefully acknowledged.