

ACCEPTED MANUSCRIPT • OPEN ACCESS

Synergies and trade-offs between energy-efficient urbanization and health

To cite this article before publication: Sohail Ahmad *et al* 2017 *Environ. Res. Lett.* in press <https://doi.org/10.1088/1748-9326/aa9281>

Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is © 2017 The Author(s). Published by IOP Publishing Ltd.

As the Version of Record of this article is going to be / has been published on a gold open access basis under a CC BY 3.0 licence, this Accepted Manuscript is available for reuse under a CC BY 3.0 licence immediately.

Everyone is permitted to use all or part of the original content in this article, provided that they adhere to all the terms of the licence <https://creativecommons.org/licenses/by/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions may be required. All third party content is fully copyright protected and is not published on a gold open access basis under a CC BY licence, unless that is specifically stated in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

1
2
3 Synergies and trade-offs between energy-efficient urbanization and health
4
5
6

7 Sohail Ahmad^{1,2}, Shonali Pachauri³, Felix Creutzig^{1,2}
8
9

10 ¹Mercator Research Institute on Global Commons and Climate Change (MCC)
11 Torgauer Straße 12-15 10829 Berlin Germany
12
13

14
15 ²Department Economics of Climate Change, Technische Universität Berlin, 10623 Berlin, Germany
16
17

18
19 ³International Institute for Applied Systems Analysis (IIASA) - Schlossplatz 1 - A-2361 Laxenburg,
20 Austria
21
22

23 Emails: SA (ahmad@mcc-berlin.net); SP (pachauri@iiasa.ac.at); FC (creutzig@mcc-berlin.net)
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Synergies and trade-offs between energy-efficient urbanization and health

Abstract. Energy-efficient urbanization and public health pose major development challenges for India. While both issues are intensively studied, their interaction is not well understood. Here we explore the relationship between urban infrastructures, public health, and household-related emissions, identifying potential synergies and trade-offs of specific interventions by analyzing nationally representative household surveys from 2005 and 2012. Our analysis confirms previous characterizations of the environmental-health transition, but also points to an important role of energy use and urbanization as modifiers of this transition. We find that non-motorized transport may prove a sweet spot for development, as its use is associated with lower emissions and better public health in cities. Urbanization and improved access to basic services correlate with lower short-term morbidity (STM), such as fever, cough and diarrhea. Our analysis suggests that a 10% increase in urbanization from current levels and concurrent improvement in access to modern cooking and clean water could lower STM for 2.4 million people. This would be associated with a modest increase in electricity related emissions of 84 ktCO₂e annually. Promoting energy-efficient mobility systems, for instance by a 10% increase in bicycling could lower chronic diseases, like diabetes and cardio-vascular diseases, for 0.3 million people while also abating emissions. These findings provide empirical evidence to validate that energy-efficient and sustainable urbanization can address both public health and climate change challenges simultaneously.

Keywords: sustainable development; morbidity; environmental health transition; India; energy-efficient urbanization

1
2
3 India is projected to add 400 million new urbanites to its existing 410 million by 2050, doubling its
4 urban population within one generation (UN-DESA, 2014). Urbanization offers the chance of a better
5 life for many, providing improved access to infrastructure and living conditions (UN Habitat, 2009).
6 Yet, when public policy lags behind urbanization, as is currently the case in India and other emerging
7 economies, it results in a new set of challenges. Where urban growth proceeds unplanned, dismal
8 living conditions and inadequate infrastructure lead to growing inequities and an urbanization of
9 poverty (McGranahan & Satterthwaite, 2014; Ravallion, Chen, & Sangraula, 2007). In 2011, an
10 estimated 17.2 percent of urban Indians were living in slums under detrimental public health
11 conditions (Census of India, 2011). Improving health and wellbeing of these populations requires
12 rapid infrastructure development and extending modern services and amenities to all. However,
13 rapid expansion of infrastructure in cities and rising personal incomes can, in turn, result in growing
14 environmental footprints and associated health impacts. On average, household greenhouse gas
15 (GHG) and pollutant emissions from urban areas already exceed those from rural areas (Donglan,
16 Dequn, & Peng, 2010; Krey et al., 2012; O'Neill, Ren, Jiang, & Dalton, 2012).

17
18
19
20
21
22 Nonetheless, urban infrastructures are also at the center of demand-side action for climate change
23 mitigation (Creutzig, Agoston, et al., 2016; Creutzig, Fernandez, et al., 2016) and provide
24 opportunities for improving social stability and economic well-being (Bongardt, Breithaupt, &
25 Creutzig, 2010). Of total household emissions in urban India, three-fourth are from electricity and
26 private transport energy use (Ahmad, Baiocchi, & Creutzig, 2015). At the same time, a significant
27 fraction of urban populations in developing countries today thus face multiple overlapping
28 environmental health risks and opportunities concurrently (Kjellstrom et al., 2007; Marcotullio & Lee,
29 2003).

30
31
32 Energy, transport, climate, and building policies all offer significant potential to improve the
33 sustainability of cities (Creutzig, 2016; Grubler et al., 2012; Lucon et al., 2014; Ribeiro et al., 2012).
34 But understanding of the potential health co-benefits and risks of these policies remains limited,
35 especially at the level of individuals and households. The few studies that assess the health burdens
36 of specific transport and energy policies do so at an aggregate level (Wilkinson et al., 2009;
37 Woodcock et al., 2009). Yet understanding of the differential health outcomes and vulnerabilities of
38 urban populations in developing countries and the multiple energy, transport and infrastructural
39 correlates and confounders of these has not received adequate attention. Here we contribute to
40 improving this understanding empirically using microdata from two rounds of the most recently
41 available nationally representative longitudinal surveys from India (Desai, Vanneman, & National
42 Council of Applied Economic Research, 2010, 2015). We assess if there is a sweet spot in specific
43 urban development measures that maybe associated with lower GHG emissions and better public
44 health. We focus on understanding differences in traditional disease, labeled short term morbidity,
45 and modern disease, labeled major morbidity, prevalence among Indians, as well as their residential
46 and transport energy spending that correlate with emissions. We find significant differences in
47 energy and transport spending and morbidity patterns among rural and urban households, about half
48 of which can be attributed to differences in socioeconomic conditions and endowments. Our
49 analyses suggest that access to clean cooking, water, and improved sanitation is correlated with
50 lower short term morbidity. At the same time access to active (non-motorized) and public transit
51 mobility options is associated with lower chronic disease morbidity, as well as more efficient energy
52 use and lower emissions. Thus sustainable infrastructure development in cities can be an effective
53 means to achieve public health as well as local and global environmental objectives and goals. This
54 requires multi-sector integrated approaches to urban policies and planning to realize the greatest
55 gains to human and environmental health and to improve livability in cities.
56
57
58
59
60

Methods

This section presents the data employed and describes the health measures, energy and transport spending based emissions measures, and other explanatory variables used in this study. We also present the statistical methods and tests for model validation applied in our analysis.

Data Sources

This paper uses the Indian Human Development Surveys (IHDS) I (2004-05) and II (2011-12), both of which are nationally representative and multi-topic surveys (Desai et al., 2010, 2015). While IHDS I surveyed 41,554 households (215,754 individuals), IHDS II re-interviewed 83% of the original households surveyed in IHDS I, as well as additional samples, so that a total of 42,152 households (204,568 individuals) were surveyed in this round. The samples are nationally representative, spread across 33 (now 34) states and union territories over rural and urban areas, covering questions on health, education, employment, expenditures, economic status, marriage, fertility, gender relations, and social capital.

Health Measures

The health measures covered by the surveys are based on information collected at the individual level, where individuals are classified as either having short-term morbidity (STM), or major morbidity (MM), or neither. An individual is assumed to have STM also referred to as traditional diseases or communicable diseases if s/he had fever, cough, or diarrhea in the last 30 days. An individual is assumed to have MM also referred to as modern diseases or non-communicable diseases if s/he has been diagnosed with high blood pressure, heart disease, diabetes, or asthma ever. Fig S1 presents distribution of these morbidities.

Energy and Transport Spending based Emissions Measures

The dataset includes detailed household consumption expenditures over the past 30 days, including data on electricity spending, and private transport spending (expenditures on diesel, petrol, CNG, and maintenance of owned vehicle). These spendings are a proxy for household energy use. These spendings may vary slightly spatially and in scale of consumption (for electricity), given differences in prices. The mean spending on electricity and private transport in 2012 are 58 ₹/capita and 76 ₹/capita respectively. Employing the data on household consumption expenditure on electricity and private transport, we estimate emissions from electricity and private transport using the methodology described in Box S1. The annual per capita emissions from electricity and private transport are estimated to be 0.282 tCO₂e and 0.052 tCO₂e respectively in 2012, corresponding to 21% of economy-wide per capita emissions (including the burning of fossil fuels and the manufacture of cement) (World Bank, 2012).

Explanatory Variables

We group these variables into four categories – built environment, basic services, energy and transport, and other control variables that consist of socioeconomic and demographics characteristics of the households. The built environment consists of location of settlements (megacity, urban (non-megacity), and rural), basic services include modern stove, piped water, flush toilet, separate kitchen, and overcrowding (i.e. three or more persons living in a room). Here, individuals living in the six largest metropolitan areas (Mumbai, Delhi, Kolkata, Chennai, Bangalore, and Hyderabad) are defined as megacity dwellers. The geographical boundaries include census “urban agglomeration” for each of six areas with the exception of Delhi, where Gurgaon district (Haryana) and Gautam Buddha district (UP) are also included. The variables related to energy and transport consist of electricity spending, private-transport spending, and vehicular ownership. The variables related to socioeconomic and demographic characteristics include household size, age, gender, highest adult education level (male and female), and income. Tables, S1-S3, present

1
2
3 descriptive statistics of these variables, and Table S4 presents the correlation coefficients among the
4 variables of interest.

5 6 *Statistical Methods*

7
8 Multivariate regressions are employed to understand the factors underlying emissions from
9 electricity and private transport based energy spending at the household level, and health (likelihood
10 of having short term and major morbidity) at the individual level, following other empirical studies
11 (Ahmad et al., 2015; Lenzen et al., 2006). Some of the household-level attributes (i.e. sanitation
12 facilities) are assigned to the individual-level, and vice versa. We perform panel regressions using
13 2005 and 2012 data to get robust estimates.

14
15 For emissions (electricity and private transportation), which are continuous variables, we use linear
16 models, whereas for the prevalence of STM (or MM) relative to neither, we use logit models using
17 Stata 14.1 (Stata Corp., College Station TX, USA). Pooled OLS regressions are run to estimate constant
18 coefficients, under the usual assumption for cross-sectional analysis that regressors are uncorrelated
19 with the error term. To account for individual heterogeneity, we estimate fixed and random effect
20 panel models (Wooldridge, 2012).

21
22 To decide between the fixed or random effects model, we run a Hausman test (Greene, 2011). This
23 basically tests whether the unique errors are correlated with the regressors, assuming under the null
24 hypothesis that they are not. Based on the Hausmann test, we conclude that the coefficients
25 estimated by the efficient random effects are not the same as the ones estimated by the consistent
26 fixed effects estimator. Therefore, it is not appropriate to use the random effects model. In addition,
27 we test for time fixed effects, a joint test to see if the dummies for years (2005 and 2012) are equal
28 to 0. In our case, we reject the null hypothesis that the coefficients for all years are jointly equal to
29 0, and therefore retain time fixed effects in the model.

30
31 Using the results from our multivariate regression models, we also estimate the changes in our
32 dependent variables - emissions and prevalence of morbidity - resulting from a change in some of
33 our key explanatory variables, like urbanization in a comparative static analysis (for details of the
34 methods employed for these computations see Box S2).

35 36 37 38 39 40 *Descriptive statistics*

41
42 Our investigation of the patterns of individual morbidity and energy spending at a household-level in
43 India shows that the prevalence of MM is higher whereas that of STM is lower in urban centers and
44 for households with higher incomes in line with the environmental health transition literature (Smith
45 & Ezzati, 2005). Per capita electricity and private transport energy consumption (both continuous
46 variables that are a proxy for emissions) are also higher for households with higher income and that
47 live in urban areas (Fig 1). Between 2005 and 2012, both morbidity and energy consumption have
48 increased, albeit energy consumption has increased at a faster pace.

49
50 Other infrastructural characteristics are also correlated with STM, MM, and spending on electricity
51 and transport (Fig 2). Households with higher STM live in dwellings that have poorer quality drinking
52 water and sanitation facilities (Fig 2 and Table S1). In contrast, households that use better quality
53 cooking fuels, have higher electric spending, and that use motorized vehicles for personal
54 transportation (i.e. own more two-wheelers and four-wheelers) have higher MM. Notably,
55 households that own vehicles have several times higher MM as well as energy consumption.
56
57
58
59
60





Fig 1. Morbidity prevalence and energy consumption by income (bottom panel), settlement type (middle panel) and over time (top panel) in Indian households. All graphs use 2012 survey data, except for the top panel, which relies on survey data from both rounds.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60






Fig 2. Comparing prevalence of morbidity and energy spending (times) to the national average, by basic services, vehicle ownership and living space in India for 2005 and 2012. The national averages: STM (15%), MM (5%), electricity spending 524₹/capita/year, and transport spending 880₹/capita/year (spending in 2005 ₹).

Multivariate regression results

To analyze these differences further and test the statistical significance of various household characteristics in affecting traditional disease (STM) and modern disease (MM) prevalence, as well as electricity and transport energy spending, we present here the fixed effect model results organized by the following categories of explanatory variables: built environment, basic services, energy and transport, and socio-economic and demographic characteristics of households (Table 1 & 2).

Built Environment

We find that inhabitants of megacities have lower odds (0.68 times) of STM than rural inhabitants. But unlike other findings from low-income countries (van der Sande et al., 2000) we do not observe statistically significant higher MM in cities, after controlling for other variables as shown in Table 1. Moreover, we do not find statistically significant differences in STM and MM between smaller cities (urban category) and rural inhabitants, contrary to the hypothesis that urban inhabitants have lower incidence of STM and higher incidence of MM (Agarwal, 2011; Gupta, Arnold, & Lhungdim, 2009). Households in smaller cities spend more on transport and electricity than rural households. After controlling for other variables, including income, we find electricity spending in megacities is lower than in other areas, suggesting possible efficiencies of scale (Bettencourt, Lobo, Helbing, Kühnert, & West, 2007). The lower electricity spending among megacity inhabitants could also be explained by other household characteristics e.g., family work pattern and density of human settlement (Makido, Dhakal, & Yamagata, 2012), which are not controlled for here. Individuals in megacities have lower

1
2
3 traditional disease prevalence and household electricity spending (use), but higher utilization of
4 private transport, and consequently higher emissions of air pollutants and greenhouse gases than
5 those living in other urban areas, when controlling for all other variables.
6
7

8 *Basic Services*

9
10 Access to basic services, specifically to modern stoves, piped water, and flush toilet, lower the odds
11 of STM, but have no significant effect on the odds of MM. Our results echo previous findings that
12 show that switching from traditional to modern non-solid cooking fuels brings about large reductions
13 in household smoke, therefore improving health (Wilkinson et al., 2009). Households with access to
14 these basic services also show higher expenditures on energy. However, households with access to
15 piped water spend 4.2% less on electricity, ceteris paribus.
16
17

18 Housing space provides the immediate environment where individuals spend two-thirds of their time
19 (Brasche & Bischof, 2005). It also provides an environment for the household economy. Therefore, it
20 is likely to play an important role in overall morbidity outcomes (Krieger & Higgins, 2002; Shaw,
21 2004) as well as energy spending, required for lighting and space conditioning. We examine the
22 relationship between dwelling space using two proxy variables – separate kitchen and room per
23 capita – on morbidity and energy spending. Dwelling units with a separate kitchen are less exposed
24 to smoke, resulting in reduced disease prevalence. Previous studies have shown that lower room per
25 capita, resulting in inadequate space or overcrowding, is associated with a lack of privacy and stress,
26 which contribute to both physical and mental illness, transmission of tuberculosis and respiratory
27 infections (Krieger & Higgins, 2002), and increased incidents of accidents.
28
29

30 Our results reveal that inhabitants living in dwelling units with a separate kitchen have lower STM
31 ($p<0.01$) and MM ($p<0.1$) compared to those without a separate kitchen. We also find that STM and
32 MM are positively associated with room per capita (Fig. 2). These findings are against intuition, as
33 more space ought to reduce STM. The average room per capita in urban smaller city households
34 (0.59) is higher than in megacities (0.53) and rural (0.52) households. It is likely that households with
35 higher room per capita are located in peripheral urban areas, which could have poorer access to
36 certain municipal services, which are not controlled for here, e.g., solid waste management, that
37 could be associated with higher STM. Moreover living in peripheral urban areas might be associated
38 with longer commute times (more exposure to air pollution) and/or more sedentary lifestyles that
39 could also be related to higher prevalence of MM.
40
41
42

43 *Electricity and Transport*

44
45 Increased electricity spending is correlated with decreasing STM, whereas private-transport spending
46 is positively correlated with increasing MM. Interestingly, we do not find a statistically significant
47 influence of electricity spending on MM and transport spending on STM, ceteris paribus. These
48 findings suggest that adequate access to electricity maybe associated with lower STM, whereas the
49 use of non-motorized personal transport maybe associated with lower MM.
50
51

52 Owning a bicycle is associated with lower STM as well as MM ($p<0.1$), everything else held constant.
53 This findings complement our earlier one that indicates a correlation between transport spending
54 and MM. Ownership of vehicles also explains variations in electricity and transport spending. We find
55 that households with a motorcycle, compared to those without one, spend 11% and 102% more on
56 electricity and transport, respectively, ceteris paribus. Similarly, households with a car, compared to
57 those without one, spend 7% and 63% more on electricity and transport, respectively.
58
59

60 *Socioeconomic and Demographic Variables*

Our study controls for socioeconomic and demographic characteristics that partially explain prevalence of STM and MM, as well as energy spending. Notably, we find higher income is associated

with lower STM but is unrelated with MM. Moreover, we find electricity and transport spending is inelastic in income, as has been shown in previous studies (Ahmad & Puppim de Oliveira, 2016; Lenzen et al., 2006). A 10% increase in income is associated with 0.7% increase in electricity spending and 1.5% increase in transport spending. Thus, given similar increases in income, we find transport spending rises twice as fast as electricity spending.

Higher education levels among female members is related to higher prevalence of MM, according to our analysis. Possibly, changing activity patterns that accompany higher educational attainment might explain this finding. We also find that with age STM is lower but MM is higher. This suggests that women and seniors might deserve special attention in designing policies to moderate morbidity incidences.

Table 1 Likelihood of the prevalence of morbidity at the individual-level in India 2005 and 2012. *** p<0.01, ** p<0.05, * p<0.1; standard errors in parentheses.

Variables	STM	MM
Megacity (ref: rural)	0.686*** (0.0625)	0.697 (0.155)
Urban (ref: rural)	1.079 (0.0866)	0.951 (0.144)
Kitchen separate	0.914*** (0.0201)	0.903* (0.0509)
Stove modern	0.932** (0.0288)	1.082 (0.0675)
Piped water	0.927*** (0.0267)	1.053 (0.0665)
Flush toilet	0.946** (0.0244)	1.105* (0.0583)
Room pc	1.164*** (0.0426)	1.185** (0.0789)
Electricity pc (log)	0.974*** (0.00760)	0.982 (0.0193)
Transport pc (log)	0.994 (0.00616)	1.036** (0.0144)
Cycle (ref: Ø)	0.949** (0.0218)	0.913* (0.0460)
Motor cycle (ref: Ø)	1.064** (0.0329)	1.029 (0.0649)
Car (ref: Ø)	1.007 (0.0640)	1.130 (0.120)
Household size	0.914*** (0.0049)	0.969** (0.0123)
Age	0.890*** (0.00552)	1.081*** (0.0176)
Age^2	1.001*** (4.54e-05)	1.000*** (0.000127)
Female	0.793 (0.136)	0.733 (0.371)
Highest edu male	1.006 (0.00365)	0.993 (0.00816)
Highest edu female	0.997 (0.00344)	1.013* (0.00735)
Income pc (log)	0.958*** (0.0121)	0.957 (0.0267)
Year12 (ref:05)	1.792*** (0.0661)	2.947*** (0.196)
Observations	54,116	15,322
R-squared		
F statistics	1953.74***	3079.52***
Number of Individuals	27,058	7,661
Individual FE	Yes	Yes

Table 2 Determinants of energy spending at the individual-level in India 2005 and 2012. *** p<0.01, ** p<0.05, * p<0.1; standard errors in parentheses.

Variables	Electricity PC (log)	Private transport PC (log)
Megacity (ref: rural)	-0.080**	0.347***


	(0.0374)	(0.0441)
Urban (ref: rural)	0.063**	0.194***
	(0.0265)	(0.0366)
Kitchen separate	0.141***	0.043***
	(0.00857)	(0.0104)
Stove modern	0.075***	0.128***
	(0.0103)	(0.0142)
Piped water	-0.041***	-0.003
	(0.0104)	(0.0134)
Flush toilet	0.035***	0.102***
	(0.00873)	(0.0120)
Room pc	0.311***	0.126***
	(0.0126)	(0.0171)
Cycle (ref: Ø)	0.090***	0.055***
	(0.00827)	(0.0106)
Motor cycle (ref: Ø)	0.112***	1.025***
	(0.0103)	(0.0135)
Car (ref: Ø)	0.073***	0.630***
	(0.0183)	(0.0274)
Household size	-0.018***	-0.024***
	(0.00180)	(0.00216)
Age	0.003	0.007**
	(0.00226)	(0.00286)
Age^2	-1.22e-05	-2.25e-05
	(1.62e-05)	(2.10e-05)
Female	-0.054	0.048
	(0.0617)	(0.0776)
Highest edu male	0.011***	0.017***
	(0.00132)	(0.00169)
Highest edu female	0.002	0.004**
	(0.00117)	(0.00156)
Income pc (log)	0.073***	0.151***
	(0.00447)	(0.00601)
Year12 (ref:05)	0.522***	0.515***
	(0.0129)	(0.0165)
Constant	1.192***	0.529***
	(0.0717)	(0.0912)
Observations	267,560	267,683
R-squared	0.153	0.192
F statistics	1249.86***	1652.72***
Number of Individuals	147,552	147,563
Individual FE	Yes	Yes

Comparative Static Analysis


We assess the potential implications for public health and energy use of specific interventions that have an important relationship with these outcomes based on a comparative static analysis. The potential effects on health and energy of specific interventions are determined assuming a change in the value of one factor, while all other independent variables are held constant. These results provide further insight on the relative importance of various factors that we find are associated with morbidity and energy use (Fig 3). All the interventions we tested are associated with lower STM, albeit with a minor increase in electricity emissions, whereas only a shift from private transport to public transport and cycling is associated with lower MM significantly. Hence, we find both potential tradeoffs and synergies between reductions in morbidity and GHG emissions from energy use (Fig 3).

As traditional diseases remain widespread ($\approx 16\%$ of population), our analysis suggests that structural interventions may have a large potential to lower STM. We find that a 10% increase in urban population (with 2012 data as the baseline) would be associated with lower STM for 0.55 million people. At the same time, this magnitude of urbanization is also associated with an increase in household electricity emissions by 28 ktCO₂/year. Alternatively, greater provision of basic services, such as a separate kitchen, modern stove, piped water, and flush toilet by a similar magnitude of 10% above 2012 levels, could lower STM by as much as 1.8 million, while increasing electricity emissions by about 56 ktCO₂/year.

Lowering private transport spending, through an associated shift from private motorized to public transit modes and non-motorized transport (e.g., bicycling) could be associated with lower major morbidity as well as energy emissions. Specifically, we estimate that an increase of 10% in bicycle ownership could lower major morbidity for 0.29 million people, and private transport emissions from households by as much as 1.5 ktCO₂/year.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



1
2
3 Fig 3. Changes in morbidity and emissions due to selected interventions in India, represented by a 10% change
4 relative to the 2012 national average values (dotted lines represent results that are not statistically significant
5 in our models). The ranges represent the 95% confidence intervals of parameter estimates.
6
7
8

9 Discussion and Conclusions

10 Our analysis of these nexus issues provides empirical evidence that energy-efficient and sustainable
11 urbanization can address both public health and climate change challenges simultaneously. Three
12 specific findings emerge from our empirical study that can inform sustainable urbanization policies:
13 the provision of non-motorized transport (NMT) may be a sweet spot for sustainable development
14 that is associated with better health and lower GHG emissions; urbanization and better access to
15 household infrastructure maybe a means to lower short-term morbidity; other socio-economic
16 developments and built environment interventions, except in the case of the transport sector,
17 remain rather unrelated with major morbidity prevalence.
18
19

20 The promotion of non-motorized and public transit options, through its affect on private transport
21 spending and bicycle use, is associated with lower major morbidity as well as transport related
22 emissions. However, trends in India and other developing economies indicate that the share of
23 bicycles is declining and private motorized vehicle ownership is rising with economic development
24 ((Pucher, Korattyswaropam, Mittal, & Ittyerah, 2005), and also see table S5). Beyond higher modern
25 disease prevalence, private motorized transport also augments the risk of road traffic injuries, air and
26 noise pollution, which pose a major global public health challenge (Garg & Hyder, 2006; Sharma,
27 2008). Hence, the provision of efficient and clean transport systems – through a combination of high-
28 quality mass transit and safe bicycle infrastructures (Bongardt et al., 2013) – could be highly
29 beneficial for both public health and climate change mitigation.
30
31

32 Mitigation effects of cycling are rather low in absolute numbers, and one order of magnitude smaller
33 than emission increases in electricity required to increase access to basic services. Nonetheless, the
34 effect is prospectively relevant, as the transport share of emissions increase with development and
35 structural change (Schäfer, 2005). Moreover, incentivizing car transport also leads to sprawled urban
36 form and long-term lock-in into car dependency and transport energy use (Borck & Brueckner, 2016;
37 Creutzig, 2014; Seto et al., 2016). Neither too dense slums, nor too sprawled suburbs help India
38 towards sustainable urbanization. Instead medium-dense suburbs with public transit and bicycle
39 access can best negotiate the trade-offs associated with urban form (Lohrey & Creutzig, 2016).
40
41

42 Other than these transport related interventions, our study did not find any evidence of the effect of
43 other socio-economic or built environment related factors on major morbidity, which has increased
44 significantly between 2005 and 2012. This also suggests that solutions to reduce major morbidity
45 might lie in measures beyond those studied here, such as better diets and physical activity.
46
47

48 Our study also provides empirical evidence of the potential role of urbanization and access to basic
49 amenities on short term morbidity. We find that interventions to improve access to clean water and
50 sanitation systems are strongly associated with lower STM as opposed to urbanization alone (Fig 3).
51

52 In contrast to other studies, we rely on microdata from nationally representative household surveys
53 to analyze the prevalence of morbidity at an individual level, and transport and energy related
54 emissions at a household level, while controlling for socio-economic and built environment related
55 variables. Methodologically, we use cross-sectional and panel regressions to identify significant
56 relationships, and provide evidence of potential interventions that may improve quality of life in
57 cities. Some caveats of our analysis are the relatively short time span of seven years (2005 and 2012)
58 between the two surveys employed, and the use of subjective measurements of morbidity as
59 captured in the surveys. Future work can build on this analysis by taking a wider systems perspective
60 and uncovering the underlying causal mechanisms behind rising morbidity and emissions in cities.
The availability of longer panel series and better data, such as objective measurement of individuals'

1
2
3 health status rather than reported health status alone, could also inform more spatially
4 differentiated analysis, and allow for other important health and emissions drivers such as nutrition
5 (food demand) and physical activity patterns to be incorporated. In this study, transported-related
6 information was also limited to the variables included in the survey. In the future more detailed
7 information such as on individuals' travelling pattern (typical transportation mode and distance)
8 could be of great use in carrying out more detailed assessments.
9

10
11 Our results have important implications for policy, particularly for the nexus between energy-
12 efficient urbanization, climate change mitigation, and sustainable development. With urbanization,
13 directed energy use from cities is expected to more than triple between 2005 and 2050 globally
14 (Creutzig, Baiocchi, Bierkandt, Pichler, & Seto, 2015). However, smart urbanization strategies, such as
15 those that rely on higher transport fuel prices, could reduce energy demand by 25% (Creutzig et al.,
16 2015). For no world region is this result more relevant than for India that is expecting the highest
17 absolute urbanization among all countries. Our analysis of households in India shows that a shift in
18 transport spending from private motorized means to clean mass transit and more bicycles could also
19 benefit public health by reducing the prevalence of modern diseases like diabetes. Thus, higher taxes
20 on gasoline and diesel to finance clean mass transit, such as BRT systems, could benefit public health
21 and the climate. The opposite is likely to be true for electricity, however. We find that an increase in
22 electricity spending is associated with greater utilization of clean cooking and water infrastructures,
23 and a significantly lower traditional disease prevalence. Hence, a blunt tax increase on electricity for
24 climate change mitigation might hinder the achievement of other sustainable development goals,
25 and in particular, improved public health. Significantly, previous research suggests that expanding
26 electricity access to households has a relatively marginal contribution to national greenhouse gas
27 emissions increases (Pachauri, 2014). On the other hand, climate mitigation policies in South Asia
28 that lead to higher fossil fuel costs could slow down clean cooking fuel uptake if not compensated for
29 by other social protection measures (Cameron et al., 2016). Climate policies hence need to shield the
30 poor and be cognizant of the potential public health benefits of higher energy and electricity use,
31 especially when starting from very low levels.
32
33
34
35
36
37

38 **Acknowledgments:** We would like to thanks participants at the third annual IHDS user conference
39 (Neemrana Fort Palace, Rajasthan, 16-18 March, 2016) for their comments and suggestions on earlier
40 draft of the manuscript. SA acknowledges the Alexander von Humboldt Foundation and the Federal
41 Ministry for Education and Research (Germany) for the research fellowship.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- Agarwal, S. (2011). The state of urban health in India; comparing the poorest quartile to the rest of the urban population in selected states and cities. *Environment and urbanization*, 23(1), 13-28.
- Ahmad, S., Baiocchi, G., & Creutzig, F. (2015). CO2 Emissions from Direct Energy Use of Urban Households in India. *Environmental science & technology*, 49(19), 11312-11320.
- Ahmad, S., & Puppim de Oliveira, J. A. (2016). Determinants of urban mobility in India: Lessons for promoting sustainable and inclusive urban transportation in developing countries. *Transport Policy*, 50, 106-114.
- Bettencourt, L. M., Lobo, J., Helbing, D., Kühnert, C., & West, G. B. (2007). Growth, innovation, scaling, and the pace of life in cities. *Proceedings of the national academy of sciences*, 104(17), 7301-7306.
- Bongardt, D., Breithaupt, M., & Creutzig, F. (2010). *Beyond the fossil city: Towards low carbon transport and green growth*. Paper presented at the Fifth Regional EST Forum.
- Bongardt, D., Creutzig, F., Hüging, H., Sakamoto, K., Bakker, S., Gota, S., & Böhler-Baedeker, S. (2013). *Low-carbon land transport: policy handbook*: Routledge.
- Borck, R., & Brueckner, J. K. (2016). Optimal energy taxation in cities.
- Brasche, S., & Bischof, W. (2005). Daily time spent indoors in German homes – Baseline data for the assessment of indoor exposure of German occupants. *International Journal of Hygiene and Environmental Health*, 208(4), 247-253. doi: <http://dx.doi.org/10.1016/j.ijheh.2005.03.003>
- Cameron, C., Pachauri, S., Rao, N. D., McCollum, D., Rogelj, J., & Riahi, K. (2016). Policy trade-offs between climate mitigation and clean cook-stove access in South Asia. *Nature Energy*, 1, 15010.
- Census of India. (2011). *Primary Census Abstract for Slum*. Delhi: Retrieved from http://www.censusindia.gov.in/pca/final_pca.aspx.
- Creutzig, F. (2014). How fuel prices determine public transport infrastructure, modal shares and urban form. *Urban Climate*, 10, Part 1, 63-76. doi: <http://dx.doi.org/10.1016/j.uclim.2014.09.003>
- Creutzig, F. (2016). Evolving narratives of low-carbon futures in transportation. *Transport Reviews*, 36(3), 341-360.
- Creutzig, F., Agoston, P., Minx, J. C., Canadell, J. G., Andrew, R. M., Le Quéré, C., . . . Dhakal, S. (2016). Urban infrastructure choices structure climate solutions. *Nature Climate Change*, 6(12), 1054-1056.
- Creutzig, F., Baiocchi, G., Bierkandt, R., Pichler, P.-P., & Seto, K. C. (2015). Global typology of urban energy use and potentials for an urbanization mitigation wedge. *Proceedings of the national academy of sciences*, 112(20), 6283-6288.
- Creutzig, F., Fernandez, B., Haberl, H., Khosla, R., Mulugetta, Y., & Seto, K. C. (2016). Beyond Technology: Demand-Side Solutions to Climate Change Mitigation. *Annual Review of Environment and Resources*, 41(1).
- Desai, S., Vanneman, R., & National Council of Applied Economic Research. (2010). *India Human Development Survey (IHDS), 2005*. Retrieved from: <http://doi.org/10.3886/ICPSR22626.v11>
- Desai, S., Vanneman, R., & National Council of Applied Economic Research. (2015). *India Human Development Survey-II (IHDS-II), 2011-12*. Retrieved from: <http://doi.org/10.3886/ICPSR36151.v2>
- Donglan, Z., Dequn, Z., & Peng, Z. (2010). Driving forces of residential CO2 emissions in urban and rural China: An index decomposition analysis. *Energy Policy*, 38(7), 3377-3383.
- Garg, N., & Hyder, A. A. (2006). Exploring the relationship between development and road traffic injuries: a case study from India. *The European Journal of Public Health*, 16(5), 487-491.
- Greene, W. (2011). *Econometric Analysis* (7th ed.). NJ: Prentice Hall, Upper Saddle River.
- Grubler, A., Bai, X., Buettner, T., Dhakal, S., Fisk, D. J., Ichinose, T., . . . Schulz, N. B. (2012). Urban energy systems.

- 1
2
3 Gupta, K., Arnold, F., & Lhungdim, H. (2009). Health and living conditions in eight Indian cities.
4 National Family Health Survey (NFHS-3) India 2005-06.
- 5 Kjestrom, T., Friel, S., Dixon, J., Corvalan, C., Rehfuess, E., Campbell-Lendrum, D., . . . Bartram, J.
6 (2007). Urban environmental health hazards and health equity. *Journal of Urban Health*,
7 84(1), 86-97.
- 8 Krey, V., O'Neill, B. C., van Ruijven, B., Chaturvedi, V., Daioglou, V., Eom, J., . . . Ren, X. (2012). Urban
9 and rural energy use and carbon dioxide emissions in Asia. *Energy Economics*, 34, S272-S283.
- 10 Krieger, J., & Higgins, D. L. (2002). Housing and Health: Time Again for Public Health Action. *American*
11 *Journal of Public Health*, 92(5), 758-768.
- 12 Lenzen, M., Wier, M., Cohen, C., Hayami, H., Pachauri, S., & Schaeffer, R. (2006). A comparative
13 multivariate analysis of household energy requirements in Australia, Brazil, Denmark, India
14 and Japan. *Energy*, 31(2), 181-207.
- 15 Lohrey, S., & Creutzig, F. (2016). A 'sustainability window' of urban form. *Transportation Research*
16 *Part D: Transport and Environment*, 45, 96-111. doi:
17 <http://dx.doi.org/10.1016/j.trd.2015.09.004>
- 18 Lucon, O., Üрге-Vorsatz, D., Ahmed, A. Z., Akbari, H., Bertoldi, P., Cabeza, L., . . . Jiang, Y. (2014).
19 Buildings.
- 20 Makido, Y., Dhakal, S., & Yamagata, Y. (2012). Relationship between urban form and CO2 emissions:
21 Evidence from fifty Japanese cities. *Urban Climate*, 2, 55-67. doi:
22 <http://dx.doi.org/10.1016/j.uclim.2012.10.006>
- 23 Marcotullio, P., & Lee, Y.-s. (2003). Urban environmental transitions and urban transportation
24 systems: A comparison of the North American and Asian experiences. *International*
25 *Development Planning Review*, 25(4), 325-354.
- 26 McGranahan, G., & Satterthwaite, D. (2014). *Urbanisation: Concepts and Trends*: IIED.
- 27 Metz, B., Solomon, S., Kuijpers, L., Andersen, S. O., Davidson, O., Pons, J., . . . Meyer, L. (2005).
28 *Safeguarding the Ozone Layer and the Global Climate System: Issues related to*
29 *hydrofluorocarbons and perfluorocarbons*. New York: Cambridge University Press.
- 30 NSSO. (2012). *Consumer Expenditure -Type 1*. Delhi: Ministry of Statistics and Programme
31 Implementation, Government of India.
- 32 O'Neill, B. C., Ren, X., Jiang, L., & Dalton, M. (2012). The effect of urbanization on energy use in India
33 and China in the iPETS model. *Energy Economics*, 34, S339-S345.
- 34 Pachauri, S. (2014). Household electricity access a trivial contributor to CO2 emissions growth in
35 India. *Nature Clim. Change*, 4(12), 1073-1076. doi: 10.1038/nclimate2414
- 36 Pucher, J., Korattyswaropam, N., Mittal, N., & Ittyerah, N. (2005). Urban transport crisis in India.
37 *Transport Policy*, 12(3), 185-198.
- 38 Ravallion, M., Chen, S., & Sangraula, P. (2007). New evidence on the urbanization of global poverty.
39 *Population and Development Review*, 33(4), 667-701.
- 40 Ribeiro, S. K., Meza, M. J. F., Creuzigt, F., Dubeaux, C., Hupe, J., & Kobayashi, S. (2012). Energy end-
41 use: transport *Global Energy Assessment*: Cambridge University Press.
- 42 Schäfer, A. (2005). Structural change in energy use. *Energy Policy*, 33(4), 429-437.
- 43 Seto, K. C., Davis, S. J., Mitchell, R. B., Stokes, E. C., Unruh, G., & Üрге-Vorsatz, D. (2016). Carbon lock-
44 in: Types, causes, and policy implications. *Annual Review of Environment and Resources*, 41,
45 425-452.
- 46 Sharma, B. (2008). Road traffic injuries: a major global public health crisis. *Public health*, 122(12),
47 1399-1406.
- 48 Shaw, M. (2004). Housing and public health. *Annu. Rev. Public Health*, 25, 397-418.
- 49 Smith, K. R., & Ezzati, M. (2005). How environmental health risks change with development: the
50 epidemiologic and environmental risk transitions revisited. *Annu. Rev. Environ. Resour.*, 30,
51 291-333.
- 52 UN-DESA. (2014). *World Urbanization Prospects: The 2014 Revision*, , CD-ROM Edition.
- 53 UN Habitat. (2009). *Planning Sustainable Cities: Global Report on Human Settlements 2009*. London:
54 Earthscan.
- 55
56
57
58
59
60

- 1
2
3 van der Sande, M. A., Milligan, P. J., Nyan, O. A., Rowley, J. T., Banya, W. A., Ceesay, S. M., . . .
4 Walraven, G. E. (2000). Blood pressure patterns and cardiovascular risk factors in rural and
5 urban gambian communities. *Journal of human hypertension*, 14(8), 489-496. doi:
6 10.1038/sj.jhh.1001050
7
8 Wilkinson, P., Smith, K. R., Davies, M., Adair, H., Armstrong, B. G., Barrett, M., . . . Oreszczyn, T.
9 (2009). Public health benefits of strategies to reduce greenhouse-gas emissions: household
10 energy. *The Lancet*, 374(9705), 1917-1929.
11
12 Woodcock, J., Edwards, P., Tonne, C., Armstrong, B. G., Ashiru, O., Banister, D., . . . Cohen, A. (2009).
13 Public health benefits of strategies to reduce greenhouse-gas emissions: urban land
14 transport. *The Lancet*, 374(9705), 1930-1943.
15
16 Wooldridge, J. (2012). *Introductory econometrics: A modern approach*: Cengage Learning.
17
18 World Bank. (2012). CO2 emissions (metric tons per capita). World Development Indicators
19 <http://data.worldbank.org/indicator/EN.ATM.CO2E.PC?locations=IN>
20
21 World Energy Council. (2014). Energy Efficiency Indicators. *CO2 emissions of transport per capita*
22 Retrieved 18 April, 2017, from [https://www.wec-indicators.enerdata.eu/transport-co2-](https://www.wec-indicators.enerdata.eu/transport-co2-emissions-per-capita.html)
23 [emissions-per-capita.html](https://www.wec-indicators.enerdata.eu/transport-co2-emissions-per-capita.html)
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60