

**UCC Library and UCC researchers have made this item openly available.
 Please [let us know](#) how this has helped you. Thanks!**

Title	Plasmin activity in Manchega ewe milk: The effect of lactation, parity and health of the udder, and its influence on milk composition and rennet coagulation
Author(s)	Caballero-Villalobos, J.; Garzón, A. I.; Martínez Marín, A. L.; Arias, R.; Ciocia, Felicia; McSweeney, Paul L. H.
Publication date	2017-10-17
Original citation	Caballero-Villalobos, J., Garzón, A. I., Martínez Marín, A. L., Arias, R., Ciocia, F. and McSweeney, P. L. H. (2017) 'Plasmin activity in Manchega ewe milk: The effect of lactation, parity and health of the udder, and its influence on milk composition and rennet coagulation', Small Ruminant Research, In Press. doi:10.1016/j.smallrumres.2017.10.005
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://www.sciencedirect.com/science/article/pii/S0921448817302778 http://dx.doi.org/10.1016/j.smallrumres.2017.10.005 Access to the full text of the published version may require a subscription.
Rights	© 2017 Elsevier B.V. This manuscript version is made available under the CC BY-NC-ND 4.0 license. http://creativecommons.org/licenses/by-nc-nd/4.0/
Embargo information	Access to this article is restricted for 12 months after publication by request of the publisher.
Embargo lift date	2018-10-17
Item downloaded from	http://hdl.handle.net/10468/4937

Downloaded on 2021-11-27T04:59:43Z

Accepted Manuscript

Title: Plasmin activity in Manchega ewe milk: The effect of lactation, parity and health of the udder, and its influence on milk composition and rennet coagulation

Authors: J. Caballero-Villalobos, A.I. Garzón, A.L. Martínez Marín, R. Arias, F. Ciocia, P.L.H. McSweeney



PII: S0921-4488(17)30277-8
DOI: <https://doi.org/10.1016/j.smallrumres.2017.10.005>
Reference: RUMIN 5573

To appear in: *Small Ruminant Research*

Received date: 7-3-2017
Revised date: 15-10-2017
Accepted date: 16-10-2017

Please cite this article as: Caballero-Villalobos, J., Garzón, A.I., Martínez Marín, A.L., Arias, R., Ciocia, F., McSweeney, P.L.H., Plasmin activity in Manchega ewe milk: The effect of lactation, parity and health of the udder, and its influence on milk composition and rennet coagulation. *Small Ruminant Research* <https://doi.org/10.1016/j.smallrumres.2017.10.005>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Plasmin activity in Manchega ewe milk: the effect of lactation, parity and health of the udder, and its influence on milk composition and rennet coagulation.

J. Caballero-Villalobos^{a,*}, A.I. Garzón^a, A.L. Martínez Marín^a, R. Arias^b, F. Ciocia^c, P.L.H. McSweeney^c

a. Departamento de Producción Animal, Universidad de Córdoba, Spain.

b. Centro Regional de Selección y Reproducción Animal (CERSYRA), Castilla-La Mancha, Spain.

c. School of Food and Nutritional Sciences, University College Cork, Ireland.

* *Corresponding author: Javier Caballero-Villalobos.*

Tel.: +34 606197367 Email address: v22cavij@uco.es

Highlights

- Plasmin activity can be used as an indicator of the health of the udder.
- Controlling plasmin activity may improve milk technological quality, reducing production costs.
- The relationship between casein and plasmin is still controversial.
- Plasmin activity has a negative impact on rennet coagulation.

Abstract

Milk from 40 Manchega ewes was collected monthly and analysed during a complete lactation (5 months). Milk samples were classified by their origin in 3 categories, termed PR (primiparous ewes), M1 (multiparous ewes with no damage of the udder in the previous lactation) and M2 (multiparous ewes with udder damage in the previous lactation). The influence on plasmin of several factors as stage of lactation, parity, somatic cell count and udder health status was studied, as well as the effect of plasmin activity on milk composition and rennet coagulation. Plasmin activity decreased throughout lactation but was not affected by parity or somatic cell count ($P>0.05$). A strong negative correlation was found between

plasmin activity and protein (especially casein), presumably due to the proteolysis of β -casein. Plasmin also worsened rennet coagulation, increasing rennet clotting time (RCT) and negatively affecting curd firmness (A_{60}), especially in multiparous ewes. However, the good health condition in the herd may have camouflaged some effects of plasmin over renneting.

Keywords: Dairy sheep; Plasmin; Udder health; Rennet coagulation.

1. Introduction

Proteolysis potentially affects all dairy products (Saint-Denis et al., 2001). This results in a lower quality of the products, the development of bitter flavours in UHT milks, a decrease in cheese yield and a degradation of caseins during storage and ripening (Mara et al., 1998).

Plasmin is an alkaline serine proteinase (Bastian and Brown, 1996). This enzyme is present in milk as its zymogen plasminogen, which activates to plasmin when somatic cell counts exceed 500×10^3 cells/ml. Milk contains the complete plasmin system: plasmin (PL), plasminogen (PG) and a complex structure of plasminogen activators (PA), plasminogen activator inhibitors (PAI) and plasmin inhibitors (PI) (Ismail et al., 2006; Silanikove et al., 2013; White et al., 1995). Both PA and PAI are known to be locally produced by mammary epithelial cells in the mammary gland (Heegard et al., 1994).

The PL system plays an important role in the breakdown of casein, reducing cheese yield and casein content due to the leakage of proteose-peptones into whey (Albenzio et al., 2009). Plasmin mainly attacks β -CN, α_{S2} -CN and α_{S1} -CN (susceptible in that particular order). However, κ -CN seems to resist its action, though some experiments have reported that it can be affected under certain conditions (Groves et al., 1998). An increase of plasmin activity in bovine milk has been described due to udder infections or advanced stages of lactation (Bastian et al., 1991; Politis et al., 1989). On the other hand, results in literature

regarding the behaviour of plasmin in ovine milk are often controversial, and many differences between breeds have been evidenced (Theodorou et al. 2007). However, most authors have evidenced low levels of plasmin activity in late lactation ewe milk (Albenzio et al., 2009; Koutsouli et al., 2015).

In cheese, plasmin-induced proteolysis can contribute to the development of flavour and texture during ripening (Ismail and Nielsen, 2010). Meanwhile, other authors associated an intense plasmin activity as a cause of a development of bitter peptides (Habibi-Najafi and Lee, 1996; Sousa et al., 2001). This seems to be more frequent in high-cooked cheese varieties (Fox and Kelly, 2006). In milk (whether raw, pasteurized or UHT), to the contrary, proteolysis is the cause of undesirable effects. In severe cases, casein hydrolysis induced by plasmin may greatly affect rennet coagulation (Albenzio et al., 2005; Srinivasan and Lucey, 2002). This represents an important issue concerning the dairy sheep industry, as almost all milk production is intended for cheesemaking. Thus, proteolysis causes a reduction of the processing capacity of milk into cheese, as well as changes in its composition and the development of bitter flavours in processed products (Guerrero et al., 2003).

Controlling plasmin activity in sheep's milk could lead to an improvement of quality in the dairy industry, perhaps also reducing production costs. Therefore, proteolysis induced by this enzyme has attracted strong interest from researchers, due to its complexity and versatile effects over the quality of milk and dairy products (Ismail and Nielsen, 2010). Manchega is the most common dairy sheep breed in Spain, and milk from this breed is used to elaborate Manchego cheese. This product is the best-selling cheese variety in Spain and one of the most popular cheeses abroad (Arias et al, 2016), and since 1996 is protected by a protected designation of origin (PDO) to guarantee its quality. However, despite the importance of Manchega ewe milk, the behavior of the plasmin system in this breed has not

been yet investigated. Thus, the aim of this study is to determine the influence of several inherent factors on plasmin activity and to evaluate the impact of plasmin on Manchega milk quality.

2. Materials and methods

2.1. Animal sampling

Milk from 40 Manchega ewes was collected throughout a whole lactation (5 months). All animals belonged to the farm La Nava Del Conejo (Valdepeñas, Castilla-La Mancha, Spain), included in the *Spanish National Association of Manchega Sheep Breeders (AGRAMA)*. Ewes were classified in 3 groups, according to their udder health status in the previous lactation: PR (primiparous ewes), M1 (multiparous ewes with no damage of the udder in the previous lactation) and M2 (multiparous ewes with udder damage in the previous lactation). Samples were divided in 4 aliquots and analysed for milk composition, rennet coagulation, somatic cell count (SCC) and plasmin activity (PL).

2.1. Milk composition and somatic cell count

Major milk components (fat, crude protein, lactose, total solids, casein and urea) were directly measured using a Milko-Scan FT-6000 (*Foss Electric*, Hillerød, Denmark). To determine ash, milk samples (10 ml) were pipetted into crucibles and dried in a laboratory oven for 24 hours to obtain total solids, and then were transferred to a muffle furnace for 2 hours at 550°C. Finally, crucibles were stored in a desiccator with silica gel for around 30-45 minutes, and weighted to calculate ash content.

SCC was measured using a Fossomatic FC (*Foss Electric*, Hillerød, Denmark) and expressed as cells/ml. A logarithmic transformation was applied to SCC values and milk samples were categorized in 3 groups: low SCC ($< 1.6 \times 10^3$), mid SCC ($1.6 \times 10^3 < \text{SCC} < 2 \times 10^3$) and high SCC ($> 2 \times 10^3$).

2.3. Rennet coagulation

Samples were preheated at 32°C and renneting parameters were monitored using a Formagraph viscometer (Foss Electric, Hillerød, Denmark), based on the oscillatory motion of circular pendula immersed in milk during coagulation. The testing time of the analysis was set to 60 minutes and the measured parameters were rennet clotting time (RCT), curd firming time (k_{20}), and curd firmness after 30 and 60 minutes (A_{30} and A_{60}). To measure curd yield, curds were individually placed in centrifuge tubes, cut with a spatula and centrifuged (30 minutes, $2800 \times g$, 37°C) to separate the whey.

2.4. Plasmin activity

Plasmin activity was determined according to the procedure described by Richardson and Pearce (1981), with slight modifications. Plasmin cleaves the peptide *N*-Succinyl-L-alanyl-L-phenylalanyl-L-lysyl-7-amino-4-methyl coumarin, and releases 7-amino-4-methyl coumarin (AMC), which can be quantified spectrofluorometrically. Milk (3 ml) was added to 1 ml 0.4M trisodium citrate and centrifuged ($29,000 \times g$, 20 min, 4°C). The supernatant and 50 μ l of filtrate were added to 825 μ l 50mM tris-HCl buffer pH 7.5 and incubated for 5 min at room temperature. Reaction was initiated by adding to the mixture 225 μ l substrate (5 mg coumarin peptide in 1.33 ml dimethylsulphoxide and 5.33 ml 50mM tris-HCl buffer, pH 7.5), and fluorescence intensity (380 nm excitation, 460 nm emission) was measured at 5 min intervals over a period of 35 min. Plasmin activity was determined from the linear part of the fluorescence emission versus time curve. One unit of plasmin activity was defined as the activity necessary to release 1 nmol AMC $\text{min}^{-1} \text{ml}^{-1}$ milk under the conditions of the assay.

2.5. Statistical analysis

Data were analysed using the GLM procedure of SAS 9.1 (SAS Institute Inc., Cary, NC). A

first statistical analysis of plasmin activity (PL) included stage of lactation (SL), parity (PAR), and their interaction as fixed effects. A second analysis included SCC group, udder health status in the previous lactation (UHS), and their interaction as fixed effects. Contrast analysis was performed to compare least squares means, and statistical significance was declared at $P < 0.05$. In addition, linear correlations were calculated between PL and milk composition, and also between PL and rennet coagulation variables.

3. Results

Considering three stages in lactation (early, mid, and late), a gradual decrease in plasmin activity with the course of lactation was observed in all samples (**Table 1**). The lowest values of plasmin activity were found in late lactation, although this drop was only significant in multiparous ewes (**Table 2**).

Table 3 compares the least square means of plasmin activity as affected by SCC and udder health status category (UHS). Somatic cell counts did not affect plasmin activity, as shown in **Table 4**, where plasmin activity means between different SCC ranges are contrasted. However, regarding UHS categories, in low SCC milk, plasmin activity in M2 was significantly higher than in PR, and also tend to be higher than in M1. In high SCC milk, plasmin activity in M2 was also higher than in PR, but in this case, similar to M1.

Correlation coefficients for plasmin activity and milk composition variables in PR, M1 and M2 are presented in **Table 5**. No correlation was found in any group between plasmin and fat content, lactose or urea concentration. Regarding milk proteins, there was a strong negative correlation between plasmin and casein content, which diminished as plasmin activity increased, especially in multiparous ewes, regardless of their clinical history. Average values for plasmin activity and casein content were 1.05 units/ml and 4.85%, respectively. Ash content was shown to be very related with plasmin activity, as significant

positive correlations were found in multiparous ewes (both M1 and M2). Lastly, there seemed to be an increase of pH with plasmin activity in primiparous ewes (PR).

Table 6 presents the correlation coefficients for plasmin activity and the renneting parameters of milk in PR, M1 and M2 ewes. In general, rennet coagulation of milk worsens as plasmin activity increases. This is evident specially in milk from M2, since plasmin activity in this category is significantly higher than in the rest. Therefore, in this group, an increase of plasmin activity is the cause of high rennet clotting times and a decrease in curd firmness.

4. Discussion

4.1. Effect of stage of lactation (SL) and parity (PAR) on plasmin activity

Although no big differences were found, the high plasmin activity recorded in the beginning of lactation, as well as its decreasing trend, agree with the results obtained in Comisana ewes by Albenzio et al. (2009, 2005, 2004) and Caroprese et al. (2007), and in different greek breeds (Karagouniko, Chios and Boutsiko) by Koutsouli et al. (2015) and Theodorou et al. (2007), who measured the lowest plasmin values in late lactation milk. Contrastingly, several studies performed on bovine milk (Bastian and Brown, 1996; Ismail and Nielsen, 2010) and milk from other Italian ewe breeds (Bianchi et al., 2004; Sevi et al., 2004) reported a different pattern in plasmin activity, where the highest values were generally recorded in the end of lactation. Politis et al. (1989) suggested that this increase of plasmin activity observed in dairy cows during late lactation is due to an increase of the activation rate of plasminogen to plasmin, rather than to a higher influx of plasmin to the mammary gland. Recent works have discussed that this presence of plasminogen in milk is probably due to a transcellular route of passage from blood (Silanikove et al., 2016). Other authors related

the increase of plasmin in late lactation with the involution of the udder (Koutsouli et al., 2015) or an increase of milking interval (Castillo et al., 2008; Kelly et al., 1998). Some experiments performed with goat milk also reported an increase of plasmin activity throughout lactation (Cortellino et al., 2006).

No significant effect of parity was found, although most of the studies performed in different species of domestic ruminants reported that plasmin activity increased with parity and age of the animals (Bastian and Brown, 1996; Battacone et al., 2005; Koutsouli et al., 2015).

4.2. Effect of somatic cell count and udder health status on plasmin activity

Although previous results from studies performed with Manchega ewes suggested an increase of casein breakdown due to the amount of somatic cells in milk (Caballero-Villalobos et al., 2015), plasmin activity in the present study was found similar in all UHS categories, despite variations in SCC. This agrees with the results published by Bianchi et al. (2004) for Sardinian ewes, and Koutsouli et al. (2015) for Greek sheep breeds, who reported that SCC did not seem to affect plasmin activity. Albenzio et al. (2009) reported that in healthy ewes with $SCC < 600 \times 10^3$ cells/ml, the plasmin system was not affected, which is consistent with the average values of SCC obtained in the present study ($\approx 230 \times 10^3$ cells/ml). Other authors found that plasmin activity increased with SCC (Battacone et al., 2005; Leitner et al., 2004; Theodorou et al. 2007). Furthermore, experiments performed in Manchega ewes (Castillo et al., 2008) found a positive correlation between plasmin and somatic cells in milk samples with very low SCC (175×10^3 cells/ml), supporting the hypothesis that the influence of SCC over the plasmin-plasminogen system may not follow a specific pattern. Theodorou et al. (2007) suggested that these mixed results may be due

to variations between breeds and the different methods used to measure plasmin. However, SCC is not the only variable for predicting PL evolution in milk, as PL activity is affected by a complex network of molecular interactions between enzyme activators and inhibitors (Albenzio et al., 2009).

Results from the present study also suggest that the levels of plasmin in Manchega milk are affected by the sanitary conditions of the udder in the previous lactation. Differences between groups were evidenced in high and low somatic cell count categories, as plasmin activity values increased with a previous deterioration of the udder health status.

4.3. Influence of plasmin on milk composition

No influence of plasmin activity on fat concentration was found, which agrees with Koutsouli et al. (2015), who reported that in Greek ewe breeds plasmin did not seem to affect fat content in milk.

Despite that some authors did not find an effect of plasmin on levels of casein (Albenzio et al., 2004), and others have found a positive correlation (Baldi et al., 1996; Bianchi et al., 2004), most of the references found in literature have reported similar results to those obtained in the present study (Jaeggi et al., 2003; Leitner et al., 2004; Politis and Ng Kwai Hang, 1989). Numerous authors have related this negative correlation with the proteolytic action of plasmin on β -CN (Leitner et al., 2004; Moatsou, 2010; Nielsen, 2002). Thus, this could also explain the negative correlation between plasmin activity and crude protein observed in PR and M2. In this case, the effect of proteolysis would not be so evident, as the measurement involves total protein and not only casein. This might prove why some recent works have not found a clear decrease in crude protein as plasmin activity increased

(Koutsouli et al., 2015). However, the results found in the literature are often controversial, as other authors have revealed a positive correlation between plasmin and crude protein. Proteolysis in low SCC milk seems to be dominated by the action of plasmin. However, as SCC increases, the relative significance of plasmin decreases, while the relative activity of other indigenous and microbial enzymes increases (Albenzio et al., 2005; Kelly et al., 2006; Santillo et al., 2009). Therefore, the further research on other indigenous enzymes should be considered to explain more clearly the effect of proteolysis in milk.

Although some authors have reported a negative correlation between lactose and plasmin activity in Sarda and Assaf breeds (Battacone et al., 2005; Leitner et al., 2004), no correlation has been found in this study for Manchega ewes.

There was a relatively slight correlation between plasmin activity and total solids, probably due to the already mentioned cleavage of casein. Meanwhile, ash content seemed to increase along with plasmin activity, but no references were found in literature to help explain this finding.

Lastly, an influence of pH over plasmin activity was only evident in PR ewes. Some authors, such as Battacone et al. (2005) reported a positive correlation between plasmin activity and pH, although others did not find an effect of plasmin on the native pH of milk (Koutsouli et al., 2015).

4.4. Influence of plasmin on rennet coagulation

Results describing a deterioration of milk coagulation (evidenced mainly in M2) agree with those obtained by Albenzio et al. (2004), Battacone et al., (2005) and Mara et al. (1998).

Plasmin mainly cleaves β -CN into γ -CN, although it can also attack α_{S2} -CN and α_{S1} -CN (Fox and Kelly, 2006). Some authors like Okigbo et al. (1985) have described a low concentration of β -CN, alpha α_{S1} -CN and κ -CN, as well as a higher content of γ -casein and other casein fragments in milk that coagulated poorly. Therefore, the decrease of casein concentration due to proteolysis in ewes with previous udder damage may explain this deterioration of milk technological properties. According to Battacone et al. (2005), the worsening of rennet coagulation parameters is more related to disorders of permeability in the mammary gland than to casein breakdown by proteolysis. However, as previously mentioned, there were no effects of plasmin over lactose that could imply alterations of the milk-blood barrier. Due to all these reasons, the relationship between casein hydrolysis and rennet coagulation needs to be further investigated in greater detail.

On the other hand, Leiber et al. (2005) and Srinivasan and Lucey (2002) found that, normally, an increase in plasmin activity negatively affected renneting parameters and curd yield. Nevertheless, other authors did not find a clear correlation between plasmin and rennet coagulation properties of milk. However, most of the experiments found in literature concerning plasmin activity describe addition of different plasmin concentrations to milk, to establish different levels or ranges of activity (Mara et al., 1998). Thus, several authors have reported that high levels of plasmin induce casein hydrolysis, affecting milk coagulation. However, in experiments performed on native raw milk, plasmin levels have been reported to be lower. This suggests that, although there was some impact of plasmin on renneting parameters in M2 ewes, in the rest of categories this effect might not be so evident at native concentrations of the enzyme.

In addition, Bastian et al. (1991) found that when SCC did not exceed 300×10^3 cells/ml, as in the animal population studied in this experiment, plasmin levels remained relatively low, and no further correlation between plasmin activity and renneting properties was found. This

might certainly explain the low impact of plasmin on milk coagulation in PR and M1.

Conclusions

In Manchega sheep, plasmin activity decreased throughout lactation, and its highest values were measured in ewes that suffered udder infections in the previous lactation. Therefore, regardless the health condition of the animals in the beginning of lactation, there seems to be a residual enzymatic activity persisting as a response to a previous infection.

In addition, in ewes with a previous udder infection, plasmin activity had a negative impact on rennet coagulation, probably due to casein breakdown. In the rest of the animals this effect was lighter. However, the good health condition of the herd (reflected by the low somatic cell counts measured) seems to camouflage the possible effects of plasmin on rennet coagulation.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

This project was funded by the Government of Spain (Ministry of Education, Culture and Sport / Ministry of Economy and Competitiveness), the industrial partner Agrométodos S.A.,

Santander Universidades and ceiA3 (Agri-food Science International Campus of Excellence), through the research program “*Doctores en Empresas*”.

The authors wish to thank Antonio Figueroa for his invaluable help during the development of the project.

References

Albenzio, M., Santillo, A., Caroprese, M., d’Angelo, F., Marino, R., Sevi, A., 2009. Role of endogenous enzymes in proteolysis of sheep milk. *J. Dairy Sci.* 92, 79–86.

Albenzio, M., Caroprese, M., Santillo, A., Marino, R., Muscio, A., Sevi, A., 2005. Proteolytic patterns and plasmin activity in ewes’ milk as affected by somatic cell count and stage of lactation. *J. Dairy Res.* 72, 86–92.

Albenzio, M., Caroprese, M., Santillo, A., Marino, R., Taibi, L., Sevi, A., 2004. Effects of somatic cell count and stage of lactation on the plasmin activity and cheese-making properties of ewe milk. *J. Dairy Sci.* 87, 533–42.

Arias, R., Gallego, R., Altares, S., Garzón, A., Romero, J., Jiménez, L., Oliete, B., Arias, C., Caballero, J., Martínez, A., Núñez, N., García, A., Ramón, M., Montoro, V. and Pérez-Guzmán, M.D. 2016. Quality of milk in Manchego sheep flocks. A review. *Arch. Zootec.* 65(251), 469–473.

Baldi, A., Savoini, G., Cheli, F., Fantuz, F., Senatore, E., Bertocch, L., Politis, I., 1996. Changes in plasmin-plasminogen-plasminogen activator system in milk from Italian Friesian herds. *Int. Dairy J.* 6, 1045–1053.

Bastian, E.D., Brown, R.J., 1996. Plasmin in milk and dairy products: an update. *Int. Dairy*

J. 6, 435–457.

Bastian, E.D., Brown, R.J., Ernstrom, C.A., 1991. Plasmin activity and milk coagulation. *J. Dairy Sci.* 74, 3677–3685.

Battacone, G., Cannas, E.A., Mazzette, A., Dimauro, C., Enne, G., 2005. Why does the increase of plasmin worsen the coagulation properties of milk in dairy sheep? *Ital. J. Anim. Sci.* 4, 342–344.

Bianchi, L., Bolla, A., Budelli, E., Caroli, A., Casoli, C., Pauselli, M., Duranti, E., 2004. Effect of udder health status and lactation phase on the characteristics of Sardinian ewe milk. *J. Dairy Sci.* 87, 2401–2408.

Caballero-Villalobos, J., Garzón A.I., Oliete, B., Arias, R., Jiménez, L., Núñez, N., Martínez, A.L., 2015. Relationship of somatic cell count and composition and coagulation properties of ewes milk. *Mljekarstvo* 65(2), 138–143.

Caroprese, M., Schena, L., Marzano, A., Marino, R., Sevi, A., Albenzio, M., 2007. Contribution of macrophages to plasmin activity in ewe bulk milk. *Ital. J. Anim. Sci.* 6(1), 545–547.

Castillo, V., Such, X., Caja, G., Casals, R., Albanell, E., Salama, A.A.K., 2008. Effect of milking interval on milk secretion and mammary tight junction permeability in dairy ewes. *J. Dairy Sci.* 91, 2610–2619.

Cortellino, G., Locci, F., Rampilli, M., 2006. An investigation of the plasmin-plasminogen system in caprine milk and cheese. *Int. Dairy J.* 16, 619–622.

Fox, P.F., Kelly, A.L., 2006. Indigenous enzymes in milk: overview and historical aspects - Part 1. *Int. Dairy J.* 16, 500–516.

Groves, M.L., Wickham, E.D., Farrell, H.M., 1998. Environmental effects on disulfide

- bonding patterns of bovine kappa-casein. *J. Protein Chem.* 17, 73–84.
- Guerrero, L., Román, S., Pacheco, L., 2003. Proteolysis during cold storage of refrigerated raw milk . Effect of proteolytic enzymes on casein integrity. *Rev. Científica FCV-LUZ* XIII, 187–192.
- Habibi-Najafi, M.B., Lee, B.H., 1996. Bitterness in cheese: a review. *Crit. Rev. Food Sci. Nutr.* 36, 397–411.
- Heegard, C.E., White, J.H., Zavizion, B., Turner, J.D., Politis, I., 1994. Production of various forms of plasminogen activator and plasminogen activator inhibitor by cultured mammary epithelial cells. *J. Dairy Sci.* 77, 2949–2958.
- Ismail, B., Choi, L.H., Were, L.M., Nielsen, S.S., 2006. Activity and nature of plasminogen activators associated with the casein micelle. *J. Dairy Sci.* 89, 3285–95.
- Ismail, B., Nielsen, S.S., 2010. Invited review: Plasmin protease in milk: current knowledge and relevance to dairy industry. *J. Dairy Sci.* 93, 4999–5009.
- Jaeggi, J.J., Govindasamy-Lucey, S., Berger, Y.M., Johnson, M.E., McKusick, B.C., Thomas, D.L., Wendorff, W.L., 2003. Hard ewe's milk cheese manufactured from milk of three different groups of somatic cell counts. *J. Dairy Sci.* 86, 3082–3089.
- Kelly, A.L., O'Flaherty, F., Fox, P.F., 2006. Indigenous proteolytic enzymes in milk: a brief overview of the present state of knowledge. *Int. Dairy J.* 6, 563–572.
- Kelly, A.L., Reid, S., Joyce, P., Meaney, W.J., Foley, J., 1998. Effect of decreased milking frequency of cows in late lactation on milk somatic cell count, polymorphonuclear leucocyte numbers, composition and proteolytic activity. *J. Dairy Res.* 65, 365–73.
- Koutsouli, P., Patounas, G., Massouras, T., Bizelis, I., Politis, I., 2015. Plasmin-plasminogen system and milk coagulation properties of two Greek dairy sheep breeds. *Small Rumin.*

Res. 124, 89–94.

Leiber, F., Nigg, D., Kunz, C., Scheeder, M.R.L., Wettstein, H.R., Kreuzer, M., 2005. Protein composition, plasmin activity and cheesemaking properties of cows' milk produced at two altitudes from hay of lowland and high-alpine origins. *J. Dairy Res.* 72, 65–74.

Leitner, G., Chaffer, M., Shamay, A., Shapiro, F., Merin, U., Ezra, E., Saran, A., Silanikove, N., 2004. Changes in milk composition as affected by subclinical mastitis in sheep. *J. Dairy Sci.* 87, 46–52.

Mara, O., Roupie, C., Duffy, A., Kelly, A.L., 1998. The curd-forming properties of milk as affected by the action of plasmin. *Int. Dairy J.* 8, 807–812.

Moatsou, G., 2010. Indigenous enzymatic activities in ovine and caprine milks. *Int. J. Dairy Technol.* 63(1), 16–31.

Nielsen, S.S., 2002. Plasmin system and microbial proteases in milk: characteristics, roles, and relationship. *J. Agric. Food Chem* 50, 6628–6634.

Okigbo, L.M., Richardson, G.H., Brown, R.J. and Ernstrom, C.A., 1985. Casein composition of cow's milk of different chymosin coagulation properties. *J. Dairy Sci.* 68, 1887–1892.

Politis, I., Lachance, E., Block, E., and Turner, J.D., 1989. Plasmin and plasminogen in bovine milk: a relationship with involution? *J. Dairy Res.* 72, 900-906.

Politis, I., Ng Kwai Hang, K.F., 1989. Environmental factors affecting plasmin activity in milk. *J. Dairy Sci.* 72, 1713–1718.

Richardson, B.C., Pearce, K.N., 1981. The determination of plasmin in dairy products. *New Zeal. J. Dairy Sci. Technol.* 16, 209–220.

Saint-Denis, B.T., Humbert, G., Gaillard, J.L., 2001. Enzymatic assays for native plasmin, plasminogen and plasminogen activators in bovine milk. *J. Dairy Res.* 68, 437–49.

- Santillo, A., Kelly, A.L., Palermo, C., Sevi, A., Albenzio, M., 2009. Role of indigenous enzymes in proteolysis of casein in caprine milk. *Int. Dairy J.* 19, 655–660.
- Sevi, A., Albenzio, M., Marino, R., Santillo, A., Muscio, A., 2004. Effects of lambing season and stage of lactation on ewe milk quality. *Small Rumin. Res.* 51, 251–259.
- Silanikove, N., 2016. Transcellular route as the most probable explanation for the presence of plasminogen in mammal's milk. *J. Theor. Biol.* 395, 221–226.
- Silanikove, N., Shapiro, F., Merin, U., and Leitner, G., 2013. Tissue-type plasminogen activator and plasminogen embedded in casein rule its degradation under physiological situations: manipulation with casein hydrolysate. *J. Dairy Res.* 80, 227–232.
- Sousa, M.J., Ardö, Y., McSweeney, P.L.H., 2001. Advances in the study of proteolysis during cheese ripening. *Int. Dairy J.* 11, 327–345.
- Srinivasan, M., Lucey, J.A., 2002. Effects of added plasmin on the formation and rheological properties of rennet-induced skim milk gels. *J. Dairy Sci.* 85, 1070–1078.
- Theodorou, G., Kominakis, a, Rogdakis, E., Politis, I., 2007. Factors affecting the plasmin-plasminogen system in milk obtained from three Greek dairy sheep breeds with major differences in milk production capacity. *J. Dairy Sci.* 90, 3263–9.
- White J.H., Zavizion, B., O'Harea, K., Gilmore, I., Guoa, M.R., Kindstedt, P. and Politis, I., 1995. Distribution of plasminogen activator in different fractions of bovine milk. *J. Dairy Res.* 62 115–122.

Table 1. Plasmin activity (least squares means) as determined by stage of lactation (SL) and parity (PAR).

	EARLY						SEM	P		
	LACT.		MID LACT.		LATE LACT.			SL	PAR	SL x PAR
	MULT		MULT		MULT					
	PRIM	I	PRIM	I	PRIM	I				
PL (u/ml)	0.76	1.69 ^a	0.71	1.08 ^{ab}	0.46	0.83 ^b	0.070	0.313	0.076	0.692

Within each PAR group, means without a common superscript are statistically different at $P < 0.05$

Table 2. Contrast analysis comparing plasmin activity among different stages of lactation (SL) in primiparous and multiparous ewes.

	Early×Mid	Early×Late	Mid×Late
Primiparous	0.935	0.640	0.699
Multiparous	0.097	0.036*	0.545

An asterisk indicates significant differences at $P < 0.05$

Table 3. Plasmin activity (least squares means) according to somatic cell count (SCC) and udder health status (UHS).

	LOW SCC			MID SCC			HIGH SCC			SEM	P		
	PR	M1	M2	PR	M1	M2	PR	M1	M2		SCC	UHS	SCC x UHS
PL	0.76	0.90	1.44	0.6	0.9	1.0	0.64	1.23	1.29	0.07	0.00		
(u/ml)	a	ab	b	1	4	7	a	b	b	0	0.562	0.761	
											3		

Within each SCC category, means without a common superscript are statistically different at $P < 0.05$

PR = primiparous ewes.

M1 = multiparous ewes with no damage of the udder in the previous lactation.

M2 = multiparous ewes with udder damage in the previous lactation.

Table 4. Contrast analysis comparing plasmin activity throughout different somatic cell count (SCC) ranges in all udder health status (UHS) categories.

	Low×Mid	Low×High	Mid×High
PR	0.628	0.747	0.925
M1	0.913	0.320	0.430
M2	0.151	0.517	0.376

An asterisk indicates significant differences at $P < 0.05$

PR = primiparous ewes.

M1 = multiparous ewes with no damage of the udder in the previous lactation.

M2 = multiparous ewes with udder damage in the previous lactation.

Table 5. Correlation between plasmin activity and milk composition.

	PR	M1	M2
Fat	-0.1863	-0.0789	-0.1274
Crude protein	-0.2158*	-0.0056	-0.1593*
Total solids	-0.2054	-0.0942	-0.1527*
Lactose	0.1752	0.061	0.0979
Casein	-0.2539*	-0.4333***	-0.4442***
Ash	-0.1127	0.3237**	0.1024***
Urea	-0.0389	0.1674	0.0871
Native pH	0.3845*	0.1974	0.4968

* P<0.05 ; ** P<0.01 ; *** P<0.001

PR = Primiparous ewes.

M1 = multiparous ewes with no damage of the udder in the previous lactation.

M2 = multiparous ewes with udder damage in the previous lactation.

Table 6. Correlation between plasmin activity and rennet coagulation variables.

	PR	M1	M2
RCT	0.0914	0.0296	0.1685*
k ₂₀	-0.0857	-0.0073	0.137
RCT + k ₂₀	0.0587	0.0223	0.1928*
A ₃₀	-0.1907	-0.0037	-0.1731*
A ₆₀	-0.0136	-0.3006**	-0.214**
Curd yield	-0.2785**	0.26*	0.0429

* P<0.05 ; ** P<0.01 ; *** P<0.001

PR = Primiparous ewes.

M1 = multiparous ewes with no damage of the udder in the previous lactation.

M2 = multiparous ewes with udder damage in the previous lactation.