



Virginia Commonwealth University
VCU Scholars Compass

Hepatobiliary Cancers: Pathobiology and
Translational Advances

Dept. of Pathology

2017

Yap Functions Via TEAD Mediated Transcriptional Activation in Hepatoblastoma Pathogenesis


junyan tao 8598903
junyantao2010@gmail.com

pin liu
ucsf, Pin.Liu@ucsf.edu

jie zhang
ucsf, Jie.Zhang@ucsf.edu

See next page for additional authors

Follow this and additional works at: http://scholarscompass.vcu.edu/hepa_cancers

 Part of the [Cancer Biology Commons](#), and the [Medicine and Health Sciences Commons](#)

© The Author(s)

Downloaded from

http://scholarscompass.vcu.edu/hepa_cancers/2

This Abstract Accepted for Presentation is brought to you for free and open access by the Dept. of Pathology at VCU Scholars Compass. It has been accepted for inclusion in Hepatobiliary Cancers: Pathobiology and Translational Advances by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

Authors

junyan tao 8598903, pin liu, jie zhang, sucha singh, xin chen, Na zhan, and Satdarshan (Paul) Singh Monga

Yap Functions Via TEAD Mediated Transcriptional Activation in Hepatoblastoma Pathogenesis

Pin Liu^{1,2}, Junyan Tao^{3,4,5}, Jie Zhang^{1,2}, Sucha Singh^{3,4,5}, Na Zhan^{3,4,6}, Satdarshan P. S. Monga^{3,4,5}, Xin Chen^{1,2}

¹Department of Bioengineering and Therapeutic Sciences and ²Liver Center, University California, San Francisco, CA; ³Department of Pathology, ⁴Pittsburgh Liver Research Center, and ⁵University of Pittsburgh, School of Medicine and University of Pittsburgh Medical Center, Pittsburgh, PA, USA; ⁶Department of pathology, Renmin Hospital of Wuhan University, Wuhan, China.

Hepatoblastoma (HB) is the most common pediatric liver tumor. Activation of Wnt/ β -catenin pathway is considered to be the driver genetic event in HB. In addition to Wnt/ β -catenin, activation of Yap is found in a majority of HBs, and high Yap transcriptional activity was recently found to be associated with high risk HB. Importantly, our studies demonstrated the coordinated activation of β -catenin and Yap led to HB formation in mice. However, how Yap functions to promote HB formation remains unknown. Yap has been shown to have transcriptional activity as well as non-transcriptional based functions. Here we investigated whether TEAD mediated transcription is the major mechanism by which Yap promotes HB formation. Specifically, we blocked Yap transcriptional activity via dominant negative form of TEAD2 (dnTEAD2). We found that overexpression of dnTEAD2 inhibited HB cell growth in culture and blocked Yap/ β -catenin induced HB formation in mice. We generated TEAD2-VP16 fusion construct which allows TEAD dependent transcriptional activation independent of Yap. When TEAD2-VP16 was co-expressed with activated β -catenin into mice, it led to HB formation. Further analysis demonstrated that TEAD2-VP16/ β -catenin HBs shared similar molecular features with Yap/ β -catenin tumors. As there are multiple TEADs (TEAD1, 2, 3 and 4) which share redundant functions, we tested the contribution of each TEAD in regulating Yap activity in HB cells. We found that silencing of TEAD4 consistently reduced Yap downstream gene expression, supporting that TEAD4 may be the major Yap partner in HB development. In summary, our studies support that Yap predominantly regulates HB development via TEAD4 mediated transcriptional activation,

and blocking Yap/TEAD interaction may represent a novel therapeutic strategy against HB.

Keywords: Hepatoblastoma (HB); Yap; TEAD2-VP16; TEAD4