



Title	Ice core records of monoterpene- and isoprene-SOA tracers from Aurora Peak in Alaska since 1660s: Implication for climate change variability in the North Pacific Rim
Author(s)	Pokhrel, Ambarish; Kawamura, Kimitaka; Ono, Kaori; Seki, Osamu; Fu, Pingqing; Matoba, Sumio; Shiraiwa, Takayuki
Citation	Atmospheric environment, 130, 105-112 https://doi.org/10.1016/j.atmosenv.2015.09.063
Issue Date	2015-09-28
Doc URL	http://hdl.handle.net/2115/67195
Rights	© 2015. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	Pokhrel.pdf



[Instructions for use](#)



Title	Ice core records of monoterpene- and isoprene-SOA tracers from Aurora Peak in Alaska since 1660s: Implication for climate change variability in the North Pacific Rim
Author(s)	Pokhrel, Ambarish; Kawamura, Kimitaka; Ono, Kaori; Seki, Osamu; Fu, Pingqing; Matoba, Sumio; Shiraiwa, Takayuki
Citation	Atmospheric environment, 130: 105-112
Issue Date	2015-09-28
Doc URL	http://hdl.handle.net/2115/67195
Right	© 2015. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	Pokhrel et al., AE -.pdf



[Instructions for use](#)

1 Revised to Atmospheric Environment (Special Issue: Chemical Characterization of SOA), accepted

2

3 **Ice core records of monoterpene- and isoprene-SOA tracers from Aurora Peak in Alaska**
4 **since 1660s: Implication for climate change variability in the North Pacific Rim**

5

6 Ambarish Pokhrel^{1,2}, Kimitaka Kawamura^{1,*}, Kaori Ono¹, Osamu Seki¹, Pingqing Fu^{1,3},

7 Sumio Matoba¹, Takayuki Shiraiwa¹

8

9

10 ¹ Institute of Low Temperature Science, Hokkaido University, Sapporo, Japan

11 ² Graduate School of Environmental Science, Hokkaido University, Sapporo, Japan

12 ³ LAPC, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029,
13 China

14

15 *Corresponding author

16 Email address: kawamura@lowtem.hokudai.ac.jp (K. Kawamura)

17

18

19 **Abstract**

20 Monoterpene and isoprene secondary organic aerosol (SOA) tracers are reported for the
21 first time in an Alaskan ice core to better understand the biological source strength before and
22 after the industrial revolution in the Northern Hemisphere. We found significantly high
23 concentrations of monoterpene- and isoprene-SOA tracers (e.g., pinic, pinonic, and 2-
24 methylglyceric acids, 2-methylthreitol and 2-methylerythritol) in the ice core, which show
25 historical trends with good correlation to each other since 1660s. They show positive
26 correlations with sugar compounds (e.g., mannitol, fructose, glucose, inositol and sucrose), and
27 anti-correlations with α -dicarbonyls (glyoxal and methylglyoxal) and fatty acids (e.g., C_{18:1}) in
28 the same ice core. These results suggest similar sources and transport pathways for
29 monoterpene- and isoprene-SOA tracers. In addition, we found that concentrations of C₅-
30 alkene triols (e.g., 3-methyl-2,3,4-trihydroxy-1-butene, *cis*-2-methyl 1,3,4-trihydroxy-1-butene
31 and *trans*-2-methyl-1,3,4-trihydroxy-1-butene) in the ice core have increased after the Great
32 Pacific Climate Shift (late 1970s). They show positive correlations with α -dicarbonyls and
33 fatty acids (e.g., C_{18:1}) in the ice core, suggesting that enhanced oceanic emissions of biogenic
34 organic compounds through the marine boundary layer are recorded in the ice core from
35 Alaska. Photochemical oxidation process for these monoterpene- and isoprene-/sesquiterpene-
36 SOA tracers are suggested to be linked with the periodicity of multi-decadal climate
37 oscillations and retreat of sea ice in the Northern Hemisphere.

38

39 Keywords: Monoterpenes, isoprene, SOA tracers, ice core, Alaska, multi-decadal climate
40 oscillation, climate change

41 1. Introduction

42 Organic aerosols (OA) are an important fraction of atmospheric fine particles. They can
43 alter the earth radiation budget directly by scattering sunlight, and indirectly by acting as cloud
44 condensation nuclei (CCN) and ice nuclei (IN) (*Aalto et al., 2001; Kanakidou et al., 2005*). OA
45 are derived from anthropogenic and biogenic sources via primary emissions and secondary
46 photochemical oxidation of various precursors (*Claeys et al., 2004; Guenther et al., 2006;*
47 *Kunwar and Kawamura 2014a, b*). The former can be classified as primary organic aerosols
48 (POA) whereas the latter as secondary organic aerosols (SOA). POA are emitted from plant
49 debris, fungal spore, fossil fuel combustion, biomass burning, and soil particles (*Guenther et*
50 *al., 2006; Fu et al., 2014*) whereas SOA are photochemically produced by homogeneous
51 (*Claeys et al., 2004*) and heterogeneous (*Limbeck et al., 2003*) oxidations of biogenic volatile
52 organic compounds (BVOCs) and anthropogenic VOCs.

53 Globally, annual emissions of BVOCs are estimated to be 1150 TgC/yr, where isoprene
54 (C_5H_8) and monoterpenes ($C_{10}H_{16}$) consist 44%, and 11%, respectively. In contrast, emissions
55 of anthropogenic VOCs (110 TgC/yr) are one order of magnitude smaller than those of BVOC;
56 contribution of aromatic hydrocarbon is estimated to be only 13% of total anthropogenic VOCs
57 (*Guenther et al. 1995, references therein*). Particularly, terrestrial and oceanic phytoplankton's
58 isoprene is the largest source of VOCs, contributing ~50% of total global emission of 309–706
59 TgC/yr (*Acosta Navarro et al., 2014, references therein*). In contrast, α -/ β -pinenes can
60 contribute 35% of total global emissions of BVOCs (*Griffin et al., 1999*). These BVOCs could
61 contribute significantly to the formation of SOA in the troposphere.

62 Isoprene- and monoterpene-oxidation products have been studied using field- (*Claeys*
63 *et al., 2004; Fu et al., 2011*) and laboratory-based measurements (*Claeys et al., 2007;*
64 *Hallquist et al., 2009; Surratt et al., 2010; Noziere et al., 2015*). Being different from the
65 previous assumption made by Limbeck et al. (2003), Claeys et al. (2004) first reported the

66 presence of 2-methyltetrols in the Amazonian aerosols as oxidation products of isoprene.
67 Based upon the laboratory work, the formation of SOA from isoprene is further confirmed
68 (*Surratt et al., 2010*). Laboratory experiments showed that the oxidation products of isoprene
69 (e.g., C₅-alkene triols and 2-methyltetrols) are enhanced in the presence of NO_x, SO₂ and/or
70 H₂SO₄ (*Surratt et al., 2006, 2010*). Moreover, in-cloud aqueous phase oxidation of isoprene is
71 suggested to significantly contribute to the SOA formation (*Carlton et al., 2009; Volkamer et*
72 *al., 2009; Hallquist et al., 2009*). Thus, these studies (i.e. field and laboratory measurement)
73 provide insight on the sources, spatial and temporal distributions and molecular evolution of
74 SOA.

75 McNeill et al. (2012) reported that organic compounds in snow/glaciers and/or in ice
76 particles could be biological in origin and deposited from the atmosphere. Isoprene- and
77 monoterpene-oxidation products are rarely studied in ice and snow, although biomass burning
78 products and photooxidation products (i.e., dicarboxylic acids) of various organic precursors
79 have been reported (*Kawamura et al., 2001; 2012a,b*). However, studies of monoterpene- and
80 isoprene-SOA in snow and glacier ice remain poorly explored.

81 Here, we studied monoterpene- and isoprene-SOA tracers in an ice core from Alaska.
82 Our objectives are to obtain the temporal profiles of monoterpene- and isoprene-SOA tracers
83 that are derived from atmospheric photooxidation of isoprene and α/β -pinenes to better
84 understand the historical changes in biological source strength of BVOCs. In this study, we
85 report for the first time monoterpene- and isoprene-SOA tracers from mountain glaciers in
86 Alaska to better understand both POA and SOA cycles in the North Pacific Rim.

87 88 **2. Samples and Analytical Procedures**

89 *2.1. Site description*

90 We collected a ~180 m deep ice core on the saddle of Aurora Peak in 2008 AD, which
91 is located in the southeast of Fairbanks (63.52°N; 146.54°W, elevation: 2,825 m) (Figure 1).

92 The ice core sample was cut into ~50 cm-long pieces and directly transported to the laboratory
93 of the Institute of Low Temperature Science (ILTS), Sapporo, Japan and stored in a dark cold
94 room at -20°C until analysis. The ice core ages were determined by annual counting of
95 hydrogen isotopes (δD) and Na^+ seasonal cycles and the age control was provided by reference
96 horizons of tritium peaks in 1963 and 1964 (*Tsushima et al., 2015*). The bottom of the sample
97 was estimated to be 343 years old, i.e., 1665 AD (*Tsushima et al., 2015*). 10-day air mass
98 backward trajectory based on Lagrangian tracking method for 1992 – 2002 has been already
99 reported for Alaskan regions in the troposphere (>300 hPa) (*Yasunari and Yamazaki, 2009*).

100 2.2. Chemical analysis

101 Total numbers of ice core sections (50 cm long) were 147, which means that sampling
102 frequency is ~40% of the 180 m deep ice core. This sampling frequency is much better than
103 that (8%) of the previous ice core studies in Greenland (*Kawamura et al., 2001*). Roughly one
104 quarter of each section was used after removing the surface 5-10 mm of the ice section. These
105 ice core samples (ca. 150 mL) were first molten and then transferred to a pear shape flask (300
106 ml) and concentrated to almost dryness using a rotary evaporator under a vacuum. The total
107 dissolved and particulate organic matters were extracted with a mixture of CH_2Cl_2/CH_3OH
108 (2:1) using ultrasonic bath. The extracts were transferred to a 1.5 mL glass vial, dried under
109 nitrogen stream and then derivatized with 99% N, O-bis-(trimethylsilyl) trifluoroacetamide
110 (BSTFA) + 1% trimethylchlorosilane (TMCS) and pyridine at 70 °C for three hours (*Fu et al.,*
111 *2010a, b; Kawamura et al., 2012b; Zhu et al., 2015*).

112 The derivatized fractions were diluted with n-hexane containing an internal standard (*n*-
113 C_{13} alkane) prior to the analysis with a gas chromatograph/mass spectrometer (GC/MS)
114 installed with a capillary column (HP-5MS, 30 m \times 0.32 mm I.D. \times 0.25 μm film thickness)
115 and split/splitless injector (*Kawamura et al., 2012b*). Monoterpene- and isoprene-SOA tracers
116 (Table 1) were determined using GC/MS method which has been reported elsewhere (*Fu et al.,*

117 2014). Triplicate analysis of real ice core sample was also conducted. Analytical errors were
118 less than 8%. Laboratory blank using pure water (150 ml), i.e., Milli-Q water, showed blank
119 levels of less than 3% of real ice core samples. All the data reported here are corrected for the
120 laboratory blank. We believe that the preservation conditions of organic compounds in the ice
121 core is fairly good and thus post-depositional microbial changes in the signals are not
122 important because no ice lense was found in the ice core studied. However, we cannot exclude
123 the possibility that some oxidants deposited in snow may react with organic species, which
124 may decompose and/or produce some organics.

125

126 3. Results and Discussion

127 3.1. Historical changes of monoterpene-SOA tracers

128 Table 1 presents ice core concentrations of monoterpene- and isoprene-SOA tracers
129 since 1660s. Pinic and pinonic acids are detected as α/β -pinene oxidation products. These
130 acids are observed in smog chamber experiments (*Glasius et al., 2000*), and are produced from
131 photooxidation of α/β -pinene in the presence of O_3 and OH radicals (*Hoffman et al., 1997*).
132 Pinic acid showed sporadic peaks in 1750 (concentration, 427 ng/L), 1786 (529), 1870 (406),
133 1875 (484), 1875, 1880 (3001), 1913 (630), 1973 (853), 1993 (352), 1998 (295) and 2005
134 (430) compared to its average concentration (157 ± 148 ng/L). Similar spikes were detected for
135 pinonic acid, except for 1980s and 1990s (Figure 2a and b). The periods for the peaks can be
136 considered as significant monoterpene oxidation periods corresponding to the formation of
137 SOA in the North Pacific region.

138 Interestingly, average concentration of pinic acid (157 ng/L) is more than double of that
139 of pinonic acid (70.6 ng/L), which contradicts the trend reported for aerosol studies from Mt.
140 Tai, East China (*Fu et al., 2010a*), tropical India (*Fu et al., 2010b*), central Greece (*Kavouras*
141 *et al., 1999*), Sierra Nevada Mountains of California (*Cahill et al., 2006*) and North Carolina

142 (*Bhat et al., 2007*). It should be noted that vapor pressure of pinonic acid is twice higher than
143 that of pinic acid (*Fu et al., 2010b*). Coniferous forest is common in southern Alaska, being
144 similar to Research Triangle Park, USA and Germany (e.g., *Kleindienst et al., 2007; Plewka et*
145 *al., 2006*). A good correlation between pinonic and pinic acids ($R = 0.93$) in the ice core
146 indicates a similar sources and formation pathways and/or similar atmospheric fate in southern
147 Alaska for 1660s-1980s (Figure 4a). Interestingly, these compounds show a decrease during
148 the 1800-1850s and after the 1920s (except for 1973) with sporadic peaks during the 1990s,
149 suggesting that source and/or source region could be shifted to this sampling site.

150 We found significant levels of 3-hydroxyglutaric acid (3-HGA), whose historical
151 trends are completely different (Figure 2c) with pinic and pinonic acids. Smog chamber
152 experiment shows that 3-HGA is an oxidation product of α -pinene under UV irradiation in the
153 presence of NO_x (*Claeys et al., 2007*). Average concentration of 3-HGA (22.4 ± 41 ng/L) is
154 several times lower than pinic and pinonic acids (Table1). It should be noted that we didn't
155 detect any β -caryophyllinic acid and related compounds. For instance, many aerosol studies
156 showed higher concentrations of β -caryophyllinic acid in late winter (e.g., *Fu et al., 2010b*).
157 Correlations (R) of 3-HGA with pinic and pinonic acids are 0.46 and 0.46, respectively. These
158 results suggest that 3-HGA involves other processes than pinic and pinonic acids over the
159 southern Alaska.

160 3.2. Historical changes of isoprene-SOA tracers

161 We detected six isoprene-SOA tracers (Table 1 and Figure 3). Isoprene is readily
162 oxidized in the atmosphere by OH, O_3 , NO_x , SO_2 and H_2SO_4 (*Carlton et al, 2009; Surratt et al.,*
163 *2010*). There are many key oxidation products of isoprene via methacrolein (a key as an
164 intermediate in SOA formation) in ambient aerosol has been found (*Noziere et al., 2015;*
165 *Hallquist et al., 2009 and references therein*). Some isoprene oxidation products have less
166 vapor pressures and thus stay in particle phase, leading to the formation of SOA (*Hoffmann et*

167 *al.*, 1997). 2-Methylglyceric acid can be formed by the oxidation of methacrolein and
168 methacrylic acid from two major gas-phase oxidation products of isoprene (*Surrat et al.*, 2007;
169 *Hu et al.*, 2008), which can be altered and/or associated with land sea breeze atmospheric
170 circulation (*Claeys et al.*, 2007; *Fu et al.*, 2010b and references therein).

171 2-Methylglyceric acid showed higher concentrations in 1750 (74.8 ng/L), 1786 (175),
172 1870 (83.4), 1875 (93.1), 1880 (73.1), 1913 (166), 1973 (144), 1977 (67.3), 1993 (230), 1998
173 (65.9), 1999 (71.8) and 2005 (282), whose concentrations are much higher than its average
174 (35.6±48 ng/L). These periods are consistent with those of 2-methylthreitol (ave. 349±383
175 ng/L) and 2-methylerythritol (692±709 ng/L) as shown in Figure 3a, b and c. Interestingly,
176 historical trends are similar to those of pinic and pinonic acids and isoprene-SOA tracers
177 (Figure 2a, b, and 3a, b, c) before 1990s. Correlation coefficients (R) of pinonic and pinic acids
178 with 2-methylglyceric acid are 0.71 and 0.83, respectively whereas that of 2-methylglyceric
179 acid and 3-HGA is 0.67 (Figure 4b, c and d). Correlation coefficients of pinonic and pinic
180 acids with 2-methylthreitol are 0.73 and 0.77, respectively (Figure 5a, b). In addition, those of
181 pinonic and pinic acids with 2-methylerythritol are 0.71 and 0.75 (Figure 5c, d) and correlation
182 coefficients (R) of 2-methylglyceric with 2-methylthreitol and 2-methylerythritol are 0.82 and
183 0.67, respectively (not shown in figure).

184 The source strength of these SOA tracers is somewhat higher before the industrial
185 revolution than in recent years in the southern Alaskan ice core (Figure 3a-c). It should be
186 noted that 3-methyl-1,2,3-butanetricarboxylic acid (MBTCA) was reported during smog
187 chamber experiments of α -pinene with UV irradiation in the presence of NO_x (*Claeys et al.*,
188 2007). But we didn't detect MBTCA from the 1660's till 2008, suggesting that anthropogenic
189 contributions are insignificant for the last 3 centuries. In contrast, C₅-alkene triols showed
190 similar historical trends each other (Figure 3d,e and f), with somewhat different trends to other
191 SOA tracers (e.g., Figure 3a, b and c). Particularly, 3-methyl-2,3,4-trihydroxy-1-butene

192 showed higher concentrations in 1786 (13.8 ng/L), 1947 (12.0), 1973 (30.2), 1989 (41.1), 1993
193 (93.5), 1998 (23.0), 1999 (31.7), 2004 (35.1), 2005 (140.7) and 2006 (46.4) compared to its
194 average concentration (6.99 ± 17 ng/L-ice). These periods are also consistent with those for the
195 reaction products of *cis* and *trans* hydroxy alkenes (e.g., Figure 3d, e and f). Correlation
196 coefficients (R) of 3-methyl-2,3,4-trihydroxy-1-butene with pinonic and pinic acids in the ice
197 core are 0.51 and 0.63, whereas those of *cis* 2-methyl-1,3,4-trihydroxy-1-butene with pinonic
198 and pinic acids are 0.49 and 0.52, respectively.

199 Moreover, correlations of 2-methylglyceric acid with 3-methyl-2,3,4-trihydroxy-1-
200 butene, *trans*-2-methyl-1,3,4-trihydroxy-1-butene and *cis* 2-methyl 1,3,4-trihydroxy-1-butene
201 are 0.83, 0.77 and 0.75, respectively. Correlations of 2-methylthreitol to 2-methylerythritol, 3-
202 methyl-2,3,4-trihydroxy-1-butene, *cis* 2-methyl 1,3,4-trihydroxy-1-butene and *trans*-2-methyl-
203 1,3,4-trihydroxy-1-butene are positive (R=0.87, 0.65, 0.54 and 0.57, respectively). Hence, we
204 consider that atmospheric fate could be strongly involved with local or regional meteorological
205 conditions with an insignificant influence of anthropogenic activities. The most prevalent SOA
206 precursors on a global scale are terpenoids emitted from vegetations (*Claeys et al., 2007*;
207 *Faiola et. al., 2014*). It should be noted that more than 100 organic compounds including many
208 numbers of SOA precursors are emitted from leaf litter and/or soil, although they are less
209 significant than direct emission from plant leaves (*Faiola et. al., 2014*). SOA formation in the
210 forest environment (i.e., from leaf litter and/or forest soil) is also important in spring to fall
211 (*Faiola et. al., 2014*). For instance, rate of particle formation was high during spring from soil
212 and leaf litter in a boreal forest of southern Finland (*Makela et al., 2000*; *Bigg et al., 2004*). In
213 addition, emissions of SOA precursors (i.e., terpenoids) are high during late spring and autumn
214 in boreal pine forest floor (e.g., *Aaltonen et al., 2011 and references therein*). These results
215 suggest the additional sources for higher spikes in an ice core.

216 *3.3. Responses of monoterpene- and isoprene-SOA tracers to tropospheric temperature*

217 We found that monoterpene- (Figure 2a-b) and isoprene-SOA tracers (Figure 3a-c)
218 inversely correlate with marine biogenic tracers, i.e., even carbon numbered fatty acids (e.g.,
219 $C_{16:0}$, $C_{18:0}$ and $C_{18:1}$) (Pokhrel *et al.*, 2015). In contrast, those SOA tracers show a positive
220 correlation with sugar compounds (e.g., mannitol, fructose, glucose, inositol and sucrose) from
221 the same ice core, suggesting the continental sources (unpublished data, Pokhrel and
222 Kawamura, 2015). Hence, emissions of isoprene and monoterpenes from vegetations could be
223 controlled by biological activity of the ocean and continents in different years, which could be
224 linked to historical variations of thermohaline circulation (Boyce *et al.*, 2010 and references
225 therein).

226 Reconstructed air temperatures from the Gulf of Alaska (GOA) using tree ring showed
227 a sharp drop in June to September since 1800 to 1875 except for 1830s (e.g., Wilson *et al.*,
228 2007b), whose historical trend is similar to lower concentrations of monoterpene- and
229 isoprene-SOA tracers (Figure 2a-b and Figure 3a-c). It should be noted that sampling sites for
230 the tree ring data are located close to coastal range of the GOA and/or south of the Brooks
231 Range at about 68 °N. This GOA temperature expresses a strong consistency with multi-
232 decadal climate oscillation and short-term atmospheric activities (e.g., ENSO/El-Nino and/or
233 Aleutian Low). Interestingly, reconstructed Northern Hemisphere annual temperature since
234 1671 based on high-latitude tree-ring data from North America (Jacoby and D'Arrigo, 1989) is
235 also similar historical trend (except for few decades). In addition, tree-ring growth data and
236 inferred temperature variability at the North American Arctic tree line (D'Arrigo *et al.*, 2009)
237 are also consistent with the trends of ice core SOA tracers.

238 Based on 10 day backward trajectory analyses for the years of 1992-2002, Yasunari
239 and Yamazaki (2009) reported that Alaskan regions can receive significant air parcels from
240 adjacent North Pacific Regions, East Asia, Eastern Russia, Siberia, the Okhotsk and Bering
241 Seas, higher latitudes of Alaskan regions, the Gulf of Alaska, Japan, Canada, the Arctic Ocean

242 and Europe in the troposphere (>300 hPa). These areas are represented by extra tropical
243 Northern Hemispheric temperature (ENHT). ENHT is the robust temperature trends for the
244 Northern Hemisphere, which is calculated with more than 25 different sampling sites from tree
245 ring proxy records of the world. ENHT shows the positive correlations (R) with the ambient
246 temperatures in European Alps (0.67), western Siberia (0.61), Mongolia (0.70), Nepal (0.49),
247 Northern Yukon (0.60), Wrangell Mountains (0.60), British Columbia (0.77), Idaho (0.41) and
248 Northern Quebec (0.42) (e.g., *Wilson et al., 2007a*).

249 Lower concentrations of monoterpene- and isoprene-SOA tracers (Figures 2 and 3)
250 during this period are further supported by historical trend of ENHT (Figure 3h). For instance,
251 decreasing trends for all these SOA tracers showed positive correlations with decreasing
252 historical trends of ENHT during 1800-1860. Particularly, correlation coefficients (R) of each
253 11 points running mean (11-RM) of pinonic, pinic and 3-HGA with 11-RM of ENHT are 0.84,
254 0.86 and 0.54, respectively. Similarly, correlation coefficients (R) of 11-RM of each isoprene-
255 SOA tracers (e.g., Figure 3a - f) with 11-RM of ENHT are 0.84, 0.95, 0.94, 0.82, 0.89, and
256 0.67, respectively. It should be noted that NH temperature departure and reconstructed solar
257 irradiance are well correlated with many climate periodicity and non-periodicity cycles in the
258 NH, which are reported elsewhere (*Wang et al., 2012; Lean et al., 2005 and references*
259 *therein*).

260 Aleutian Low (AL) can easily alter air parcel flow and heat exchange between the
261 Arctic and mid-latitudinal region (*Wang et al., 2012*). *Mantua et al. (1997)* pointed out that the
262 AL was significantly correlated with the Pacific Decadal Oscillation (PDO), which can be
263 described as a long-lived El Niño-like pattern of Pacific climate variability. For instance, the
264 warm phase of PDO corresponds to the strong AL, while the cold phase of PDO corresponds to
265 the weak AL. The AL would have mixed continental and marine air mass, and more drive
266 prevailing winds from the northwestern Pacific regions to northeastward in North America,

267 which can intensify the positive North Pacific Index (NPI) over American west (*Trouet and*
268 *Taylor, 2010*). It represents the region of 30°N-65°N, 160°E-140°W for decadal variations in
269 the atmospheric circulation (<https://climatedataguide.ucar.edu>). Moreover, on a decadal scale,
270 the AL has been consistently strong and drifted eastward since 1970s (*Wang et al, 2012*).

271 These monoterpene- and isoprene-SOA tracers (Figure 2a-b and Figure 3a-c,
272 respectively) showed decreased trends since 1920s, which are similar to the trend of NPI.
273 Historical trends of C₅-alkene triols (Figure 3d-f) showed anti-correlation with those of other
274 compounds. Particularly, large-scale atmospheric circulation (or relatively local) is associated
275 with the AL system in the Bering Sea (or Alaskan regions) (*Sasaki and Minobe, 2005*). If this
276 is the case, it is important to understand the historical variations of atmospheric circulation
277 over the Bering Sea, Gulf of Alaska and Alaskan regions. The climatic oscillations can
278 represent the atmospheric transport in the lower latitudes. The effect of climatic oscillations
279 can be observed using NPI (*Trouet and Taylor, 2010*). For instance, relations between 7-RM of
280 C₅-alkene triols, i.e., 3-methyl-2,3,4-trihydroxy-1-butene, cis 2-methyl 1,3,4-trihydroxy-1-
281 butene and trans-2-methyl-1,3,4-trihydroxy-1-butene (Figure 3d-f), anti-correlate with 7-RM
282 of NPI since 1899-2007 (R = 0.31, 0.32 and 0.29, respectively). These results suggest that
283 decadal variability (or atmospheric stability) may be involved with the alteration in the
284 concentrations of SOA tracers.

285 *Tsushima et al. (2015)* reported that concentrations of Na⁺ in the same ice core
286 increased after the 1970s. This increase could be resulted from changes in the position and/or
287 strength of winter storms in the Gulf of Alaska, which is associated with the PDO shift in 1976.
288 We obtained higher concentrations of all the compounds around 1975 (Figures 2 and 3), being
289 similar to Na⁺ cycle of same ice core, rapid change of sea ice concentrations (SIC) and
290 temporal wind pattern (TW) over the Bering Sea (*Sasaki and Minobe, 2005*). Moreover,
291 historical trends C₅-alkene triols (Figure 3d-f) are similar to changes of SIC and TW over the

292 Bering Sea for spring season after 1950s. It should be noted that SOA formation in forest
293 environment is important during spring and fall season compared to summer season (*Faiola et*
294 *al., 2014*). *Makela et al.* (2000) and *Bigg et al.* (2004) reported that particle formation rate was
295 high during spring season in the boreal forest of southern Finland. SOA precursors and/or
296 terpenoids are also high during late spring and autumn in boreal pine forest floor (*Aaltonen et*
297 *al., 2011 and references therein*).

298 In contrast, all other compounds in the ice core (Figure 2a-c and 3a-c) exhibit similar
299 trends with wintertime SIC and TW over the Bering Sea during the same periods. These results
300 indicate that different pattern of wind circulations can alter the temporal trends of these
301 compounds in ice core, further suggesting that the first case is linked with local circulation
302 over the North Pacific Rim compared to regional wind circulation for the latter case. These
303 results may suggest that climatic variations could be recorded in the ice core profile of SOA
304 tracers. Similarly, we can explain another process for variable concentrations of SOA tracers
305 since 1660s. For instance, we found lower concentrations of SOA-tracers around 1650-1700s
306 and 1800-1850s, which are known as cooling period of Maunder and Dalton minima and a part
307 of Little Ice Age (Figure 3h, i), respectively (*Krivova, et al., 2010*), suggesting lower emission
308 of isoprene and monoterpenes during the cold periods. Our ice core records of SOA tracers can
309 be further supported by air-sea and oceanic heat fluxes of the northern Gulf of Alaska (*Janout*
310 *et al., 2013*). Net heat flux and/or heat flux anomalies in winter are significantly increased
311 during 1975-2010 (except around 1998) compared to 1950-1975 (*Janout et al., 2013*), which is
312 consistent with historical trends of C₅-alkene triols.

313 These considerations suggest that concentrations of SOA tracers are controlled by
314 temperature, pressure and wind field of the Gulf of Alaska and/or Northern North Pacific Rim.
315 These three parameters alter the atmospheric transport (e.g., the AL can alter the atmospheric
316 transport), which are correlated with ENHT and NPI elsewhere (NPI has negative correlation

317 with C₅-alkenes) (*Wilson et al., 2007a,b; Wang et al., 2012 and references therein*). We found
318 that C₅-alkene triols positively correlate with ENHT during the 1750s to 1980s. Particularly,
319 correlation coefficients (R) between 5 points running mean (5-RM) of C₅-alkene triols (Figure
320 3d, e and f) with ENHT are 0.73, 0.80 and 0.78 for individual species (3-methyl-2,3,4-
321 trihydroxy-1-butene, cis 2-methyl 1,3,4-trihydroxy-1-butene and trans-2-methyl-1,3,4-
322 trihydroxy-1-butene), respectively. It is likely that BVOC emissions could be associated with
323 type and density of vegetation in landscape, temperature and radiation, having a significant
324 impact on SOA formation (*Pokhrel, 2015*).

325 In addition, C₅-alkene triols have increasing trends of the concentrations since 1920s,
326 which is similar to sea surface temperature (SST) over the past century (*Boyce et al., 2010 and*
327 *references therein*). Hence, these increasing trends of C₅-alkene triols (see Figure 3d-f) could
328 be involved with climatic and oceanographic variability. Particularly, an increased SST over
329 the past century since 1920s (except for few points) is similar to historical trends of SOA
330 tracers, suggesting that an enhanced ocean warming could cause a restructuring of marine
331 ecosystems and/or likely ocean circulation (*Boyce et al., 2010; and references therein*),
332 followed by intensified production of SOA in southern Alaska. In the same ways, we may
333 explain other historical trends since 1660s, which could be altered by complex mechanisms of
334 thermohaline circulation and productivity of marine biota, which are reported elsewhere (*Boyce*
335 *et al., 2010; and references therein*).

336 Moreover, C₅-alkene triols have positive correlations with α -dicarbonyls (i.e., glyoxal
337 and methylglyoxal) and even-carbon numbered low molecular weight fatty acids (e.g., C_{16:0}
338 and C_{18:0}) including unsaturated fatty acids (C_{18:1}), which are originated from marine
339 phytoplankton (*Pokhrel et al., 2015*). Increased historical trends of C₅-alkenes after 1920s are
340 somewhat similar to the reconstructed solar total irradiance (Figure 3i) (*Krivova et al., 2010;*
341 *Ball et al., 2012*). In addition, historical concentrations of C₅-alkene triols are

342 changed/increased significantly after the Great Pacific Climate Shift. In contrast, other
343 compounds (Figure 2a-c and 3a-c) show negative correlation with α -dicarbonyls and fatty
344 acids including oleic acid (e.g., C_{16:0}, C_{18:0} and C_{18:1}). But they have positive correlations with
345 sugar compounds (e.g., mannitol, fructose, glucose, inositol and sucrose) from same ice core
346 sample (unpublished data, Pokhrel and Kawamura, 2015). These results all suggest that an
347 increased biogenic emission capacity and/or source could be changed, being recorded in a clear
348 historical profile of dual sources; marine phytoplankton and terrestrial plants.

349

350 **4. Summary and Conclusions**

351 We report for the first time historical records of monoterpene- and isoprene-derived
352 secondary organic aerosols (SOA) tracers in glacier ice core from Aurora Peak in Alaska since
353 1660s, which could be derived from dual biogenic sources (marine and continental). Positive
354 correlations of monoterpene tracers with 2-methylglyceric acid and 2-methyltetrols are found
355 to be strong, indicating the shared source regions and transport. In contrast, correlations of
356 monoterpene tracers with C₅-alkene triols are weak, suggesting the different source regions and
357 transport pathways.

358 Ice core concentrations of monoterpene- and isoprene-SOA tracers showed lower levels
359 during 1800-1860, which are similar to temperature trends of the Gulf of Alaska and showed a
360 positive correlation with extra-tropical Northern Hemisphere temperature (ENHT). In addition,
361 C₅-alkene triols showed positive correlations with α -dicarbonyls, ENHT and fatty acids, and
362 their historical trends are increased after the Great Pacific Climate Shift. In contrast, all other
363 compounds (i.e., monoterpene-SOA tracers) have negative correlation with α -dicarbonyls and
364 fatty acids including oleic acid (C_{18:1}). But they have positive correlations with sugar
365 compounds (e.g., mannitol, fructose, glucose, inositol and sucrose) in the same ice core.

366 These results suggest that source could be changed significantly after the Great Pacific
367 Climate Shift and exhibit a clear historical dual sources likely to be marine boundary layer.
368 This study reveals that ice core records of SOA tracers and atmospheric oxidizing capability or
369 emission of monoterpenes and isoprene are associated with multi-decadal climate oscillation
370 periodicity in the North Pacific Rim, which are associated with the Northern Hemispheric
371 temperature signals.

372

373 **Acknowledgements**

374 This study was partly supported by the Japan Society for the Promotion of Science (JSPS)
375 through grand-in-aid Nos. 19340137 and 24221001. We also acknowledge the support from the
376 Institute of Low Temperature Science, Hokkaido University for the ice core program.

377 **References**

- 378 Aalto, P., Hameri, K., Becker, E., Weber, R., Salm, J., Makela, J. M., Hoell, C., O'Dowd, C. D.,
379 Karlsson, H., Hansson, H. C., Vakeva, M., Koponen, I. K., Buzorius, G., Kulmala, M.,
380 2001. Physical characterization of aerosol particles during nucleation events. *Tellus B.* 53,
381 344-358.
- 382 Aaltonen, H., Pumpanen, J., Pihlatie, M., Hakola, H., Hellen, H., Kulmala, L., Vesala, T., Back,
383 J., 2011. Boreal pine forest floor biogenic volatile organic compound emissions peak in
384 early summer and autumn. *Agric. For. Meteorol.* 151, 682-691.
- 385 Acosta Navarro, J. C., Smolander, S., Struthers, H., Zorita, E., Ekman, A. M., Kaplan, J.,
386 Guenther, A., Arneeth, A., Riipinen, I., 2014. Global emissions of terpenoid VOCs from
387 terrestrial vegetation in the last millennium. *J. Geophys. Res.* 119, 6867-6885.
- 388 Ball, W. T., Unruh, Y. C., Krivova, N. A., Solanki, S., Wenzler, T., Mortlock, D. J., Jaffe, A.
389 H., 2012. Reconstruction of total solar irradiance 1974-2009. *Astronomy & Astrophysics.*
390 541, id.A27. doi:10.1051/0004-6361/201118702.
- 391 Bhat, S., Fraser, M. P., 2007. Primary source attribution and analysis of α -pinene
392 photooxidation products in Duke Forest, North Carolina. *Atmos. Environ.* 41, 2958-2966.
- 393 Bigg, E. K., 2004. Gas emissions from soil and leaf litter as a source of new particle formation.
394 *Atmos. Res.* 70, 33-42.
- 395 Boyce, D. G., Lewis, M. R., Worm, B., 2010. Global phytoplankton decline over the past century.
396 *Nature.* 466, 591-596. doi: 10.1038/nature09268.
- 397 Cahill, T. M., Seaman, V. Y., Charles, M. J., Holzinger, R., Goldstein, A. H., 2006. Secondary
398 organic aerosols formed from oxidation of biogenic volatile organic compounds in the
399 Sierra Nevada Mountains of California. *J. Geophys. Res. Atmos.* 111, D16312.
400 doi:10.1029/2006JD007178.
- 401 Carlton, G., Wiedinmyer, C., Kroll, J. H., 2009. A review of secondary organic aerosol (SOA)
402 formation from isoprene. *Atmos. Chem. Phys.* 9, 4987-5005.

- 403 Claeys, M., Graham, B., Vas, G., Wang, W., Vermeylen, R., Pashynska, V., Cafmeyer, J.,
404 Guyon, P., Andreae, M., Artaxo, P., Maenhaut, W., 2004. Formation of secondary
405 organic aerosols through photooxidation of isoprene. *Science*. 303, 1173-1176.
- 406 Claeys, M., Szmigielski, R., Kourtchev, I., Veken, P. V. D., Vermeylen, R., Maenhaut, W.,
407 Jaoui, M., Kleindienst, T. E., Lewandowski, M., Offenberg, J. H., Edney, E. O., 2007.
408 Hydroxydicarboxylic Acids: Markers for Secondary Organic Aerosol from the
409 Photooxidation of α -Pinene. *Environ. Sci. Technol.* 41, 1628-1634.
- 410 D'Arrigo, R., Jacoby, G., Buckley, B., Sakulich, J., Frank, D., Wilson, R., Curtis, A.,
411 Anchukaitis, K., 2009. Tree growth and inferred temperature variability at the North
412 American Arctic tree line. *Global and Planetary Change*. 65, 71–82. doi:
413 10.1016/j.gloplacha.2008.10.011.
- 414 Faiola, C. L., VanderSchelden, G. S., Wen, M., Elloy, F. C., Cobos, D. R., Watts, R. J., Jobson,
415 B. T., VanReken, T. M., 2014. SOA Formation Potential of Emissions from Soil and Leaf
416 Litter. *Environ. Sci. & Technol.* 48, 938-946.
- 417 Fu, P., Kawamura, K., Kanaya, Y., Wang, Z., 2010a. Contributions of biogenic volatile organic
418 compounds to the formation of secondary organic aerosols in the troposphere over Mt.
419 Tai, Central East China. *Atmos. Environ.* 44, 4817-4826.
- 420 Fu, P. Q., Kawamura, K., Pavuluri, C. M., Swaminathan, T., Chen, J., 2010b. Molecular
421 characterization of urban organic aerosol in tropical India: contributions of primary
422 emissions and secondary photooxidation. *Atmos. Chem. Phys.* 10, 2663-2689.
- 423 Fu, P., Kawamura, K., Miura, K., 2011. Molecular Characteristics of Marine Organic Aerosols
424 Collected during a Round-the-World Cruise. *J. Geophys. Res. Atmos.* 116, D13302.
425 doi:10.1029/2011JD015604.
- 426 Fu, P., Kawamura, K., Chen, J., Miyazaki, Y., 2014. Secondary Production of Organic
427 Aerosols from Biogenic VOCs over Mt. Fuji, Japan. *Environ. Sci. & Technol.* 48, 8491-
428 8497.
- 429 Glasius, M., Lahaniati, M., Calogiron, A., Di Bella, D., Jensen, N. R., Hjorth, J., Kotzias, D.,
430 Larsen, B. R., 2000. Carboxylic acids in secondary aerosols from oxidation of cyclic
431 monoterpenes by ozone. *Environ. Sci. & Technol.* 34, 1001-1010.
- 432 Griffin, R. J., Cocker, D. R., Seinfeld, J. H., Dabdub, D., 1999. Estimate of global atmospheric
433 organic aerosol from oxidation of biogenic hydrocarbons. *Geophys. Res. Lett.*, 26, 2721-
434 2724.
- 435 Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger,
436 L., Lerdau, M., McKay, W., 1995. A global model of natural volatile organic compound
437 emissions. *J. Geophys. Res.* 100, 8873–8892.
- 438 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., Geron, C., 2006. Estimates of
439 global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and
440 Aerosols from Nature). *Atmos. Chem. Phys.* 6, 3181-3210.
- 441 Hallquist, M., Wenger, J. C., Baltensperger, U. 2009. The formation, properties and impact of
442 secondary organic aerosol: current and emerging issues. *Atmos. Chem. Phys.* 9, 5155-
443 5235.
- 444 Hoffmann, T., Odum, J. R., Bowman, F., Collins, D., Klockow, D., Flagan, R. C., Seinfeld, J.
445 H., 1997. Formation of organic aerosols from the oxidation of biogenic hydrocarbons. *J.*
446 *Atmos. Chem.* 26, 189-222.
- 447 Hu, D., Bian, Q., Li, T. W. Y., Lau, A. K. H., Yu, J. Z., 2008. Contributions of isoprene,
448 monoterpenes, β -caryophyllene, and toluene to secondary organic aerosols in Hong Kong
449 during the summer of 2006. *J. Geophys. Res.* 113, D22206. doi: 10.1029/2008JD010437.
- 450 Jacoby, G. C., D'Arrigo, R., 1989. Reconstructed northern hemisphere annual temperature
451 since 1671 based on high-latitude tree-ring data from North America. *Clim. Change*. 14,
452 39-59.

- 453 Janout, M. A., Weingartner, T. J., Stabenro, P. J., 2013. Air-sea and oceanic heat flux
454 contributions to the heat budget of the northern Gulf of Alaska shelf. *J. Geophys. Res.*
455 *Oceans*. 118. doi: 10.1002/jgrc.20095.
- 456 Jaoui, M., Lewandowski, M., Kleindienst, T. E., Offenberg, J. H., Edney, E. O., 2007. β -
457 caryophyllinic acid: An atmospheric tracer for β -caryophyllene secondary organic aerosol.
458 *Geophys. Res. Lett.* 34, L05816. doi: 10.1029/2006GL028827.
- 459 Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van
460 Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P.,
461 Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L.,
462 Tsigaridis, K., Vignati, E., Stephanou, E. G., Wilson, J., 2005. Organic aerosol and global
463 climate modelling: a review. *Atmos. Chem. and Phys.* 5, 1053e1123.
- 464 Kavouras, I. G., Mihalopoulos, N., Stephanou, E. G., 1999. Formation and gas particle
465 partitioning of monoterpene photo-oxidation products over forests. *Geophys. Res. Lett.*
466 26, 55-58.
- 467 Kawamura, K., Yokoyama, K., Fujii, Y., Watanabe, O., 2001. A Greenland ice core record of
468 low molecular weight dicarboxylic acids, ketocarboxylic acids, and dicarbonyls: A trend
469 from Little Ice Age to the present (1540 to 1989 A.D.). *J. Geophys. Res.* D1, 106, 1331-
470 1345.
- 471 Kawamura, K., Matsumoto, K., Tachibana, E., Aoki, K., 2012a. Low molecular weight (C1-
472 C10) monocarboxylic acids, dissolved organic carbon and major inorganic ions in alpine
473 snow pit sequence from a high mountain site, central Japan. *Atmos. Environ.* 62, 272-280.
- 474 Kawamura K., Izawa Y., Mochida M. and Shiraiwa T. (2012b), Ice core records of biomass
475 burning tracers (levoglucosan and dehydroabietic, vanillic and p-hydroxybenzoic acids)
476 and total organic carbon for past 300 years in the Kamchatka Peninsula, Northeast Asia,
477 *Geochimica et Cosmochimica Acta*, 99, 317-329.
- 478 Kleindienst, T. E., Jaoui, M., Lewandowski, M., Offenberg, J. H., Lewis, C. W., Bhawe, P. V.,
479 Edney, E. O., 2007. Estimates of the contributions of biogenic and anthropogenic
480 hydrocarbons to secondary organic aerosol at a southeastern US location. *Atmos. Environ.*
481 41, 8288 – 8300, doi:10.1016/j.atmosenv.2007.06.045.
- 482 Krivova, N. A., Vieira, L. E. A., Solanki, S. K., 2010. Reconstruction of solar spectral
483 irradiance since the Maunder minimum. *J. Geophys. Res. Space Physics*. 115, A12112.
484 doi: 10.1029/2010JA015431.
- 485 Kunwar, B., Kawamura, K., 2014b. Seasonal distributions and sources of low molecular
486 weight dicarboxylic acids, ω -oxocarboxylic acids, pyruvic acid, α -dicarbonyls and fatty
487 acids in ambient aerosols from subtropical Okinawa in the western Pacific Rim. *Environ.*
488 *Chem.* doi: org/10.1071/EN14097.
- 489 Kunwar, B., Kawamura, K., 2014a. One-year observations of carbonaceous and nitrogenous
490 components and major ions in the aerosols from subtropical Okinawa Island, an outflow
491 region of Asian dusts. *Atmos. Chem. Phys.* 14, 1819-1836, 2014. doi: 10.5194/acp-14-
492 1819-2014.
- 493 Lean, J., Rottman, G., Harder, J., Kopp, G., 2005. *SORCE* contributions to new understanding
494 of global change and solar variability. *Solar Physics*. 230, 27-53.
- 495 Limbeck, A., Kulmala, M., Puxbaum, H., 2003. Secondary organic aerosol formation in the
496 atmosphere via heterogeneous reaction of gases isoprene on acidic particles. *Geophys.*
497 *Res. Lett.* 30, doi: 10.1029/2003Glo177738.
- 498 Makela, J. M., Maso, M. D., Pirjola, L., Keronen, P., Laakso, L., Kulmala, M., Laaksonen, A.,
499 2000. Characteristics of the atmospheric particle formation events observed at a boreal
500 forest site in southern Finland. *Boreal Environ. Res.* 5, 299–313.

- 501 Mantua, N. J., Hare, S. R., Zhang, Y., Wallace, J. M., Francis, R. C., 1997. A Pacific decadal
502 climate oscillation with impacts on salmon. *Bulletin of the American Meteorological*
503 *Society*. 78, pp 1069-1079.
- 504 Matsunaga, S. N., Kato, S., Yoshino, A., Greenberg, J. P., Kajii, Y., and Guenther, A. B., 2005.
505 Gas-aerosol partitioning of semi volatile carbonyls in polluted atmosphere in Hachioji,
506 Tokyo. *Geophys. Res. Lett.* 32, L11805. doi: 10.1029/2004GL021893.
- 507 McNeill, V. F., Grannas, A. M., Abbatt, J. P. D., Ammann, M., Ariya, P., Bartels-Rausch, T.,
508 Domine, F., Donaldson, D. J., Guzman, M. I., Heger, D., Kahan, T. F., Klan, K., Masclin,
509 S., Toubin, C., Voisin, D., 2012. Organics in environmental ices: sources, chemistry, and
510 impacts. *Atmos. Chem. Phys.* 12, 9653–9678. doi: 10.5194/acp-12-9653-2012.
- 511 Noziere, B., Kalberer, M., Claeys, M., Allan, J., D’Anna, B., Decesari, S., Finessi, E., Glasius,
512 M., Grgić, I., Hamilton, J. F., Hoffmann, T., Iinuma, Y., Jaoui, M., Kahnt, A., Kampf, C.
513 F., Kourtchev, I., Maenhaut, W., Marsden, M., Saarikoski, S., Schnelle-Kreis, J., Surratt,
514 J. D., Szidat, S., Szmigielski, R., Wisthaler, A., 2015. The Molecular Identification of
515 Organic Compounds in the Atmosphere: State of the Art and Challenges. *Chem. Rev.* 115,
516 3919–3983. doi: 10.1021/cr5003485.
- 517 Plewaka, A., Gnauk, T., Brüggemann, E., Herrmann, H., 2006. Biogenic contributions to the
518 chemical compositions of airborne particles in a coniferous forest in Germany. *Atmos.*
519 *Environ.* 40, S103-S115.
- 520 Pokhrel, A., Kawamura, K., Seki, O., Matoba, S., Shiraiwa, T., 2015. Ice core profiles of
521 saturated fatty acids (C_{12:0} - C_{30:0}) and oleic acid (C_{18:1}) from southern Alaska since 1734
522 AD: A link to climate change in the Northern Hemisphere. *Atmos. Environ.* 100, 202-209.
- 523 Pokhrel, A., 2015. Studies on ice core records of dicarboxylic acids, w-oxocarboxylic acids,
524 pyruvic acid, a-diacarbonyls and fatty acids from southern Alaska since 1665 Ad: A link to
525 climate change in the Northern Hemisphere. Ph. D thesis, Hokkaido University,
526 HUSCAP 11786. <http://hdl.handle.net/2115/59331>.
- 527 Sasaki, Y. N., Minobe, S., 2005. Seasonally dependent interannual variability of sea ice in the
528 Bering Sea and its relation to atmospheric fluctuations. *J. Geophys. Res.* 110, C05011.
529 doi: 10.1029/2004JC002486.
- 530 Surratt, J. D., Lewandowski, M., Offenberg, J. H., Jaoui, M., Kleindienst, T. E., Edney, E. O.,
531 Seinfeld, J. H., 2007. Effect of acidity on secondary organic aerosol formation from
532 isoprene. *Environ. Sci. Technol.* 41, 5363 – 5369. doi:10.1021/es0704176.
- 533 Surratt, J. D., Chan, A. W. H., Eddingsaasa, N. C., Chan, M. N., Loza, C. L., Kwan, A. J.,
534 Hersey, S. P., Flagan, R. C., Wennberg, P. O., Seinfeld, J. H., 2010. Reactive
535 intermediates revealed in secondary organic aerosol formation from isoprene. *PNAS.* 107,
536 15, 6640–6645.
- 537 Trouet, V., Taylor, A., 2010. Multi-century variability in the Pacific North American (PNA)
538 circulation pattern reconstructed from tree rings. *Climate Dynamics*. doi:10.1007/s00382-
539 009-0605-9.
- 540 Tsushima, A., Matoba, S., Shiraiwa, T., Okamoto, S., Sasaki, H., Solie, D. J., Yoshikawa, K.,
541 2015. Reconstruction of recent climate change in Alaska from the Aurora Peak ice core,
542 central Alaska. *Clim. Past.* 11, 217-226. doi:10.5194/cp-11-217-2015, 2015.
- 543 Volkamer, R., Ziemann, P. J., and Molina, M. J., 2009. Secondary Organic Aerosol Formation
544 from Acetylene (C₂H₂): seed effect on SOA yields due to organic photochemistry in the
545 aerosol aqueous phase. *Atmos. Chem. Phys.* 9, 1907–1928.
- 546 Wang, P. X., Julian, X. L., Wang, H. Z., Wang, Y. K., Sun, X. J., 2012. Circulation indices of
547 the Aleutian low pressure system: Definitions and relationships to climate anomalies in the
548 northern hemisphere. *Adv. Atmos. Sci.* 29(5), 1111-1118. doi: 10.1007/s00376-012-1196-
549 7.

- 550 Wilson, R., D'Arrigo, R., Buckley, B., Buntgen, U., Esper, J., Frank, D., Luckman, B., Payette,
551 S., Vose, R., Youngblut, D., 2007a. A matter of divergence: Tracking recent warming at
552 hemispheric scales using tree ring data. *J. Geophys. Res.* 112, D17103. doi:
553 10.1029/2006JD008318.
- 554 Wilson, R., Wiles, G., D'Arrigo, R., Zweck, C., 2007b. Cycles and shifts: 1,300 years of multi-
555 decadal temperature variability in the Gulf of Alaska. *Clim. Dyn.* 28:425–440. doi:
556 10.1007/s00382-006-0194-9.
- 557 Yasunari, T. J. Yamazaki, K., 2009. Origins of air mass over an Alaskan glacier and
558 implications for ice core studies in the North Pacific regions. *SOLA*. 5, 077-080. doi:
559 10.2151/sola.2009-020.
- 560 Zhu, C., Kawamura, K., Kunwar, B., 2015. Effect of biomass burning over the western North
561 Pacific Rim: wintertime maxima of anhydrosugars in ambient aerosols from Okinawa.
562 *Atmos. Chem. Phys.* 15, 1–15. doi: 10.5194/acp-15-1-2015.
563
564


565 Table 1. Concentrations of monoterpene and isoprene secondary organic tracers in ice core
 566 samples from southeast Alaska in 1665 – 2008.

Name	Concentrations (ng/L)			
	Ave.	Min.	Max.	SD
<i>α/β-Pinene-SOA tracers</i>				
Pinic acid	157	BDL	853	148
Pinonic acid	70.6	BDL	300	68
3-HGA	22.4	BDL	234	41
<i>Isoprene-SOA tracers</i>				
2-methylglyceric acid	35.6	BDL	282	48
2-methylthreitol	349	BDL	1740	383
2-methylerythritol	692	BDL	3509	709
3-methyl-2,3,4-trihydroxy-1-butene	6.99	BDL	141	17
cis-2-methyl-1,3,4-trihydroxy-1-butene	23.1	BDL	384	52
trans-2-methyl-1,3,4-trihydroxy-1-butene	36.5	BDL	786	98

567 BDL= Below detection limit (0.001 ng/g-ice)

568

569



570

571 Figure 1. Map showing the geographical location of Aurora Peak of Alaska, from which 180-
572 meter long ice core sample was drilled in 2008 (*Pokhrel et al., 2014*).

573

574

575


576

577

578


579

580




581
 582 Figure 2. Concentration changes of (a) pinic acid, (b) pinonic acid (c) 3-hydroxyglutaric acid
 583 (3-HGA) and (d) average concentration changes of monoterpene tracers (a)-(c) in the Alaska
 584 ice core records collected from Aurora peak.
 585

586




587

588 Figure 3. Concentration changes of (a) 2-methylglyceric (b) 2-methylthreitol (c) 2-methylerythritol (d) cis-
 589 methyl 1,3,4-trihydroxy-1-butene (e) 3-methyl-2,3,4-trihydroxy-1-butene (f) trans-2-methyl-1,3,4-trihydroxy-1-
 590 butene and (g) total annual average concentrations of all these isoprene SOA tracers (a – f) in the Alaska ice core
 591 records collected from Aurora peak (h) extra tropical Northern Hemispheric temperature (ENHT) departure, e.g.,
 592 Wilson et al. (2007a) and (i) 7 years running mean of reconstructed solar total irradiance (STI) from IPCC AR5
 593 based on Krivova et al. (2010) and Ball et al. (2012).
 594



595
 596 Figure 4. Correlation between the concentrations of (a) pinonic and pinic (b) pinonic acid and
 597 2-methylglyceric (e) pinic acid and 2-methylglyceric and (f) 2-methylglyceric and 3-
 598 hydroxyglutaric acid (3-HGA) in the Alaska ice core records collected from saddle of Aurora
 599 Peak.



600
601 Figure 5. Correlation between the concentrations of (a) pinonic and 2-methylthreitol (b) pinic
602 and 2-methylthreitol, (c) pinonic and 2-methylerythritol, and (d) pinic and 2-methylerythritol
603 in the Alaskan ice core records collected from saddle of Aurora Peak.
604

605 **Highlights**

606 1. Monoterpene-SOA tracers in south Alaskan ice core are associated with continental sources.

607 (80 with space)

608 2. Isoprene-SOA tracers in the ice core are derived from both continental and marine sources.

609 (75 with space)

610 3. Monoterpene- and isoprene-SOA tracers are linked with climate oscillations of Aleutian Low.

611 (80 with space)