



Title	Sulfuriflexus mobilis gen. nov., sp nov., a sulfur-oxidizing bacterium isolated from a brackish lake sediment
Author(s)	Kojima, Hisaya; Fukui, Manabu
Citation	International journal of systematic and evolutionary microbiology, 66, 3515-3518 https://doi.org/10.1099/ijsem.0.001227
Issue Date	2016-09-01
Doc URL	http://hdl.handle.net/2115/67091
Type	article (author version)
File Information	20170106.pdf



[Instructions for use](#)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17

Sulfuriflexus mobilis gen. nov., sp. nov., a sulfur-oxidizing
bacterium isolated from a brackish lake sediment

Hisaya Kojima* and Manabu Fukui

The Institute of Low Temperature Science, Hokkaido University. Kita-19, Nishi-8,
Kita-ku, Sapporo 060-0819, Japan



*Corresponding author.
E-mail: kojimah@pop.lowtem.hokudai.ac.jp
Phone: +81-11-706-5460
Fax: +81-11-706-5460

Running head: *Sulfuriflexus mobilis* gen. nov., sp. nov.

Subject category: New taxa: *Proteobacteria*

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of
strain is LC131141.

18 Summary

19 A chemolithotrophic sulfur-oxidizing bacterium, strain aks1^T was isolated from
20 sediment of a brackish lake in Japan. The cells were curved rod-shaped and Gram-
21 stain-negative. The G+C content of genomic DNA was 53 mol%. The major
22 components in the cellular fatty acid profile were C_{16:0} and summed feature 3 (C_{16:1}
23 _{ω7c} and/or C_{16:1}_{ω6c}). As electron donor for chemolithoautotrophic growth, strain
24 aks1^T oxidized thiosulfate, sulfide, and elemental sulfur. The strain could utilize oxygen
25 and nitrate as an electron acceptor for thiosulfate oxidation. Growth was observed at a
26 temperature range of 5–34°C, with optimum growth at 30–32°C. Growth of the strain
27 was observed at a pH range of 6.4–8.7. Phylogenetic analysis based on 16S rRNA gene
28 indicated that the strain is related to members of the family *Granulosicoccaceae* within
29 the order *Chromatiales*, with sequence similarities around 92%. On the basis of its
30 phylogenetic and phenotypic properties, the strain aks1^T (= DSM 102939^T = NBRC
31 111889^T) is proposed as type strain of a new species of a novel genus, *Sulfuriflexus*
32 *mobilis* gen. nov., sp. nov.

33

34 The original description of the order *Chromatiales* contains 3 families,
35 *Chromatiaceae*, *Ectothiorhodospiraceae*, and *Halothiobacillaceae* (Imhoff, 2005), and
36 4 families, *Granulosicoccaceae* (Lee *et al.*, 2007), *Thioalkalispiraceae* (Mori *et al.*,
37 2011), *Wenzhouxiangellaceae* (Wang *et al.*, 2015), and *Woeseiaceae* (Du *et al.*, 2016)
38 have been added in the order. In the present study, a novel chemolithoautotrophic
39 sulfur-oxidizing bacterium, strain aks1^T, was isolated and characterized to be proposed
40 as type species of a new genus in this order.

41

42 The strain aks1^T was isolated from sediment of a brackish lake in Japan, Lake Akkeshi.
43 Throughout this study, a bicarbonate-buffered defined medium was used as basal
44 medium. To prepare the medium, following constituents (l⁻¹) were dissolved in distilled
45 water and then autoclaved: 20 g NaCl, 3 g MgCl₂·6H₂O, 0.3 g MgSO₄·7H₂O, 0.1 g
46 CaCl₂·2H₂O, 0.1 g NH₄Cl, 0.1 g KH₂PO₄, 0.1 g KCl. After cooling down to room
47 temperature, 1 ml trace element solution, 1 ml selenite-tungstate solution, 30 ml
48 NaHCO₃ solution, and 1 ml vitamin mixture solution (DSM 141) were aseptically added
49 to the main body of medium. The solutions of trace element, selenite-tungstate and
50 NaHCO₃ were prepared as described previously (Widdel & Bak, 1992). Before
51 dispensing into culture containers, the pH of the medium was adjusted to 7.0–7.2. The

52 enrichment culture was established with the basal medium supplemented with elemental
53 sulfur (ca. 0.5 g l⁻¹). After 11 times transfer to fresh medium of the same composition
54 (1–2%), the sole electron donor was changed to 20 mM Na₂S₂O₃. Finally, strain was
55 isolated in pure culture by repeated serial dilution in the medium supplemented with
56 Na₂S₂O₃. Purity of the isolate was checked by microscopy and sequencing of the 16S
57 rRNA gene fragments amplified with using some PCR primer pairs, as described
58 previously (Higashioka *et al.*, 2012).

59

60 For the characterization of the strain, the basal medium supplemented with 20 mM
61 Na₂S₂O₃ was used unless otherwise specified. Culturing experiments were performed in
62 bottles closed with rubber stoppers, and the bottles were incubated without shaking at
63 30°C unless otherwise specified.

64 The Gram-stain test was conducted with a kit (Fluka), and oxidase activity was tested
65 by using an oxidase test reagent (bioMérieux). Catalase activity was assessed by
66 pouring 3% H₂O₂ solution onto a pellet of cells. The genomic G+C content of the DNA
67 was determined with the HPLC methods (Katayama-Fujimura *et al.*, 1984), using a kit
68 (Yamasa Shoyu). Fatty acid profile of the strain was analyzed at the Techno Suruga Co.
69 Ltd (Shizuoka, Japan), by using the Sherlock Microbial Identification System Version

70 6.0 (MIDI) with database TSBA6.

71 Effects of temperature on growth were examined by culturing at various temperatures
72 (0, 5, 8, 10, 13, 15, 18, 22, 25, 28, 30, 32, 34, 36, and 37°C). To examine effects of salt
73 concentration, strain aks1 was cultured in modified media with varying concentrations
74 of NaCl (0.0, 0.5, 1.0, 2.0, 3.0, 4.0, 4.5 and 5.0% w/v) and lowered concentration of
75 $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ (0.2 g l^{-1}). Effect of pH on the growth was tested with a method previously
76 described (Kojima *et al.*, 2015), with a modification for the NaCl-requiring strain. The
77 test was performed with media containing 2.0% (w/v) NaCl, and tested pH and
78 buffering reagents were as follows; pH 5.9, 6.3, 6.4, 6.5, 6.8 and 7.2 with MES; pH 6.9,
79 7.0, 7.2, 7.3, 7.4 and 7.6 with PIPES; pH 7.7, 7.8, 7.9 and 8.2 with MOPS; pH 8.2, 8.4,
80 8.6, and 8.7 with Tricine; pH 8.9, 9.4, 9.6, and 10.0 with CHES.

81 Utilization of electron donors was tested in the basal medium supplemented with one
82 of the substances listed later. The utilization of electron acceptor was tested with
83 $\text{Na}_2\text{S}_2\text{O}_3$ (10 mM) as an electron donor, in the same medium under anoxic conditions
84 (headspace of the bottles was filled with 4:1 mixture of N_2/CO_2). Heterotrophic growth
85 in complex liquid media was tested under oxic conditions, for Marine Broth 2216 (MB;
86 Difco) and following media all supplemented with 2% NaCl; R2A (Daigo), diluted
87 (1/10) R2A, NB (Difco), and TSB (OXOID).

88 For phylogenetic analysis, 16S rRNA gene was amplified by PCR using the primer
89 pair 27F and 1492R (Lane, 1991) and then sequenced with a BigDye Terminator v3.1
90 Cycle Sequencing kit (Applied Biosystems). For the resulting sequence, phylogenetic
91 analysis was conducted using the program MEGA version 5.22 (Tamura *et al.*, 2011).

92

93 Cells of aks1^T were motile curved rods, 0.9–6.0 μm long and 0.3–0.5 μm wide (Fig.
94 1). Strain aks1^T was Gram-stain-negative, and tests of catalase and oxidase resulted in
95 negative and positive respectively. The G+C content of the genomic DNA assessed by
96 the HPLC-based method was 53 mol%. In the cellular fatty acid profile, major
97 components were summed feature 3 (C_{16:1ω7c} and/or C_{16:1ω6c}; 57.3 %) and C_{16:0}
98 (32.2%). The other fatty acids detected were summed feature 9 (isoC_{17:1ω7c} and/or
99 C_{16:0} 10-methyl; 5.2 %), summed feature 8 (C_{18:1ω7c} and/or C_{18:1ω6c}; 1.7 %), C_{10:0}
100 3-OH (1.3%), C_{18:0} (0.8%), C_{14:0} (0.6%), C_{12:0} 3-OH (0.4%), C_{17:0} (0.3%) and C_{10:0}
101 (0.2%).

102 Growth of strain aks1^T was observed over a temperature range between 5°C and 34°C,
103 with optimal growth at 30–32°C. Growth of the strain was observed at a pH range of
104 6.4–8.7, and optimum growth was observed at pH range of 6.8–8.3. Growth was
105 observed in medium with 1–4% NaCl with an optimum of 3%.

106 The isolate grew chemolithotrophically on thiosulfate (20 mM), sulfide (2 mM) and
107 elemental sulfur (0.5 g l⁻¹). Hydrogen gas (Air/H₂; 4:1, v/v; 125 kPa total pressure),
108 tetrathionate (20 mM) and sulfite (5 mM) did not support autotrophic growth of the
109 strain. The following organic substrates did not support growth of strain aks1^T: pyruvate
110 (5 mM), lactate (5 mM), acetate (5 mM), methanol (5 mM), succinate (2.5 mM),
111 fumarate (2.5 mM), butyrate (2.5 mM), isobutyrate (2.5 mM), ethanol (2.5 mM),
112 formate (5 mM), lactose (2.5 mM), glucose (2.5 mM), xylose (2.5 mM). The strain
113 exhibited no growth on MB or other complex media tested. Nitrate (20 mM) supported
114 anaerobic growth of strain as sole electron acceptor for thiosulfate oxidation.

115 Among characterized strains of species with validly published names, *Thiopfundum*
116 *hispidum* gsp61^T showed the highest sequence similarity to strain aks1^T (93%), followed
117 by the type strains of *Granulosicoccus* species (92%) and some other bacteria in the
118 order *Chromatiales*. By constructing phylogenetic trees, it was revealed that strain aks1^T
119 forms a monophyletic cluster with *Granulosicoccus* species (Fig. 2, Fig S1). The
120 methods of neighbor-joining and minimum-evolution generated trees of identical
121 topology (Fig. 2), but a different tree was obtained with the maximum-likelihood
122 method (Fig. S1). In all trees, aks1^T represents a sister group of the genus
123 *Granulosicoccus*, and they form a cluster with the genera *Thioalkalispira* and

124 *Thiohalophilus*. The genus *Granulosicoccus* is the sole genus in the family
125 *Granulosicoccaceae*, whereas *Thioalkalispira* and *Thiohalophilus* belong to the family
126 *Thioalkalispiraceae*. The other genus of the family *Thioalkalispiraceae*, genus
127 *Thiopfundum* was positioned apart from these genera (Fig. 2, Fig S1). This
128 phylogenetic isolation of *Thiopfundum* from the other genera was also shown in
129 phylogenetic trees previously constructed (Mori & Suzuki 2014; Mori *et al.*, 2015),
130 suggesting that the family *Thioalkalispiraceae* is not monophyletic. As pointed out
131 previously, it is difficult to clarify phylogenetic relationships among the families in the
132 order *Chromatiales* (Mori *et al.*, 2015), and reclassification of some taxa may be
133 required in the future. However, it seems reasonable to place the strain aks1^T in the
134 family *Granulosicoccaceae* at this point.

135 Differential properties of strain aks1^T and related genera are summarized in Table 1. In
136 contrast to heterotrophic *Granulosicoccus* species, growth of strains aks1^T was not
137 observed in complex media including the MB, or synthetic medium supplemented with
138 organic substrates. Differences between strains aks1^T and *Granulosicoccus* species are
139 apparent in cell morphology, oxygen requirement, catalase activity (Table 1). On the
140 basis of its distinct phenotypic properties and isolated phylogenetic position, strain
141 aks1^T is proposed to be assigned to a new species of a novel genus in the family

142 *Granulosicoccaceae*, with the name *Sulfuriflexus mobilis* gen. nov., sp. nov.

143

144 **Description of *Sulfuriflexus* gen. nov.**

145 *Sulfuriflexus* (Sul.fu.ri.fle'xus. L. neut. n. *sulfur* sulfur; L. masc. n. *flexus*, a bending;

146 N.L. masc. n. *Sulfuriflexus* sulfur-oxidizing bending).

147 Cells are motile and Gram-stain-negative. Grow chemolithoautotrophically by the

148 oxidation of inorganic sulfur compounds. As determined by 16S rRNA gene sequence

149 analysis, belonging to family *Granulosicoccaceae*. The type species is *Sulfuriflexus*

150 *mobilis*.

151

152 **Description of *Sulfuriflexus mobilis* sp. nov.**

153 *Sulfuriflexus mobilis* (mo'bi.lis. L. masc. adj. *mobilis*, movable, motile).

154 Cells are curved rod-shaped, 0.9–6.0 μm in length and 0.3–0.5 μm in width.

155 Autotrophic growth occurs with oxidation of thiosulfate, sulfide and elemental sulfur.

156 Oxidase-positive and catalase-negative. Growth occurs at temperatures 5–34°C, with

157 optimum growth at 30–32°C. The pH range for growth is 6.4–8.7. The G+C content of

158 genomic DNA is 53 mol%. Major cellular fatty acids are C_{16:0} and summed feature 3

159 (C_{16:1 ω 7c} and/or C_{16:1 ω 6c}). The type strain aks1^T (= DSM 102939^T = NBRC 111889^T)

160 was isolated from sediment of a brackish lake in Japan (Lake Akkeshi).

161

162 **ACKNOWLEDGMENTS**

163 We thank R. Tokizawa and A. Shinohara, Hokkaido University, for their
164 technical assistance. This study was supported by JSPS KAKENHI Grant Number
165 15K07209 to H. Kojima.

166

167

168 **REFERENCES**

169 **Baek, K., Choi, A., Kang, I., Im, M., & Cho, J. C. (2014).** *Granulosicoccus marinus*
170 sp. nov., isolated from Antarctic seawater, and emended description of the genus
171 *Granulosicoccus*. *Int J Syst Evol Microbiol* **64**, 4103–4108.

172

173 **Du, Z-J., Wang, Z-J, Zhao, J-X, Chen & G-J. (2016).** *Woeseia oceani* gen. nov., sp.
174 nov., a chemoheterotrophic member of the order *Chromatiales*, and proposal of
175 *Woeseiaceae* fam. nov. *Int J Syst Evol Microbiol* **66**, 107–112.

176

177 **Higashioka, Y., Kojima, H., & Fukui, M. (2012).** Isolation and characterization of

178 novel sulfate-reducing bacterium capable of anaerobic degradation of p-xylene. *Microb*
179 *Environ* **27**, 273–277.

180

181 **Imhoff, J.F. (2005).** Order I. *Chromatiales* ord. nov. In Bergey's Manual of Systematic
182 Bacteriology, vol. 2 (The Proteobacteria), part B (The Gammaproteobacteria), 2nd edn,
183 pp. 1–3. Edited by Brenner D. J., Krieg N. R., Staley J. T., Garrity G. M.. New York:
184 Springer.

185

186 **Katayama-Fujimura, Y., Komatsu, Y., Kuraishi, H. & Kaneko, T. (1984).**
187 Estimation of DNA base composition by high-performance liquid chromatography of its
188 Nuclease PI hydrolysate. *Agric Bio. Chem* **48**, 3169–3172.

189

190 **Kojima, H. Shinohara, A. & Fukui, M. (2015).** *Sulfurifustis variabilis* gen. nov., sp.
191 nov., a novel sulfur oxidizer isolated from a lake, and proposal of *Acidiferrobacteraceae*
192 fam. nov. and *Acidiferrobacterales* ord. nov. *Int J Syst Evol Microbiol* **65**, 3709–3713

193

194 **Kurilenko, V.V., Christen, R., Zhukova, N.V., Kalinovskaya, N.I., Mikhailov, V.V.,**
195 **Crawford, R.J., & Ivanova, E.P. (2010).** *Granulosicoccus coccooides* sp. nov., isolated

196 from leaves of seagrass (*Zostera marina*). *Int J Syst Evol Microbiol* **60**, 972–976.

197

198 **Lane, D. J. (1991)**. 16S/23S rRNA sequencing. In *Nucleic Acid Techniques in Bacterial*
199 *Systematics*, chapter 6, pp. 115–175. Edited by E. Stackebrandt & M. Goodfellow.
200 Chichester: Wiley.

201

202 **Lee, K., Lee, H.K., Choi, T.-H., Kim, K.M., & Cho J.-C. (2007)**. *Granulosicoccaceae*
203 fam. nov., to include *Granulosicoccus antarcticus* gen. nov., sp. nov., a
204 non-phototrophic, obligately aerobic chemoheterotroph in the order *Chromatiales*,
205 isolated from Antarctic seawater. *J Microbiol Biotechnol* **17**, 1483–1490.

206

207 **Mori, K., Suzuki, K., Urabe, T., Sugihara, M., Tanaka, K., Hamada, M., & Hanada,**
208 **S. (2011)**. *Thiopfundum hispidum* sp. nov., an obligately chemolithoautotrophic
209 sulfur-oxidizing gammaproteobacterium isolated from the hydrothermal field on Suiyo
210 Seamount, and proposal of *Thioalkalispiraceae* fam. nov. in the order *Chromatiales*. *Int*
211 *J Syst Evol Microbiol* **61**, 2412–2418.

212

213 **Mori, K. & Suzuki, K. (2014)**. The family *Thioalkalispiraceae*. In *The Prokaryotes*,

214 4th edn, pp. 199–223. Edited by Rosenberg E., DeLong E. F., Lory S., Stackebrandt E.,
215 Thompson F. Berlin: Springer.

216

217 **Mori, K., Suzuki, K., Yamaguchi, K., Urabe, T. & Hanada, S. (2015).** *Thiogranum*

218 *longum* gen. nov., sp. nov., an obligately chemolithoautotrophic, sulfur-oxidizing

219 bacterium of the family *Ectothiorhodospiraceae* isolated from a deep-sea hydrothermal

220 field, and an emended description of the genus *Thiohalomonas*. *Int J Syst Evol*

221 *Microbiol* **65**, 235–241.

222

223 **Park, S., Jung, Y. T., Won, S. M., Park, J. M., & Yoon, J. H. (2014).** *Granulosicoccus*

224 *undariae* sp. nov., a member of the family *Granulosicoccaceae* isolated from a brown

225 algae reservoir and emended description of the genus *Granulosicoccus*. *Antonie van*

226 *Leeuwenhoek* **106**, 845–852.

227

228 **Sorokin, D. Y., Tourova, T. P., Kolganova, T. V., Sjollema, K. A. & Kuenen, J. G.**

229 **(2002).** *Thioalkalispira microaerophila* gen. nov., sp. nov., a novel lithoautotrophic,

230 sulfur-oxidizing bacterium from a soda lake. *Int J Syst Evol Microbiol* **52**, 2175–2182.

231

232 **Sorokin, D. Y., Tourova, T. P., Bezsoudnova, E. Y., Pol, A. & Muyzer, G. (2007).**
233 Denitrification in a binary culture and thiocyanate metabolism in *Thiohalophilus*
234 *thiocyanoxidans* gen. nov. sp. nov. - a moderately halophilic chemolithoautotrophic
235 sulfur-oxidizing Gammaproteobacterium from hypersaline lakes. *Arch Microbiol* **187**,
236 441–450
237
238 **Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011).**
239 MEGA5: Molecular evolutionary genetics analysis using maximum likelihood,
240 evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* **28**,
241 2731–2739.
242
243 **Wang, G., Tang, M., Li, T., Dai, S., Wu, H., Chen, C., He, H., Fan, J., Xiang, W. &**
244 **Li, X. (2015).** *Wenzhouxiangella marina* gen. nov, sp. nov, a marine bacterium from the
245 culture broth of *Picochlorum* sp. 122, and proposal of *Wenzhouxiangellaceae* fam. nov.
246 in the order *Chromatiales*. *Antonie van Leeuwenhoek* **107**, 1625–1632.
247
248 **Widdel, F. & Bak, F. (1992).** Gram-negative mesotrophic sulfate-reducing bacteria. In
249 *The Prokaryotes* 2nd edn, vol. 4, pp. 3352–3378. Edited by A. Balows, H. G. Trüper, M.

251 Table 1. Differential properties of strain aks1^T and related genera. Genera: 1,
 252 *Granulosicoccus*; 2, *Thioalkalispira*; 3, *Thiohalophilus*. The properties of
 253 *Granulosicoccus* were compiled from the description of type strains representing 4
 254 species in this genus, *G. antarcticus* IMCC3135^T (Lee *et al.*, 2007), *G. coccoides* Z
 255 271^T (Kurilenko *et al.*, 2010), *G. marinus* IMCC3490^T (Baek *et al.*, 2014) and *G.*
 256 *undariae* W-BA3^T (Park *et al.*, 2014). Those of *Thioalkalispira* and *Thiohalophilus* are
 257 from Sorokin *et al.*, 2002 and Sorokin *et al.*, 2007, respectively.

258

Characteristics	aks1	Genera		
		1	2	3
Cell morphology	Curved rod	Coccoid	Spiral rod	Rod
Motility	+	+/-*	+	-
Heterotrophic growth	-	+	-	-
Anaerobic growth	+	-	-	+
Catalase	-	+	+	ND

259 *One of the 4 species is non-motile and the others are motile.

260

261

262 Figure legends

263

264 Fig. 1 Phase-contrast micrograph of strain aks1^T. Bar, 5 μm.

265

266 Fig. 2 Minimum-evolution tree showing the phylogenetic position of aks1^T within the

267 order *Chromatiales* based on the 16S rRNA gene sequence analysis. This tree was

268 constructed using *ca.* 1200 sites. *Sulfuricaulis limicola* and *Acidiferrobacter*

269 *thiooxydans* are included as outgroup species. Neighbor-joining method yielded a tree

270 of identical topology. Numbers on nodes represent percentage values of 1000 bootstrap

271 resampling (values less than 50 are not shown).

272




Fig.1




Fig.2

Supplementary material

***Sulfuriflexus mobilis* gen. nov., sp. nov., a sulfur-oxidizing bacterium isolated from
a brackish lake sediment**

Hisaya Kojima* and Manabu Fukui

The Institute of Low Temperature Science, Hokkaido University. Kita-19, Nishi-8,
Kita-ku, Sapporo 060-0819, Japan

*Corresponding author. E-mail: kojimah@pop.lowtem.hokudai.ac.jp




Fig S1. Maximum-likelihood tree showing the phylogenetic position of of aks1^T within the order *Chromatiales* based on the 16S rRNA gene sequence analysis.