

1 **BOLDNESS IS FOR ROOKIES: PRE-FIGHT BOLDNESS AND FIGHTING SUCCESS IN**
2 **A SEA ANEMONE**

3 **Sarah M. Lane* & Mark Briffa**

4 Marine biology and ecology research centre, Plymouth University, Drake Circus, Plymouth,
5 UK, PL4 8AA

6 *Correspondence: Sarah M. Lane, Davy 620, Marine biology and ecology research centre,
7 Plymouth University, Drake Circus, Plymouth, UK, PL4 8AA

8 E-mail: sarah.lane@plymouth.ac.uk

9

10 **Fighting experience (specifically winning or losing a fight) can significantly alter boldness,**
11 **a component of resource holding potential (RHP). Previous studies have shown that both**
12 **the repeatability of boldness and mean-level boldness can be affected by fighting**
13 **experience and that these effects are strongest in the recipients of agonistic behaviour.**
14 **However, whether these post-fight changes in boldness impact future contest success and**
15 **whether subsequent contests further affect boldness remains unknown. Furthermore,**
16 **little is known about the effects of the specific tactics used within a fight (within-fight**
17 **experience) and how these might influence future fight performance and boldness. Here,**
18 **we investigate the relationship between fighting success and boldness (measured as**
19 **recovery time when startled) across repeated contests in the beadlet sea anemone *Actinia***
20 ***equina*, measuring boldness at 5 occasions before, between and after two contests. We**
21 **found that boldness (both repeatability and mean-level) was generally robust to the**

22 effects of fighting experience, with the exception of a decrease in the immediate boldness
23 of losers after their second fight. Furthermore, we found that while pre-fight boldness
24 significantly predicted fighting success and the level of aggression used in an individual's
25 first fight, it did not predict victory or aggression in the second fight. Our findings thus
26 indicate that different traits may be important in determining fighting success in
27 consecutive fights and moreover that fighting experience may alter which traits contribute
28 to an individual's RHP.

29

30 **Keywords:** *Actinia equina*; boldness; fighting success; injuries; prior fighting experience;
31 resource holding potential

32

33 INTRODUCTION

34 Resource holding potential (RHP) is comprised of a multitude of traits – including weapon
35 size, body size, strength and endurance – all of which combine to determine an individual's
36 ability to win a fight (Parker, 1974). Recently, RHP has been shown to be influenced not only
37 by morphological and physiological traits but also by consistent between-individual
38 differences in behaviour (personality traits), namely boldness (Barlow, Rogers, & Fraley,
39 1986; reviewed in Briffa, Sneddon, & Wilson, 2015). Boldness can be measured in different
40 ways depending upon the species and context of interest. The most common measures
41 include; exploratory behaviour in a novel environment (high/low), investigation of novel
42 objects (readily/slowly) and recovery time when startled (fast/slow) (Briffa, Rundle, & Fryer,
43 2008). As a component of RHP, boldness significantly affects the fighting success of
44 individuals, for instance in the beadlet sea anemone *Actinia equina* - in which boldness has

45 been measured as recovery time when startled (referred to hereafter as startle response) -
46 bolder individuals have been shown to inflict a higher number of attacks and win more
47 fights than their shyer counterparts (Rudin & Briffa, 2012). However bolder is not always
48 better. In the asymmetric contests of the hermit crab *Pagurus bernhardus*, an attacker's
49 chance of winning is not influenced by its boldness (startle response duration) but shyer
50 individuals are better able to defend their shells from eviction (Courtene-Jones & Briffa,
51 2014).

52 The experience of winning or losing a fight can significantly alter traits contributing
53 to RHP which can in turn affect behaviour and success in subsequent contests (Hsu & Wolf,
54 2001; Rutte, Taborsky, & Brinkhof, 2006). The effect of fighting experience on boldness has
55 thus far been investigated in only a handful of studies (willingness to approach a novel
56 object - Frost, Winrow-Giffen, Ashley, & Sneddon, 2007; startle response duration- Rudin &
57 Briffa, 2012; Courtene-Jones & Briffa, 2014), the results of which demonstrate that fighting
58 experience (specifically, winning or losing a fight) significantly affects both the repeatability
59 of boldness and mean-level boldness, with the most extreme effects being seen in the
60 recipients of agonistic behaviour, i.e. losers and defenders (Courtene-Jones & Briffa, 2014;
61 Rudin & Briffa, 2012). For example, in *P. bernhardus*, defenders showed a significant
62 reduction in the repeatability of boldness (measured as startle response) after fighting,
63 while the boldness of attackers remained stable across situations (Courtene-Jones & Briffa,
64 2014). Furthermore the mean-level post-fight boldness of defenders varied with the
65 intensity of agonistic behaviours they were subjected to during the fight. In *A. equina*, pre-
66 fight boldness was highly repeatable for both eventual winners and losers but the post-fight
67 boldness of losers was not repeatable at all. Losers also showed a significant reduction in

68 mean-level boldness (Rudin & Briffa, 2012). Although these studies provide evidence that
69 fighting experience (specifically winning or losing a fight) can significantly affect the
70 consistency and level of boldness, measures of pre-fight boldness were only compared with
71 one or two post-fight measures, and as such it is unclear how long the observed post-fight
72 changes in boldness persist. Moreover, if boldness contributes to RHP, then post-fight
73 changes in boldness could alter the potential to win subsequent fights.

74 However, it is not just the outcome of a fight that can affect an individual's future
75 contest behaviour/success, but also what happens during a fight (referred to hereafter as
76 within-fight experience). Within-fight experience can vary in terms of outcome (winning or
77 losing – as discussed above), level of aggression/escalation, tactics employed, duration and
78 injury (both receiving and inflicting injuries; Lane & Briffa, 2017). Injuries can significantly
79 affect subsequent contest performance by reducing fighting ability. For example, in blue
80 crabs *Callinectes sapidus* Rathbun (Smith, 1992) and stomatopods *Gonodactylus bredini*
81 (Berzins & Caldwell, 1983), injury affects an individual's ability to retain possession of
82 females and territories respectively, with injured individuals losing to intact opponents.
83 Injury has also been shown to interact with correlates of RHP to determine fighting success.
84 For instance in the jumping spider *Trite planiceps*, body size is a major predictor of fight
85 outcome in intact individuals, with larger individuals being more likely to win as the size
86 difference between opponents increases. However, this size advantage diminishes when an
87 individual is injured (Taylor & Jackson, 2003), the most injured rival being more likely to lose
88 regardless of size difference. While these studies all indicate that injury can have a
89 significant effect on subsequent fighting success, all three were carried out on individuals
90 who had been injured in ways other than through fighting itself (autotomy of unknown

91 cause – Smith, 1992; Taylor & Jackson, 2003; surgically injured – Berzins & Caldwell, 1983),
92 and thus the direct effect of injuries sustained in fights on future contest performance
93 remains unclear.

94 Although previous studies have compared the effects of winning and losing fights
95 against individuals that have not fought, little is known about the effects of the specific
96 tactics used within a fight and how these might influence future fight performance and
97 boldness. Understanding whether there is a link between aggressive performance and
98 repeatable behaviour is important because it has been suggested that both fighting
99 behaviour and consistent behavioural differences between individuals can be explained by
100 negative frequency dependent selection (e.g. Wolf & Weissing, 2010). Here, we investigate
101 the relationship between boldness and within-fight experience in the beadlet sea anemone
102 *Actinia equina*, where the outcomes of the fight can be win, lose or draw (i.e. no clear
103 winner). Although lacking a centralised nervous system, *A. equina* possess weapons in the
104 form of specialised stinging structures called acrorhagi which contain high concentrations of
105 stinging cells (nematocytes) and are used during fights with conspecifics (Williams, 1978;
106 Brace, Pavey, & Quickie, 1979; Bigger, 1982). As mentioned above, boldness is a known
107 component of RHP in *A. equina* and has previously been shown to be affected by winning or
108 losing a fight (Rudin & Briffa, 2012). Thus far however, post-fight boldness has been
109 measured only once and thus the extent to which fighting experience affects boldness in the
110 longer term, the extent to which it affects the repeatability of boldness, and hence the
111 effect that these changes could have on future fights, is unclear. Although not all contests in
112 *A. equina* result in injuries, when fights do escalate, anemones drag inflated acrorhagi along
113 the body column of their opponent, leaving behind nematocyte-filled ectoderm which rips

114 off from their acrorhagi (referred to hereafter as 'peels'). These assaults injure the attacker
115 as well as the recipient and thus are potentially costly to both contestants (Lane & Briffa,
116 2017). In this study we aimed to examine (i) how boldness contributes to fighting success
117 across multiple contests, (ii) how post-fight changes in boldness affect subsequent contest
118 success (iii) if and how subsequent contests further affect boldness and (iv) how injury state
119 and contest outcome (of focal and opponent) influence the effect of boldness on
120 subsequent fighting success. We therefore measured boldness before, between and after
121 two staged contests using startle response duration as our index of boldness (startle
122 response duration has previously been shown to provide highly repeatable measures of
123 boldness in *A. equina*, Bigger 1982; Briffa & Greenaway, 2011; Rudin & Briffa, 2012).

124

125 MATERIALS AND METHODS

126 *Anemone collection and startle-response measures*

127 *Actinia equina* (N= 126) were collected from Portwrinkle (Cornwall, UK; grid reference: SX
128 357539) on 4 collection trips carried out between December 2015 and June 2016 (an
129 average of 38 anemones collected on each trip) and taken back to the lab within 1-2 hours
130 of collection. As in previous studies investigating aggression in *A. equina*, only anemones of
131 the red/brown colour morph were collected. The red/brown morph has previously been
132 shown to exhibit higher levels of aggression than anemones of the green/orange morphs
133 found lower down on the shore (Manuel, 1988). Once in the lab, anemones were
134 individually housed in plastic tanks (23 x 16 x 17.5cm) containing 700ml of filtered, aerated
135 seawater and maintained in a controlled temperature room at 15°C ± 0.5°C. Throughout the

136 experiment, anemones were fed *ad libitum* aquaria marine fish flakes every 2-3 days and
137 seawater was changed every 7 days.

138 The first startle-response test ('pre-fight 1') was conducted 7-14 days after collection
139 from the shore, allowing the anemones time to habituate to the laboratory environment
140 and attach their pedal discs to the side of their tank. The test was carried out by discharging
141 a 5ml syringe full of seawater directly into the oral disc at a range of approximately 2cm
142 (Briffa & Greenaway, 2011), causing the anemone to retract its tentacles. The anemone's
143 response was calculated from the time the stimulus was applied until the point at which the
144 anemone had re-opened fully to match its pre-stimulus state. Photographs were taken
145 immediately before the stimulus was applied in order to accurately identify this state. The
146 response was timed using a stopwatch and converted into seconds prior to analysis.
147 Anemones were observed for a maximum of one hour after the stimulus was applied. If an
148 anemone failed to reopen within the hour, no startle response time was recorded. This
149 process was repeated early morning and late afternoon (with at least 6 hours between the
150 morning and afternoon tests) for 2 days before the first fight and one last time the morning
151 of the fight. It was then further repeated 5 times after the first fight to obtain 'between-
152 fight' measures of startle response and again 5 times after both fights in order to obtain
153 'after-fight' measures (15 measures per individual - see figure 1 for details). Between-fight
154 measures of startle response were treated as post-fight startle responses with respect to
155 fight 1 and pre-fight startle responses with respect to fight 2 (figure 1). The total number of
156 startle response observations for each situation was as follows: Pre-fight = 231 (82
157 anemones); Between-fights = 234 (78 anemones); After-fights=220 (78 anemones).

158

159 *Staging contests*

160 On the morning of day 2, the anemones were dislodged from their position on the tank
161 surface and their tanks lined with stones for them to attach to. The sides of the tank were
162 also lined with a thin layer of removable plastic in case the anemones re-adhered to the
163 tank walls. On the afternoon of day 3, anemones were randomly paired and placed into the
164 centre of a clean tank containing 700ml of aerated and filtered seawater. The anemones
165 were positioned such that they were in contact with one another, which stimulates them to
166 fight over territory. This was defined as the beginning of the fight and fights were
167 considered concluded when one individual (the loser) either moved away from its opponent
168 by an approximate distance of one pedal disc (estimated visually) or retracted its tentacles
169 completely for at least 10 mins (Rudin & Briffa, 2011; 2012). If both opponents performed
170 these retreating behaviours, the outcome of the fight was classified as a draw. Contest
171 duration was then back-calculated from the time of initial contact to the time at which the
172 loser first began to move away from its opponent or first retracted its tentacles completely.
173 At the end of the contest, the number of acrorhagial peels inflicted on each opponent was
174 counted and the fights were classified into two escalation categories –‘no peel’ or ‘peel’ –
175 depending on whether or not peels were inflicted. Anemones that failed to fight were
176 removed from the experiment.

177 In order to observe the effects of within-fight experience and post-fight changes in
178 boldness on future contest success and behaviour, a second fight was staged on the
179 afternoon of day 10. In order to investigate how the prior contest outcomes of the focal and
180 opponent interact to affect subsequent fights, individuals from the first fights were paired
181 according to their victory status (winner or loser – individuals who drew in their first fight

182 were excluded from the rest of the study [$N= 6$]) in a fully orthogonal design. Individuals
183 were randomly allocated as either focal (F) or opponent (O) for the second fight based on
184 their first fight ID, i.e. the pair of anemones that fought in first fight 1 were allocated as focal
185 individuals in the second fight while the anemone pair that fought in first fight 2 were
186 allocated as opponents in the second fight and so on. This resulted in a combination of four
187 pairings of focal and opponent individuals: - winner (F) - loser (O) ($N = 10$ pairs); loser (F) -
188 winner (O) ($N = 9$ pairs); winner (F) - winner (O) ($N = 11$ pairs); loser (F) - loser (O) ($N = 11$
189 pairs) (A total of 41 focals and 41 opponents). Individuals were never re-paired with the
190 same opponent from their previous fight. Fights were then staged as outlined above, but
191 this time data were only taken for focal individuals within pairs. As before, anemones that
192 failed to fight were removed from the experiment.

193 After both contests had taken place, the minimum and maximum pedal disc
194 diameter of each anemone was measured using callipers to the nearest 0.1 mm. As pedal
195 disc shape is often elliptical, body size was then calculated as the average of the maximum
196 and minimum diameter (Brace & Quicke, 1986). A small piece of pedal disc tissue
197 (approximately 1cm x 1cm) was removed using a scalpel and stored in 100% molecular
198 grade ethanol for genetic analysis at a later date.

199

200 *Ethical Note*

201 After use in this study all anemones were returned to the collection site at Portwrinkle. No
202 licences or permits were required for this study.

203 *Statistical analyses*

204 *Calculating and comparing repeatability of startle response duration*

205 To determine the repeatability of startle response duration across fight outcomes (winner,
206 loser, draw), fight types (peel, no peel) and situations (pre-fight, post-fight), we conducted
207 Bayesian Markov chain Monte Carlo (MCMC) generalised linear-mixed models (using R
208 package MCMCglmm, Hadfield, 2010) to calculate repeatability (R) and its confidence
209 intervals separately for each round of fights (first, second) (Royauté, Buddle, & Vincent,
210 2015). To determine the effect of fight outcome and situation (pre- or post- fight) on R , we
211 first created an outcome x situation interaction variable containing all possible combinations
212 of outcome and situation. We then fitted a linear mixed model (lmm) with outcome,
213 situation and the outcome x situation interaction variable as fixed effects and the
214 interaction variable and ID as random effects. To determine the combined effect of fight
215 outcome, fight type and situation on R , we created an outcome x situation x fight type
216 interaction variable containing all possible combinations of outcome, situation and fight
217 type. We then fitted an lmm with outcome, situation, fight type and the interaction variable
218 as fixed effects alongside the interaction variable and ID as random effects. For both models,
219 we then extracted the situation and outcome (along with fight type for model 2) specific
220 posterior mode variance components (between individuals / G-structure, within individuals
221 / R-structure) and from these calculated posterior mode values for R . We then compared R
222 values by calculating differences in repeatability (ΔR) across outcomes, fight types and
223 situations. Differences were deemed to be significant if the 95% CIs of ΔR did not span zero.
224 See appendix for more details on this approach.

225 *Fighting experience and mean-level boldness*

226 To investigate the relationships between boldness and within-fight experience we
227 conducted a series of generalised linear-mixed models (glmm) using the R package lme4
228 (Bates, Maechler, Bolker, & Walker, 2015). Minimal adequate models were reached based
229 on significance values gained from model comparison using likelihood ratio tests.

230 As our response data consisted of a mixture of binary, categorical and continuous
231 variables, the type of model and error family changed depending on the nature of the
232 response variable. For models with binary variables (fight outcome and fight type) as the
233 response variable, a binomial error family was used in a glmm. Fixed effects included in
234 these models were average pre-fight boldness, number of peels received and number of
235 peels inflicted. Homoscedasticity was checked for by visual inspection of the model residuals.
236 When examining the effect of first fight outcome on second fight outcome, first fight
237 outcome was also included as a fixed effect. First fight ID was included in all models as a
238 random effect to control for pseudoreplication resulting from taking two data points from
239 the same fight. This was not necessary for the second fight as data points were only taken
240 from one (focal) individual per fight in this second contest. Individual ID was also included as
241 a random effect in models with startle response duration as the response variable since
242 there were multiple startle response durations for each individual. For models with
243 continuous response variables (startle response duration, number of peels inflicted, number
244 of peels received), a linear mixed model (lmm) was used. Due to the non-normal distribution
245 of startle response duration, this variable was log₁₀ transformed before analysis. When
246 analysing the effect of fight outcome on post-fight startle response duration, fight outcome,
247 fight type and situation were included as fixed effects in the model. When analysing the
248 effect of boldness and fighting experience on the number of peels inflicted and received in

249 the second fight, average pre-fight boldness was included as a fixed effect. Relative size
250 difference (RSD) between opponents was calculated following Briffa, Elwood, & Dick, 1998
251 ($RSD = 1 - (\text{opponent size} / \text{focal size})$) and was included as a covariate in all models.

252

253 RESULTS

254 *Pre-fight boldness as a predictor of fight outcome*

255 In the first fight, average pre-fight boldness had a significant effect on fighting success ($\chi^2 =$
256 4.37, $P = 0.037$), with bolder individuals winning more fights than shyer individuals (figure
257 2a). Average pre-fight boldness also predicted whether or not an individual received ($\chi^2 =$
258 4.28, $P = 0.039$) or inflicted peels ($\chi^2 = 8.59$, $P = 0.003$) in the first fight, with bolder
259 individuals being more likely on average to inflict and receive peels than their shyer
260 counterparts. In the second fight however, average pre-fight boldness (i.e. the startle
261 responses recorded between the first and second fight) did not significantly predict the
262 chance of victory ($\chi^2 = 1.01$, $P = 0.31$) or the likelihood of receiving ($\chi^2 = 0.53$, $P = 0.47$) or
263 inflicting peels ($\chi^2 = 0.37$, $P = 0.54$), but rather predicted whether a fight ended in a clear
264 outcome or in a draw ($\chi^2 = 3.91$, $P = 0.048$) (figure 2b), such that focal anemones that drew
265 their second fight had longer startle responses prior to this fight than focal anemones that
266 won or lost it. In order to check whether the loss of correlation between pre-fight boldness
267 and the likelihood of inflicting peels was driving the breakdown of the link between pre-fight
268 boldness and fighting success, we analysed second fights separately according to fight type.
269 For both types of second fight (those that involved peels and those that did not) we again
270 found that pre-fight boldness significantly affected whether a fight ended in a draw or a

271 clear outcome (No Peels: $X^2 = 4.51$, $P = 0.03$; Peels: $X^2 = 3.63$, $P = 0.005$) but not whether an
272 individual won or lost (No Peels: $X^2 = 0.37$; Peels: $X^2 = 2.93$, $P = 0.087$). There was no
273 significant effect of pre-fight boldness on the number of peels received (1st fight: $X^2 = 1.25$,
274 $P = 0.26$; 2nd fight: $X^2 = 1.69$, $P = 0.19$) or inflicted (1st fight: $X^2 = 0.05$, $P = 0.82$; 2nd fight: $X^2 =$
275 $2e-04$, $P = 0.99$) in either fight. Relative size difference had no effect on either first ($X^2 = 1.48$,
276 $P = 0.22$) or second fight outcome ($X^2 = 0.19$, $P = 0.67$).

277

278 *Effect of fighting experience on boldness – repeatability and mean-level*

279 Boldness (measured as startle response duration) was found to be significantly repeatable
280 across all individuals, situations (pre-fight, between-fight and post-fight) and fights (1st and
281 2nd fight) regardless of fight outcome (table 1). Furthermore, there was no significant effect
282 of fight type on the repeatability of boldness in any of the three situations irrespective of
283 victory status. The only exception was a significant difference in the repeatability of post-
284 fight boldness after the second fight between winners of 'no peel' fights and winners of
285 fights involving peels. The post-fight boldness of individuals who won 'no peel' fights was
286 significantly more repeatable than that of winners of fights involving peels (table 1).

287 First fight outcome had no effect on average ($X^2 = 1.15$, $P = 0.28$) or immediate (i.e.
288 the first startle response in the sequence) post-fight boldness ($X^2 = 1.72$, $P = 0.19$). Second
289 fight outcome on the other hand had a significant effect on immediate post-fight boldness.
290 Anemones that lost their second fight significantly increased their startle response in the
291 first 24 hours post-fight ($X^2 = 8.65$, $P = 0.01$) (Figure 3). However this significant increase did
292 not persist past these first 24 hours, with no significant difference in losers' average post-

293 fight startle response ($\chi^2 = 3.99, P = 0.14$). The number of peels received in a fight had no
294 effect on average post-fight boldness in either the first ($\chi^2 = 0.03, P = 0.87$) or the second
295 fight ($\chi^2 = 0.21, P = 0.65$).

296

297 *Effect of within-fight experience on second fight*

298 There was no significant effect of first fight outcome on second fight outcome ($\chi^2 = 0.13, P =$
299 0.722). There was also no effect of opponent victory status (i.e. whether they had won or
300 lost the first fight) on the second fight outcome of focal individuals ($\chi^2 = 0.17, P = 0.68$), nor
301 was there a significant interaction between focal and opponent status ($\chi^2 = 1.84, P = 0.18$).
302 The number of peels inflicted in the first fight had no effect on second fight outcome ($\chi^2 =$
303 $0.52, P = 0.47$) and although there was a trend between the number of peels received in the
304 first fight and second fight outcome, this trend was not statistically significant ($\chi^2 = 3.18, P =$
305 0.07).

306

307 DISCUSSION

308 Resource holding potential is defined as a phenotypic trait that will increase the likelihood
309 of victory in a contest. In contrast to motivational state, which should vary from encounter
310 to encounter, RHP traits might be subject to post-fight change but they should be relatively
311 stable between episodes of fighting. Although resource value dependent changes in
312 motivation within a fight can drive changes in startle response duration (Elwood & Briffa
313 2001), this index of boldness is consistent between fights in several species. Since
314 individuals that are bolder outside of a fight situation show a higher probability of winning

315 compared to shyer individuals, boldness appears to be an RHP component in many species.
316 For example, consistent pre-fight boldness been shown to predict subsequent fighting
317 success in the beadlet sea anemone *Actinia equina* (Rudin & Briffa, 2012). However, our
318 findings indicate that in *A.equina*, boldness may only determine fighting success in an
319 individual's first fight. We found that in the first fight, pre-fight boldness determined
320 whether an individual won or lost, while in the second fight, boldness no longer influenced
321 an individual's victory but rather whether the fight ended in a draw or a clear outcome. Thus,
322 although consistent boldness appears to act as an RHP trait (determining the chance of
323 victory in a subsequent fight) its influence appears to vary with recent experience.

324 In agreement with previous work (Rudin & Briffa, 2012), we found that pre-fight
325 boldness significantly predicted whether an individual won or lost a fight, bolder individuals
326 winning more fights on average than shyer individuals. However, this effect was only
327 present in the first fight, not the second fight. In the second fight, pre-fight boldness did not
328 predict fighting success per se but rather whether a fight ended in a clear outcome or a
329 draw, with shyer individuals drawing more often than bolder individuals. Pre-fight boldness
330 also had differential effects on the injury state of individuals across the two fights. In the
331 first fight, as well as predicting fighting success, pre-fight boldness predicted whether or not
332 an individual inflicted or received peels. Bolder individuals were not only more likely to
333 inflict peels (boldness has previously been found to covary with aggressiveness –
334 Huntingford, 1976; Rudin & Briffa, 2012) but were also more likely to receive peels than
335 their shyer counterparts. In the second fight, however, this correlation between boldness
336 and injurious fighting was absent. This suggests that while shyer individuals were more likely
337 to lose their first fight, they were less likely to become injured in the process. Low boldness

338 also appears to have advantages in other examples of fighting. In the hermit crab *Pagurus*
339 *bernhardus*, for example, shy individuals are better able to resist eviction from their shells
340 when attacked (Courtene-Jones & Briffa, 2014). In the case of hermit crabs the advantage of
341 being shy (at least when playing a defender-role) is obvious since these individuals are more
342 likely to win. In the present example of anemones the advantages of shy behaviour are less
343 clear, since it was the bold individuals rather than the shy ones that were more likely to win.
344 Perhaps then, these differences between fight-outcomes for bold and shy anemones
345 represent alternative strategies; bold individuals are more likely to win territory but at the
346 cost of injuries, whereas shy individuals avoid injuries but at the cost of losing a territory.
347 Such a scenario has clear parallels with the predictions of the classic Hawk-Dove game
348 (Maynard Smith and Price 1973, Maynard Smith and Parker 1976), which predicts a stable
349 mix (i.e. a mixed evolutionarily stable strategy, ESS) of injurious and non-injurious fighting
350 strategies if the costs of injury are (on average) greater than the value of the contested
351 resource. Such a mix can arise in two ways, either through a mixture of consistently hawkish
352 and dove-like individuals, or through a population of individuals that fluctuate between both
353 strategies with the proportion of time playing each governed by the ratio of costs to
354 resource value. Indeed, the actual agonistic behaviour of the anemones in this study
355 indicates a relatively stable mix of injurious and non-injurious fighting since the proportion
356 of fights with peels did not differ between the first and second fights. On the other hand the
357 link between boldness and fighting tactics does not appear to be stable since the effect of
358 pre-fight boldness on the likelihood of inflicting or receiving injury in the first fight was
359 absent in the second fight.

360

361 It is important to note here that while we refer to these fights as ‘first’ and ‘second’,
362 the anemones used were collected from the wild, and are likely to have experienced fights
363 prior to this experiment. However, none of the anemones possessed any sign of injury from
364 recent fights when collected from the shore and all individuals were housed for 7-14 days
365 before being fought. Thus while these may well not be their true first and second fights, any
366 experience effects leftover from prior contests in the wild would very likely have dissipated
367 by the time this experiment was run. Furthermore, we found no effect of prior fighting
368 experience (winning or losing, receiving or inflicting injuries) on second fight outcome or
369 behaviour, an absence which could possibly be due to the length of time between the first
370 and second fight in our study being too long. It has previously been shown that the effects
371 of fighting experience can last for very specific amounts of time, for example when male
372 broad-horned flour beetles *Gnathocerus cornutus* lose fights, they exhibit behavioural
373 changes which last for exactly four days after the fight, returning to their pre-fight state on
374 day 5 (Okada, Yamane, & Miyatake, 2010). However, as information on the duration of
375 fighting experience effects in anemones is currently lacking we cannot make any conclusions
376 about timing effects in *A. equina*.

377

378 Although boldness is regarded as a highly repeatable behaviour in many species,
379 previous studies have shown that both the repeatability of boldness and mean-level
380 boldness can be significantly affected by fighting experience, especially in recipients of
381 agonistic behaviour (Frost, Winrow-Giffen, Ashley, & Sneddon, 2007; Rudin & Briffa, 2012;
382 Courtene-Jones & Briffa, 2014; Briffa, Sneddon, & Wilson, 2015). However, the results of our
383 study illustrate that in *A. equina* boldness is generally robust to the effects of fighting

384 experience. We found that boldness was significantly repeatable regardless of when it was
385 measured (pre-fight, between-fights, post-fight) and furthermore that the repeatability of
386 boldness was generally unaffected by an individual's within-fight experience (i.e. whether it
387 won or lost, engaged in fights with or without peels). The only instance in which the
388 repeatability of boldness was apparently altered was seen in winners after the second fight.
389 Individuals who won fights involving peels exhibited significantly lower boldness
390 repeatability than individuals who had won fights not involving peels, however there was no
391 significant change in boldness repeatability within these groups across the two situations
392 (i.e. between pre- and post-fight measures). This result may signify costs associated with
393 competing in an injurious fight, especially as both inflicting and receiving injuries may pose
394 costs in *A. equina* (Lane & Briffa, 2017). Mean-level boldness also appeared generally robust
395 to the effects of fighting experience, the only exception being a significant decrease in the
396 immediate boldness of losers after the second fight. As previous work has found losers to be
397 more susceptible to the effects of fighting experience on boldness (Frost, Winrow-Giffen,
398 Ashley, & Sneddon, 2007; Rudin & Briffa, 2012; Courtene-Jones & Briffa, 2014), this result is
399 perhaps not surprising. However, why the first fight did not elicit a similar response in losers
400 is unclear. It is possible that there may be a cumulative effect of fighting experience on the
401 boldness of losers, but we did not detect any such effect in our study.

402 Fighting experience can have significant effects on traits that contribute to an
403 individual's resource holding potential, which can in turn affect behaviour and success in
404 subsequent contests (Hsu & Wolf, 2001; Rutte, Taborsky, & Brinkhof, 2006). For example,
405 losing a fight can cause a reduction in RHP via injury or the physiological costs of fighting,
406 causing individuals who have lost a fight to be more likely to lose the next fight they enter, a

407 phenomenon known as the loser effect (winner effects also exist). While previous studies
408 have shown that fighting experience can significantly impact boldness, we have found the
409 opposite, that in *A. equina*, fighting experience (at least initially) has very little impact on
410 boldness but does affect the importance of boldness as an RHP trait, specifically the link
411 between boldness and fighting tactics that was present in the first fight was absent in the
412 second fight. Thus our study indicates that fighting experience can not only change an
413 individual's RHP (as shown elsewhere) but may also impact which traits contribute to RHP as
414 well. One example of such phenomena has been seen in the New Zealand jumping spider
415 *Trite planiceps*, in which the size advantage gained from being substantially bigger than your
416 opponent is lost once an individual is injured, relative injury level becoming more important
417 in determining fight outcome than relative size (Taylor & Jackson, 2003).

418 Here we have shown that despite being a highly repeatable trait, robust to the
419 effects of fighting experience, boldness does not consistently predict fighting success in *A.*
420 *equina*, determining first fight but not second fight success. Our findings suggest that
421 different traits may be important in determining fighting success in consecutive fights and
422 moreover that fighting experience may alter which traits contribute to an individual's RHP.

423

424 **Acknowledgments**

425 We thank Ann Torr for help with collecting anemones. This study was funded by a BBSRC
426 grant awarded to M.B. (grant no. BB/ M019772/1).

427

428

429 **REFERENCES**

430 Barlow, G., Rogers, W., & Fraley, N. (1986) Do Midas cichlids win through prowess or daring?
431 It depends. *Behavioural Ecology & Sociobiology*, 19, 1-8.

432

433 Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015) Fitting linear mixed-effects models
434 using lme4. *Journal of Statistical Software*, 67, 1-48. (doi: 10.18637/jss.v067.i01)

435

436 Bigger, C. H. (1982) The cellular basis of the aggressive acrorhagial response of sea-
437 anemones. *Journal of Morphology*, 173, 259-278.

438

439 Brace, R. C., Pavey, J., & Quicke, D. L. J. (1979) Intraspecific aggression in the colour morphs
440 of the anemone *Actinia equina*: the convention governing dominance ranking. *Animal*
441 *Behaviour*, 27, 553-561.

442

443 Brace, R. C., & Quicke, L. J. (1986) Dynamics of colonization by the beadlet anemone, *Actinia*
444 *equina*. *Journal of the Marine Biological Association of the United Kingdom*, 66, 21-47.

445

446 Briffa, M, Elwood, R. W., & Dick, J. T. A. (1998) Analysis of repeated signals during shell fights
447 in the hermit crab *Pagurus bernhardus*. *Proceedings of the Royal Society B*, 265, 1467-1474.

448

449 Briffa, M., Rundle, S. D., & Fryer, A. (2008) Comparing the strength of behavioural plasticity
450 and consistency across situations: animal personalities in the hermit crab *Pagurus*
451 *bernhardus*. *Proceedings of the Royal Society B*, 275, 1305-1311. (doi:
452 10.1098/rspb.2008.0025)

453

454 Briffa, M., & Greenaway, J. (2011) High *In situ* repeatability of behaviour indicates animal
455 personality in the beadlet anemone *Actinia equina* (Cnidaria). *PLoS One*, 6, e21963. (doi:
456 10.1371/journal.pone.0021963)

457

458 Briffa, M., Sneddon, L. U., & Wilson, A. J. (2015) Animal personality as a cause and
459 consequence of contest behaviour. *Biology Letters*, 11, 20141007. (doi:
460 10.1098/rsbl.2014.1007)

461

462 Courtene-Jones, W., & Briffa, M. (2014) Boldness and asymmetric contests : role- and
463 outcome-dependent effects of fighting in hermit crabs. *Behavioural Ecology*, 25, 1073-1082.
464 (doi: 10.1093/beheco/aru085)

465

466 Elwood, R.W. & Briffa, M. (2001) Information gathering and communication during agonistic
467 encounters: A case study of hermit crabs. *Advances in the Study of Behavior*, 30, 53-97.

468

469 Frost, A., Winrow-Giffen, A., Ashley, P. J., & Sneddon, L. U. (2007) Plasticity in animal
470 personality traits: does prior experience alter the degree of boldness? *Proceedings of the*
471 *Royal Society B*, 274, 333-339. (doi: 10.1098/rspb.2006.3751)

472

473 Hadfield, J. D. (2010) MCMC Methods for Multi-Response Generalized Linear Mixed Models:
474 The MCMCglmm R Package. *Journal of Statistical Software*, 33, 1-22. (URL: -
475 [http://www.jstatsoft.org/v33/i02/.](http://www.jstatsoft.org/v33/i02/))

476

477 Hsu, Y., & Wolf, L. L. (2001) The winner and loser effect: what fighting behaviours are
478 influenced? *Animal Behaviour*, 61, 777-786. (doi: 10.1006/anbe.2000.1650)

479

480 Lane, S. M. & Briffa, M. (2017) The price of attack: rethinking damage costs in animal
481 contests. *Animal Behaviour*, 126, 23-229.

482

483 Manuel, R. L. (1988) *British Anthozoa*. Academic Press, London, UK.

484

485 Maynard Smith, J., & Parker, G. A. (1976) The logic of asymmetric contests. *Animal*
486 *Behaviour*, 24, 159-175.

487

488 Maynard Smith, J., & Price, G. R. (1973) The logic of animal conflict. *Nature*, 246, 15-18.

489

490 Okada, K., Yamane, T., & Miyatake, T. (2010) Ejaculatory strategies associated with
491 experience of losing. *Biology Letters*, *6*, 593-596.

492

493 Parker, G. A. (1974) Assessment strategy and the evolution of fighting behaviour. *Journal of*
494 *Theoretical Biology*, *47*, 223-243. (doi: 10.1016/0022-5193(74)90111-8)

495

496 Royauté, R., Buddle, C. M., & Vincent, C. (2015) Under the influence : sublethal exposure to
497 an insecticide affects personality expression in a jumping spider. *Functional Ecology*, *29*,
498 962-970. (doi: 10.1111/1365-2435.12413)

499

500 Rudin, F. S., & Briffa, M. (2011) The logical polyp: assessments and decisions during contests
501 in the beadlet anemone *Actinia equina*. *Behavioural Ecology*, *22*, 1278-1285. (doi:
502 10.1093/beheco/arr125)

503

504 Rudin, F. S., & Briffa, M. (2012) Is boldness a resource-holding potential trait? Fighting
505 prowess and changes in startle response in the sea anemone *Actinia equina*. *Proceedings of*
506 *the Royal Society B*, *279*, 1904-1910. (doi: 10.1098/rspb.2011.2418)

507

508 Rutte, C., Taborsky, M., Brinkhof, & M. W. G. (2006) What sets the odds of winning and
509 losing? *Trends in Ecology and Evolution*, 21, 16-21. (doi: 10.1016/j.tree.2005.10.014)

510

511 Williams, R. B. (1978) Some recent observations on acrorhagi of sea anemones. *Journal of*
512 *the Marine Biological Association UK*, 80, 719-724.

513

514 Wolf, M., & Weissing, F. J. (2010) An explanatory framework for adaptive personality
515 difference. *Philosophical Transactions of the Royal Society: B*, 365, 3965-3968. (doi:
516 10.1098/rstb.2010.0215)

517

518

519

520

521

522

523

524

525

526

527

528 **APPENDIX**

529 **Model specifications for comparing repeatability of boldness between fight outcomes and**
530 **situations**

531 We first split data into 'first fight' (first fight outcome; pre-fight boldness; between-fights
532 boldness) and 'second fight' (second fight outcome; between-fights boldness; after-fights
533 boldness), analysing these two sets separately in the following way.

534 We began by creating an outcome x situation interaction variable (referred to hereafter as
535 outcome_sit), accounting for every possible combination of outcome and situation. We then
536 used an inverse-wishart prior ($V=\text{diag}(n)$, $\nu=n$), where n is the number of behavioural
537 variables being considered and ν is the degree of belief parameter (Hadfield 2010) to
538 incorporate all combinations of fight outcome and situation, resulting in a 6x6 matrix. We
539 specified a Markov Chain Monte Carlo (MCMC) glmm with 50,000 iterations, a 30,000
540 iteration burn-in and a thinning level of 10. This yielded an MCMC sample size of 2,000 and
541 autocorrelation of <0.1 in all instances. We used a poisson error family due to the count
542 nature of our response variable (startle response duration) and included outcome, situation
543 and outcome_sit as fixed effects along with outcome_sit and ID as random effects. We then
544 extracted the outcome and situation specific posterior variance components (between
545 individuals/ G-structure/R-structure) from this model and used these to calculate posterior
546 mode values for repeatability (R) for each outcome_sit combination. Finally, we compared
547 R values by calculating differences in repeatability (ΔR) across outcomes and situations.

548

549 **Model specifications for comparing repeatability of boldness between fight outcomes,**
550 **fight types and situations**

551 We again split our data into 'first fight' and 'second fight' and analysed these two sets
552 separately.

553 We created an outcome x fight type x situation interaction variable (referred to hereafter as
554 outcome_type_sit), accounting for every possible combination of outcome, fight type and
555 situation. We again used an inverse-wishart prior to create a 12x12 matrix incorporating all
556 combinations of outcome, fight type and situation. We then specified an MCMCglmm with
557 500,000 iterations, a 300,000 iteration burn-in and a thinning level of 10. This yielded an
558 MCMC sample size of 20,000 and autocorrelation of <0.1 in all instances. We again used a
559 poisson error family to account for the count nature of our response variable (startle
560 response duration) and this time included outcome, situation, fight type and
561 outcome_type_sit as fixed effects alongside outcome_type_sit and ID as random effects.
562 We then extracted the outcome and situation specific posterior variance components
563 (between individuals/ G-structure/R-structure) from this model and used these to calculate
564 posterior mode values for repeatability (R) for each outcome _sit combination. Finally, we
565 compared R values by calculating differences in repeatability (ΔR) across outcomes and
566 situations.

567

568

569

570

571 **Table 1** Repeatability of startle response duration (boldness) \pm 95% confidence intervals
572 Repeatability of startle response duration (boldness) \pm 95% confidence intervals (CIs) for
573 each combination of fight outcome and situation and each combination of fight outcome,
574 situation and fight type along with differences in repeatability (ΔR) between outcomes,
575 situations and fight types. Significant values (if the 95% CIs crossed zero) are indicated in
576 bold.
577

	Pre-fight 1	Between-fights (post fight 1)	ΔR (Pre-fight – between-fights)	Pre-fight 2 (between fights)	After fights	ΔR (between-fights – after-fights)
Winners	0.32 [0.18,0.51]	0.25 [0.15,0.46]	-0.02 [-0.28,0.18]	0.35 [0.15,0.46]	0.36 [0.18,0.64]	0.07 [-0.020,0.43]
Losers	0.20 [0.11,0.26]	0.22 [0.13,0.41]	0.002 [-0.15,0.25]	0.22 [0.13,0.41]	0.26 [0.14,0.59]	-0.22 [-0.48,0.16]
Drawers	-	-	-	0.42 [0.19,0.68]	0.24 [0.11,0.49]	-0.13 [-0.44,0.18]
ΔR (Winners – Losers)	-0.08 [-0.32,0.11]	-0.02 [-0.23,0.19]	-	-0.02 [-0.23,0.19]	0.09 [-0.28,0.38]	-
ΔR (Winners –Drawers)	-	-	-	0.19 [-0.17,0.46]	0.12 [-0.19,0.43]	-
ΔR (Losers-Drawers)	-	-	-	0.15 [-0.16,0.47]	-0.03[-0.35,0.28]	-
No Peels: Winners	0.28 [0.15,0.51]	0.21 [0.11,0.43]	-0.04 [-0.29,0.21]	0.27 [0.11,0.69]	0.78 [0.40,0.96]	0.35 [-0.09,0.74]
No Peels: Losers	0.29 [0.15,0.51]	0.19 [0.10,0.39]	0.03 [-0.16,0.33]	0.26 [0.10,0.66]	0.31 [0.10,0.71]	0.050 [-0.43,0.46]
No Peels: Drawers	-	-	-	0.49 [0.22,0.75]	0.29 [0.13,0.57]	-0.19 [-0.50,0.20]
ΔR (No Peels: Winners – Losers)	0.20[-0.16,0.34]	0.20 [-0.30,0.20]	-	0.01 [-0.41,0.46]	0.35 [-0.10,0.75]	-
ΔR (No Peels: Winners – Drawers)	-	-	-	-0.18 [-0.51,0.30]	0.45 [-0.02,0.71]	-
ΔR (No Peels: Losers – Drawers)	-	-	-	-0.17[-0.61,0.34]	0.05 [-0.44,0.51]	-
Peels: Winners	0.42 [0.22,0.69]	0.57 [0.25,0.74]	0.10 [-0.32,0.41]	0.34 [0.13,0.60]	0.22 [0.10,0.54]	-0.05 [-0.40,0.29]
Peels: Losers	0.34 [0.14,0.59]	0.26 [0.14,0.54]	0.004 [-0.36,0.27]	0.31 [0.13,0.63]	0.38 [0.15,0.71]	-0.08 [-0.48,0.39]
Peels: Drawers	-	-	-	0.48 [0.12,0.88]	0.25 [0.22,0.75]	-0.17 [-0.65,0.44]
ΔR (Peels: Winners –Losers)	0.20 [-0.22,0.44]	0.20 [-0.13,0.52]	-	0.001 [-0.36,0.36]	-0.09 [-0.49,0.26]	-
ΔR (Peels: Winners –Drawers)	-	-	-	-0.13 [-0.58,0.34]	-0.03 [-0.54,0.35]	-
ΔR (Peels: Losers – Drawers)	-	-	-	-0.16 [-0.51,0.29]	0.04 [-0.34,0.45]	-
ΔR (Peels: Winners – No Peels: Winners)	0.19 [-0.13,0.47]	0.22 [-0.06,0.52]	-	-0.05 [-0.38,0.42]	0.47 [0.006,0.74]	-
ΔR (Peels: Losers – No Peels: Losers)	-0.11 [-0.41,0.13]	-0.009 [-0.30,0.27]	-	0.006 [-0.40,0.40]	0.078 [-0.34,0.45]	-
ΔR (Peels: Drawers – No Peels: Drawers)	-	-	-	0.04 [-0.47,0.49]	0.05 [-0.49,0.38]	-

578

579


580

581

582

583

584




585

586

587


Figure 1 Overview of the experimental structure. Between-fight startle responses (SR) were treated as post-fight responses with respect to fight 1 and pre-fight responses with respect to fight 2.



597

598

Figure 2 Effect of mean (\pm SE) pre-fight startle response time on (a) first fight outcome and (b) second fight outcome. Asterisks indicate significant differences within each panel.



599 **Figure 3** Mean (\pm SE) startle response duration immediately before and immediately after
 600 the second fight.

601