CALIBRATION OF A FAST FIELD CYCLING RELAXOMETER FOR ULTRA-LOW FIELD MEASUREMENTS OF *T*₁-DISPERSION

Vasileios Zampetoulas, Lionel M. Broche and David J. Lurie

Aberdeen Biomedical Imaging Centre, University of Aberdeen, Foresterhill, Aberdeen, AB25 2ZD, Scotland, United Kingdom www.ffc-mri.org

In Fast-Field Cycling (FFC) MRI the field switches to different levels, allowing for the measurement of NMR parameters such as T_1 over a range of fields [1].

For the successful implementation of the ultra-low field FFC techniques on a commercial benchtop FFC relaxometer (Stelar S.r.l., Italy) the stray magnetic fields need to be compensated. These are determined with the application of FFC measurements in a range of fields close to zero. During this process correction fields of varying magnitude and orientation are applied by the relaxometer while the magnetisation precesses around a stray component of unknown direction with the frequency of precession determined by the magnitude of the transverse and longitudinal fields. The successful compensation of the stray fields is shown as a minimum in the frequency of precession (Figure 1a) [2].

After the successful calibration of the relaxometer FC methods at ultra-low fields can be applied, including extension of the T_1 -dispersion curve acquired to the region of μ T (Figure 1b).




FIGURE 1. (a) Precession frequency against the longitudinal correction field applied. The minimum is observed in the range of 6 to 12 μ T. For values larger than 22 μ T magnetisation decays exponentially as in conventional FFC experiments thus no precession is observed (b) R_1 -dispersion curve (where $R_1=1/T_1$) of a sample of MnCl₂ acquired before and after the calibration, with the range of minimum ¹H Larmor frequencies extending to 100 Hz (corresponding to a field of 2.3 μ T).

Aknowledgements

The author acknowledges funding from the EPSRC through the Centre for Doctoral Training in Integrated Magnetic Resonance.

References

[1] Kimmich R, Anoardo E., Prog. NMR Spectrosc., 44, 257-320 (2004).

[2] Anoardo E, Ferrante GM, Appl. Magn. Reson., 24, 85-96 (2003).

AMPERE NMR School, Zakopane, Poland, June 2014, p92.