

CRYSTALLOGRAPHIC
COMMUNICATIONS

ISSN 2056-9890

Crystal structure of 9-methacryloylanthracene

Aditya Agrahari,^a Patrick O. Wagers,^b Steven M. Schildcrout,^c John Masnovi^{a*} and Wiley J. Youngs^b^aDepartment of Chemistry, Cleveland State University, Cleveland OH 44115, USA, ^bDepartment of Chemistry, University of Akron, Akron OH 44325, USA, and ^cDepartment of Chemistry, Youngstown State University, Youngstown OH 44555, USA. *Correspondence e-mail: j.masnovi@csuohio.edu

Received 20 September 2014

Accepted 26 February 2015


Edited by W. T. A. Harrison, University of Aberdeen, Scotland

Keywords: crystal structure; aldol; methylation; acetylanthracene; stereochemistry**CCDC reference:** 1051418**Supporting information:** this article has supporting information at journals.iucr.org/e

In the title compound, C₁₈H₁₄O, with systematic name 1-(anthracen-9-yl)-2-methylprop-2-en-1-one, the ketonic C atom lies 0.2030 (16) Å out of the anthryl-ring-system plane. The dihedral angle between the planes of the anthryl and methacryloyl moieties is 88.30 (3)° and the stereochemistry about the Csp²–Csp² bond in the side chain is *transoid*. In the crystal, the end rings of the anthryl units in adjacent molecules associate in parallel-planar orientations [shortest centroid–centroid distance = 3.6320 (7) Å]. A weak hydrogen bond is observed between an aromatic H atom and the O atom of a molecule displaced by translation in the *a*-axis direction, forming sheets of parallel-planar anthryl groups packing in this direction.


1. Chemical Context

Enolizable aldehydes react with formaldehyde in strong aqueous base to form polyols, whereas ketones usually react to form polyhydroxyketones (Davidson & Bogert, 1935; Vik *et al.*, 1973; Weissmermel & Arpe, 1997; Wittcoff *et al.*, 2013). Therefore, the observed methylation of 9-acetylanthracene by formaldehyde with alcoholic potassium carbonate (see Scheme below) is remarkable in that the reaction occurs with weak base in a non-aqueous medium by reduction of formaldehyde to form the methyl group (Pande *et al.*, 1998). Consequently, we obtained an X-ray structure determination to confirm the identity of the isolated product, 9-methacryloylanthracene or 1-(9-anthryl)-2-methyl-2-propen-1-one.



2. Structural commentary

The crystal structure (Fig. 1) establishes the material to be the α -methylated aldol condensation product. Bond distances and valence angles agree well between the observed and the calculated structures. The anthryl ring system is essentially planar, as is the methacryloyl substituent (excepting the hydrogen atoms of the methyl group), whereas the calculated structure shows a slight deviation, about 7°, of the methacryloyl skeleton from planarity. The substituted C atom (C9) of the anthryl group also lies in the plane of the substituent, deviating by only 0.002 (2) Å. However, this C atom is puckered, so that the carbonyl C atom resides 0.2030 (16) Å out of the anthryl plane. This puckering is



OPEN ACCESS




Figure 1
ORTEP (30% probability ellipsoids) plot of the title compound showing the atom-labeling scheme.

absent in the calculated structure. The planes of the anthryl and methacryloyl moieties are nearly perpendicular with a dihedral angle of 88.30 (3)° (but about 12° from perpendicular in the calculated structure). This general orientation is demanded by the close intramolecular approach of the methacryloyl group to the *peri*-H atoms (H1 and H8), but packing effects may also contribute to deciding the exact angle since that calculated for the energy minimum differs by about 10° from that observed. The observed positioning is not quite




Figure 2
Views parallel to the planes of both the anthryl and the methacryloyl moieties (top) and parallel to the methacryloyl but perpendicular to the anthryl with H atoms omitted for clarity (bottom).

Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$C3-H3\cdots O^i$	0.95	2.48	3.3747 (16)	157

Symmetry code: (i) $x - 1, y, z$.

symmetrical, with C11 being slightly closer (0.018 Å) to H1 than to H8. Similar geometries are found in 9-acetyl-anthracene, with a dihedral angle of 88.70 (3)° (Andersson *et al.*, 1984) and in 9-(bromoacetyl)anthracene, with a dihedral angle of 74.2 (1)° (Kubo *et al.*, 2007). Unfavorable non-bonded interactions in the present structure are likely the reason that the methyl group, which is bulkier than the methylene group, projects away from the anthryl moiety, making the stereochemistry of the C11–C12 bond *transoid*. The puckering observed at C9 would partially relieve these unfavorable steric interactions occurring about this position.

3. Supramolecular features

Intermolecular close contacts between large aromatic groups in the solid state often involve π – π stacking interactions involving parallel planar associations (Główska *et al.*, 1999). This motif is observed here as well, with the anthryl rings displaced and stacking alternately with those of neighboring molecules (Fig. 2). The centroid–centroid separations are 3.6320 (7) and 3.7532 (7) and 3.7807 (8) Å. The methacryloyl substituent prevents such interactions involving the central ring of the anthryl moiety. A weak hydrogen bond is observed (Fig. 3) between an aromatic H atom (H3) and the O atom of a molecule displaced by translation in the *a*-axis direction




Figure 3
A fragment of a [100] hydrogen-bonded chain of molecules in the crystal showing the intermolecular O...H close contact (dotted line).

(Table 1), resulting in the formation of anthryl groups packing in parallel-planar sheets in this direction.

4. Synthesis and crystallization

Refluxing 9-acetylanthracene (1.0 g), paraformaldehyde (273 mg), and potassium carbonate (942 mg) in 3.0 ml ethanol afforded 80 mg product which eluted first from an alumina column with 10% ethyl acetate–hexane and was crystallized from chloroform–hexane in the form of colorless plates.

5. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. The H atoms were placed in calculated positions and refined as riding atoms, with C–H = 0.95 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C-alkene and C-aromatic})$, and C–H = 0.98 Å and $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{C-methyl})$.

6. Calculations

Density-functional theoretical computations were performed using *Gaussian* software (Frisch *et al.*, 2010) through the Ohio Supercomputing Center (in Columbus OH) with Zhao and Truhlar's hybrid meta exchange-correlation functional, M06-2X, (Choe, 2012; Huh & Choe, 2010; Zhao & Truhlar, 2008), which is parameterized for non-metallic systems with non-covalent π – π interactions for accurate modelling of intramolecular dispersion effects. The basis set used is 6-31+G(d). To obtain the geometry at the global minimum potential energy, optimization was based on the minimum-energy conformation from a two-torsion MM2 plot (*ChemBio3D Ultra 12.0*; www.CambridgeSoft.com) using rotations about the C9–C11 and C11–C12 single bonds. The M06-2X structure has all vibrational frequencies positive, verifying that it is at a potential-energy minimum. Calculated values for geometrical parameters in the optimized isolated molecule are given in the Supporting information.

Acknowledgements

The authors would like to thank the Graduate College and Chemistry Department at Cleveland State University for support, the Ohio Supercomputing Center for a grant of computer time, and the National Science Foundation (CHE-0840446) for funds used to purchase the Bruker APEXII DUO X-ray diffractometer used in this research.

Table 2
Experimental details.

Crystal data	
Chemical formula	C ₁₈ H ₁₄ O
M_r	246.29
Crystal system, space group	Triclinic, $P\bar{1}$
Temperature (K)	100
a, b, c (Å)	8.7602 (5), 9.1784 (5), 9.2032 (5)
α, β, γ (°)	67.206 (2), 71.670 (3), 75.195 (2)
V (Å ³)	639.98 (6)
Z	2
Radiation type	Mo $K\alpha$
μ (mm ⁻¹)	0.08
Crystal size (mm)	0.21 × 0.17 × 0.05
Data collection	
Diffractometer	Bruker APEXII CCD
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 1997)
$T_{\text{min}}, T_{\text{max}}$	0.984, 0.996
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	12757, 2590, 2348
R_{int}	0.026
$(\sin \theta/\lambda)_{\text{max}}$ (Å ⁻¹)	0.623
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.036, 0.106, 1.06
No. of reflections	2590
No. of parameters	173
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å ⁻³)	0.26, -0.21

Computer programs: *APEX2* and *SAINT* (Bruker, 1997), *SHELXS97* and *SHELXTL* (Sheldrick, 2008).

References

- Andersson, K., Becker, H. D., Engelhardt, L. M., Hansen, L. & White, A. H. (1984). *Aust. J. Chem.* **37**, 1337–1340.
- Bruker (1997). *APEX2, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Choe, S. J. (2012). *Bull. Korean Chem. Soc.* **33**, 2861–2866.
- Davidson, D. & Bogert, M. T. (1935). *J. Am. Chem. Soc.* **57**, 905–905.
- Frisch, M. J. *et al.* (2010). *Gaussian 09*. Revision C.01. Gaussian Inc., Wallingford, CT, USA.
- Główska, M. L., Martynowski, D. & Kozłowska, K. (1999). *J. Mol. Struct.* **474**, 81–89.
- Huh, D. S. & Choe, S. J. (2010). *J. Porphyrins Phthalocyanines*, **14**, 592–604.
- Kubo, K., Watanabe, K. & Sakurai, T. (2007). *Acta Cryst.* **E63**, o1300–o1301.
- Pande, P. P., Joshi, G. C. & Mathela, C. S. (1998). *Synth. Commun.* **28**, 4193–4200.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Vik, J. E., Kierkegaard, C., Pappas, J., Skaarup, S., Aaltonen, R. & Swahn, C. G. (1973). *Acta Chem. Scand.* **27**, 251–257.
- Weissermel, K. & Arpe, H.-J. (1997). In *Industrial Organic Chemistry*. Weinheim: VCH Verlagsgesellschaft.
- Wittcoff, H. A., Reuben, B. G. & Plotkin, J. S. (2013). In *Industrial Organic Chemicals*, 2nd ed. Hoboken, NJ: Wiley.
- Zhao, Y. & Truhlar, D. G. (2008). *Theor. Chem. Acc.* **120**, 215–241.

supporting information

Acta Cryst. (2015). E71, 357-359 [doi:10.1107/S2056989015004090]

Crystal structure of 9-methacryloylanthracene

Aditya Agrahari, Patrick O. Wagers, Steven M. Schildcrout, John Masnovi and Wiley J. Youngs

Computing details

Data collection: *APEX2* (Bruker, 1997); cell refinement: *SAINT* (Bruker, 1997); data reduction: *SAINT* (Bruker, 1997); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXS97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

1-(Anthracen-9-yl)-2-methylprop-2-en-1-one

Crystal data

$C_{18}H_{14}O$

$M_r = 246.29$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 8.7602$ (5) Å

$b = 9.1784$ (5) Å

$c = 9.2032$ (5) Å

$\alpha = 67.206$ (2)°

$\beta = 71.670$ (3)°

$\gamma = 75.195$ (2)°

$V = 639.98$ (6) Å³

$Z = 2$

$F(000) = 260$

$D_x = 1.278$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9953 reflections

$\theta = 2.5$ – 28.4 °

$\mu = 0.08$ mm⁻¹

$T = 100$ K

Plate, colourless

$0.21 \times 0.17 \times 0.05$ mm

Data collection

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 1997)

$T_{\min} = 0.984$, $T_{\max} = 0.996$

12757 measured reflections

2590 independent reflections

2348 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\max} = 26.3$ °, $\theta_{\min} = 2.4$ °

$h = -10 \rightarrow 10$

$k = -11 \rightarrow 11$

$l = -11 \rightarrow 11$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.036$

$wR(F^2) = 0.106$

$S = 1.06$

2590 reflections

173 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0479P)^2 + 0.2652P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.26$ e Å⁻³

$\Delta\rho_{\min} = -0.21$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O	0.88771 (10)	0.57102 (9)	0.67719 (10)	0.0221 (2)
C1	0.49492 (14)	0.63847 (13)	0.64670 (14)	0.0176 (2)
C2	0.35652 (15)	0.59642 (13)	0.64748 (15)	0.0203 (3)
C3	0.21748 (14)	0.59128 (14)	0.77952 (15)	0.0214 (3)
C4	0.22034 (14)	0.63126 (13)	0.90640 (14)	0.0202 (3)
C4a	0.36214 (13)	0.67831 (13)	0.91022 (14)	0.0164 (2)
C5	0.51116 (15)	0.80977 (14)	1.17279 (14)	0.0206 (3)
C6	0.64792 (16)	0.85245 (14)	1.17433 (14)	0.0229 (3)
C7	0.78992 (15)	0.85161 (14)	1.04616 (15)	0.0223 (3)
C8	0.79096 (14)	0.80876 (13)	0.91974 (14)	0.0183 (3)
C8a	0.64945 (13)	0.76406 (12)	0.91228 (13)	0.0153 (2)
C9	0.64571 (13)	0.71974 (12)	0.78388 (13)	0.0148 (2)
C9a	0.50386 (13)	0.67960 (12)	0.77832 (13)	0.0152 (2)
C10	0.36743 (14)	0.71921 (13)	1.03939 (13)	0.0179 (3)
C10a	0.50634 (14)	0.76402 (13)	1.04251 (13)	0.0166 (2)
C11	0.80055 (13)	0.70042 (13)	0.65638 (13)	0.0150 (2)
C12	0.84479 (14)	0.83810 (13)	0.50679 (13)	0.0178 (2)
C13	0.74810 (16)	0.97827 (14)	0.48462 (15)	0.0242 (3)
C14	1.00033 (15)	0.80523 (15)	0.38767 (15)	0.0264 (3)
H1	0.5869	0.6405	0.5575	0.021*
H2	0.3528	0.5701	0.5586	0.024*
H3	0.1223	0.5599	0.7793	0.026*
H4	0.1265	0.6278	0.9939	0.024*
H5	0.4172	0.8103	1.2596	0.025*
H6	0.6486	0.8830	1.2616	0.027*
H7	0.8851	0.8812	1.0487	0.027*
H8	0.8871	0.8086	0.8353	0.022*
H10	0.2741	0.7164	1.1273	0.022*
H13A	0.6505	0.9906	0.5644	0.029*
H13B	0.7765	1.0662	0.3890	0.029*
H14A	1.0155	0.9009	0.2910	0.040*
H14B	0.9950	0.7164	0.3564	0.040*
H14C	1.0920	0.7770	0.4379	0.040*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O	0.0204 (4)	0.0189 (4)	0.0224 (4)	0.0011 (3)	-0.0021 (3)	-0.0070 (3)
C1	0.0187 (5)	0.0149 (5)	0.0175 (5)	-0.0013 (4)	-0.0045 (4)	-0.0042 (4)
C2	0.0244 (6)	0.0157 (5)	0.0228 (6)	-0.0011 (4)	-0.0109 (5)	-0.0057 (4)
C3	0.0177 (6)	0.0166 (5)	0.0285 (6)	-0.0039 (4)	-0.0105 (5)	-0.0015 (5)
C4	0.0148 (5)	0.0174 (5)	0.0218 (6)	-0.0025 (4)	-0.0032 (4)	-0.0007 (4)
C4a	0.0153 (5)	0.0118 (5)	0.0171 (5)	-0.0006 (4)	-0.0038 (4)	-0.0007 (4)
C5	0.0265 (6)	0.0168 (5)	0.0146 (5)	-0.0013 (5)	-0.0018 (5)	-0.0050 (4)
C6	0.0351 (7)	0.0191 (6)	0.0170 (6)	-0.0030 (5)	-0.0085 (5)	-0.0076 (5)
C7	0.0264 (6)	0.0193 (6)	0.0241 (6)	-0.0051 (5)	-0.0101 (5)	-0.0063 (5)
C8	0.0184 (6)	0.0168 (5)	0.0188 (6)	-0.0025 (4)	-0.0038 (4)	-0.0057 (4)
C8a	0.0173 (5)	0.0116 (5)	0.0149 (5)	-0.0007 (4)	-0.0040 (4)	-0.0030 (4)
C9	0.0158 (5)	0.0112 (5)	0.0141 (5)	-0.0009 (4)	-0.0030 (4)	-0.0018 (4)
C9a	0.0159 (5)	0.0109 (5)	0.0155 (5)	-0.0004 (4)	-0.0042 (4)	-0.0018 (4)
C10	0.0166 (5)	0.0153 (5)	0.0146 (5)	-0.0002 (4)	0.0004 (4)	-0.0021 (4)
C10a	0.0201 (6)	0.0122 (5)	0.0136 (5)	0.0000 (4)	-0.0031 (4)	-0.0025 (4)
C11	0.0146 (5)	0.0180 (5)	0.0155 (5)	-0.0037 (4)	-0.0041 (4)	-0.0076 (4)
C12	0.0193 (6)	0.0200 (6)	0.0154 (5)	-0.0071 (4)	-0.0017 (4)	-0.0068 (4)
C13	0.0299 (6)	0.0190 (6)	0.0189 (6)	-0.0054 (5)	-0.0016 (5)	-0.0037 (5)
C14	0.0249 (6)	0.0269 (6)	0.0215 (6)	-0.0077 (5)	0.0039 (5)	-0.0070 (5)

Geometric parameters (\AA , $^\circ$)

O—C11	1.2176 (13)	C9—C8a	1.4024 (15)
C1—C2	1.3603 (17)	C9—C9a	1.4040 (16)
C1—H1	0.9500	C9—C11	1.5117 (14)
C2—C3	1.4201 (17)	C9a—C1	1.4300 (16)
C2—H2	0.9500	C9a—C4a	1.4365 (15)
C3—C4	1.3615 (17)	C10—C10a	1.3920 (17)
C3—H3	0.9500	C10—H10	0.9500
C4—C4a	1.4297 (16)	C10a—C5	1.4307 (16)
C4—H4	0.9500	C10a—C8a	1.4363 (15)
C4a—C10	1.3948 (16)	C11—C12	1.4864 (15)
C5—C6	1.3579 (18)	C12—C13	1.3277 (17)
C5—H5	0.9500	C12—C14	1.5020 (16)
C6—C7	1.4207 (17)	C13—H13A	0.9500
C6—H6	0.9500	C13—H13B	0.9500
C7—C8	1.3618 (16)	C14—H14A	0.9800
C7—H7	0.9500	C14—H14B	0.9800
C8—C8a	1.4301 (16)	C14—H14C	0.9800
C8—H8	0.9500		
C1—C9a—C4a	118.42 (10)	C9—C8a—C10a	119.11 (10)
C2—C1—C9a	120.88 (11)	C8—C8a—C10a	118.27 (10)
C2—C1—H1	119.6	C8a—C9—C9a	121.28 (10)
C9a—C1—H1	119.6	C8a—C9—C11	119.66 (10)

C1—C2—C3	120.84 (11)	C9a—C9—C11	118.84 (10)
C1—C2—H2	119.6	C9—C9a—C1	122.57 (10)
C3—C2—H2	119.6	C9—C9a—C4a	119.02 (10)
C4—C3—C2	120.12 (11)	C10a—C10—C4a	121.54 (10)
C4—C3—H3	119.9	C10a—C10—H10	119.2
C2—C3—H3	119.9	C4a—C10—H10	119.2
C3—C4—C4a	121.14 (11)	C10—C10a—C5	121.77 (10)
C3—C4—H4	119.4	C10—C10a—C8a	119.51 (10)
C4a—C4—H4	119.4	C5—C10a—C8a	118.72 (11)
C10—C4a—C4	121.93 (10)	O—C11—C12	120.54 (10)
C10—C4a—C9a	119.49 (10)	O—C11—C9	119.34 (10)
C4—C4a—C9a	118.57 (10)	C12—C11—C9	120.11 (9)
C6—C5—C10a	121.02 (11)	C13—C12—C11	120.05 (10)
C6—C5—H5	119.5	C13—C12—C14	124.29 (11)
C10a—C5—H5	119.5	C11—C12—C14	115.66 (10)
C5—C6—C7	120.35 (11)	C12—C13—H13A	120.0
C5—C6—H6	119.8	C12—C13—H13B	120.0
C7—C6—H6	119.8	H13A—C13—H13B	120.0
C8—C7—C6	120.66 (11)	C12—C14—H14A	109.5
C8—C7—H7	119.7	C12—C14—H14B	109.5
C6—C7—H7	119.7	H14A—C14—H14B	109.5
C7—C8—C8a	120.98 (11)	C12—C14—H14C	109.5
C7—C8—H8	119.5	H14A—C14—H14C	109.5
C8a—C8—H8	119.5	H14B—C14—H14C	109.5
C9—C8a—C8	122.61 (10)		
C9a—C1—C2—C3	-0.39 (17)	C9a—C9—C11—O	-85.73 (13)
C1—C2—C3—C4	1.10 (17)	C8a—C9—C11—C12	-91.47 (12)
C2—C3—C4—C4a	-0.25 (17)	C9a—C9—C11—C12	93.86 (12)
C3—C4—C4a—C10	-179.81 (10)	C9—C9a—C1—C2	178.37 (10)
C3—C4—C4a—C9a	-1.24 (16)	C4a—C9a—C1—C2	-1.12 (16)
C4—C4a—C10—C10a	179.63 (10)	C9—C9a—C4a—C10	0.99 (15)
C9a—C4a—C10—C10a	1.08 (16)	C1—C9a—C4a—C10	-179.50 (10)
C10a—C5—C6—C7	-0.34 (18)	C9—C9a—C4a—C4	-177.61 (9)
C5—C6—C7—C8	0.28 (18)	C1—C9a—C4a—C4	1.90 (15)
C6—C7—C8—C8a	0.20 (18)	C4a—C10—C10a—C5	178.48 (10)
C7—C8—C8a—C9	179.79 (10)	C4a—C10—C10a—C8a	-1.78 (16)
C7—C8—C8a—C10a	-0.58 (16)	C10—C10a—C5—C6	179.69 (10)
C9a—C9—C8a—C8	-178.68 (9)	C8a—C10a—C5—C6	-0.05 (17)
C11—C9—C8a—C8	6.78 (16)	C10—C10a—C8a—C9	0.39 (16)
C9a—C9—C8a—C10a	1.70 (16)	C5—C10a—C8a—C9	-179.85 (9)
C11—C9—C8a—C10a	-172.84 (9)	C10—C10a—C8a—C8	-179.25 (10)
C8a—C9—C9a—C1	178.13 (9)	C5—C10a—C8a—C8	0.51 (15)
C11—C9—C9a—C1	-7.29 (16)	O—C11—C12—C13	179.37 (11)
C8a—C9—C9a—C4a	-2.38 (16)	C9—C11—C12—C13	-0.21 (16)
C11—C9—C9a—C4a	172.20 (9)	O—C11—C12—C14	0.10 (15)
C8a—C9—C11—O	88.94 (13)	C9—C11—C12—C14	-179.48 (10)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C3—H3 \cdots O ⁱ	0.95	2.48	3.3747 (16)	157

Symmetry code: (i) $x-1, y, z$.*Geometric parameters* (Å, °)

Calculated values using RM06-2X/6-31+G(d) for optimized isolated molecule given in square brackets

O—C11	1.2176 (13)	[1.217]
C1—C2	1.3603 (17)	[1.365]
C1—H1	0.9500	[1.086]
C2—C3	1.4201 (17)	[1.428]
C2—H2	0.9500	[1.086]
C3—C4	1.3615 (17)	[1.364]
C3—H3	0.9500	[1.086]
C4—C4a	1.4297 (16)	[1.432]
C4—H4	0.9500	[1.087]
C4a—C10	1.3948 (16)	[1.396]
C5—C6	1.3579 (18)	[1.364]
C5—H5	0.9500	[1.087]
C6—C7	1.4207 (17)	[1.428]
C6—H6	0.9500	[1.086]
C7—C8	1.3618 (16)	[1.365]
C7—H7	0.9500	[1.086]
C8—C8a	1.4301 (16)	[1.434]
C8—H8	0.9500	[1.087]
C9—C8a	1.4024 (15)	[1.404]
C9—C9a	1.4040 (16)	[1.404]
C9—C11	1.5117 (14)	[1.509]
C9a—C1	1.4300 (16)	[1.434]
C9a—C4a	1.4365 (15)	[1.437]
C10—C10a	1.3920 (17)	[1.396]
C10—H10	0.9500	[1.089]
C10a—C5	1.4307 (16)	[1.432]
C10a—C8a	1.4363 (15)	[1.436]
C11—C12	1.4864 (15)	[1.497]
C12—C13	1.3277 (17)	[1.339]
C12—C14	1.5020 (16)	[1.502]
C13—H13A	0.9500	[1.087]
C13—H13B	0.9500	[1.087]
C14—H14A	0.9500	[1.093]
C14—H14B	0.9500	[1.095]
C14—H14C	0.9500	[1.095]
C2—C1—C9a	120.88 (11)	[120.8]
C2—C1—H1	119.6	[120.0]
C9a—C1—H1	119.6	[119.1]

C1—C2—C3	120.84 (11)	[120.8]
C1—C2—H2	119.6	[119.9]
C3—C2—H2	119.6	[119.4]
C2—C3—C4	120.12 (11)	[120.2]
C2—C3—H3	119.9	[119.5]
C4—C3—H3	119.9	[120.3]
C3—C4—C4a	121.14 (11)	[120.9]
C3—C4—H4	119.4	[120.8]
C4a—C4—H4	119.4	[118.3]
C10—C4a—C4	121.93 (10)	[121.6]
C10—C4a—C9a	119.49 (10)	[119.5]
C4—C4a—C9a	118.57 (10)	[119.0]
C6—C5—C10a	121.02 (11)	[120.9]
C6—C5—H5	119.5	[120.8]
C10a—C5—H5	119.5	[118.3]
C5—C6—C7	120.35 (11)	[120.2]
C5—C6—H6	119.8	[120.3]
C7—C6—H6	119.8	[119.5]
C6—C7—C8	120.66 (11)	[120.7]
C6—C7—H7	119.7	[119.4]
C8—C7—H7	119.7	[119.9]
C7—C8—C8a	120.98 (11)	[120.9]
C7—C8—H8	119.5	[119.9]
C8a—C8—H8	119.5	[119.2]
C8—C8a—C9	122.61 (10)	[122.5]
C8—C8a—C10a	118.27 (10)	[118.3]
C9—C8a—C10a	119.11 (10)	[119.2]
C8a—C9—C9a	121.28 (10)	[121.2]
C8a—C9—C11	119.66 (10)	[119.7]
C9a—C9—C11	118.84 (10)	[119.0]
C1—C9a—C4a	118.42 (10)	[118.4]
C1—C9a—C9	122.57 (10)	[122.5]
C4a—C9a—C9	119.02 (10)	[119.1]
C4a—C10—C10a	121.54 (10)	[121.5]
C4a—C10—H10	119.2	[119.2]
C10a—C10—H10	119.2	[119.2]
C5—C10a—C10	121.77 (10)	[121.6]
C8a—C10a—C10	119.51 (10)	[119.4]
C5—C10a—C8a	118.72 (11)	[119.0]
O—C11—C9	119.34 (10)	[120.5]
O—C11—C12	120.54 (10)	[120.2]
C9—C11—C12	120.11 (9)	[119.3]
C11—C12—C13	120.05 (10)	[120.4]
C11—C12—C14	115.66 (10)	[115.4]
C13—C12—C14	124.29 (11)	[124.2]
C12—C13—H13A	120.0	[121.8]
C12—C13—H13B	120.0	[121.0]
H13A—C13—H13B	120.0	[117.2]

C12—C14—H14A	109.5	[110.9]
C12—C14—H14B	109.5	[110.6]
C12—C14—H14C	109.5	[110.6]
H14A—C14—H14B	109.5	[109.2]
H14A—C14—H14C	109.5	[109.1]
H14B—C14—H14C	109.5	[106.5]
C9a—C1—C2—C3	-0.39 (17)	[-0.1]
C1—C2—C3—C4	1.10 (17)	[-0.1]
C2—C3—C4—C4a	-0.25 (17)	[0.2]
C3—C4—C4a—C10	-179.81 (10)	[179.7]
C3—C4—C4a—C9a	-1.24 (16)	[-0.2]
C4—C4a—C10—C10a	179.63 (10)	[-179.9]
C9a—C4a—C10—C10a	1.08 (16)	[0.0]
C10a—C5—C6—C7	-0.34 (18)	[-0.2]
C5—C6—C7—C8	0.28 (18)	[0.3]
C6—C7—C8—C8a	0.20 (18)	[0.0]
C7—C8—C8a—C9	179.79 (10)	[-179.9]
C7—C8—C8a—C10a	-0.58 (16)	[-0.3]
C9a—C9—C8a—C8	-178.68 (9)	[179.0]
C11—C9—C8a—C8	6.78 (16)	[0.2]
C9a—C9—C8a—C10a	1.70 (16)	[-0.6]
C11—C9—C8a—C10a	-172.84 (9)	[-179.4]
C8a—C9—C9a—C1	178.13 (9)	[-179.8]
C11—C9—C9a—C1	-7.29 (16)	[-1.1]
C8a—C9—C9a—C4a	-2.38 (16)	[1.1]
C11—C9—C9a—C4a	172.20 (9)	[179.8]
C8a—C9—C11—O	88.94 (13)	[102.3]
C9a—C9—C11—O	-85.73 (13)	[-76.5]
C8a—C9—C11—C12	-91.47 (12)	[-78.8]
C9a—C9—C11—C12	93.86 (12)	[102.4]
C9—C9a—C1—C2	178.37 (10)	[-179.0]
C4a—C9a—C1—C2	-1.12 (16)	[0.1]
C9—C9a—C4a—C10	0.99 (15)	[-0.7]
C1—C9a—C4a—C10	-179.50 (10)	[-179.9]
C9—C9a—C4a—C4	-177.61 (9)	[179.1]
C1—C9a—C4a—C4	1.90 (15)	[0.0]
C4a—C10—C10a—C5	178.48 (10)	[-179.7]
C4a—C10—C10a—C8a	-1.78 (16)	[0.4]
C10—C10a—C5—C6	179.69 (10)	[180.0]
C8a—C10a—C5—C6	-0.05 (17)	[-0.1]
C10—C10a—C8a—C9	0.39 (16)	[-0.1]
C5—C10a—C8a—C9	-179.85 (9)	[180.0]
C10—C10a—C8a—C8	-179.25 (10)	[-179.7]
C5—C10a—C8a—C8	0.51 (15)	[0.4]
O—C11—C12—C13	179.37 (11)	[172.7]
C9—C11—C12—C13	-0.21 (16)	[-6.1]
O—C11—C12—C14	0.10 (15)	[-5.8]

C9—C11—C12—C14

-179.48 (10)

[175.3]
