



Fitness, J; Floyd, S; Warndorff, DK; Sichali, L; Mwaungulu, L; Crampin, AC; Fine, PE; Hill, AV (2004) LARGE-SCALE CANDIDATE GENE STUDY OF LEPROSY SUSCEPTIBILITY IN THE KARONGA DISTRICT OF NORTHERN MALAWI. *The American journal of tropical medicine and hygiene*, 71 (3). pp. 330-40. ISSN 0002-9637

Downloaded from: <http://researchonline.lshtm.ac.uk/14334/>

DOI:

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: Copyright the publishers

LARGE-SCALE CANDIDATE GENE STUDY OF LEPROSY SUSCEPTIBILITY IN THE KARONGA DISTRICT OF NORTHERN MALAWI

JODENE FITNESS, SIAN FLOYD, DAVID K. WARNDORFF, LIFTED SICHALI, LORREN MWAUNGULU, AMELIA C. CRAMPIN, PAUL E. M. FINE, AND ADRIAN V. S. HILL

Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, United Kingdom; Department of Tropical Hygiene, London School of Hygiene and Tropical Medicine, London, United Kingdom; Karonga Prevention Study, Chilumba, Karonga District, Malawi

Abstract. We present a large case-control candidate gene study of leprosy susceptibility. Thirty-eight polymorphic sites from 13 genes were investigated for their role in susceptibility to leprosy by comparing 270 cases with 452 controls in Karonga district, northern Malawi. Homozygotes for a silent T→C change in codon 352 of the vitamin D receptor gene appeared to be at high risk (odds ratio [OR] = 4.3, 95% confidence interval [CI] = 1.6–11.4, $P = 0.004$), while homozygotes for the McCoy b blood group defining variant K1590E in exon 29 of the complement receptor 1 (formerly CD35) gene appeared to be protected (OR = 0.3, 95% CI = 0.1–0.8, $P = 0.02$). Borderline evidence for association with leprosy susceptibility was found for seven polymorphic sites in an additional six genes. Some of these apparent associations may be false-positive results from multiple comparisons, and several associations suggested by studies in other populations were not replicated here. These data provide evidence of inter-population heterogeneity in leprosy susceptibility.

INTRODUCTION

Leprosy is a chronic disabling disease attributable to infection with *Mycobacterium leprae*. It has a broad clinical spectrum: at one pole is paucibacillary (PB) leprosy, characterized by a low bacterial count, strong cell-mediated immunity, and localized disease. At the other pole, multibacillary (MB) leprosy is characterized by a high bacterial count, poor cell-mediated immunity and strong humoral immunity, with progressive and disseminated disease. The natural history of leprosy is poorly understood, but family studies, segregation analyses, and twin studies provide evidence that, in addition to environmental and exposure factors, host genetic factors influence leprosy susceptibility.^{1–4}

Elucidation of genetic determinants of host susceptibility to leprosy could facilitate the development of better preventative and therapeutic strategies. The earliest strong evidence pointed towards the major histocompatibility complex region on chromosome 6.^{5,6} More recently susceptibility loci have been mapped using linkage analysis of multi-case families to chromosomes 10p13 and 20p12 in populations in south India and to chromosome 6q25 and, again, 10p13, in Vietnam.^{4,7,8} The linkage on chromosome 10p13 appears to be mainly with tuberculoid leprosy and the chromosome 6q25 linkage with leprosy *per se*. Association studies in various populations have indicated that several other non-HLA genes may also influence either leprosy susceptibility or the type of leprosy that develops upon infection.³ In general, these association studies have been carried out on relatively small sample sizes, and some of the reported associations may represent chance findings, particularly in studies that examined numerous variants of multiple genes. A major objective of the present study was to re-evaluate reported associations in an independent population.

The Karonga Prevention Study (KPS), a long-term epidemiologic study of mycobacterial disease in the Karonga District of northern Malawi, provides an excellent opportunity to attempt to replicate previously reported leprosy associations as well as to investigate *de novo* hypotheses. A case-control association study was thus designed to investigate a large number of putative susceptibility genes (38 polymorphisms

from 13 genes) in this population. The genotyping was performed blind to the disease status of the samples.

Polymorphisms in tumor necrosis factor (TNF, formerly TNF α), the vitamin D receptor (VDR), toll-like receptor 2 (TLR2), and lymphotoxin alpha (LTA, formerly TNF β) are associated with leprosy in other populations and were included here in an attempt to replicate these findings.³ An interleukin-10 (IL10) promoter that has recently been suggested to be associated with leprosy susceptibility in Brazil, and other polymorphisms that may influence expression of the immunomodulatory genes IL10 and TLR2 were included (Segal S and others, unpublished data).⁹ Polymorphisms in the genes encoding solute carrier family 11, member 1 (SLC11A1, formerly NRAMP1), mannose-binding lectin (MBL2, formerly MBP or MBL) and interferon- γ (IFNG) were investigated because they have been reported to be associated, in at least one population, with susceptibility to a related mycobacterial disease, tuberculosis.¹⁰ The toll-like receptor 4 (TLR4), IL10, and intercellular adhesion molecule 1 (ICAM-1, formerly CD54) loci contain polymorphisms that are associated with susceptibility to other infectious diseases, and it was speculated that these genes might represent general infectious disease susceptibility loci.^{11–13} The caspase recruitment domain family, member 15 (CARD15, formerly NOD2) and CR1 were selected as functional candidates because of their roles in innate immunity and potential role in mycobacterial disease. The chemokine (C-C motif) ligand 3 (CCL3, formerly MIP1 α) was selected as a functional candidate due to its involvement in T helper cell 1 (TH1) differentiation.

METHODS

The study protocol was approved by the National Health Sciences Research Committee of Malawi and by the Ethics Committee of the London School of Hygiene and Tropical Medicine.

Recruitment and sample collection. Cases were ascertained within the context of the KPS, a long-term epidemiologic study of mycobacterial disease in the Karonga District, a rural area of northern Malawi. The population and basic

field methods of the KPS have been described in detail elsewhere.^{14,15} Leprosy cases were identified in the context of total population surveys during the 1980s, and through “enhanced” passive detection (examination of all individuals who appeared at health centers) in the 1990s, in addition to self-reporting. Individuals were examined first by paramedical leprosy control assistants (LCAs), and suspects referred to project Medical Officers for review examination and (4-mm punch) biopsy. Diagnostic certainty was defined in terms of clinical, skin slit smear, and histopathologic data.¹⁶ Cases were defined as PB or MB based upon an algorithm that took into account all clinical, bacteriologic (slit skin smear), biopsy, and historical record information. The algorithm implies a conservative definition of MB (e.g., biopsy or slit skin smear bacterial index > 1).

Recruitment for this study began in 1996. Since leprosy incidence had decreased dramatically in the district, cases were selected retrospectively from project data files. Only cases that met strict diagnostic criteria were included (N and M categories as stipulated elsewhere¹⁶). Individuals were eligible for selection as controls if they were resident in the district during the 1986–1989 population survey, and had no history of either tuberculosis or leprosy. Controls were matched to cases by age (15–19, 20–24, 25–29, 30–34, 35–44, 45–54, 55–64, and > 65 years old), sex, and geographic area (residence within 1 km of the case). First- or second-degree relatives of cases were not eligible for selection as controls. Each case was matched to up to three controls.

Cases and controls were visited in their homes by field teams consisting of an LCA and an interviewer. After having

the study explained to them, those who consented were examined and invited to provide a blood or saliva sample for genetic testing (those who were willing to provide blood were told their blood group, and offered syphilis tests with treatment and also human immunodeficiency virus [HIV] counseling and testing if they so desired). All cases ascertained since 1987 have been offered voluntary counseling and HIV testing at the time of initial registration as part of a continuous case control study of leprosy, tuberculosis, and HIV.¹⁷ Blood (7.5 mL) was collected into EDTA and placed in vaccine carriers with wet ice until it reached the project laboratory within three days of collection, where cells were separated and stored at –70°C. The DNA was separated using Nucleon kits (Scotlab, Coatbridge, United Kingdom) following the manufacturer’s instructions, and shipped at 4°C to the Wellcome Trust Center for Human Genetics in Oxford, United Kingdom for analysis.

Genotyping. Genotyping was performed blind to disease status. A variety of genotyping methods were used to investigate 38 polymorphisms from 13 candidate genes (Tables 1 and 2). Four CARD15 polymorphisms were genotyped using polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP), the IFN- γ +874 single nucleotide polymorphism (SNP) was typed using amplification refractory mutation system PCR, and the CARD15 and IFNG products were analyzed using agarose gel electrophoresis.^{18,19} Seven polymorphisms in SLC11A1, LTA, CCL3, and TLR2 were genotyped by fluorescence PCR followed by electrophoresis and detection using an ABI fluorescent DNA sequencer (Applied Biosystems, Foster City, CA) as de-

TABLE 1
Oligonucleotides and conditions used for non-LDR genotyping*

Gene	Location, polymorphism Amino acid change (Restriction enzyme)	Genotyping method (% agarose gel)	PCR primers	PCR temperature product length
CARD15 (NOD2)	Exon 4, 802C→T	PCR-RFLP (3%)	F: CAGTCTCGCTTCCTCAGTACC	55°C
	P268S (<i>Bam</i> HI)		R: AGTGTCCGCATCGTCATTG	187 bp
	Exon 4, 2104C→T	PCR-RFLP (3%)	F: TTCCTGGCAGGGCTGTGTGCTcGG	55°C
	R702W (<i>Msp</i> I)		R: GGATGGAGTGGAAAGTGTCTTG	139 bp
	Exon 8, 2722G→C	PCR-RFLP (3%)	F: AGGCCACTCTGGGATTGAG	55°C
	G908R (<i>Hha</i> I)		R: GTGATCACCCAAGGCTTCAG	196 bp
Exon 11, 3020insC	PCR-RFLP (4%)	F: GGCAGAAGCCCTCCTGCAGGgCC	55°C	
1007fs (<i>Apa</i> I)		R: CCTCAAAATCTTGCCATTCC	152 bp	
CCL3 (MIP1- α)	Promoter, microsatellite (TA) _n at -906	Fluorescence PCR	F: FAM-AAGGCATGTATTTCGAAGC R: CTGACCCAGCATCGTTTA	58°C 327–337 bp
IFNG	Intron 1 874T→A	ARMS PCR (2%)	IFNGT: TTCTTACAACACAAAATCAAATCT IFNGA: TTCTTACAACACAAAATCAAATCA IFNGCOM: TCAACAAAGCTGATACTCCA HGHR: GCCTTCCCAACCATTCCTTA HGHR: TCACGGATTCTGTGTGTGTTTC	62°C, 50°C 461 bp 426 bp
LTA (TNF- β)	5'UTR, microsatellite (AC/GT) _n at -3.5 kb	Fluorescence PCR	F: GCCCTAGATTTCATCCAGCCACAG R: HEX-CCTCTCTCCCCTGCAACACACA	58°C, 50°C 93–119 bp
SLC11A1 (NRAMP1)	Promoter, microsatellite (GT) _n	Fluorescence PCR	F: FAM-TTCTGTGCCCTCCCAAGTTAGC R: ACTCGCATTAGGCCAACGAG	52°C 180–220 bp
	Exon 2, 9 bp del	Fluorescence PCR	F: GTGACAAAGGGTCCCAAC R: TET-CGGTTTTGTGTCTGGGAT	60°C 143, 132 bp
	3' UTR, TGTG ins/del	Fluorescence PCR	F: GCATCTCCCAATTCATGGT R: TET-AACTGTCCCCTCTATCCTG	60°C 240, 244 bp
	3' UTR CAAA ins/del	Fluorescence PCR	F: TET-CCTAGCGCAGCCATGTGATTACCC R: AGCCTGTGTCCC GCCCAAGTCCT	58°C 234, 238 bp
	Intron 2, microsatellite	Fluorescence PCR	F: TATCCCCATTTCATTCGTTCCATT R: FAM-ACCCCAAGACCCACCC	58°C 213–243 bp

* LDR = ligation detection reaction; PCR = polymerase chain reaction; CARD15 = caspase recruitment family, member 15; RFLP = restriction fragment length polymorphism; bp = basepairs; CCL3 = chemokine (C-C motif) ligand 3; IFNG = interferon- γ ; ARMS = amplification refractory mutation system; LTA = lymphotoxin α ; UTR = untranslated region; TNF- β = tumor necrosis factor β ; SLC11A1 = solute carrier family 11, member 1; del = deletion; ins = insertion; TLR2 = toll-like receptor 2; fs = frameshift.

TABLE 2
Oligonucleotides and conditions required for ligation detection reaction (LDR) genotyping*

Gene	SNP location Base change Amino acid change (SNP name) [†]	PCR primer sequences	PCR temperature [MgCl ₂] and product length	LDR probe sequences [‡]	LDR product length [§] and reaction temperature
CRI (CD35)	Exon 19	F: TGGGTTGGATCTTTCCCATG R: CCTGGTTCCAGCAAGGATACA	63°C 2 mM	P1: FAM-aaattaaaaatTGGTGCATGTGATGCACAGACATCCAG P2: HEX-taaattaaaaatTGGTGCATGTGATGCACAGACATCCAT P3: GTTGGATCAGAACTCAACTATTTCTGTACTACAGaaaataaaaa	81 bp 82 bp 70°C
	Exon 28	F: TAAAAAATAAGCTGTTTTACCACTACTC R: CCTCACACCAGCAAAAGTC	61°C 1 mM 476 bp	P1: FAM-atacagccctccccctCGGTGTATTTCTACTAATA P2: HEX-ataaagccccctccccctCGGTGTATTTCTACTAATG P3: AATGCACAGCTCCAGAAAGTTGAAAATGCAAtat	71 bp 70 bp 68°C
	Exon 29	F: TAAAAAATAAGCTGTTTTACCACTACTC R: CCTCACACCAGCAAAAGTC	61°C 1 mM	P1: HEX-GAAACAGAGTTCCTTTTCCCTCACTGAGATA P2: FAM-AAACAGAGTTCCTTTTCCCTCACTGAGATCG P3: TCAGATTTAGATGTCAGCCCGGGTTTGTGTC	68°C
	Exon 29	F: TAAAAAATAAGCTGTTTTACCACTACTC R: CCTCACACCAGCAAAAGTC	61°C 1 mM	P1: HEX-GAGGAAAAGAACTCCTGTTCTGTTACTCC P2: FAM-GTAGGAAAAGAACTCCTGTTTCTGTTACTCT P3: AATGCAATTTTCAACTTCTGGAGCTGTGCATT	65 bp 67 bp 68°C
	Exon 33	F: AAGCGCACAGTCAACAGGTCAC R: GGAGGTAGTTCTGCTCTGAC	63°C 2 mM	P1: FAM-taaaaataataataaCATCCCTGCACAAGTGACCC P2: HEX-ataaaaaataataataaCATCCCTGCACAAGTGACCC P3: TCATGGGAATGGGTTGGAGTtaaaaaaataaaaa	72 bp 73 bp 64°C
	P1827R	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	53°C 1 mM	P1: FAM-CAGCACCTCTGTGACCCAGCCCAT P2: TET-aAGCACCTCTGTGACCCAGCCCAA P3: GTTGTGGCAATAGAGACCCCGTTGCCattt	55 bp 55 bp 64-68°C
	Exon 2	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	1 mM 130 bp		
	Exon 2	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	1 mM 130 bp		
	Exon 2	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	1 mM 130 bp		
	Exon 2	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	1 mM 130 bp		
ICAM1	Exon 2	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	1 mM 130 bp		
	Exon 2	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	1 mM 130 bp		
	Exon 2	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	1 mM 130 bp		
	Exon 2	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	1 mM 130 bp		
	Exon 2	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	1 mM 130 bp		
	Exon 2	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	1 mM 130 bp		
	Exon 2	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	1 mM 130 bp		
	Exon 2	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	1 mM 130 bp		
	Exon 2	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	1 mM 130 bp		
	Exon 2	F: TGTCCCCCTCAAAAAGTCATCCTG R: TTTCCAGGAGGAGCAACTCCTT	1 mM 130 bp		
IL10	Promoter	F: CTGGCTCCCTTACCTTCTACACA R: TGGGCTAAAATATCCTCAAAGTTCC	60°C 1.5 mM	P1: FAM-CAAAGACAACACTACTAAGGCTTCTTTGGAA P2: TET-CCAAGACAACACTACTAAGGCTTCTTTGGAG P3: GGGAAAGTAGGATAGGTAAGGAAAGTAAGGAC	68 bp 68 bp 64-68°C
	Promoter	F: CTGGCTCCCTTACCTTCTACACA R: TGGGCTAAAATATCCTCAAAGTTCC	60°C 1.5 mM	P1: FAM-GCAGTGGTACCCCTTGTACAGGTGATTAAT P2: TET-GCAGTGGTACCCCTTGTACAGGTGATTAAC P3: ATCTCTGTGCTCAGTTTGTCTCACTATAAATAAGAGACG	71 bp 71 bp 64-68°C
	Promoter	F: CTGGCTCCCTTACCTTCTACACA R: TGGGCTAAAATATCCTCAAAGTTCC	60°C 1.5 mM	P1: FAM-AAACACATCCTGTGACCCCGCTGTGTA P2: TET-ttCACATCCTGTGACCCCGCTGTGTC P3: CTGTAGGAAGCCAGTCTCTGAAAGTAAAATGGAA	60 bp 60 bp 64-68°C
	Promoter	F: CTGGCTCCCTTACCTTCTACACA R: TGGGCTAAAATATCCTCAAAGTTCC	60°C 1.5 mM	P1: FAM-ttCGGCTTCCAGGCAAAAGATGGGC P2: TET-AAACGCTTCCAGGCAAAAGATGGGT P3: GTGATGGCACCAGGGGAGAAAAGGGGG	52 bp 52 bp 73°C
	Promoter	F: CTGGCTCCCTTACCTTCTACACA R: TGGGCTAAAATATCCTCAAAGTTCC	60°C 1.5 mM	P1: FAM-aaaaaaaCCAGCAAAAGATGGGCTGATGG P2: TET-aaaaaaaCCAGGCAAAAGATGGGCTGATGG P3: CACCAAGGAGAAAAGGGGAAACCAGGaat	62 bp 62 bp 64°C
	Promoter	F: CTGGCTCCCTTACCTTCTACACA R: TGGGCTAAAATATCCTCAAAGTTCC	60°C 1.5 mM	P1: FAM-ttatttaaATGGGCTGTGATGCCACCAAGG P2: TET-ttatttaaATGGGCTGTGATGCCACCAAGG P3: AGAAAAGGGGAAACCAGGCAAAAGGaaat	58 bp 55 bp 64°C
	Promoter	F: CTGGCTCCCTTACCTTCTACACA R: TGGGCTAAAATATCCTCAAAGTTCC	60°C 1.5 mM	P1: FAM-AAATTCGATTTAGCATTTAGACTACTACCTCGATGA P2: HEX-TTCCGATTTAGCATTTAGACTACTACCTCGATGG P3: TATTATTGACTTTAATTTGTTGACAAATGTTCTTCATTTTCC	83 bp 81 bp 54°C
	Promoter	F: CTGGCTCCCTTACCTTCTACACA R: TGGGCTAAAATATCCTCAAAGTTCC	60°C 1.5 mM	P1: FAM-attcaAAAAGGAAAGCTGAGAAAGTGAAGAAAAGT P2: TET-attcaAAAAGGAAAGCTGAGAAAGTGAAGAAAAGC P3: CAGGCTCTGGAGGGGGGaaaaataataata	71 bp 71 bp 64°C
	Promoter	F: CTGGCTCCCTTACCTTCTACACA R: TGGGCTAAAATATCCTCAAAGTTCC	60°C 1.5 mM		
	Promoter	F: CTGGCTCCCTTACCTTCTACACA R: TGGGCTAAAATATCCTCAAAGTTCC	60°C 1.5 mM		
MBL2 (MBP, MBL)	Exon 1	F: GCACCCAGATTTAGGACAGAG R: CAGGCAGTTTCTCTGGAAAGG	58°C 2 mM	P1: FAM-ttCGGCTTCCAGGCAAAAGATGGGC P2: TET-AAACGCTTCCAGGCAAAAGATGGGT P3: GTGATGGCACCAGGGGAGAAAAGGGGG	52 bp 52 bp 73°C
	R52C (D variant)	F: GCACCCAGATTTAGGACAGAG R: CAGGCAGTTTCTCTGGAAAGG	339 bp 58°C		62 bp 62 bp
	Exon 1	F: GCACCCAGATTTAGGACAGAG R: CAGGCAGTTTCTCTGGAAAGG	339 bp 58°C		64°C
	G54D (B variant)	F: GCACCCAGATTTAGGACAGAG R: CAGGCAGTTTCTCTGGAAAGG	339 bp 58°C		58 bp
	Exon 1	F: GCACCCAGATTTAGGACAGAG R: CAGGCAGTTTCTCTGGAAAGG	2 mM 339 bp		55 bp 64°C
	G57E (C variant)	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	55°C 2.5 mM		83 bp 81 bp
	Exon 4	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	55°C 2.5 mM		54°C
	896A→G	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	440 bp 58°C		71 bp
	D299G	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	58°C 1 mM		71 bp
	Promoter	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	1 mM 1,240 bp		64°C
TLR4	Exon 4	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	55°C 2.5 mM		83 bp 81 bp
	896A→G	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	55°C 2.5 mM		54°C
	D299G	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	440 bp 58°C		71 bp
	Promoter	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	58°C 1 mM		71 bp
	-1031T→C	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	1 mM 1,240 bp		64°C
	Exon 4	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	55°C 2.5 mM		83 bp 81 bp
	896A→G	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	55°C 2.5 mM		54°C
	D299G	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	440 bp 58°C		71 bp
	Promoter	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	58°C 1 mM		71 bp
	-1031T→C	F: CTGGCTGGTTTAGAAGTCC R: CCATCCGAAATTAAGAAAAGT	1 mM 1,240 bp		64°C

Promoter -863C→A	F: GTGGGGAGAACAAAAGGATAAG R: GATGTGGCGTCTGAGGGTTGTTTT	58°C 1 mM 1,240 bp	P1: FAM-taaaaataGAAAAGTCGAGTAATGGGACCCCCC P2: TET-taaaaataGAAAAGTCGAGTAATGGGACCCCCC P3: CTTAAACGAGACAGGGCCCAATGAGAGGGMaataaaat	69 bp 69 bp 64°C
Promoter -857C→A	F: GTGGGGAGAACAAAAGGATAAG R: GATGTGGCGTCTGAGGGTTGTTTT	58°C 1 mM 1,240 bp	P1: FAM-aataaataAGTCGAGTAATGGGACCCCMCTTAAAC P2: TET-aataaataAGTCGAGTAATGGGACCCCMCTTAAAC P3: GAAGACAGGCCCAATGAGAGGGCCCaataat	65 bp 65 bp 64°C
Promoter -376G→A	F: GTGGGGAGAACAAAAGGATAAG R: GATGTGGCGTCTGAGGGTTGTTTT	58°C 1 mM 1,240 bp	P1: FAM-gttctatctttttctctgcatcctctgtctggaaa P2: TET-gttctatctttttctctgcatcctctgtctggaaa P3: TTAGAAGGAAAACAGCCACAGACCTGGTCTCC	63 bp 63 bp 64°C
Promoter -308G→A (TNF*1→TNF*2)	F: GTGGGGAGAACAAAAGGATAAG R: GATGTGGCGTCTGAGGGTTGTTTT	58°C 1 mM 1,240 bp	P1: FAM-aaAGGCAATAGGTTTTGAGGGGCATGA P2: TET-agAGGCAATAGGTTTTGAGGGGCATGA P3: GGACGGGTTTCAGCTCCAGGGGaaaa	54 bp 53 bp 64°C
Promoter -238G→A	F: GTGGGGAGAACAAAAGGATAAG R: GATGTGGCGTCTGAGGGTTGTTTT	58°C 1 mM 1,240 bp	P1: FAM-aCCAGAAAGACCCCTCGGAATCG P2: TET-aCCAGAAAGACCCCTCGGAATCG P3: GAGCAGGAGGATGGGGAGTGTGAGG	51 bp 51 bp 64°C
Exon 1 C insertion at 70	F: GTGGGGAGAACAAAAGGATAAG R: GATGTGGCGTCTGAGGGTTGTTTT	58°C 1 mM 1,240 bp	P1: FAM-taatatGACCAGCTAAAGGGGAGAGAAAGCAACTACAGAC P2: TET-taatatCCAGCTAAAGGGGAGAGAAAGCAACTACAGACC P3: CCCCCTGAAAACAAACCCCTCAGACCaataaaaaa	75 bp 75 bp 64°C
Intron 8 G→T	F: CTGGGGAGCGGGAGTATGAAGGA R: GGGTGGCGCAGCGGATGTA	58°C 1 mM	P1: FAM-gggtggTGGGATTTGAGCAGTGAAGT P2: HEX-aggtggTGGGATTTGAGCAGTGAAGT P3: GCCCAGCTGAGAGCTCTGTGCCCTTaaaaaaaTaaaaaaa	67 bp 67 bp 72°C
Intron 8 C→T	F: CTGGGGAGCGGGAGTATGAAGGA R: GGGTGGCGCAGCGGATGTA	1,100 bp 58°C 1 mM	P1: FAM-CAAGAGCAGAGCTGAGTATTTGGGAAATGT P2: HEX-aaAGAGCAGAGCTGAGTATTTGGGAAATGC P3: GCAGCCCTCTCTGTGGCCCAAGattaaaaaaaata	65 bp 65 bp 72°C
Exon 9 T→C	F: CTGGGGAGCGGGAGTATGAAGGA R: GGGTGGCGCAGCGGATGTA	58°C 1 mM	P1: FAM-TGCAGGACGGCCGGCTGATT P2: HEX-agCAGGACGGCCGGCTGATC P3: GAGGCCATCCAGGACCGCCCTCattataataataataaaaaa	70 bp 70 bp 72°C
352 silent (Taq I; T→t)		1,100 bp		

* CR = complement receptor; ICAMI = intercellular adhesion molecule 1; IL10 = interleukin-10; MBL = mannose-binding lectin; VDR = vitamin D receptor; SNP = single nucleotide polymorphism. For definition of other abbreviations, see Table 1.

† SNP name used by previous investigators.

‡ Non-specific polymucleotides, added to LDR probes to obtain desired ligation product lengths, are shown in lower case. P1 = allelic probe 1, P2 = allelic probe 2, P3 = common probe.

§ Apparent LDR product lengths obtained using an ABI 3700 with Rox 500 size standards may be less than the absolute length of the LDR product. This discrepancy is not observed using an ABI 373 with GS-550 size standards.

scribed later in this report (Segal S and others, unpublished data).^{20,21} The PCR primers, conditions, and restriction enzymes used are shown in Table 1.

Direct sequence analysis. Direct sequence analysis was used to determine whether the TLR2 R677W, R753Q, and CARD15 R790Q variants were present in the Malawi population^{13,22} (National Center for Biotechnology Information SNP database, Bethesda, MD). The PCR primers and thermocycles were TLR2 exon 3 586-basepair (bp) segment: 5'-gcgtggccagcaggttcag-3', 5'-gggacctccaggtaggtcttg-3' (thermocycles: one cycle at 94°C for four minutes; 36 cycles at 94°C for 30 seconds, 58°C for 40 seconds, and 72°C for 60 seconds; and one cycle at 72°C for two minutes) and CARD15 exon 4 509-bp segment: 5'-gccgagccgcacaacctt-3', 5'-gtgctccccatacctgaaca-3' (thermocycles: one cycle at 94°C for five minutes; 35 cycles at 60°C for 30 seconds, and 72°C for 45 seconds; and one cycle at 72°C for four minutes). The PCR (2 × 25 µL) and sequencing (20 µL) reaction were performed using standard conditions. Sequence analysis was performed using an ABI 3700 fluorescent DNA sequencer. Each segment was sequenced in both directions.

Genotyping using the post-PCR multiplexed ligation detection reaction (LDR). The LDR was used to genotype the remaining 23 SNPs located in seven genes.²³ Briefly, fragments containing the SNPs of interest were PCR-amplified using appropriate primers and conditions (Table 2). Amplifications were performed in a total volume of 15 µL and contained 10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1–2.5 mM MgCl₂, 200 µM of each dNTP, 0.2 µM of each primer, approximately 50 ng of DNA, and 0.5 units of Ampliqaq Gold polymerase (Perkin-Elmer, Boston, MA). To prevent polymerase extension during the subsequent LDR procedure, residual polymerase activity was removed by incubation with 1:10 volume of 1 mg/mL of proteinase K in 50 mM EDTA at 37°C for 30 minutes and 55°C for 10 minutes. The proteinase K was inactivated by incubation at 99°C for 10 minutes.

A set of three probes consisting of two fluorescently labeled allelic and one common probe was designed for each of the SNPs investigated using the LDR (Table 2). The common probes (100 pmol of each) were 5' phosphorylated in a volume of 100 µL using 10 units of T4 polynucleotide kinase and 1 mM ATP in 70 mM Tris-HCl buffer, pH 7.6, containing 10 mM MgCl₂ and 5 mM dithiothreitol. After incubation at 37°C for 45 minutes, the T4 kinase was inactivated by the addition of 20 µg of proteinase K in 100 µL of TE buffer (10 mM Tris-HCl, 5 mM EDTA, pH 8.0) and incubating at 37°C for 30 minutes. The proteinase K was inactivated by incubation at 99°C for 10 minutes.

The multiplexed LDR was performed in 15-µL volumes of 20 mM Tris-HCl buffer, pH 7.6, containing 25 mM KCl, 10 mM MgCl₂, 1 mM NAD⁺, 10 mM dithiothreitol, 0.1% Triton X-100, 10 nM (200 fmol) of each LDR probe, 2 µL of PCR product, and 2 units of *Taq* DNA ligase. The reactions were prepared on ice and then initiated by placing on PCR blocks pre-heated to 95°C. An initial one-minute denaturation at 95°C was followed by 15 thermal cycles of denaturation at 95°C for 15 seconds, and annealing and ligation at the appropriate temperature (Table 2) for four minutes. The activity of the ligase was stopped after 15 cycles by cooling to 4°C and adding 2 µL of 100 mM EDTA.

Electrophoresis and detection of multiplexed and/or pooled LDR or fluorescent PCR products was performed on

an ABI 373 or ABI 3700 fluorescent DNA sequencer. For the ABI 3700, 1 µL of LDR product was added to 10 µL of Hi-Di formamide and 0.01 µL of 8 nM ABI Genescan-500 Rox size standards (Applied Biosystems). For the ABI373, 2 µL of LDR product was added to 1.5 µL of 5:1 deionized formamide, 25 mM EDTA with blue dextran (1 mg/mL), and 0.5 µL of fluorescently labeled size standards (GeneScan-350 tetramethylrhodamine [TAMRA]). After denaturation, the product was subjected to electrophoresis on 6% polyacrylamide gels. Product sizes were calculated relative to the standard with Genescan 3.5 (Perkin Elmer Applied Biosystems, Warrington, United Kingdom) using the second order least squares method. Automated allele calling of LDR products was performed using Genotyper 2.5 (Perkin Elmer Applied Biosystems).

Statistical analysis. For each polymorphism, the percentage of individuals with a particular genotype was calculated separately for cases and controls. Conditional logistic regression was used to quantify the association between disease and genotype, with and without controlling for ethnic group, and with and without restriction to PB cases. The percentage of cases with a particular genotype was also calculated separately for PB and MB cases.

RESULTS

Recruitment. A total of 334 cases and 702 controls were selected from the database and sought in the field. Of the 334 cases, 22 had left the district and 34 had died by the time they were sought for genetics testing, 7 were alive but neither a blood nor a buccal swab specimen were provided, and for one case data were not available for a matched control. This gave a total of 270 cases that could be included in analyses, 26 of which were MB. Of the 702 controls that were sought, 74 had left the district and 90 had died by the time they were sought for genetics testing, 26 were alive but neither a blood nor a buccal swab specimen were provided, and for 60 data on the matched case were not available. Thus, 452 controls were included in these analyses. There were one or two controls for each case. Blood specimens were available for all except eight individuals, who provided buccal swabs.

Direct sequence analysis. Sequence analysis of 83 individuals (including cases and controls) indicated that the TLR2 R677W and R753Q variants were not present, or were very rare, in northern Malawi. Likewise, the CARD15 SNP R790Q was not found upon sequencing of 63 individuals. Accordingly, further genotyping of these SNPs was not performed.

Genotyping results. The PCR-RFLP analysis of more than 150 Malawians indicated that the CARD15 variants P268S, R702W, G908R, and 1007fs were not present in this population. The TNF-1031 variant was also not found. The MBL2 R52C, MBL2 G54D, TNF + 70, and -857 variants were very rare in Karonga District, occurring with an allele frequency of < 1.0%. No homozygotes of these variants were identified.

Association study results for leprosy susceptibility. The genotype frequencies and odds ratios (ORs) of the 26 variants that were sufficiently polymorphic for leprosy association analysis are shown in Table 3, with ORs controlled for ethnic group. The results shown combine PB and MB cases, but all results were unchanged when restricted to PB cases. For most polymorphisms, there was no evidence of an association with

TABLE 3

Genotype frequencies for all leprosy (paucibacillary and multibacillary)* cases and controls, and odds ratios (ORS) controlled for age, sex, area of residence, and ethnic group†

Gene	Polymorphism	Genotype	Case		Control		OR (95% confidence interval)
			%	n	%	n	
CCL3 (MIP1- α)	Promoter microsatellite (TA) _n at -906	331/331	41	54	46	131	
		331/327	31	40	25	71	1.4 (0.7-2.7), <i>P</i> = 0.37
		331/333	16	21	18	52	0.9 (0.4-2.0), <i>P</i> = 0.79
		Other	12	16	11	31	1.0 (0.4-2.9), <i>P</i> = 0.94
					131	285	(110, 146)‡
CR1	Exon 19 3093G→T Q1022H	g/g	79	131	81	151	
		g/t	20	33	17	31	1.1 (0.5-2.1), <i>P</i> = 0.86
		t/t	1	2	2	4	0.4 (0.06-2.6), <i>P</i> = 0.33
				166		186	(136, 136)
	Exon 28 4795A→G K1590E (<i>McC^a</i> → <i>McC^b</i>)	a/a	65	164	60	238	
		a/g	32	80	34	135	0.8 (0.5-1.1), <i>P</i> = 0.18
		g/g	3	7	6	26	0.3 (0.1-0.8), <i>P</i> = 0.02
				251		399	(245, 374)
	Exon 29 4828G→A R1601G (<i>SI1</i> → <i>SI2</i>)	g/g	45	114	49	196	
		g/a	45	114	40	157	1.3 (0.9-1.9), <i>P</i> = 0.12
		a/a	10	24	11	42	1.0 (0.5-1.9), <i>P</i> = 0.99
				252		395	(246, 371)
	Exon 29 4870A→G I1615V	g/g	76	192	79	311	
		g/a	23	57	20.5	81	1.3 (0.9-2.0), <i>P</i> = 0.21
a/a		1	3	0.5	2	1.7 (0.3-11.1), <i>P</i> = 0.58	
			252		394	(246, 370)	
Exon 33 5507C→G P1827R	c/c	71	128	79	172		
	c/g	28	51	18	40	1.4 (0.8-2.5), <i>P</i> = 0.21	
	g/g	1	2	3	7	0.2 (0.01-1.7), <i>P</i> = 0.12	
			181		219	(153, 168)	
ICAM1	Exon 2 179A→T K29M (<i>ICAM-1^{ref}</i> → <i>ICAM-1^{Kilifi}</i>)	a/a	54	112	55	134	
		a/t	32	68	31	76	1.1 (0.6-1.7), <i>P</i> = 0.83
		t/t	14	29	14	35	1.0 (0.5-2.0), <i>P</i> = 0.93
				209		245	(189, 208)
IFNG	Intron 1 874T→A	a/a	67	157	66	265	
		a/t	28	67	27	109	1.1 (0.7-1.6), <i>P</i> = 0.72
		t/t	5	12	7	28	0.8 (0.4-1.8), <i>P</i> = 0.59
				236		402	
IL10	Promoter -592C→A	c/c	46	97	43	157	
		c/a	38	81	47	169	0.7 (0.5-1.1), <i>P</i> = 0.13
		a/a	16	34	10	36	1.7 (1.0-3.3), <i>P</i> = 0.06
				212		362	(203, 296)
	Promoter -819C→T	c/c	44	95	42	147	
		c/t	41	88	48	170	0.8 (0.5-1.2), <i>P</i> = 0.23
		t/t	15	32	10	36	1.6 (0.9-3.0), <i>P</i> = 0.11
			215		353	(204, 293)	
	Promoter -1082A→G	a/a	45	86	43	149	
		a/g	44	84	44	154	1.0 (0.6-1.5), <i>P</i> = 0.96
g/g		11	21	13	46	0.7 (0.4-1.4), <i>P</i> = 0.30	
			191		349	(183, 262)	
LTA (TNF- β)	5'UTR microsatellite (AC/GT) _n at -3.5 kb§	101 absent	89	164	89	296	
		101 1 copy	10	18	11	36	0.9 (0.5-1.7), <i>P</i> = 0.80
		101 2 copies	1	2	0.3	1	
				184		333	
		103 absent	52	96	59	196	
		103 1 copy	45	83	36	121	1.2 (0.8-1.8), <i>P</i> = 0.42
		103 2 copies	3	5	5	16	
				184		333	
		105 absent	49	90	58	192	
		105 1 copy	42	78	37	124	1.6 (1.1-2.4), <i>P</i> = 0.03
		105 2 copies	9	16	5	17	
				184		333	
		111 absent	95	174	90	301	
		111 1 copy	5	10	10	32	0.4 (0.2-1.1), <i>P</i> = 0.07¶
111 2 copies	0	0	0	0			
		184		333	(163, 228)		
MBL2 (MBL)	Exon 1 239G→A G57E (C variant)	g/g	63	146	65	169	
		g/a	32	73	30	78	1.1 (0.7-1.7), <i>P</i> = 0.63
		a/a	5	12	5	14	0.9 (0.4-2.1), <i>P</i> = 0.73
				231		261	(224, 238)

TABLE 3
Continued

Gene	Polymorphism	Genotype	Case		Control		OR (95% confidence interval)
			%	n	%	n	
SLC11A1 (NRAMP1)	Promoter microsatellite (GT) _n (199 = allele 3, 201 = allele 2)	199/199	63	158	56	237	
		199/201	30	75	36	153	0.7 (0.5–1.0), <i>P</i> = 0.06
		201/201	4	11	6	25	0.6 (0.3–1.4), <i>P</i> = 0.24
		Other	2	5	2	8	1.4 (0.4–4.3), <i>P</i> = 0.60 (247, 388)
	Exon 2 9 bp del	Ins/ins	93	228	91	258	
		del present	7	16	9	25	0.7 (0.3–1.4), <i>P</i> = 0.31 (236, 261)
	3'UTR TGTG ins/del	Ins/ins	47	122	50	203	
		Ins/del	43	112	41	164	1.1 (0.8–1.5), <i>P</i> = 0.64
		del/del	9	24	9	35	1.2 (0.7–2.1), <i>P</i> = 0.58 (248, 391)
	3'UTR CAAA ins/del	del/del	41	106	45	192	
		Ins/del	45	117	44	191	1.1 (0.8–1.6), <i>P</i> = 0.48
		Ins/ins	14	35	11	46	1.5 (0.9–2.6), <i>P</i> = 0.14 (257, 410)
TLR2	Intron 2 microsatellite#	216 absent	90	188	88	332	
		216 1 copy	10	22	12	45	0.7 (0.4–1.3), <i>P</i> = 0.25
		216 2 copies	0	0	0.5	2	
				210		379	
		222 absent	78	164	79	299	
		222 1 copy	21	44	19	74	0.9 (0.6–1.5), <i>P</i> = 0.73
		222 2 copies	1	2	2	6	
				210		379	
		224 absent	37	78	40	151	
		224 1 copy	52	108	50	188	1.0 (0.7–1.4), <i>P</i> = 0.91
		224 2 copies	11	24	10	40	
				210		379	
TLR4	Exon 4 896A→G D299G	a/a	95	223	91	262	
		g/a	5	12	9	25	0.5 (0.2–1.1), <i>P</i> = 0.08
		g/g	0	0	0.4	1	
				235		288	(243, 383)
TNF	Promoter –238G→A	g/g	85	206	84	237	
		g/a	15	36	15	43	0.9 (0.5–1.5), <i>P</i> = 0.71
		a/a	0.4	1	1	3	1.4 (0.1–16.6), <i>P</i> = 0.80 (235, 262)
				243		283	
	Promoter –308G→A (TNF*1→TNF*2)	g/g	80	173	78	201	
		g/a	19	42	20	51	1.5 (0.9–2.5), <i>P</i> = 0.15
		a/a	1	1	2	6	0.2 (0.02–1.8), <i>P</i> = 0.15 (195, 223)
	Promoter –376G→A	g/g	88	211	86	242	
		g/a	11	27	13	38	0.8 (0.4–1.4), <i>P</i> = 0.37
		a/a	0.4	1	1	2	1.3 (0.1–15.8), <i>P</i> = 0.83 (231, 257)
	Promoter –863C→A	c/c	75	177	75	212	
		a/c	22	51	24	68	0.9 (0.6–1.5), <i>P</i> = 0.77
a/a		3	8	1	3	4.0 (0.8–19.7), <i>P</i> = 0.09 (228, 254)	
VDR	Intron 8 G→T (Apa I; a→A)	t/t	50	85	45	147	
		t/g	43	73	48	158	0.7 (0.5–1.2), <i>P</i> = 0.21
		g/g	7	11	7	23	1.5 (0.6–4.0), <i>P</i> = 0.43 (159, 235)
	Intron 8 C→T (Bsm I; b→B)	c/c	62	104	60	197	
		c/t	28	47	35	115	0.8 (0.5–1.3), <i>P</i> = 0.35
		t/t	10	17	5	18	2.1 (0.9–5.0), <i>P</i> = 0.08 (158, 235)
	Exon 9 T→C (codon 352) (Taq I; T→t)	t/t	53	131	57	228	
		t/c	41	101	40	158	1.1 (0.8–1.7), <i>P</i> = 0.54
		c/c	6	15	3	12	4.3 (1.6–11.4), <i>P</i> = 0.004 (240, 375)

* The results did not change when analysis was restricted to paucibacillary cases and their matched controls.

† For definitions of abbreviations, see Tables 1 and 2.

‡ The number of cases and controls used in the logistic regression.

§ ORs are for one copy of the allele, and are with respect to baseline of pairs of alleles not including any of 101, 103, 105, and 111.

¶ *P* = 0.03, test for heterogeneity among LTA alleles.

ORs are for one copy of the allele, and are with respect to baseline of pairs of alleles not including any of 216, 222, 224, and 226.

leprosy susceptibility; however, for TNF -376 and -238, the 95% confidence intervals (CIs) were too wide to eliminate this. The polymorphisms ICAM1 K29M, MBL2 G57E, SLC11A1 exon 2 9-bp deletion, TGTG insertion/deletion, CAAA insertion/deletion, CR1 R1601G, IFNG +874, CCL3 microsatellite, TLR2 intron 2 microsatellite, and IL10 -1082 were not found to be associated with leprosy susceptibility. Borderline evidence of association with leprosy susceptibility was found for six polymorphisms: TNF -863, SLC11A1 promoter microsatellite, IL10 -592, TLR4 D299G, VDR *Bsm* I, and the LTA microsatellite 111-bp allele. Two SNPs were found to be associated with leprosy susceptibility with a 5% level of statistical significance without correcting for multiple comparisons: homozygotes (tt) for a silent T→C change in codon 352 of the VDR gene are susceptible (OR = 4.3, 95% CI = 1.6–11.4, $P = 0.004$), while homozygotes for the CR1 variant K1590E are protected (OR = 0.3, 95% CI = 0.1–0.8, $P = 0.02$). Presence of the LTA microsatellite 105-bp allele, but not the overall distribution of alleles at this microsatellite locus, was also associated with leprosy susceptibility (OR = 1.6, 95% CI = 1.1–2.4, $P = 0.03$).

Association study results for leprosy type. Comparison of genotype frequency between MB and PB cases usually indicated no differences. However, there was suggestive evidence that presence of the LTA 101-bp microsatellite allele is associated with MB leprosy ($\chi^2 = 11.4$, $P = 0.003$) and borderline evidence that the TLR2 224-bp microsatellite allele ($\chi^2 = 6.3$, $P = 0.042$) genotype frequencies may differ between leprosy types. However, since only 26 MB cases were included in this study these results could be due to sampling error.

DISCUSSION

Of the 38 polymorphisms investigated, 12 were insufficiently polymorphic for analysis and 2 showed evidence of association with leprosy susceptibility in this population. These two are discussed first.

The nuclear hormone receptor VDR mediates the many immunomodulatory effects of the active form of vitamin D ($1\alpha,25(\text{OH})_2\text{D}_3$), which include suppression of cytokine synthesis, immunoglobulin production, and lymphocyte proliferation. Epidemiologic evidence suggests that vitamin D deficiency may be associated with susceptibility to tuberculosis and vitamin D impairs growth of *M. tuberculosis* in human macrophage and monocytic cell lines, while vitamin D-containing medications have been useful in treating some leprosy cases.^{10,24} Homozygosity for a *Taq* I polymorphism (tt) in the ligand binding domain of VDR was associated with PB leprosy in Bengali Indians and resistance to tuberculosis in The Gambia.^{10,25} In the Bengali leprosy study the TT genotype was associated with MB leprosy and the Tt genotype was associated with leprosy susceptibility *per se*. In Karonga, where the leprosy cases are predominantly (90%) PB, the tt genotype was associated with susceptibility to leprosy *per se*. However, the apparent association in Karonga should be interpreted with caution. The expected frequency of tt homozygotes based on the measured t allele frequency in controls is 5%. The observed frequency in cases is 6% and in controls 3%. Thus, much of the apparent difference relates to a deviation of the control frequency from Hardy-Weinberg equilibrium, which may be a chance finding. In Mali, no associa-

tion was found for the VDR genotype with either lepromatous or tuberculoid leprosy or leprosy *per se* (Meisner S, unpublished data). It is not clear whether the silent T/t polymorphism has a direct effect on function or expression of VDR, and it seems more likely that reported disease associations, if real, relate to linkage disequilibrium with a flanking functional polymorphism.^{26,27}

On phagocytic cells, CR1 mediates the adherence and phagocytosis of complement-opsonized pathogens. Since *M. leprae* and *M. tuberculosis* can both gain entry into macrophages using this mechanism, it was hypothesized that variants in CR1 may influence susceptibility to mycobacterial disease.^{28,29} Five polymorphisms that encode amino acid changes in the extracellular domains of the protein and may affect either ligand binding or CR1 stability were investigated.^{30,31} These included the variants K1590E and R1601G, which underlie the Knops blood group McCoy ($McC^a \rightarrow McC^b$) and Swain-Langley ($SI1 \rightarrow SI2$) polymorphisms, respectively. Both the McC^b and $SI2$ alleles are more prevalent in African populations than in Caucasian, Asian, or Hispanic Americans, and it has been suggested that they may confer a selective advantage against infectious disease.³² These alleles were less common in Malawi ($McC^b = 0.23$, $SI2 = 0.31$) than in the west African Gambian population ($McC^b = 0.39$, $SI2 = 0.80$), but more common than in non-African populations ($McC^b \leq 0.025$, $SI2 \leq 0.03$). Homozygosity for the McC^b allele was associated with protection against leprosy in Malawi; however, the biologic mechanism underlying this finding is unclear.

Four other genes, TNF, LTA, IL-10, and TLR2, which have been reported to be associated with leprosy susceptibility elsewhere, failed to show association of the relevant polymorphisms in Karonga.³

The proinflammatory cytokine TNF influences several aspects of the immune response, including macrophage activation (important for killing of mycobacteria) and granuloma formation (important for containment of mycobacterial infections), and can also cause tissue damage and apoptosis. Several promoter polymorphisms exist which may be involved in regulation of TNF expression and such variants have been associated with both autoimmune and infectious diseases.^{33–35} A G→A polymorphism (TNF*2) at -308, which may be associated with increased TNF expression, was strongly linked to and associated with resistance to leprosy in Brazil.^{6,9,36,37} Contrary to these findings, the same variant has been associated with MB leprosy susceptibility in Bengali Indians, indicating, perhaps, that different aspects of TNF may be important in controlling the outcome of leprosy infection in different populations, perhaps due to the presence of different *M. leprae* strains, or a modified cytokine milieu resulting from presence of varying infectious pathogens or differences in genetic backgrounds.³⁸ We found no evidence that the -308 variant has a role in leprosy susceptibility in Malawi. Six other TNF polymorphisms were investigated in this study, but the -1031, +70 and -857 variants were very rare and thus could not be tested for any association. There was no evidence that the -376 polymorphism, which upregulates TNF expression by introducing a binding site for the OCT-1 transcription factor, and is associated with severe malaria, was associated with leprosy susceptibility in Malawi, although the confidence limits were too wide to eliminate this possibility.³³ Consistent with a recent Brazilian leprosy study, the -238

SNP, which may produce increased TNF transcription, was not found to be associated with leprosy in Malawi.^{9,39} The -863 SNP is located in an NF- κ B regulatory site and results in reduced p50-p50 enhancement of expression.⁴⁰ There was borderline evidence that this variant may be associated with leprosy susceptibility in Malawi, which is consistent with the findings that higher levels of TNF production may be protective against leprosy in Brazil.

Located close to the TNF gene, LTA encodes a cytokine that is produced by lymphocytes and natural killer cells and which has pleiotropic immunomodulatory effects. A haplotype that includes an intronic LTA SNP is associated with leprosy susceptibility in Brazil.⁶ In Malawi, the 105-bp allele may be weakly associated with leprosy susceptibility, while the 101-bp allele may influence leprosy type. It is possible that these associations are due to linkage disequilibrium with other polymorphisms either in another gene in the HLA region, or within LTA, or they may just be chance findings.²⁰ However, further investigation of possible associations, if any, between LTA and susceptibility both to leprosy *per se*, and to leprosy type is warranted.

The cytokine IL10 has pleiotropic effects in immunoregulation and inflammation, including the inhibition of TH1 cytokine secretion and T cell proliferation. Polymorphisms in the promoter influence the amount of IL10 production.^{10,41,42} It was speculated that by suppressing inflammation, high IL10 production might increase mycobacterial susceptibility. Three promoter polymorphisms in IL10 were investigated in Malawi and borderline evidence was found that homozygotes for the -592 A variant may have increased susceptibility to leprosy. Recently, the -819 T variant has been reported to be weakly associated with leprosy susceptibility in Brazil.⁹ These two variants exhibit linkage disequilibrium and usually occur together on a haplotype (-1082 A, -1892 T, -592 A) that is associated with increased IL10 production. In Brazil, homozygosity for the -592 A variant is not significantly associated with susceptibility, but there is a trend in this direction. Thus, it is possible that the ATA haplotype may be associated with leprosy susceptibility in both these populations but much larger studies would be required to assess this.

Mammalian toll-like receptors recognize and react to a variety of bacterial cell wall components to activate direct antimicrobial mechanisms and to facilitate transcription of genes that regulate the adaptive inflammatory immune response. TLR2, probably acting as a heterodimer with other TLRs, recognizes components from a variety of microbial pathogens including mycobacterial lipoproteins and lipoarabinomannan. *Mycobacterium leprae* can activate both TLR2-TLR1 heterodimers and TLR2 homodimers.⁴³ TLR4 is the receptor for lipopolysaccharide (LPS), a major cell wall component of gram-negative bacteria, but it seems that both TLR2 and TLR4 can mediate cellular activation in response to components of gram-positive *Mycobacterium tuberculosis*.⁴⁴ An Arg299Gly polymorphism in the extracellular domain results in impaired TLR4-mediated LPS induced signaling.⁴⁵ Heterozygosity is associated with hypo-responsiveness to inhaled LPS. Thus, it appears counter-intuitive that although only borderline significant, heterozygosity for the rare 299Gly allele was increased among controls.⁴⁶ It would be of interest to determine whether this genotype is associated with protection in other populations.

Three TLR2 polymorphisms were also investigated. Se-

quencing indicated that Arg677Trp and Arg753Gln, identified as rare variants in other populations, are extremely rare in Malawi and variation in a microsatellite located in intron 2 of the TLR2 gene was not associated with leprosy susceptibility.^{13,22}

Mannose-binding lectin binds to mannose and N-acetylglucosamine groups on a variety of bacteria, resulting in complement activation and opsonophagocytosis. Any one of three exon 1 point mutations (R52C, G54D, and G57E) reduces serum MBL concentrations. Since low serum levels of MBL may be associated with recurrent infections in young children and yet low-producing alleles are maintained in all populations, as for ICAM1, it has been proposed that balancing selection might exist.⁴⁷ One hypothesis is that low MBL levels may protect against mycobacterial disease by limiting phagocytotic entry of mycobacteria into host cells.⁴⁸ Consistent with studies of other African populations, the R52C and G54D polymorphisms were found only rarely in Malawi. The G57E (C) variant had an allele frequency of 20% and was not associated with leprosy susceptibility.

Interferon- γ is an essential and central regulator of the human response to infection. The intronic +874 T/A SNP lies in a putative binding site for the transcription factor NF- κ B. The T allele, which may be associated with high IFN- γ production has recently been found to be associated with resistance to tuberculosis in Sicilians and South Africans.^{10,19,49,50} This SNP has not previously been investigated for an association with leprosy susceptibility and was not found to be associated with leprosy in Malawi. This could be because the lower allele frequency in Malawi (20% compared with 50%) reduces the power to detect such an association because the tuberculosis association is due to another polymorphism that is in linkage disequilibrium with the +874 SNP in Sicily but not in Malawi, or because this allele does not affect leprosy susceptibility in this population.

The SLC11A1 gene encodes a protein that may have a direct effect on the survival of mycobacteria within macrophage phagosomes, but may also have pleiotropic effects that include regulation of the TH1:TH2 balance of the adaptive immune response to infection. Since the mouse homolog Slc11a1 protects against some mycobacterial infections and SLC11A1 variants have been associated with human tuberculosis susceptibility, SLC11A1 has long been a candidate for leprosy susceptibility. However, although a haplotype segregation study suggested this gene may influence leprosy susceptibility in Vietnam,⁵¹ other studies have found no evidence for association and/or linkage with susceptibility.^{25,51-54} However, variants of this gene have been associated with leprosy type in Mali and linked with the size of the Mitsuda response in southeast Asians.^{53,55} For three of the four SLC11A1 polymorphisms examined in the Malawi population, there was no evidence of an association with leprosy susceptibility. The fourth, a promoter microsatellite, showed borderline evidence that the putative lower-expressing allele 201 (allele 2) may be protective against leprosy in this population. This contrasts with the reported association of this allele with tuberculosis susceptibility in The Gambia.^{10,56}

The aim of this study was to better understand the genetic factors that underlie host immune responses to leprosy. We find evidence that variants in the VDR and CR1 genes may influence leprosy susceptibility in the Karonga District of northern Malawi, and it is possible that variants of some of

the others genes studied may also play a minor role. The sibling risk ratio for leprosy in this population appears to be of the order of two, and thus the genes and modest associations studied here appear insufficient to account for the overall genetic component identified.⁵⁷ Therefore, it seems likely that there are several other host genes that contribute to a complex network of gene products that may synergize or counteract each other to influence leprosy susceptibility. Interaction of strain, environment, nutrition, presence and nature of concurrent background infections, varying linkage disequilibrium, and allele frequencies may limit the usefulness of extrapolating these findings to other populations. Indeed, a major conclusion from this large-scale study is that many of the associations suggested by studies in other populations are not found in this Malawian population. Our data urge caution in the interpretation of infectious disease genetic associations from single-population analyses and support the re-assessment of positive findings in replication studies.

Received August 28, 2003. Accepted for publication January 19, 2004.

Acknowledgments: We thank the many field, laboratory, and data management staff of the KPS who have carried out the leprosy work in Karonga District since 1979, the people of Karonga District, and the Ministry of Health and Population and the National Health Sciences Research Committee of Malawi for their encouragement of the KPS over many years. We also thank past and present Wellcome Trust Centre for Human Genetics (WTCHG) researchers and collaborators including Peter Zimmerman, Patricia Ramaley, and Graham Cooke for their role in developing the genotyping methods used here, as well as WTCHG core facilities staff, and Kerrie Tosh, Christophe Aucan, and Dr. Branwen Hennig, for their assistance.

Financial support: The leprosy work of the KPS has been funded since its inception by the British Leprosy Relief Association, with contributions from the International Federation of Anti-Leprosy Associations. This study was supported by the Wellcome Trust. Adrian V. S. Hill is a Wellcome Trust Principal Research Fellow.

Authors' addresses: Jodene Fitness, School of Biological Sciences, Victoria University of Wellington, PO Box 600, Wellington, New Zealand, E-mail: jodene.fitness@vuw.ac.nz. Sian Floyd and Paul E. M. Fine, Department of Tropical Hygiene, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, United Kingdom, E-mails: sian.floyd@lshtm.ac.uk and paul.fine@lshtm.ac.uk. David K. Warndorff, Lifted Sichali, Lorren Mwaungulu, and Amelia C. Crampin, Karonga Prevention Study, PO Box 46, Chilumba, Karonga District, Malawi. Adrian V. S. Hill, Wellcome Trust Centre for Human Genetics, Roosevelt Drive, Oxford OX3 7BN, United Kingdom, E-mail: adrian.hill@molecular-medicine.oxford.ac.uk.

REFERENCES

1. Fine PE, 1988. Implications of genetics for the epidemiology and control of leprosy. *Philos Trans R Soc Lond B Biol Sci* 321: 365–376.
2. Chakravarti MR, Vogel F, 1973. A twin study on leprosy. *Top Hum Genet* 1: 1–123.
3. Fitness J, Tosh K, Hill AV, 2002. Genetics of susceptibility to leprosy. *Genes Immun* 3: 441–453.
4. Tosh K, Meisner S, Siddiqui MR, Balakrishnan K, Ghei S, Golding M, Sengupta U, Pitchappan R, Hill AV, 2002. A region of chromosome 20 is linked to leprosy susceptibility in a south Indian population. *J Infect Dis* 186: 1190–1193.
5. Todd JR, West BC, McDonald JC, 1990. Human leukocyte antigen and leprosy: study in northern Louisiana and review. *Rev Infect Dis* 12: 63–74.
6. Shaw MA, Donaldson IJ, Collins A, Peacock CS, Lins-Lainson Z, Shaw JJ, Ramos F, Silveira F, Blackwell JM, 2001. Association and linkage of leprosy phenotypes with HLA class II and tumour necrosis factor genes. *Genes Immun* 2: 196–204.
7. Siddiqui MR, Meisner S, Tosh K, Balakrishnan K, Ghei S, Fisher SE, Golding M, Shanker Narayan NP, Sitaraman T, Sengupta U, Pitchappan R, Hill AV, 2001. A major susceptibility locus for leprosy in India maps to chromosome 10p13. *Nat Genet* 27: 439–441.
8. Mira MT, Alcais A, van Thuc N, Thai VH, Huong NT, Ba NN, Verner A, Hudson TJ, Abel L, Schurr E, 2003. Chromosome 6q25 is linked to susceptibility to leprosy in a Vietnamese population. *Nat Genet* 33: 412–415.
9. Santos AR, Suffys PN, Vanderborcht PR, Moraes MO, Vieira LM, Cabello PH, Bakker AM, Matos HJ, Huizinga TW, Ottenhoff TH, Sampaio EP, Sarno EN, 2002. Role of tumor necrosis factor- α and interleukin-10 promoter gene polymorphisms in leprosy. *J Infect Dis* 186: 1687–1691.
10. Fitness J, Floyd S, Warndorff DK, Sichali L, Malema S, Crampin AC, Fine PEM, Hill AVS, 2004. Large-scale candidate gene study of tuberculosis susceptibility in the Karonga district of northern Malawi. *Am J Trop Med Hyg* 70: (in press).
11. Fernandez-Reyes D, Craig AG, Kyes SA, Peshu N, Snow RW, Berendt AR, Marsh K, Newbold CI, 1997. A high frequency African coding polymorphism in the N-terminal domain of ICAM-1 predisposing to cerebral malaria in Kenya. *Hum Mol Genet* 6: 1357–1360.
12. Shin HD, Winkler C, Stephens JC, Bream J, Young H, Goedert JJ, O'Brien TR, Vlahov D, Buchbinder S, Giorgi J, Rinaldo C, Donfield S, Willoughby A, O'Brien SJ, Smith MW, 2000. Genetic restriction of HIV-1 pathogenesis to AIDS by promoter alleles of IL10. *Proc Natl Acad Sci U S A* 97: 14467–14472.
13. Lorenz E, Mira JP, Cornish KL, Arbour NC, Schwartz DA, 2000. A novel polymorphism in the toll-like receptor 2 gene and its potential association with staphylococcal infection. *Infect Immun* 68: 6398–6401.
14. Ponnighaus JM, Fine PE, Sterne JA, Bliss L, Wilson RJ, Malema SS, Kileta S, 1994. Incidence rates of leprosy in Karonga District, northern Malawi: patterns by age, sex, BCG status and classification. *Int J Lepr Other Mycobact Dis* 62: 10–23.
15. Ponnighaus JM, Fine PE, Bliss L, Sliney IJ, Bradley DJ, Rees RJ, 1987. The Lepa Evaluation Project (LEP), an epidemiological study of leprosy in northern Malawi. I. Methods. *Lepr Rev* 58: 359–375.
16. Ponnighaus JM, Fine PE, Bliss L, 1987. Certainty levels in the diagnosis of leprosy. *Int J Lepr Other Mycobact Dis* 55: 454–462.
17. Ponnighaus JM, Mwanjasi LJ, Fine PE, Shaw MA, Turner AC, Oxborrow SM, Lucas SB, Jenkins PA, Sterne JA, Bliss L, 1991. Is HIV infection a risk factor for leprosy? *Int J Lepr Other Mycobact Dis* 59: 221–228.
18. van Heel DA, McGovern DP, Cardon LR, Dechairo BM, Lench NJ, Carey AH, Jewell DP, 2002. Fine mapping of the IBD1 locus did not identify Crohn disease-associated NOD2 variants: implications for complex disease genetics. *Am J Med Genet* 111: 253–259.
19. Pravica V, Perrey C, Stevens A, Lee JH, Hutchinson IV, 2000. A single nucleotide polymorphism in the first intron of the human IFN- γ gene: absolute correlation with a polymorphic CA microsatellite marker of high IFN- γ production. *Hum Immunol* 61: 863–866.
20. Jongeneel CV, Briant L, Udalova IA, Sevin A, Nedospasov SA, Cambon-Thomsen A, 1991. Extensive genetic polymorphism in the human tumor necrosis factor region and relation to extended HLA haplotypes. *Proc Natl Acad Sci U S A* 88: 9717–9721.
21. Al-Sharif FM, Makki RF, Ollier WE, Hajeer AH, 1999. A new microsatellite marker within the promoter region of the MIP-1A gene. *Immunogenetics* 49: 740–741.
22. Kang TJ, Chae GT, 2001. Detection of Toll-like receptor 2 (TLR2) mutation in the lepromatous leprosy patients. *FEMS Immunol Med Microbiol* 31: 53–58.
23. Day DJ, Speiser PW, White PC, Barany F, 1995. Detection of steroid 21-hydroxylase alleles using gene-specific PCR and a multiplexed ligation detection reaction. *Genomics* 29: 152–162.
24. Rogers L, 1921. Chaulmoogra oil in leprosy and tuberculosis. *Lancet* 1: 1178–1180.

25. Roy S, Frodsham A, Saha B, Hazra SK, Mascie-Taylor CG, Hill AV, 1999. Association of vitamin D receptor genotype with leprosy type. *J Infect Dis* 179: 187–191.
26. Verbeek W, Gombart AF, Shiohara M, Campbell M, Koeffler HP, 1997. Vitamin D receptor: no evidence for allele-specific mRNA stability in cells which are heterozygous for the Taq I restriction enzyme polymorphism. *Biochem Biophys Res Commun* 238: 77–80.
27. Mocharla H, Butch AW, Pappas AA, Flick JT, Weinstein RS, De Togni P, Jilka RL, Roberson PK, Parfitt AM, Manolagas SC, 1997. Quantification of vitamin D receptor mRNA by competitive polymerase chain reaction in PBMC: lack of correspondence with common allelic variants. *J Bone Miner Res* 12: 726–733.
28. Schlesinger LS, Horwitz MA, 1990. Phagocytosis of leprosy bacilli is mediated by complement receptors CR1 and CR3 on human monocytes and complement component C3 in serum. *J Clin Invest* 85: 1304–1314.
29. Hirsch CS, Ellner JJ, Russell DG, Rich EA, 1994. Complement receptor-mediated uptake and tumor necrosis factor- α -mediated growth inhibition of *Mycobacterium tuberculosis* by human alveolar macrophages. *J Immunol* 152: 743–753.
30. Moulds JM, Zimmerman PA, Doumbo OK, Kassambara L, Sagarra I, Diallo DA, Atkinson JP, Krych-Goldberg M, Hauhart RE, Hourcade DE, McNamara DT, Birmingham DJ, Rowe JA, Moulds JJ, Miller LH, 2001. Molecular identification of Knops blood group polymorphisms found in long homologous region D of complement receptor 1. *Blood* 97: 2879–2885.
31. Xiang L, Rundles JR, Hamilton DR, Wilson JG, 1999. Quantitative alleles of CR1: coding sequence analysis and comparison of haplotypes in two ethnic groups. *J Immunol* 163: 4939–4945.
32. Zimmerman PA, Fitness J, Moulds JM, McNamara DT, Kasehagen LJ, Rowe JA, Hill AV, 2003. CR1 Knops blood group alleles are not associated with severe malaria in the Gambia. *Genes Immun* 4: 368–373.
33. Knight JC, Udalova I, Hill AV, Greenwood BM, Peshu N, Marsh K, Kwiatkowski D, 1999. A polymorphism that affects OCT-1 binding to the TNF promoter region is associated with severe malaria. *Nat Genet* 22: 145–150.
34. Cabrera M, Shaw MA, Sharples C, Williams H, Castes M, Convit J, Blackwell JM, 1995. Polymorphism in tumor necrosis factor genes associated with mucocutaneous leishmaniasis. *J Exp Med* 182: 1259–1264.
35. Sashio H, Tamura K, Ito R, Yamamoto Y, Bamba H, Kosaka T, Fukui S, Sawada K, Fukuda Y, Satomi M, Shimoyama T, Furuyama J, 2002. Polymorphisms of the TNF gene and the TNF receptor superfamily member 1B gene are associated with susceptibility to ulcerative colitis and Crohn's disease, respectively. *Immunogenetics* 53: 1020–1027.
36. Santos AR, Almeida AS, Suffys PN, Moraes MO, Filho VF, Mattos HJ, Nery JA, Cabello PH, Sampaio EP, Sarno EN, 2000. Tumor necrosis factor promoter polymorphism (TNF2) seems to protect against development of severe forms of leprosy in a pilot study in Brazilian patients. *Int J Lepr Other Mycobact Dis* 68: 325–327.
37. Moraes MO, Duppre NC, Suffys PN, Santos AR, Almeida AS, Nery JA, Sampaio EP, Sarno EN, 2001. Tumor necrosis factor- α promoter polymorphism TNF2 is associated with a stronger delayed-type hypersensitivity reaction in the skin of borderline tuberculoid leprosy patients. *Immunogenetics* 53: 45–47.
38. Roy S, McGuire W, Mascie-Taylor CG, Saha B, Hazra SK, Hill AV, Kwiatkowski D, 1997. Tumor necrosis factor promoter polymorphism and susceptibility to lepromatous leprosy. *J Infect Dis* 176: 530–532.
39. Bayley JP, de Rooij H, van den Elsen PJ, Huizinga TW, Verweij CL, 2001. Functional analysis of linker-scan mutants spanning the -376, -308, -244, and -238 polymorphic sites of the TNF- α promoter. *Cytokine* 14: 316–323.
40. Udalova IA, Richardson A, Denys A, Smith C, Ackerman H, Foxwell B, Kwiatkowski D, 2000. Functional consequences of a polymorphism affecting NF- κ B p50-p50 binding to the TNF promoter region. *Mol Cell Biol* 20: 9113–9119.
41. Gibson AW, Edberg JC, Wu J, Westendorp RG, Huizinga TW, Kimberly RP, 2001. Novel single nucleotide polymorphisms in the distal IL-10 promoter affect IL-10 production and enhance the risk of systemic lupus erythematosus. *J Immunol* 166: 3915–3922.
42. Rees LE, Wood NA, Gillespie KM, Lai KN, Gaston K, Mathieson PW, 2002. The interleukin-10-1082 G/A polymorphism: allele frequency in different populations and functional significance. *Cell Mol Life Sci* 59: 560–569.
43. Krutzik SR, Ochoa MT, Sieling PA, Uematsu S, Ng YW, Legaspi A, Liu PT, Cole ST, Godowski PJ, Maeda Y, Sarno EN, Norgard MV, Brennan PJ, Akira S, Rea TH, Modlin RL, 2003. Activation and regulation of Toll-like receptors 2 and 1 in human leprosy. *Nat Med* 9: 525–532.
44. Means TK, Wang S, Lien E, Yoshimura A, Golenbock DT, Fenton MJ, 1999. Human toll-like receptors mediate cellular activation by *Mycobacterium tuberculosis*. *J Immunol* 163: 3920–3927.
45. Arbour NC, Lorenz E, Schutte BC, Zabner J, Kline JN, Jones M, Frees K, Watt JL, Schwartz DA, 2000. TLR4 mutations are associated with endotoxin hyporesponsiveness in humans. *Nat Genet* 25: 187–191.
46. Lorenz E, Mira JP, Frees KL, Schwartz DA, 2002. Relevance of mutations in the TLR4 receptor in patients with gram-negative septic shock. *Arch Intern Med* 162: 1028–1032.
47. Garred P, Madsen HO, Kurtzhals JA, Lamm LU, Thiel S, Hey AS, Svejgaard A, 1992. Diallelic polymorphism may explain variations of the blood concentration of mannan-binding protein in Eskimos, but not in black Africans. *Eur J Immunogenet* 19: 403–412.
48. Garred P, Harboe M, Oettinger T, Koch C, Svejgaard A, 1994. Dual role of mannan-binding protein in infections: another case of heterosis? *Eur J Immunogenet* 21: 125–131.
49. Lio D, Marino V, Serauto A, Gioia V, Scola L, Crivello A, Forte GI, Colonna-Romano G, Candore G, Caruso C, 2002. Genotype frequencies of the +874T→A single nucleotide polymorphism in the first intron of the interferon- γ gene in a sample of Sicilian patients affected by tuberculosis. *Eur J Immunogenet* 29: 371–374.
50. Rossouw M, Nel HJ, Cooke GS, van Helden PD, Hoal EG, 2003. Association between tuberculosis and a polymorphic NF- κ B binding site in the interferon γ gene. *Lancet* 361: 1871–1872.
51. Abel L, Sanchez FO, Oberti J, Thuc NV, Hoa LV, Lap VD, Skamene E, Lagrange PH, Schurr E, 1998. Susceptibility to leprosy is linked to the human NRAMP1 gene. *J Infect Dis* 177: 133–145.
52. Shaw MA, Atkinson S, Dockrell H, Hussain R, Lins-Lainson Z, Shaw J, Ramos F, Silveira F, Mehdi SQ, Kaukab F, 1993. An RFLP map for 2q33-q37 from multicase mycobacterial and leishmanial disease families: no evidence for an Lsh/Ity/Bcg gene homologue influencing susceptibility to leprosy. *Ann Hum Genet* 57: 251–271.
53. Meisner SJ, Mucklow S, Warner G, Sow SO, Lienhardt C, Hill AV, 2001. Association of NRAMP1 polymorphism with leprosy type but not susceptibility to leprosy per se in west Africans. *Am J Trop Med Hyg* 65: 733–735.
54. Medina E, Rogerson BJ, North RJ, 1996. The Nramp1 antimicrobial resistance gene segregates independently of resistance to virulent *Mycobacterium tuberculosis*. *Immunology* 88: 479–481.
55. Alcais A, Sanchez FO, Thuc NV, Lap VD, Oberti J, Lagrange PH, Schurr E, Abel L, 2000. Granulomatous reaction to intradermal injection of lepromin (Mitsuda reaction) is linked to the human NRAMP1 gene in Vietnamese leprosy sibships. *J Infect Dis* 181: 302–308.
56. Bellamy R, Ruwende C, Corrah T, McAdam KP, Whittle HC, Hill AV, 1998. Variations in the NRAMP1 gene and susceptibility to tuberculosis in west Africans. *N Engl J Med* 338: 640–644.
57. Wallace C, Clayton D, Fine P, 2003. Estimating the relative recurrence risk ratio for leprosy in Karonga District, Malawi. *Lepr Rev* 74: 133–140.