

1 **A Systematic Review of Cost-of-Illness Studies of Multimorbidity**

2

3 Author names

4 Lili Wang², Lei Si², Fiona Cocker^{2,3}, Andrew J Palmer², Kristy Sanderson^{1,2}

5

6 Author affiliations

7 ¹School of Health Sciences, Faculty of Medicine and Health Sciences, University of East Anglia, Norwich, UK

8 ²Menzies Institute for Medical Research and University of Tasmania, Hobart, Tasmania, Australia

9 ³School of Medicine and University of Tasmania, Hobart, Tasmania, Australia

10

11 Corresponding author

12 Prof Kristy Sanderson

13 Chair in Applied Health Research, School of Health Sciences, University of East Anglia

14 Adjunct Professor, Menzies Institute for Medical Research, University of Tasmania

15 Mailing address: School of Health Sciences, University of East Anglia, Edith Cavell Building room 1.27, Norwich
16 Research Park Norwich, United Kingdom, NR4 7TJ

17 Phone Number: +44 (0) 1603 591142

18 Email Address: Kristy.Sanderson@uea.ac.uk

1 **Abstract**

2 Objectives: The economic burden of multimorbidity is considerable. This review analyzed the methods of cost-of-
3 illness (COI) studies and summarized the economic outcomes of multimorbidity.

4 Methods: A systematic review (2000-2016) was performed, which was registered with Prospero, reported according
5 to PRISMA, and used a quality checklist adapted for COI studies. The inclusion criteria were peer-reviewed COI
6 studies on multimorbidity, whereas the exclusion criterion was studies focusing on an index disease. Extracted data
7 included the definition, measure, and prevalence of multimorbidity; the number of included health conditions; the
8 age of study population; the variables used in the COI methodology; the percentage of multimorbidity vs. total costs;
9 and the average costs per capita.

10 Results: Among the 26 included articles, 14 defined multimorbidity as a simple count of 2 or more conditions.

11 Methodologies used to derive the costs were markedly different. Given different healthcare systems, OOP payments
12 of multimorbidity varied across countries. In the 17 and 12 studies with cut-offs of ≥ 2 and ≥ 3 conditions,
13 respectively, the ratios of multimorbidity to non-multimorbidity costs ranged from 2-16 and 2-10. Among the 10
14 studies that provided cost breakdowns, studies with and without a societal perspective attributed the largest
15 percentage of multimorbidity costs to social care and inpatient care/medicine, respectively.

16 Conclusion: Multimorbidity was associated with considerable economic burden. Synthesising the cost of
17 multimorbidity was challenging due to multiple definitions of multimorbidity and heterogeneity in COI methods.

18 Count method was most popular to define multimorbidity. There is consistent evidence that multimorbidity was
19 associated with higher costs.

20 Words: 250/250

21

22 Key points for decision makers:

23 1. Despite substantial methodological variations between COI studies, there is consistent evidence of considerable
24 economic burden associated with multimorbidity.

25 2. Yet pooling the costs from different studies is impossible given different environments, such as healthcare
26 systems, period of observation and perspectives.

27 3. Social care is the most important cost drivers in multimorbidity COI studies with societal perspective while
28 inpatient care/medicine in studies without societal perspective.

1 **Background**

2 The term multimorbidity refers to the presence of multiple concurrent chronic health conditions in one
3 individual without an index disease [1]. Regardless of the specific definition of multimorbidity adopted, it is
4 common [2], particularly in the elderly with prevalence estimates of 65-98% for those aged >65 years [3-5].
5 Additionally, a growing body of evidence has indicated an increasing prevalence of multimorbidity [6]. In the
6 Netherlands, Uijen and van de Lisdonk found that the prevalence of people with two or more chronic health
7 conditions increased from 12.3% to 20.5% in primary care from 1985 to 2005 [7]. In the United States, Ward found
8 that the prevalence of multimorbidity increased from 21.8% in 2001 to 25.5% in 2012 using the data from a national
9 household survey [8, 9].

10 Multimorbidity is one of the most problematic “chronic health conditions” [10] because of the escalating
11 prevalence and its far-reaching health consequences. Multimorbidity can have a drastic and lifetime impact, as it is
12 unlikely to be cured. Additionally, compared to single health conditions, multimorbidity has been related to poorer
13 health-related quality of life [11, 12], higher health service utilization [13], and negative occupational consequences
14 [14], such as productivity loss due to presenteeism (e.g., ‘continuing to work while sick’) and absenteeism.
15 Moreover, healthcare resource consumption is expected to increase not only because of the accumulation of chronic
16 health conditions but also because of interactions and synergies among health conditions present within an
17 individual [15]. Given the concurrent changes in epidemiology, the use of resources and morbidity-related costs of
18 multimorbid conditions are likely to undergo enormous changes as well, especially since uniform definition and
19 measure of multimorbidity have been lacking.

20 Some researchers have begun to summarize the associations of multimorbidity and costs. Lehnert et al.
21 reviewed the literature in 2011 which was restricted to studies of older adults only [16]. Sambamoorthi et al.
22 conducted a narrative expert review which does not meet the criteria for a systematic review, i.e. did not report use
23 of systematic review methodology, did not describe a study protocol and therefore was not registered on Prospero,
24 did not include a standardised assessment of study quality, and did not follow guidelines for reporting systematic
25 reviews (e.g. PRISMA) [17]. Our review meets all of these criteria and we believe it presents an important and
26 distinct contribution to this field. Another advantage of this review was providing the breakdown of costs. The aim
27 of this study was two-fold: we first compiled a general description of COI methods, and we subsequently

1 systematically reviewed studies on the costs of multimorbidity, analyzing the different methods used, summarizing
2 their findings on the economic impact of multimorbidity and evaluating the quality of the included COI studies.

3

4 **Methods**

5 A literature search was performed in the following electronic databases: PROSPERO, Cochrane Library
6 (including the HTA Database, DARE and Cochrane Database of Systematic Reviews), Health Economic
7 Evaluations databases (including the NHS Economic Evaluation Database (NHS EED) and Health Economic
8 Evaluations Database (HEED)), National Institute for Health and Care Excellence (NICE) Evidence Services,
9 Google Scholar, Scopus, and PubMed. The search strategy combined key words related to multimorbidity,
10 comorbidity and multiple chronic health conditions. The search was restricted to papers written in English and
11 published since 2000 up to October 2016. The inclusion criteria were peer-reviewed COI studies (including cross-
12 sectional, cohort and modeling studies); the exclusion criterion was studies focusing on an index disease. The main
13 difference between comorbidity and multimorbidity was whether an index disease was specified or not. Calculating
14 the costs without distinguishing those two situations may lead to an underestimation of the burden of multimorbidity.
15 As in “comorbidity”, allied treatments of the dominant disease might also apply to the triggered secondary diseases,
16 while in “multimorbidity”, each disease receives relatively independent treatments. Therefore, we included
17 “comorbidity” in the search terms primarily because of the interchangeable use of the terms “comorbidity” and
18 “multimorbidity” in the literature. Then, during the article screening stage, studies were excluded if they focused on
19 “an index disease”. Figure 1 illustrates the literature search and selection process and presents the reasons for study
20 exclusion. As an example, the search strategy for PubMed is shown below.

21 (((multimorbidity[Title/Abstract]) OR (multi-morbidity[Title/Abstract]) OR (comorbidity[MeSH Terms])
22 OR (co-morbidity[Title/Abstract]) OR ((multiple[Title/Abstract]) AND (chronic[Title/Abstract] OR long-
23 term[Title/Abstract] OR "long term"[Title/Abstract]) AND (illnesses[Title/Abstract] OR diseases[Title/Abstract]
24 OR conditions[Title/Abstract]))) AND ((forecasting[MeSH Terms]) OR (health expenditures[MeSH Terms]) OR
25 (spending[Title/Abstract]) OR (costs and cost analysis[MeSH Terms]) OR (cost-of-illness[Title/Abstract]) OR (cost
26 of illness[Title/Abstract])) AND English[Language] AND ("2000"[Date - Publication] : "3000"[Date - Publication]))
27 NOT (letter[Publication Type] OR news[Publication Type] OR editorial[Publication Type] OR “newspaper
28 article”[Publication Type] OR comment[Publication Type]).

1 All titles and abstracts were screened by two independent reviewers (LLW and LS), after which the full
2 texts of all potentially eligible papers were obtained and screened by the same two reviewers. For any disagreement,
3 the abstract was set aside for further evaluation. After a consensus was reached on the final sample of papers, the
4 primary reviewer (LLW) screened the reference lists of the included papers for additional papers that fulfilled the
5 inclusion criteria. This review was reported in accordance with the Preferred Reporting Items for Systematic
6 Reviews and Meta-Analyses (PRISMA) guidelines [19].

7 Formal international guidelines for quality analyses of COI studies are lacking; therefore, relevant
8 information was extracted referring to the British Medical Journal Checklist [20] for economic submissions and was
9 adapted for COI studies by Molinier et al. [21]. Equal weight was assigned to each item of the checklist, and the
10 final score was the sum of the 10 individual items. The two reviewers assessed each study separately. If there was
11 disagreement between two reviewers at this stage, the paper was discussed with reference to the aforementioned
12 COI study checklist until agreement was reached.

13 This systematic review summarized the results referring to the items of COI methods, which have been
14 described elsewhere [18, 20, 21]. The items included the definition of multimorbidity, the epidemiological approach,
15 the perspective of the study and the type of costs assessed, resource consumption and unit costs, and sensitivity
16 analyses (dimensions shown in Table 1 and Table 2).

17 Investigating subgroup heterogeneity in COI estimates represents an area for future research [22].
18 Therefore, the included studies had to be stratified and presented by different components of costs, with clear
19 explanations of the groups. To make the costs comparable, cost estimates were all converted to USD (\$), according
20 to the 2016 exchange rate for each study and each currency, with adjustments over time based on the Consumer
21 Price Index (CPI) for the original currency. Costs were reported as average annual costs (per-capita costs) unless
22 stated otherwise, because the total costs reported in the different studies varied depending on the included sample
23 sizes. The results were synthesized descriptively.

24 25 **Results**

26 A total of 7,249 studies were identified from the PubMed and Scopus literature search. After the titles,
27 abstracts and full text were screened, 19 studies remained. Then, we incorporated three studies from other databases
28 that were not identified from PubMed and Scopus. With these 22 studies, we screened the references and identified

1 four studies that had not been identified in our literature search. Finally, twenty-six studies met our criteria (shown
2 in Tables 1-3). The years of valuation ranged from 1996 to 2013. Thirteen studies were conducted in the United
3 States [23, 24, 3, 25-34], seven in Europe [13, 35-40], two in Australia [41], one each from Canada [42], Singapore
4 [43] and Taiwan [44], and two in middle- or low-income regions [40, 45]. Overall, twenty studies used a prevalence
5 approach [23, 24, 3, 25-28, 13, 35, 30, 31, 41, 37, 38, 45, 39, 46, 40, 32, 43], seven used an incidence approach [36,
6 29, 44, 42, 33, 34, 27], and only one used an economic model to estimate the lifetime costs of multimorbidity [36].
7 The studies analyzed samples ranging in size from 1,252 to 292 million [28]. Twenty-five studies specified the age
8 range of the sample [26]. Twenty-one studies calculated estimates in a population 65 years and older [23, 24, 3, 25,
9 27, 28, 13, 35, 36, 44, 30, 31, 41, 37, 38, 45, 39, 46, 40, 32, 43], eight studies included people under 18 years old
10 [23-25, 28, 36, 44, 30, 38], and three studies were conducted in children only [42, 33, 34]. The average annual cost
11 of multimorbidity per capita ranged from \$49 [40] to \$252,313 [33], showing significant variation by study.
12 Additionally, out-of-pocket (OOP) expenditures ranged from \$49 [40] to \$6,858 [27], which was lower than public
13 insurance costs. Children with three or more life-threatening complex chronic conditions in their last year of life had
14 the highest costs (\$252,313) [33].

15

16 ***Identifying multimorbidity***

17 In total, fourteen studies provided the same, clear definition of multimorbidity, i.e., the ≥ 2 simple count
18 method [23, 13, 29, 35, 30, 31, 41, 37, 38, 45, 39, 40, 32, 43]. Twelve studies estimated the costs by number,
19 including five “organ system” and seven “health condition or symptom” studies, although they did not refer to the
20 term “multimorbidity”. For other definitions of multimorbidity, COI information was very limited. Only four studies
21 accounted for the severity of health conditions when measuring multimorbidity; two of them used the Cumulative
22 Illness Rating Scale (CIRS) [35, 37]; the Clinical Risk Groups (CRG) model [36] and Rx-defined morbidity groups
23 (Rx-MG) [44] were each used only once. The number of health conditions included when identifying
24 multimorbidity ranged from 4 [32] to 259 [23, 26, 29].

25

26 ***Epidemiological approach***

1 Six studies followed an incidence-based approach [27, 29, 36, 44, 42, 33, 34], and twenty studies calculated
2 prevalence-based healthcare costs [23, 24, 3, 25-28, 13, 35, 30, 31, 41, 37, 38, 45, 39, 46, 40, 32, 43]. Lifetime costs
3 were estimated in only one study [36], and unfortunately, specific multimorbidity-related costs were unavailable.

4

5 ***Perspective of the analysis and costs assessed***

6 Three perspectives were included: eighteen studies were from the payer's perspective [23, 24, 3, 25-29, 36,
7 44, 30, 31, 41, 45, 39, 46, 40, 34], six were from healthcare providers' perspective [13, 35, 38, 32, 33, 42], and two
8 used the societal perspective [37, 43]. However, both of the studies from the societal perspective defined costs as
9 including only healthcare and social care costs. Twelve studies included both medical and non-medical expenditures
10 when quantifying direct costs [23, 24, 3, 27, 28, 31, 41, 37, 46, 40, 32, 43].

11

12 ***Estimating resource consumption***

13 Three approaches can be used to estimate resource consumption: bottom-up, top-down and econometric
14 [47]. While the top-down approach typically requires cost data as well as relative risks to calculate population-
15 attributable fractions, the bottom-up approach often requires data from multiple sources, and the econometric
16 approach often requires only a single dataset [47]. Sixteen studies gathered data on resource consumption from
17 different departments (bottom-up approach) [3, 25, 26, 13, 29, 35, 36, 44, 30, 31, 37, 38, 32, 42, 33, 34]. One used a
18 combined bottom-up and top-down approach [35]. Ten studies extracted costs from the single database, called an
19 econometric approach [23, 24, 27, 28, 41, 45, 39, 46, 40, 43]. The follow-up periods included lifetime follow-up in a
20 study that adopted an incidence-based approach [36], six years in one study [44], four years in three studies [29, 33,
21 34] and two years in one study [42].

22

23 ***Valuation of unit costs***

24 ***Sources of cost estimations***

25 Most American studies calculated costs from Medicare payments and the Medical Expenditure Panel
26 Survey (MEPS), which provided national, continuous and comparable estimates over time. An Irish study used data
27 from primary care consultations and outpatient and inpatient visits extracted from family practices [13]. One study

1 quantified indirect costs [43]. Four studies did not provide the unit costs [25, 38, 36], and one study reported the
2 unit incremental cost only [37].

3 *Discounting costs*

4 Studies with time horizon less than one or two years did not normally discount costs. In all included studies
5 in this review, costs were not discounted, even in the longitudinal studies with more than a two-year follow-up.

7 *Sensitivity analysis*

8 None of the studies analyzed or discussed the variables that had a significant impact on cost estimates.

10 *Presentation of results*

11 The results were clearly presented in most studies and were mainly well explained and consistently
12 reported in relation to the methods adopted. Three studies did not differentiate costs. Based on the key
13 methodological points, a checklist of questions was used with full explanations given for clarity (Table 3). For
14 fourteen studies, the answer to seven of ten questions was “yes”, and all the studies were scored “no” on question 9
15 “Were the major assumptions tested in a sensitivity analysis?” Questions 3 “Were direct/indirect costs sufficiently
16 disaggregated?” and 7 “Were unit costs appropriately valued?” received fewer “yes” answers than the other
17 questions. In one American study [24], the costs were sufficiently disaggregated only for single conditions, and the
18 costs of multimorbidity were presented only as additional or supplementary information.

20 **Discussion**

21 We systematically reviewed 26 COI studies on multimorbidity without restricting the studies to any
22 specific definition of multimorbidity, and this broad inclusion contributed to a comprehensive understanding of
23 multimorbidity and its economic burden. The costs of multimorbidity ranged from \$49 [40] to \$252,313 [33] annual
24 per capita and increased according to the level of multimorbidity within each study. We found a relative paucity of
25 data on the costs of multimorbidity, but the available data still provided valuable information for us to better
26 elucidate the current magnitude of the economic burden of multimorbidity. Methods were highly heterogeneous
27 producing a wide range of COI estimates. Even at the lower bounds, these costs were substantial.

28

1 ***Costly multimorbidity***

2 The proportion of costs due to multimorbidity in relation to the total costs ranged from 3.4 to 97.8%. Most
3 (n=18) estimates were 60% and above. One study with an extraordinarily low estimate (3.4%) [46], which seemed
4 inconsistent with the other studies, only evaluated three-month cases of out-of-pocket (OOP) expenditures in
5 Australia. Two factors could explain this finding. First, the respondents had a 17% higher average income than the
6 Australian general population [46] and thus underrepresented lower socioeconomic groups (including the indigenous
7 population), who are most likely to experience higher cost burdens. Second, the conditions included in the study
8 were all chronic, which required ongoing treatment [48], and the short duration of the study may not have reflected
9 all incurred costs.

10 The highest costs of multimorbidity per person occurred in the last year of life among children with life-
11 threatening conditions (\$252,313) [33]. The costs in all three studies with young respondents ranged from \$8,551
12 [34] to \$252,313 [33] and did not include direct non-medical or indirect costs. Although the childhood prevalence
13 estimates of chronic health conditions ranged from 0.22% to 44% [49], which was much lower than the 12.9% to
14 95.1% prevalence of multimorbidity in the broader age groups [50], multimorbid children and their families still
15 faced substantial financial pressure. Moreover, the included studies indicated a persistence of high costs in the
16 following years.

17

18 ***Heterogeneity of multimorbidity COI studies***

19 Three relevant perspectives of the costs of multimorbidity were included. The societal perspective,
20 including care costs, was used in two studies, but they did not account for the costs of productivity loss due to
21 multimorbidity [51], including presenteeism, absenteeism, premature retirement and death, which are responsible for
22 a substantial proportion of the financial burden [52]. Information about productivity loss, premature retirement and
23 death could be derived from the working population. Only one Australian study in this review was conducted among
24 working-age adults and included those who were not in the workforce [29]. Unemployed populations are more likely
25 to have more chronic conditions than employed groups [51]. However, that study did not estimate productivity loss,
26 which could have been addressed with the available data.

27 Six studies adopted a cohort study design, with follow-up periods ranging from two to six years. The
28 remaining twenty studies used cross-sectional data, which reflect only the time of data collection and are limited in

1 their ability to draw valid conclusions about associations or possible causality [53]. Compared to other reviews of
2 COI studies on a specific single disease, this review on multimorbidity included fewer cohort studies [54]. Data
3 collection over a long period of time is difficult and time- and cost-intensive; however, modeling designs could
4 compensate for these challenges [55]. In this review, only Carreras et al. simulated individual costs until death using
5 a stationary Markov chain under the assumption that transition probabilities were constant [36]. This approach was
6 not consistent with the nature of chronic conditions, in which health states change dynamically, and modeling of
7 chronic conditions should consider this difference [55]. However, the lifetime multimorbidity costs could be
8 reasonably predicted in this regional study.

9 Several studies did not fully describe their methods and were thus difficult to assess. This ambiguity might
10 be due to a general lack of economic awareness in the medical journals that support economic studies. A
11 community-based cohort ensures a more representative patient population, but the diagnosis of this cohort may rely
12 on self-reported data, which are certainly less precise. The studies analyzed here confirm that multimorbidity is
13 costly and suggest that the costs of multimorbidity account for a large share of the total costs (Table 2).

14 Given different healthcare systems, OOP payments varied across countries, but OOP expenditure of
15 multimorbidity is always greater than that of non-multimorbidity. For example, in China, the patients with
16 multimorbidity have higher OOP expenditure than those without multimorbidity, even among those with health
17 insurance [40]. Findings from economic studies in different countries or regions cannot be easily generalized due to
18 monetary issues; for example, different currencies have different purchasing power for the same product [57].

19

20 *Definitions of multimorbidity*

21 It is well known that there is no singular definition of multimorbidity, and the two cut-off count method is
22 generally the most broadly accepted definition used. In this review, we found that all the COI studies that provided a
23 definition of multimorbidity adopted only this method. Most of the studies that did not specifically define
24 multimorbidity also presented costs by the number of multimorbid conditions. Using the same definition increased
25 the comparability within the available COI studies.

26 The number of included health conditions used to identify multimorbidity ranged from 4, which were
27 highly prevalent, disabling or expensive conditions in an American community [32], to 259, which included all
28 conditions in clinical classification systems [29]. The costs did not increase as more conditions were included. The

1 wide variation in severity within specific conditions [56] could produce different costs. For example, children with
2 life-threatening conditions had the highest healthcare expenditure in this review [33].

3 Using the two cut-off count method to define multimorbidity, the ratios of the costs of multimorbidity to
4 non-multimorbidity ranged from 2 to 16, while the ratios ranged from 2 to 10 using the three cut-off count method.
5 Nevertheless, interpreting these quantitative results is problematic because of the different approaches used.
6 Domestic characteristics within each country or region, such as clinical practice settings and healthcare systems, also
7 affect resource consumption and unit costs. For example, medication costs can vary among studies because of the
8 use of tariffs in solidarity systems, which are not comparable to free prices in private systems.

9 The different methodologies used to identify multimorbidity led to the wide range in expenditures reported
10 above. The number and diversity of available studies on multimorbidity provide an insufficient scientific basis for
11 further explorations on multimorbidity. Therefore, it is vital to improve the methodological quality of
12 multimorbidity COI research to gain a better understanding of this common and important phenomenon. Moreover,
13 further research is needed to clarify the costs of multimorbidity from the societal perspective.

14

15 *Limitations and strengths*

16 The results of this review are limited by the nature of the studies identified. The main limitation of this
17 review is its inability to include all relevant studies. Costs were estimated in 16 countries or regions from 1996 to
18 2013. The large number of abstracts derived from the databases improved the sensitivity of our search strategy. The
19 absence of a MeSH term for multimorbidity is a clear limitation. However, adding multimorbidity-related terms
20 from previous studies to our search strategy helped circumvent this limitation. We included papers published in
21 English only, which restricted our sample to some extent. The OOP can vary widely between countries because of
22 different health insurance systems and types of diseases, therefore, we have only reported the range of OOP payment
23 in different countries. Based on the fact that multimorbidity is not prevalent in the young population, the pediatric
24 multimorbidity studies were rare, therefore, the costs of multimorbidity could not be distinguished by age and the
25 finding of pediatric studies in this review was limited.

26 Moreover, the practicality of COI studies themselves in aiding policy decision-making has been debated
27 [58, 59], and their inability to prioritize resources has been criticized as well [60, 61]. COI studies, which aim to
28 identify and measure all costs of health condition serve a different purpose than other health economic evaluations

1 (e.g., cost-benefit, cost-effectiveness, and cost-utility analyses), which aim to assess both costs and outcomes of the
2 adopted intervention/policy [62-64]. However, COI studies can provide useful information as long as they adhere to
3 standardized and acceptable methodologies [65, 66]. Furthermore, the results of COI studies have been used by
4 organizations such as the World Bank and the World Health Organization to estimate public, private and total
5 national health expenditures globally [67]. Different stakeholders can utilize COI studies for different purposes [68].
6 For example, governments can estimate the financial impact of a disease on public budgets for resource allocation
7 purposes, whereas pharmaceutical corporations can identify diseases with high management costs and direct
8 research and development investments accordingly. However, caution is warranted when using COI studies; for
9 optimal resource allocation, they should be used in combination with other thorough economic evaluations [69].

10 Despite these limitations, this review provides an overview of the range of estimates reported in recent
11 decades, and the collated evidence provides a greater understanding of the COI of multimorbidity than the results
12 provided by individual studies. Moreover, this review adds systematic evidence about the methodologies used to
13 analyze multimorbidity costs and provides insight into the reasons for the disparate results among studies. Although
14 multimorbidity complicates the findings of COI studies, this review can be useful for informing decisions about the
15 prioritization of resources [70, 71], particularly when combined with other economic assessments.

17 **Conclusion**

18 Noting the substantial methodological variations between studies, multimorbidity was associated with a
19 considerable economic burden. Although this review identified two studies estimating the costs from a societal
20 perspective, there was a consistent theme throughout the included studies that those with multimorbidity had higher
21 costs than those without multimorbidity. Future research should focus on improving the methods of estimating costs.
22 A closer agreement of definition of multimorbidity is still required to allow consistent comparisons and enhance the
23 interpretation of study findings among future studies.

26 **Compliance with Ethical Standards**

27 **Funding**

28 Lili Wang was funded by a University of Tasmania/Anhui Medical University PhD Scholarship. No other funding
29 was received for this study.

1 Conflict of interest

2 The authors, Lili Wang, Lei Si, Fiona Cocker, Andrew J Palmer and Kristy Sanderson, declare that they have no
3 competing interests.

4

5 **Author Contributions**

6 Lili Wang conceptualised the article, registered in PROSPERO, conducted the preliminary searches, screening of
7 search results, data extraction, risk of bias (quality) assessment and data analysis, and wrote the manuscript. Lei Si
8 advised on data analysis, conducted the second round screening of search results, data extraction and risk of bias
9 (quality) assessment, and helped revise the manuscript. Fiona Cocker helped revise the manuscript. Andrew J
10 Palmer assisted in conceptualising the study and revised the manuscript. Kristy Sanderson assisted in
11 conceptualising the paper and revised the protocol and manuscript. All authors gave final approval of the version to
12 be published.

13

1 **Figure 1. Flowchart illustrating the search process**

2

3




Table 1. Methodology of included cost-of-illness studies in multimorbidity

Study	Country	Perspective	Epidemiological approach	Study design	Year of valuation	Currency
Hwang et. al [23]	USA	Payer (OOP)	Prevalence	Cross-sectional	1996	USD
Garis et. al [24]	USA	Payer (public insurance)	Prevalence	Cross-sectional	1995	USD
Wolff et. al [3]	USA	Payer (public insurance)	Prevalence	Cross-sectional	1999	USD
Anderson et. al [25]	USA	Payer (public insurance)	Prevalence	Cross-sectional	1998	USD
Thorpe et. al [26]	USA	Payer (public insurance)	Prevalence	Cross-sectional	1987	USD
Thorpe et. al [26]	USA	Payer (public insurance)	Prevalence	Cross-sectional	1997	USD
Thorpe et. al [26]	USA	Payer (public insurance)	Prevalence	Cross-sectional	2002	USD
Schoenberg et. al [27]	USA	payer (OOP)	Prevalence/incidence	Cross-sectional	1998	USD
Schoenberg et. al [27]	USA	Payer (OOP)	Prevalence/incidence	Cross-sectional	2002	USD
Paez et. al [28]	USA	Payer (OOP)	Prevalence	Cross-sectional	2005	USD
Glynn et. al [13]	West of Ireland (national representative)	Health care providers	Prevalence	Cross-sectional	2009	EUR
Naessens et. al [29]	USA	Payer	Incidence	Cohort (4 years follow-up)	2007	USD
Nagl et. al [35]	Germany	Health care providers	Prevalence	Cross-sectional	2010	EUR
Carreras et. al [36]	the county of Baix Empordà in Catalonia (Spain)	Payer (public insurance)	Incidence	cohort (lifetime)	2007	EUR
Kuo et. al [44]	Taiwan	payer (public insurance)	Incidence	Cohort (6 years follow-up)	2010	USD
Lochner et. al [30]	USA	Payer (public insurance)	Prevalence	Cross-sectional	2011	USD
Machlin et. al [31]	USA	Payer (public insurance)	Prevalence	Cross-sectional	2009	USD
McRae et. al [41]	Australia	Payer (public insurance)	Prevalence	Cross-sectional	2009	AUD
Heider et. al [37]	Germany	Payer (OOP)	Prevalence	Cross-sectional	2009	EUR
Orueta et. al [38]	Basque country (region in Spain/France)	Societal	Prevalence	Cross-sectional	2011	EUR
Pati et. al [45]	India	Health care providers	Prevalence	Cross-sectional	2007	INR
Bahler et. al [39]	Switzerland	Payer	Prevalence	Cross-sectional	2013	Swiss francs
Lee et. al [40]	China	Payer (OOP)	Prevalence	Cross-sectional	2010	CNY
Lee et. al [40]	Ghana	Payer (OOP)	Prevalence	Cross-sectional	2010	GHC
Lee et. al [40]	Mexico	Payer (OOP)	Prevalence	Cross-sectional	2010	INR
Lee et. al [40]	Russia	Payer (OOP)	Prevalence	Cross-sectional	2010	MXN
Lee et. al [40]	South Africa	Payer (OOP)	Prevalence	Cross-sectional	2010	RUB
Lee et. al [40]	India	payer (OOP)	Prevalence	Cross-sectional	2010	ZAR
Meraya et. al [32]	USA	health care providers & payer	Prevalence	Cross-sectional	2011	USD
Picco et. al [43]	Singapore	societal	Prevalence	Cross-sectional	2013	SGD
Carpenter et. al [46]	Australia	payer (OOP)	Prevalence	Cross-sectional	2009	USD
Cohen et. al [42]	Canada	health care providers	Incidence	Cohort (2 years follow-up)	2005-2007	CAD
Ananth et. al [33]	USA	health care providers	Incidence	Cohort (4 years follow-up)	2012	USD
Zhong et. al [34]	USA	payer (public insurance)	Incidence	Cohort (4 years follow-up)	2004	USD

OOP, out-of-pocket; USA, United States of America; USD, United States Dollar; EUR, Euro; AUD, Australian Dollar; INR, Indian Rupee; CNY, Chinese Yuan; GHC, Ghana Cedi; MXN, Mexican Peso; RUB, Russian Rouble; ZAR, South African Rand; SGD, Singapore Dollar; CAD, Canadian Dollar.

Table 2. The definition, measure, costs of multi morbidity

Study	Definition of MM	Measure of MM	Number of included conditions	Age range (y.o.)	Prevalence of MM (%)	% (MM) of total costs	Direct costs			Average costs of MM (\$)*		MM/non-MM	
							Direct medical costs	Direct non-medical costs	Indirect costs	MM2+	MM3+	MM2+	MM3+
Hwang et. al [23]	MM2+	Count	259	0-80+	17.0	38.1	yes	no	no	1387	1733.0	3	3
Garis et. al [24]	NS	Count	9	0+	NA	NA	yes	yes	no	7938	NA	NA	NA
Wolff et. al [3]	NS	ACG	3493	65+	65(MM2+)/43(MM3+)	95.3	yes	yes	no	10627	14276.0	11	10
Anderson et. al [25]	NS	Count	NS	0+	NA	NA	yes	no	no	NA	NA	NA	NA
Thorpe et. al [26]	NS	Count	259	NS	76.4	92.2	yes	no	no	13330	14989.8	6	3
Thorpe et. al [26]	NS	Count	259	NS	80.5	95.1	yes	no	no	10950	12158.8	5	4
Thorpe et. al [26]	NS	Count	259	NS	86.2	97.2	yes	no	no	11666	12864.0	6	5
Schoenberg et. al [27]	NS	Count	8	65+	58.1	70.6	yes	yes	no	3858	4109.0	2	2
Schoenberg et. al [27]	NS	Count	8	65+	70.4	78.6	yes	yes	no	6856	7687.6	2	2
Paez et. al [28]	NS	Count	NS	0+	24(MM2+)/13(MM3+)	48.5	yes	yes	no	1844	2306.0	16	4
Glynn et. al [13]	MM2+	Count	147	50+	66.2	82.5	yes	no	no	2211	2602.0	2	2
Naessens et. al [29]	MM2+	Count	259	18-64	54.3	82.5	yes	no	no	13285	16245.0	4	4
Nagl et. al [35]	MM2+	CIRS	33	65+	86.4	94.8	yes	no	no	3778	4422.0	2	2
Carreras et. al [36]	NS	CRG model	all 857,385 ICD codes (815,227 diagnostics and 42,158 procedures)	0+	17.8	NA	yes	no	no	NA	NA	NA	NA
Kuo et. al [44]	NS	counting the number of Rx-MG	55	0-71	80	NA	yes	no	no	1045	NA	4	NA
Lochner et. al [30]	MM2+	Count	15	0+	67.3	92.6	yes	no	no	13949	NA	6	NA
Machlin et. al [31]	MM2+	Count	20	18+	25.0	60.3	yes	yes	no	11934	NA	4	NA
McRae et. al [41]	MM2+	Count	6	50+	55.8	81.0	yes	yes	no	1781	2014	2	2
Heider et. al [37]	MM2+	CIRS-G	14	57-84	NA	74.0	yes	yes	no	NA	NA	NA	NA
Orueta et. al [38]	MM2+	ACG	52	0+	23.6	63.6	yes	no	no	NA	NA	NA	NA
Pati et. al [45]	MM2+	Count	NS	18+	1.3-30.6	NA	yes	no	no	240	NA	NA	NA
Bahler et. al [39]	MM2+	Count	22	65+	76.6	94.7	yes	no	no	8233	NA	5	NA
Lee et. al [40]	MM2+	Count	9	18+		NA	yes	yes	no	655	NA	NA	NA
Lee et. al [40]	MM2+	Count	9	18+		NA	yes	yes	no	92	NA	NA	NA
Lee et. al [40]	MM2+	Count	9	18+	1.4% in 18-29 years old to 40.0% in those aged 70+ years	NA	yes	yes	no	165	NA	NA	NA
Lee et. al [40]	MM2+	Count	9	18+		NA	yes	yes	no	151	NA	NA	NA
Lee et. al [40]	MM2+	Count	9	18+		NA	yes	yes	no	49	NA	NA	NA
Lee et. al [40]	MM2+	Count	9	18+		NA	yes	yes	no	60	NA	NA	NA
Meraya et. al [32]	MM2+	Count	4	21+	100.0	NA	yes	yes	no	12317	16454	NA	NA
Picco et. al [43]	MM2+	Count	10	60+	51.5	80.7	yes	yes	no	11167	NA	2	NA
Carpenter et. al [46]	NS	Count	11	50+	71.1	3.4	yes	yes	no	4447	3415	3	2
Cohen et. al [42]	NS	Count	9 organ systems	0-16	6.7	NA	yes	no	no	36434	NA	NA	NA
Ananth et. al [33]	NS	Count	9 organ systems	0-17	66.1	88.59	yes	no	no	252313	360046	4	4
Zhong et. al [34]	NS	Count	20	1-19	17	44.84	yes	no	no	8551	15797	4	6

* All costs are in \$ (1 EUR=1.0886 USD; 1 AUD=0.762966 USD; 1 INR=0.014948 USD; 1 CHF=1.005635 USD; 1 CNY=0.143719 USD; 1 MXN=0.049118 USD; 1 RUB=0.016234 USD; 1 ZAR=0.071561 USD; 1 SGD=0.717926 USD; December 18, 2016).

ACG, Adjusted Clinical Groups; Rx-MG, Rx-defined morbidity groups; CRG, Clinical Risk Groups; CIRS, Cumulative Illness Rating Scale; CIRS-G, Cumulative Illness Rating Scale for Geriatrics; ICD, the International Classification of Diseases; MM, multimorbidity; MM2+, two-cut off count method of multimorbidity; MM3+, three-cut off count method of multimorbidity; y.o., years old; NS, not specific; NA, not available.

Table 3. Answers to the methodological questions by study

Questions/answers	All studies	Hwang et. al [23]	Garis et. al [24]	Wolff et. al [3]	Anderson et. al [25]	Thorpe et. al [26]	Schoenberg et. al [27]	Paez et. al [28]	Glynn et. al [13]	Naessens et. al [29]	Nagl et. al [35]	Carreras et. al [36]	Kuo et. al [44]	Lochner et. al [30]	Machlin et. al [31]
1 Was a clear definition of the illness given?		1	p	p	0	p	p	0	1	1	1	p	p	1	1
2 Were epidemiological sources carefully described?		1	1	1	P	0	1	1	1	0	1	p	1	1	1
3 Were direct/indirect costs sufficiently disaggregated?		1	0	0	0	0	0	1	1	0	1	0	0	0	0
4 Were activity data sources carefully described?		1	1	1	1	1	1	1	1	0	p	p	1	1	p
5 Were activity data appropriately assessed?		1	1	1	0	p	1	1	1	p	1	1	1	1	p
6 Were the sources of all cost values analytically described?		1	0	0	0	1	1	1	1	0	1	1	1	p	0
7 Were unit costs appropriately valued?		1	1	1	1	1	1	1	1	1	1	p	1	1	1
8 Were the methods adopted carefully explained?		p	1	1	p	p	1	1	1	0	1	1	1	p	p
9 Were the major assumptions tested in a sensitivity analysis?		0	0	0	0	0	0	0	0	0	0	0	0	0	0
10 Was the presentation of study results consistent with the methodology of the study?		1	1	1	1	1	1	1	1	1	1	1	1	1	1
Total score by study															
YES(1)	166	8	6	6	3	3	7	8	9	3	8	5	7	6	4
NO(0)	58	1	3	3	5	3	2	2	1	6	1	2	2	2	3
PARTIALLY(p)	37	1	1	1	2	4	1	0	0	1	1	3	1	2	3

Table 3. (Continous)

Questions/answers	All studies	McRae et. al [41]	Heider et. al [37]	Orueta et. al [38]	Pati et. al [45]	Bahler et. al [39]	Lee et. al [40]	Meraya et. al [32]	Picco et. al [43]	Carpenter et. al [46]	Cohen et. al [42]	Ananth et. al [33]	Zhong et. al [34]
1 Was a clear definition of the illness given?		1	1	1	p	1	1	1	1	p	p	p	p
2 Were epidemiological sources carefully described?		1	1	1	1	1	p	1	1	1	1	1	1
3 Were direct/indirect costs sufficiently disaggregated?		0	p	1	1	1	1	0	1	1	0	0	0
4 Were activity data sources carefully described?		1	1	1	1	1	1	p	p	1	1	1	p
5 Were activity data appropriately assessed?		1	1	1	1	1	1	1	1	1	1	1	p
6 Were the sources of all cost values analytically described?		1	p	1	p	1	1	0	1	1	p	1	1
7 Were unit costs appropriately valued?		1	1	1	0	0	0	0	1	1	1	1	1
8 Were the methods adopted carefully explained?		1	1	p	1	1	p	1	1	1	1	1	1
9 Were the major assumptions tested in a sensitivity analysis?		0	0	0	0	0	0	0	0	0	0	0	0
10 Was the presentation of study results consistent with the methodology of the study?		1	1	1	1	1	1	1	1	1	1	1	1
Total score by study													
YES(1)	166	8	7	8	6	8	6	5	8	8	6	7	6
NO(0)	58	2	1	1	2	2	2	4	1	1	2	2	2
PARTIALLY(p)	37	0					3	1					

Total score by study is the sum of answers. P, partially.

References

1. van den Akker M, Buntinx F, Knottnerus JA. Comorbidity or multimorbidity: what's in a name? A review of literature. *The European journal of general practice*. 1996;2(2):65-70.
2. Fortin M, Stewart M, Poitras M-E, Almirall J, Maddocks H. A systematic review of prevalence studies on multimorbidity: toward a more uniform methodology. *The Annals of Family Medicine*. 2012;10(2):142-51.
3. Wolff JL, Starfield B, Anderson G. Prevalence, expenditures, and complications of multiple chronic conditions in the elderly. *Archives of internal medicine*. 2002;162(20):2269-76.
4. Hoffman C, Rice D, Sung HY. Persons with chronic conditions. Their prevalence and costs. *Jama*. 1996;276(18):1473-9.
5. Fortin M, Bravo G, Hudon C, Vanasse A, Lapointe L. Prevalence of multimorbidity among adults seen in family practice. *Annals of family medicine*. 2005;3(3):223-8. doi:10.1370/afm.272.
6. Wu S-Y, Green A. Projection of chronic illness prevalence and cost inflation. Santa Monica, CA: RAND Health. 2000;18.
7. Uijen AA, van de Lisdonk EH. Multimorbidity in primary care: prevalence and trend over the last 20 years. *The European journal of general practice*. 2008;14(sup1):28-32.
8. Ward BW. Prevalence of multiple chronic conditions among US adults: estimates from the National Health Interview Survey, 2010. *Preventing chronic disease*. 2013;10.
9. Ward BW. Multiple chronic conditions among US adults: a 2012 update. *Preventing chronic disease*. 2014;11.
10. Salisbury C. Multimorbidity: redesigning health care for people who use it. *The Lancet*. 2012;380(9836):7-9.
11. Fortin M, Bravo G, Hudon C, Lapointe L, Almirall J, Dubois MF et al. Relationship between multimorbidity and health-related quality of life of patients in primary care. *Quality of life research : an international journal of quality of life aspects of treatment, care and rehabilitation*. 2006;15(1):83-91. doi:10.1007/s11136-005-8661-z.
12. Wang L, Palmer AJ, Cocker F, Sanderson K. Multimorbidity and health-related quality of life (HRQoL) in a nationally representative population sample: implications of count versus cluster method for defining multimorbidity on HRQoL. *Health and quality of life outcomes*. 2017;15(1):7.
13. Glynn LG, Valderas JM, Healy P, Burke E, Newell J, Gillespie P et al. The prevalence of multimorbidity in primary care and its effect on health care utilization and cost. *Family practice*. 2011;28(5):516-23. doi:10.1093/fampra/cmr013.

14. Loeppke R, Taitel M, Richling D, Parry T, Kessler RC, Hymel P et al. Health and productivity as a business strategy. *Journal of Occupational and Environmental Medicine*. 2007;49(7):712-21.
15. Mercer S, Salisbury C, Fortin M. *ABC of Multimorbidity*. John Wiley & Sons; 2014.
16. Lehnert T, Heider D, Leicht H, Heinrich S, Corrieri S, Lupp M et al. Review: health care utilization and costs of elderly persons with multiple chronic conditions. *Medical care research and review : MCRR*. 2011;68(4):387-420. doi:10.1177/1077558711399580.
17. Sambamoorthi U, Tan X, Deb A. Multiple chronic conditions and healthcare costs among adults. *Expert review of pharmacoeconomics & outcomes research*. 2015;15(5):823-32. doi:10.1586/14737167.2015.1091730.
18. Jo C. Cost-of-illness studies: concepts, scopes, and methods. *Clinical and molecular hepatology*. 2014;20(4):327-37.
19. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS medicine*. 2009;6(7):e1000097.
20. Drummond MF, Sculpher MJ, Claxton K, Stoddart GL, Torrance GW. *Methods for the economic evaluation of health care programmes*. Oxford university press; 2015.
21. Molinier L, Bauvin E, Combescure C, Castelli C, Rebillard X, Soulié M et al. Methodological considerations in cost of prostate cancer studies: a systematic review. *Value in Health*. 2008;11(5):878-85.
22. Onukwugha E, McRae J, Kravetz A, Varga S, Khairnar R, Mullins CD. Cost-of-illness studies: an updated review of current methods. *Pharmacoeconomics*. 2016;34(1):43-58.
23. Hwang W, Weller W, Ireys H, Anderson G. Out-of-pocket medical spending for care of chronic conditions. *Health affairs*. 2001;20(6):267-78.
24. Garis RI, Farmer KC. Examining costs of chronic conditions in a Medicaid population. *Managed care (Langhorne, Pa)*. 2002;11(8):43-50.
25. Anderson G, Horvath J. The growing burden of chronic disease in America. *Public health reports (Washington, DC : 1974)*. 2004;119(3):263-70. doi:10.1016/j.phr.2004.04.005.
26. Thorpe KE, Howard DH. The rise in spending among Medicare beneficiaries: the role of chronic disease prevalence and changes in treatment intensity. *Health Affairs*. 2006;25(5):w378-w88.
27. Schoenberg NE, Kim H, Edwards W, Fleming ST. Burden of common multiple-morbidity constellations on out-of-pocket medical expenditures among older adults. *The Gerontologist*. 2007;47(4):423-37.

28. Paez KA, Zhao L, Hwang W. Rising out-of-pocket spending for chronic conditions: a ten-year trend. *Health affairs (Project Hope)*. 2009;28(1):15-25. doi:10.1377/hlthaff.28.1.15.
29. Naessens JM, Stroebel RJ, Finnie DM, Shah ND, Wagie AE, Litchy WJ et al. Effect of multiple chronic conditions among working-age adults. *American Journal of Managed Care*. 2011;17(2):118-22.
30. Lochner KA, Goodman RA, Posner S, Parekh A. Multiple chronic conditions among Medicare beneficiaries: state-level variations in prevalence, utilization, and cost, 2011. *Medicare & medicaid research review*. 2013;3(3). doi:10.5600/mmrr.003.03.b02.
31. Machlin SR. Health care expenditures for adults with multiple treated chronic conditions: estimates from the Medical Expenditure Panel Survey, 2009. *Preventing chronic disease*. 2013;10.
32. Meraya AM, Raval AD, Sambamoorthi U. Chronic condition combinations and health care expenditures and out-of-pocket spending burden among adults, Medical Expenditure Panel Survey, 2009 and 2011. *Preventing chronic disease*. 2015;12:E12. doi:10.5888/pcd12.140388.
33. Ananth P, Melvin P, Feudtner C, Wolfe J, Berry JG. Hospital Use in the Last Year of Life for Children With Life-Threatening Complex Chronic Conditions. *Pediatrics*. 2015;136(5):938-46. doi:10.1542/peds.2015-0260.
34. Zhong W, Finnie DM, Shah ND, Wagie AE, St Sauver JL, Jacobson DJ et al. Effect of multiple chronic diseases on health care expenditures in childhood. *Journal of primary care & community health*. 2015;6(1):2-9. doi:10.1177/2150131914540916.
35. Nagl A, Witte J, Hodek J-M, Greiner W. Relationship between multimorbidity and direct healthcare costs in an advanced elderly population. *Zeitschrift für Gerontologie und Geriatrie*. 2012;45(2):146-54.
36. Carreras M, Ibern P, Coderch J, Sánchez I, Inoriza JM. Estimating lifetime healthcare costs with morbidity data. *BMC health services research*. 2013;13(1). doi:10.1186/1472-6963-13-440.
37. Heider D, Matschinger H, Müller H, Saum K-U, Quinzler R, Haefeli WE et al. Health care costs in the elderly in Germany: an analysis applying Andersen's behavioral model of health care utilization. *BMC health services research*. 2014;14(1):1.
38. Orueta JF, García-Álvarez A, García-Goñi M, Paolucci F, Nuño-Solinís R. Prevalence and costs of multimorbidity by deprivation levels in the Basque Country: A population based study using health administrative databases. *PloS one*. 2014;9(2). doi:10.1371/journal.pone.0089787.

39. Bahler C, Huber CA, Brungger B, Reich O. Multimorbidity, health care utilization and costs in an elderly community-dwelling population: a claims data based observational study. *BMC health services research*. 2015;15:23. doi:10.1186/s12913-015-0698-2.
40. Lee JT, Hamid F, Pati S, Atun R, Millett C. Impact of Noncommunicable Disease Multimorbidity on Healthcare Utilisation and Out-Of-Pocket Expenditures in Middle-Income Countries: Cross Sectional Analysis. *PloS one*. 2015;10(7):e0127199. doi:10.1371/journal.pone.0127199.
41. McRae I, Yen L, Jeon YH, Herath PM, Essue B. Multimorbidity is associated with higher out-of-pocket spending: a study of older Australians with multiple chronic conditions. *Australian journal of primary health*. 2013;19(2):144-9. doi:10.1071/py12035.
42. Cohen E, Berry JG, Camacho X, Anderson G, Wodchis W, Guttman A. Patterns and costs of health care use of children with medical complexity. *Pediatrics*. 2012;130(6):e1463-70. doi:10.1542/peds.2012-0175.
43. Picco L, Achilla E, Abdin E, Chong SA, Vaingankar JA, McCrone P et al. Economic burden of multimorbidity among older adults: impact on healthcare and societal costs. *BMC health services research*. 2016;16(1):1.
44. Kuo RN, Lai M-S. The influence of socio-economic status and multimorbidity patterns on healthcare costs: a six-year follow-up under a universal healthcare system. *International journal for equity in health*. 2013;12(1):1.
45. Pati S, Agrawal S, Swain S, Lee JT, Vellakkal S, Hussain MA et al. Non communicable disease multimorbidity and associated health care utilization and expenditures in India: cross-sectional study. *BMC health services research*. 2014;14:451. doi:10.1186/1472-6963-14-451.
46. Carpenter A, Islam MM, Yen L, McRae I. Affordability of out-of-pocket health care expenses among older Australians. *Health policy (Amsterdam, Netherlands)*. 2015;119(7):907-14. doi:10.1016/j.healthpol.2015.03.010.
47. Jo C. Cost-of-illness studies: concepts, scopes, and methods. *Clin Mol Hepatol*. 2014;20(4):327-37.
48. Coulter A, Entwistle VA, Eccles A, Ryan S, Shepperd S, Perera R. Personalised care planning for adults with chronic or long-term health conditions. *The Cochrane Library*. 2015.
49. van der Lee JH, Mokkink LB, Grootenhuys MA, Heymans HS, Offringa M. Definitions and measurement of chronic health conditions in childhood: a systematic review. *Jama*. 2007;297(24):2741-51.
50. Violan C, Foguet-Boreu Q, Flores-Mateo G, Salisbury C, Blom J, Freitag M et al. Prevalence, determinants and patterns of multimorbidity in primary care: a systematic review of observational studies. *PloS one*. 2014;9(7):e102149.

51. Australian Institute of Health and Welfare (AIHW). Chronic disease and participation in work (Cat. no. PHE 109). Canberra: AIHW 2009.
52. Loeppke R, Taitel M, Haufle V, Parry T, Kessler RC, Jinnett K. Health and productivity as a business strategy: a multiemployer study. *Journal of Occupational and Environmental Medicine*. 2009;51(4):411-28.
53. Chen H, Hailey D, Wang N, Yu P. A review of data quality assessment methods for public health information systems. *International journal of environmental research and public health*. 2014;11(5):5170-207.
54. Ng CS, Lee JY, Toh MP, Ko Y. Cost-of-illness studies of diabetes mellitus: a systematic review. *Diabetes research and clinical practice*. 2014;105(2):151-63.
55. Steimle LN, Denton BT. *Markov Decision Processes for Screening and Treatment of Chronic Diseases*. 2015.
56. Machlin SR, Taylor AK. Design, Methods, and Field Results of the 1996 Medical Expenditure Panel Survey, Medical Provider Component. US Department of Health and Human Services, Public Health Service, Agency for Healthcare Research and Quality; 2000.
57. Costa N, Derumeaux H, Rapp T, Garnault V, Ferlicq L, Gillette S et al. Methodological considerations in cost of illness studies on Alzheimer disease. *Health economics review*. 2012;2(1):1.
58. Rice D. Cost-of-illness studies: fact or fiction? *The Lancet*. 1994;344(8936):1519-20.
59. Shiell A, Gerard K, Donaldson C. Cost of illness studies: an aid to decision-making? *Health policy (Amsterdam, Netherlands)*. 1987;8(3):317-23.
60. Wiseman V, Mooney G. SOUNDING BOARD: Burden of illness estimates for priority setting: a debate revisited. *Health policy (Amsterdam, Netherlands)*. 1998;43(3):243-51.
61. Drummond M. Cost-of-illness studies. *Pharmacoeconomics*. 1992;2(1):1-4.
62. Druss BG, Marcus SC, Olfson M, Pincus HA. The most expensive medical conditions in America. *Health Affairs*. 2002;21(4):105-11.
63. Byford S, Torgerson DJ, Raftery J. Cost of illness studies. *British Medical Journal*. 2000;320:1335-.
64. Segel JE. Cost-of-illness studies—a primer. RTI-UNC Center of Excellence in Health Promotion Economics. 2006:1-39.
65. Rice DP. Cost of illness studies: what is good about them? *Injury Prevention*. 2000;6(3):177-9.
66. Ament A, Evers S. Cost of illness studies in health care: a comparison of two cases. *Health policy (Amsterdam, Netherlands)*. 1993;26(1):29-42.

67. Murray CJ, Lopez AD, Organization WH. Global comparative assessments in the health sector: disease burden, expenditures and intervention packages. 1994.
68. Angelis A, Tordrup D, Kanavos P. Socio-economic burden of rare diseases: A systematic review of cost of illness evidence. *Health policy (Amsterdam, Netherlands)*. 2015;119(7):964-79.
69. Robinson R. Economic evaluation and health care. What does it mean? *BMJ (Clinical research ed)*. 1993;307(6905):670-3.
70. Wittenberg R. The challenge of measuring multi-morbidity and its costs. *Israel journal of health policy research*. 2015;4:1. doi:10.1186/2045-4015-4-1.
71. Weinstein M, Siegel J, Gold M, Kamlet M, Russell L. *Cost-effectiveness in health and medicine*. New York: Oxford University. 1996;55.